



Attachment Two

Draft insert labeling for proposed product

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ESCITALOPRAM OXALATE ODT safely and effectively. See full prescribing information for ESCITALOPRAM OXALATE ODT.

Escitalopram Oxalate ODT (escitalopram) tablets,
for oral use Escitalopram Oxalate ODT
Initial U.S. Approval: 2002

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS
See full prescribing information for complete boxed warning.

Increased risk of suicidal thoughts and behavior in pediatric and young adult patients taking antidepressants. Closely monitor all antidepressant-treated patients for clinical worsening and emergence of suicidal thoughts and behaviors (5.1). Escitalopram Oxalate ODT is not approved for use in pediatric patients less than 7 years of age (8.4).

RECENT MAJOR CHANGES

Indications (1)	5/2023
Dosage and Administration (2.2, 2.3, 2.5)	5/2023
Dosage and Administration, Use of Escitalopram Oxalate ODT with Other MAOIs such as Linezolid or Methylene Blue (2.7) - Removed	5/2023
Warnings and Precautions (5.2, 5.7)	8/2023

INDICATIONS AND USAGE

Escitalopram Oxalate ODT is a selective serotonin reuptake inhibitor (SSRI) indicated for the:

- treatment of major depressive disorder (MDD) in adults and pediatric patients 12 years of age and older (1)
- treatment of generalized anxiety disorder (GAD) in adults and pediatric patients 7 years and older (1)

DOSAGE AND ADMINISTRATION

Indication and Population	Recommended Dosage
MDD in Adults (2.1)	Initial: 10 mg once daily Recommended: 10 mg once daily Maximum: 20 mg once daily
MDD in Pediatric Patients 12 years and older (2.1)	Initial: 10 mg once daily Recommended: 10 mg once daily Maximum: 20 mg once daily
GAD in Adults (2.2)	Initial: 10 mg once daily Recommended: 10 mg once daily Maximum: 20 mg once daily
GAD in Pediatric Patients 7 years and older (2.2)	Initial: 10 mg once daily Recommended: 10 mg once daily Maximum: 20 mg once daily

- No additional benefits were seen at 20 mg once daily (2.1)
- Administer once daily, morning or evening, with or without food (2.3)
- Elderly patients: recommended dosage is 10 mg once daily (2.4)
- Hepatic impairment: recommended dosage is 10 mg once daily (2.4, 8.6)
- When discontinuing Escitalopram Oxalate ODT, reduce dose gradually whenever possible (2.5)

DOSAGE FORMS AND STRENGTHS

- Orally Disintegrating Tablets: 5 mg, 10 mg, 15mg and 20 mg

CONTRAINDICATIONS

- Do not use MAOIs intended to treat psychiatric disorders with Escitalopram Oxalate ODT or within 14 days of stopping treatment with Escitalopram Oxalate ODT. Do not use

Escitalopram Oxalate ODT within 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start Escitalopram Oxalate ODT in a patient who is being treated with linezolid or intravenous methylene blue (4)

- Concomitant use of pimozide (4)
- Known hypersensitivity to escitalopram or citalopram or any of the inactive ingredients (4)

WARNINGS AND PRECAUTIONS

- Serotonin Syndrome: Increased risk when co-administered with other serotonergic agents but also when taken alone. If it occurs, discontinue Escitalopram Oxalate ODT and serotonergic agents and initiate supportive treatment (4, 5.2)
- Discontinuation syndrome: When discontinuing Escitalopram Oxalate ODT, reduce dosage gradually whenever possible, and monitor for discontinuation symptoms (5.3)
- Seizures: Use with caution in patients with a history of seizure (5.4)
- Activation of Mania/Hypomania: Screen patients for bipolar disorder (5.5)
- Hyponatremia: Can occur in association with syndrome of inappropriate antidiuretic hormone secretion (5.6)
- Increased Risk of Bleeding: Concomitant use of nonsteroidal anti-inflammatory drugs, aspirin, other antiplatelet drugs, warfarin and other drugs that affect coagulation may increase risk (5.7)
- Interference with Cognitive and Motor Performance: Use caution when operating machinery (5.8)
- Angle Closure Glaucoma: Angle closure glaucoma has occurred in patients with untreated anatomically narrow angles treated with antidepressants (5.9)
- Use in Patients with Concomitant Illness: Use caution in patients with diseases or conditions that produce altered metabolism or hemodynamic responses (5.10)
- Sexual Dysfunction: Escitalopram Oxalate ODT may cause symptoms of sexual dysfunction (5.11)

ADVERSE REACTIONS

Most commonly observed adverse reactions (incidence \geq 5% and at least twice the incidence of placebo patients) are: insomnia, ejaculation disorder (primarily ejaculatory delay), nausea, sweating increased, fatigue and somnolence, decreased libido, and anorgasmia (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact TBD at 1-XXX-XXX-XXXX or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Concomitant use with SSRIs, SNRIs or Tryptophan is not recommended (7)
- Use caution when concomitant use with drugs that affect Hemostasis (NSAIDs, Aspirin, Warfarin) (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: SSRI use, particularly later in pregnancy, may increase the risk for persistent pulmonary hypertension and symptoms of poor adaptation (respiratory distress, temperature instability, feeding difficulties, hypotonia, tremor, irritability) in the neonate (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

FULL PRESCRIBING INFORMATION: CONTENTS*
WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Major Depressive Disorder
- 2.2 Generalized Anxiety Disorder
- 2.3 Administration Information
- 2.4 Screen for Bipolar Disorder Prior to Starting Escitalopram Oxalate ODT
- 2.5 Recommended Dosage for Specific Populations
- 2.6 Discontinuation of Treatment with Escitalopram Oxalate ODT
- 2.7 Switching Patients to or from a Monoamine Oxidase Inhibitor (MAOI) Antidepressant

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Suicidal Thoughts and Behaviors in Adolescents and Young Adults
- 5.2 Serotonin Syndrome
- 5.3 Discontinuation Syndrome
- 5.4 Seizures
- 5.5 Activation of Mania or Hypomania
- 5.6 Hyponatremia
- 5.7 Increased Risk of Bleeding
- 5.8 Interference with Cognitive and Motor Performance
- 5.9 Angle Closure Glaucoma
- 5.10 Use in Patients with Concomitant Illness

- 5.11 Sexual Dysfunction

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Post-Marketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment

9 DRUG ABUSE AND DEPENDENCE

- 9.2 Abuse and Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

- 14.1 Major Depressive Disorder
- 14.2 Generalized Anxiety Disorder

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

Antidepressants increased the risk of suicidal thoughts and behaviors in pediatric and young adult patients in short-term studies. Closely monitor all antidepressant-treated patients for clinical worsening, and for emergence of suicidal thoughts and behaviors [see *Warnings and Precautions (5.1)*]. Escitalopram Oxalate ODT is not approved for use in pediatric patients less than 7 years of age [see *Use in Specific Populations (8.4)*].

1 INDICATIONS AND USAGE

Escitalopram Oxalate ODT is indicated for the treatment of:

- major depressive disorder (MDD) in adults and pediatric patients 12 years of age and older.
- generalized anxiety disorder (GAD) in adults and pediatric patients 7 years of age and older.

2 DOSAGE AND ADMINISTRATION

2.1 Major Depressive Disorder

Adults

The recommended dosage of Escitalopram Oxalate ODT in adults is 10 mg once daily. A fixed-dose trial of Escitalopram Oxalate ODT demonstrated the effectiveness of both 10 mg and 20 mg of Escitalopram Oxalate ODT but failed to demonstrate a greater benefit of 20 mg over 10 mg [see *Clinical Studies (14.1)*]. Depending on clinical response and tolerability, dosage may be increased to the maximum recommended dosage of 20 mg once daily at an interval of no less than 1 week.

Pediatric Patients 12 years of age and older

The recommended dosage of Escitalopram Oxalate ODT in pediatric patients 12 years of age and older is 10 mg once daily. Depending on clinical response and tolerability, dosage may be increased to the maximum recommended dosage of 20 mg once daily at an interval of no less than 3 weeks.

2.2 Generalized Anxiety Disorder

Adults

The recommended starting dosage of Escitalopram Oxalate ODT in adults is 10 mg once daily. Depending on clinical response and tolerability, dosage may be increased to the maximum recommended dosage of 20 mg once daily at an interval of no less than 1 week.

Pediatric Patients 7 years of age and older

The recommended starting dosage of Escitalopram Oxalate ODT for pediatric patients ages 7 years of age and older is 10 mg once daily. Depending on clinical response and tolerability, dosage may be increased to the maximum recommended dosage of 20 mg once daily at an interval of no less than 2 weeks.

2.3 Administration Information

- | Administer Escitalopram Oxalate ODT orally once daily, in the morning or evening, with or without food.

2.4 Screen for Bipolar Disorder Prior to Starting Escitalopram Oxalate ODT

Prior to initiating treatment with Escitalopram Oxalate ODT or another antidepressant, screen patients for a personal family history of bipolar disorder, mania, or hypomania [see *Warnings and Precautions (5.5)*].

2.5 Recommended Dosage for Specific Populations

The recommended dosage for most elderly patients and patients with hepatic impairment is 10 mg once daily [see *Use in Specific Populations* (8.5, 8.6)].

The recommended dosage for Escitalopram Oxalate ODT in adults with a creatinine clearance less than 20 mL/minute has not been determined. No dosage adjustment is necessary for patients with mild or moderate renal impairment [see *Use in Specific Populations* (8.7)].

2.6 Discontinuation of Treatment with Escitalopram Oxalate ODT

Symptoms associated with discontinuation of Escitalopram Oxalate ODT and other SSRIs and SNRIs have been reported [see *Warnings and Precautions* (5.3)]. Patients should be monitored for these symptoms when discontinuing treatment. A gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose but at a more gradual rate.

2.7 Switching Patients to or from a Monoamine Oxidase Inhibitor (MAOI) Antidepressant

At least 14 days should elapse between discontinuation of an MAOI intended to treat psychiatric disorders and initiation of therapy with Escitalopram Oxalate ODT. Conversely, at least 14 days should be allowed after stopping Escitalopram Oxalate ODT before starting an MAOI intended to treat psychiatric disorders [see *Contraindications* (4)].

3 DOSAGE FORMS AND STRENGTHS

Orally Disintegrating Tablets

5 mg: TBD will be disclosed in ANDA filing.

10 mg: TBD will be disclosed in ANDA filing.

15 mg: TBD will be disclosed in ANDA filing.

20 mg: TBD will be disclosed in ANDA filing.

4 CONTRAINDICATIONS

Escitalopram Oxalate ODT is contraindicated in patients:

- taking MAOIs with Escitalopram Oxalate ODT or within 14 days of stopping treatment with Escitalopram Oxalate ODT because of an increased risk of serotonin syndrome. The use of Escitalopram Oxalate ODT within 14 days of stopping an MAOI intended to treat psychiatric disorders is also contraindicated [see *Dosage and Administration* (2.7) and *Warnings and Precautions* (5.2)]. Starting Escitalopram Oxalate ODT in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome [see *Dosage and Administration* (2.6) and *Warnings and Precautions* (5.2)].
- taking pimozide [see *Drug Interactions* (7)].
- with a hypersensitivity to escitalopram or citalopram or any of the inactive ingredients in Escitalopram Oxalate ODT.

5 WARNINGS AND PRECAUTIONS

5.1 Suicidal Thoughts and Behaviors in Adolescents and Young Adults

In pooled analyses of placebo-controlled trials of antidepressant drugs (SSRIs and other antidepressant classes) that included approximately 77,000 adult patients and 4,500 pediatric patients, the incidence of suicidal thoughts and behaviors in the

antidepressant-treated patients aged 24 years and younger was greater than in placebo-treated patients. There was considerable variation in risk of suicidal thoughts and behaviors among drugs, but there was an increased risk identified in young patients for most drugs studied. There were differences in absolute risk of suicidal thoughts and behaviors across the different indications, with the highest incidence in patients with MDD. The drug-placebo differences in the number of cases of suicidal thoughts and behaviors per 1000 patients treated are provided in **Table 1**.

Table 1: Risk Differences of the Number of Patients of Suicidal Thoughts and Behaviors in the Pooled Placebo-Controlled Trials of Antidepressants in Pediatric and Adult Patients

Age Range	Drug-Placebo Difference in Number of Patients of Suicidal Thoughts and Behaviors per 1000 Patients Treated
	Increases Compared to Placebo
<18 years old	14 additional patients
18 to 24 years old	5 additional patients
	Decreases Compared to Placebo
25 to 64 years old	1 fewer patient
≥65 years old	6 fewer patients

It is unknown whether the risk of suicidal thoughts and behaviors in children, adolescents, and young adults extends to longer-term use, i.e., beyond four months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with MDD that antidepressants delay the recurrence of depression and that depression itself is a risk factor for suicidal thoughts and behaviors.

Monitor all antidepressant-treated patients for any indication for clinical worsening and emergence of suicidal thoughts and behaviors, especially during the initial few months of drug therapy, and at times of dosage changes. Counsel family members or caregivers of patients to monitor for changes in behavior and to alert the healthcare provider. Consider changing the therapeutic regimen, including possibly discontinuing Escitalopram Oxalate ODT, in patients whose depression is persistently worse, or who are experiencing emergent suicidal thoughts or behaviors.

5.2 Serotonin Syndrome

SSRIs, including Escitalopram Oxalate ODT, can precipitate serotonin syndrome, a potentially life-threatening condition. The risk is increased with concomitant use of other serotonergic drugs (including triptans, tricyclic antidepressants, fentanyl, meperidine, methadone, lithium, tramadol, tryptophan, buspirone, amphetamines, and St. John’s Wort) and with drugs that impair metabolism of serotonin, i.e., MAOIs [see *Contraindications (4) and Drug Interactions (7)*].

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination) seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea).

The concomitant use of Escitalopram Oxalate ODT with MAOIs is contraindicated. In addition, do not initiate Escitalopram Oxalate ODT in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue. No reports involved the administration of methylene blue by other routes (such as oral tablets or local tissue injection). If it is necessary to initiate treatment with an MAOI such as linezolid or intravenous methylene blue in a patient taking Escitalopram Oxalate ODT, discontinue Escitalopram Oxalate ODT before initiating treatment with the MAOI [see *Contraindications (4) and Dosage and Administration (2.7)*].

Monitor all patients taking Escitalopram Oxalate ODT for the emergence of serotonin syndrome. Discontinue treatment with Escitalopram Oxalate ODT and any concomitant serotonergic agents immediately if the above symptoms occur, and initiate supportive symptomatic treatment. If concomitant use of Escitalopram Oxalate ODT with other serotonergic drugs is clinically warranted, inform patients of the increased risk for serotonin syndrome and monitor for symptoms.

5.3 Discontinuation Syndrome

During marketing of Escitalopram Oxalate ODT and other SSRIs, there have been spontaneous reports of adverse reactions occurring upon discontinuation of these drugs, particularly when abrupt, including the following: dysphoric mood, irritability, agitation, dizziness, sensory disturbances (e.g., paresthesias such as electric shock sensations), anxiety, confusion, headache, lethargy, emotional lability, insomnia, and hypomania. While these events are generally self-limiting, there have been reports of serious discontinuation symptoms.

Monitor for these symptoms when discontinuing treatment with Escitalopram Oxalate ODT. A gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose but at a more gradual rate [*see Dosage and Administration (2.6)*].

5.4 Seizures

Although anticonvulsant effects of racemic citalopram have been observed in animal studies, Escitalopram Oxalate ODT has not been systematically evaluated in patients with a seizure disorder. These patients were excluded from clinical studies during the product's premarketing testing. In clinical trials of Escitalopram Oxalate ODT, cases of convulsion have been reported in association with Escitalopram Oxalate ODT treatment. Like other drugs effective in the treatment of major depressive disorder, Escitalopram Oxalate ODT should be introduced with care in patients with a history of seizure disorder.

5.5 Activation of Mania or Hypomania

In patients with bipolar disorder, treating a depressive episode with Escitalopram Oxalate ODT or another antidepressant may precipitate a mixed/manic episode. In placebo-controlled trials of Escitalopram Oxalate ODT in major depressive disorder, activation of mania/hypomania was reported in one (0.1%) of 715 patients treated with Escitalopram Oxalate ODT and in none of the 592 patients treated with placebo. One additional case of hypomania has been reported in association with Escitalopram Oxalate ODT treatment. Activation of mania/hypomania has also been reported in a small proportion of patients with major affective disorders treated with racemic citalopram and other marketed drugs effective in the treatment of major depressive disorder. Prior to initiating treatment with Escitalopram Oxalate ODT, screen patients for any personal or family history of bipolar disorder, mania, or hypomania [*see Dosage and Administration (2.4)*].

5.6 Hyponatremia

Hyponatremia may occur as a result of treatment with SSRIs, including Escitalopram Oxalate ODT. In many cases, this hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH), and was reversible when Escitalopram Oxalate ODT was discontinued. Cases with serum sodium lower than 110 mmol/L have been reported. Elderly patients may be at greater risk of developing hyponatremia with SSRIs and SNRIs. Also, patients taking diuretics or who are otherwise volume depleted may be at greater risk [*see Use in Specific Populations (8.5)*]. Consider discontinuation of Escitalopram Oxalate ODT in patients with symptomatic hyponatremia and appropriate medical intervention should be instituted.

Signs and symptoms of hyponatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which may lead to falls. Signs and symptoms associated with more severe and/or acute cases have included hallucination, syncope, seizure, coma, respiratory arrest, and death.

5.7 Increased Risk of Bleeding

Drugs that interfere with serotonin reuptake inhibition, including Escitalopram Oxalate ODT, increase the risk of bleeding events. Concomitant use of aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), other antiplatelet drugs, warfarin, and other anticoagulants may add to

the risk. Case reports and epidemiological studies (case-control and cohort design) have demonstrated an association between use of drugs that interfere with serotonin reuptake and the occurrence of gastrointestinal bleeding. Based on data from the published observational studies, exposure to SSRIs, particularly in the month before delivery, has been associated with a less than 2-fold increase in the risk of postpartum hemorrhage [see *Use in Specific Populations (8.1)*]. Bleeding events related to drugs that interfere with serotonin reuptake have ranged from ecchymoses, hematomas, epistaxis, and petechiae to life-threatening hemorrhages.

Inform patients about the increased risk of bleeding associated with the concomitant use of Escitalopram Oxalate ODT and antiplatelet agents or anticoagulants. For patients taking warfarin, carefully monitor the international normalized ratio [see *Drug Interactions (7)*].

5.8 Interference with Cognitive and Motor Performance

In a study in normal volunteers, Escitalopram Oxalate ODT 10 mg daily did not produce impairment of intellectual function or psychomotor performance. Because any psychoactive drug may impair judgment, thinking, or motor skills, however, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that Escitalopram Oxalate ODT therapy does not affect their ability to engage in such activities.

5.9 Angle Closure Glaucoma

The pupillary dilation that occurs following use of many antidepressant drugs, including Escitalopram Oxalate ODT, may trigger an angle closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy.

5.10 Use in Patients with Concomitant Illness

Clinical experience with Escitalopram Oxalate ODT in patients with certain concomitant systemic illnesses is limited. Caution is advisable in using Escitalopram Oxalate ODT in patients with diseases or conditions that produce altered metabolism or hemodynamic responses.

Escitalopram Oxalate ODT has not been systematically evaluated in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were generally excluded from clinical studies during the product's premarketing testing.

In subjects with hepatic impairment, clearance of racemic citalopram was decreased and plasma concentrations were increased. The recommended dose of Escitalopram Oxalate ODT in hepatically impaired patients is 10 mg daily [see *Dosage and Administration (2.5)* and *Use in Specific Populations (8.6)*].

Because escitalopram is extensively metabolized, excretion of unchanged drug in urine is a minor route of elimination. Until adequate numbers of patients with severe renal impairment have been evaluated during chronic treatment with Escitalopram Oxalate ODT, however, it should be used with caution in such patients [see *Dosage and Administration (2.5)* and *Use in Specific Populations (8.7)*].

5.11 Sexual Dysfunction

Use of SSRIs, including Escitalopram Oxalate ODT, may cause symptoms of sexual dysfunction [see *Adverse Reactions (6.1)*]. In male patients, SSRI use may result in ejaculatory delay or failure, decreased libido, and erectile dysfunction. In female patients, SSRI use may result in decreased libido and delayed or absent orgasm.

It is important for prescribers to inquire about sexual function prior to initiation of Escitalopram Oxalate ODT and to inquire specifically about changes in sexual function during treatment, because sexual function may not be spontaneously reported. When evaluating changes in sexual function, obtaining a detailed history (including timing of symptom onset) is important because sexual symptoms may have other causes, including the underlying psychiatric disorder. Discuss potential management strategies to support patients in making informed decisions about treatment.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Suicidal thoughts and behaviors in adolescents and young adults [*see Warnings and Precautions (5.1)*]
- Serotonin syndrome [*see Warnings and Precautions (5.2)*]
- Discontinuation syndrome [*see Warnings and Precautions (5.3)*]
- Seizures [*see Warnings and Precautions (5.4)*]
- Activation of mania or hypomania [*see Warnings and Precautions (5.5)*]
- Hyponatremia [*see Warnings and Precautions (5.6)*]
- Increased Risk of Bleeding [*see Warnings and Precautions (5.7)*]
- Interference with Cognitive and Motor Performance [*see Warnings and Precautions (5.8)*]
- Angle-closure glaucoma [*see Warnings and Precautions (5.9)*]
- Use in Patients with Concomitant Illness [*see Warnings and Precautions (5.10)*]
- Sexual Dysfunction [*see Warnings and Precautions (5.11)*]

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Clinical Trial Data Sources

Adults

Adverse reactions information for Escitalopram Oxalate ODT was collected from 715 patients with major depressive disorder who were exposed to escitalopram and from 592 patients who were exposed to placebo in double-blind, placebo-controlled trials. An additional 284 patients with major depressive disorder were newly exposed to escitalopram in open-label trials. The adverse reaction information for Escitalopram Oxalate ODT in patients with GAD was collected from 429 patients exposed to escitalopram and from 427 patients exposed to placebo in double-blind, placebo-controlled trials.

Adverse reactions during exposure were obtained primarily by general inquiry and recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse reactions without first grouping similar types of reactions into a smaller number of standardized event categories. In the tables and tabulations that follow, standard World Health Organization (WHO) terminology has been used to classify reported adverse reactions.

The stated frequencies of adverse reactions represent the proportion of individuals who experienced, at least once, a treatment emergent adverse event of the type listed. An event was considered treatment-emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation.

Pediatric Patients

Adverse reaction information for pediatric patients was collected in double-blind placebo-controlled studies in 576 pediatric patients 6 to 17 years of age, (286 Escitalopram Oxalate ODT, 290 placebo) with major depressive disorder and in 273 pediatric patients 7 to 17 years of age (137 Escitalopram Oxalate ODT, 136 placebo) with generalized anxiety disorder.

The safety and effectiveness of Escitalopram Oxalate ODT have not been established in pediatric patients less than 12 years of age with MDD or less than 7 years of age with GAD.

Adverse Reactions Associated with Discontinuation of Treatment

Major Depressive Disorder

Adults

Among the 715 depressed patients who received Escitalopram Oxalate ODT in placebo-controlled trials, 6% discontinued treatment due to an adverse event, as compared to 2% of 592 patients receiving placebo. In two fixed-dose studies, the rate of discontinuation for adverse reactions in patients receiving 10 mg/day Escitalopram Oxalate ODT was not significantly different from the rate of discontinuation for adverse reactions in patients receiving placebo. The rate of discontinuation for adverse reactions in patients assigned to a fixed dose of 20 mg/day Escitalopram Oxalate ODT was 10%, which was significantly different from the rate of discontinuation for adverse reactions in patients receiving 10 mg/day Escitalopram Oxalate ODT (4%) and placebo (3%). Adverse reactions that were associated with the discontinuation of at least 1% of patients treated with Escitalopram Oxalate ODT, and for which the rate was at least twice that of placebo, were nausea (2%) and ejaculation disorder (2% of male patients).

Pediatric Patients

Adverse reactions in pediatric patients 6 to 17 years of age were associated with discontinuation of 3.5% of 286 patients receiving Escitalopram Oxalate ODT and 1% of 290 patients receiving placebo. The most common adverse reaction (incidence at least 1% for Escitalopram Oxalate ODT and greater than placebo) associated with discontinuation was insomnia (1% Escitalopram Oxalate ODT, 0% placebo).

The safety and effectiveness of Escitalopram Oxalate ODT have not been established in pediatric patients less than 12 years of age with MDD.

Generalized Anxiety Disorder

Adults

Among the 429 GAD patients who received Escitalopram Oxalate ODT 10 to 20 mg/day in placebo-controlled trials, 8% discontinued treatment due to an adverse event, as compared to 4% of 427 patients receiving placebo. Adverse reactions that were associated with the discontinuation of at least 1% of patients treated with Escitalopram Oxalate ODT, and for which the rate was at least twice the placebo rate, were nausea (2%), insomnia (1%), and fatigue (1%).

Pediatric Patients

Adverse reactions in pediatric patients 7 to 17 years were associated with discontinuation of 2.9% of 137 patients receiving Escitalopram Oxalate ODT and 1.5% of 136 patients receiving placebo. The most common adverse reaction (incidence at least 1% for Escitalopram Oxalate ODT and greater than placebo) associated with discontinuation was activation syndrome (1% Escitalopram Oxalate ODT, 0% placebo), intentional self injury (1% Escitalopram Oxalate ODT, 0% placebo), epistaxis (1% Escitalopram Oxalate ODT, 0% placebo), and nausea (1% Escitalopram Oxalate ODT, 0% placebo).

Incidence of Adverse Reactions in Placebo-Controlled Clinical Trials

Major Depressive Disorder

Adults

The most commonly observed adverse reactions in Escitalopram Oxalate ODT patients (incidence of approximately 5% or greater and approximately twice the incidence in placebo patients) were insomnia, ejaculation disorder (primarily ejaculatory delay), nausea, sweating increased, fatigue, and somnolence.

Table 2 enumerates the incidence, rounded to the nearest percent, of adverse reactions that occurred among 715 depressed patients who received Escitalopram Oxalate ODT at doses ranging from 10 to 20 mg/day in placebo-controlled trials. Reactions included are those occurring in 2% or more of patients treated with Escitalopram Oxalate ODT and for which the incidence in patients treated with Escitalopram Oxalate ODT was greater than the incidence in placebo-treated patients.

TABLE 2		
Adverse Reactions observed with a frequency of $\geq 2\%$ and greater than placebo for Major Depressive Disorder (Adults)		
<u>Adverse Reaction</u>	<u>Escitalopram Oxalate ODT</u>	<u>Placebo</u>
	(N=715)	(N=592)
	%	%
Autonomic Nervous System Disorders		
Dry Mouth	6%	5%
Sweating Increased	5%	2%
Central & Peripheral Nervous System Disorders		
Dizziness	5%	3%
Gastrointestinal Disorders		
Nausea	15%	7%
Diarrhea	8%	5%
Constipation	3%	1%
Indigestion	3%	1%
Abdominal Pain	2%	1%
General		
Influenza-like Symptoms	5%	4%
Fatigue	5%	2%
Psychiatric Disorders		
Insomnia	9%	4%
Somnolence	6%	2%
Appetite Decreased	3%	1%
Libido Decreased	3%	1%
Respiratory System Disorders		
Rhinitis	5%	4%
Sinusitis	3%	2%
Urogenital		
Ejaculation Disorder ^{1,2}	9%	<1%
Impotence ²	3%	<1%
Anorgasmia ³	2%	<1%

¹Primarily ejaculatory delay.

²Denominator used was for males only (N=225 Escitalopram Oxalate ODT; N=188 placebo).

³Denominator used was for females only (N=490 Escitalopram Oxalate ODT; N=404 placebo).

Pediatric Patients

The overall profile of adverse reactions in pediatric patients 6 to 17 years in major depressive disorder was generally similar to that seen in adult studies, as shown in Table 2. However, the following adverse reactions (excluding those which appear in Table 2 and those for which the coded terms were uninformative or misleading) were reported at an incidence of at least 2% for Escitalopram Oxalate ODT and greater than placebo: back pain, urinary tract infection, vomiting, and nasal congestion.

The safety and effectiveness of Escitalopram Oxalate ODT have not been established in pediatric patients less than 12 years of age with MDD.

Generalized Anxiety Disorder

Adults

The most commonly observed adverse reactions in Escitalopram Oxalate ODT patients (incidence of approximately 5% or greater and approximately twice the incidence in placebo patients) were nausea, ejaculation disorder (primarily ejaculatory delay), insomnia, fatigue, decreased libido, and anorgasmia.

Table 3 enumerates the incidence, rounded to the nearest percent of treatment-emergent adverse reactions that occurred among 429 GAD patients who received Escitalopram Oxalate ODT 10 to 20 mg/day in placebo-controlled trials. Reactions included are those occurring in 2% or more of patients treated with Escitalopram Oxalate ODT and for which the incidence in patients treated with Escitalopram Oxalate ODT was greater than the incidence in placebo-treated patients.

TABLE 3		
Adverse Reactions Observed with a Frequency of $\geq 2\%$ and $>$ placebo for Generalized Anxiety Disorder (Adults)		
<u>Adverse Reactions</u>	<u>Escitalopram Oxalate ODT</u>	<u>Placebo</u>
	(N=429)	(N=427)
	%	%
Autonomic Nervous System Disorders		
Dry Mouth	9%	5%
Sweating Increased	4%	1%
Central & Peripheral Nervous System Disorders		
Headache	24%	17%
Paresthesia	2%	1%
Gastrointestinal Disorders		
Nausea	18%	8%
Diarrhea	8%	6%
Constipation	5%	4%
Indigestion	3%	2%
Vomiting	3%	1%
Abdominal Pain	2%	1%
Flatulence	2%	1%
Toothache	2%	0%
General		
Fatigue	8%	2%
Influenza-like Symptoms	5%	4%
Musculoskeletal System Disorder		
Neck/Shoulder Pain	3%	1%
Psychiatric Disorders		
Somnolence	13%	7%

Insomnia	12%	6%
Libido Decreased	7%	2%
Dreaming Abnormal	3%	2%
Appetite Decreased	3%	1%
Lethargy	3%	1%
Respiratory System Disorders		
Yawning	2%	1%
Urogenital		
Ejaculation Disorder ^{1,2}	14%	2%
Anorgasmia ³	6%	<1%
Menstrual Disorder	2%	1%

¹Primarily ejaculatory delay.

²Denominator used was for males only (N=182 Escitalopram Oxalate ODT; N=195 placebo).

³Denominator used was for females only (N=247 Escitalopram Oxalate ODT; N=232 placebo).

Pediatric Patients

The overall profile of adverse reactions in pediatric patients 7 to 17 years in generalized anxiety disorder was generally similar to that seen in adult studies, as shown in Table 3. However, the following adverse reactions (excluding those which appear in Table 3) were reported at an incidence of at least 2% for Escitalopram Oxalate ODT and greater than placebo: dizziness (3% Escitalopram Oxalate ODT and 2% placebo), nasopharyngitis (3% Escitalopram Oxalate ODT and 1% placebo), abdominal discomfort (3% Escitalopram Oxalate ODT and 1% placebo), anxiety (3% Escitalopram Oxalate ODT and 1% placebo), irritability (2% Escitalopram Oxalate ODT and 1% placebo), and anger (2% Escitalopram Oxalate ODT and 0% placebo).

Dose Dependency of Adverse Reactions

The potential dose dependency of common adverse reactions (defined as an incidence rate of $\geq 5\%$ in either the 10 mg or 20 mg Escitalopram Oxalate ODT groups) was examined on the basis of the combined incidence of adverse reactions in two fixed-dose trials. The overall incidence rates of adverse reactions in 10 mg Escitalopram Oxalate ODT-treated patients (66%) was similar to that of the placebo-treated patients (61%), while the incidence rate in 20 mg/day Escitalopram Oxalate ODT-treated patients was greater (86%). Table 4 shows common adverse reactions that occurred in the 20 mg/day Escitalopram Oxalate ODT group with an incidence that was approximately twice that of the 10 mg/day Escitalopram Oxalate ODT group and approximately twice that of the placebo group.

Adverse Reaction	Placebo (N=311)	10 mg/day Escitalopram Oxalate ODT (N=310)	20 mg/day Escitalopram Oxalate ODT (N=125)
Insomnia	4%	7%	14%
Diarrhea	5%	6%	14%
Dry Mouth	3%	4%	9%
Somnolence	1%	4%	9%
Dizziness	2%	4%	7%
Sweating Increased	<1%	3%	8%
Constipation	1%	3%	6%
Fatigue	2%	2%	6%

Indigestion	1%	2%	6%
--------------------	----	----	----

Male and Female Sexual Dysfunction with SSRIs

Although changes in sexual desire, sexual performance, and sexual satisfaction often occur as manifestations of a psychiatric disorder, they may also be a consequence of pharmacologic treatment. In particular, some evidence suggests that SSRIs can cause such untoward sexual experiences.

Reliable estimates of the incidence and severity of untoward experiences involving sexual desire, performance, and satisfaction are difficult to obtain, however, in part because patients and physicians may be reluctant to discuss them. Accordingly, estimates of the incidence of untoward sexual experience and performance cited in product labeling are likely to underestimate their actual incidence.

TABLE 5		
Incidence of Sexual Side Effects in Placebo-Controlled Clinical Trials		
Adverse Event	Escitalopram Oxalate ODT	Placebo
	In Males Only	
	(N=407)	(N=383)
Ejaculation Disorder (primarily ejaculatory delay)	12%	1%
Libido Decreased	6%	2%
Impotence	2%	<1%
	In Females Only	
	(N=737)	(N=636)
Libido Decreased	3%	1%
Anorgasmia	3%	<1%

There are no adequately designed studies examining sexual dysfunction with escitalopram treatment.

Priapism has been reported with all SSRIs.

While it is difficult to know the precise risk of sexual dysfunction associated with the use of SSRIs, physicians should routinely inquire about such possible side effects.

Vital Sign Changes

Escitalopram Oxalate ODT and placebo groups were compared with respect to (1) mean change from baseline in vital signs (pulse, systolic blood pressure, and diastolic blood pressure) and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses did not reveal any clinically important changes in vital signs associated with Escitalopram Oxalate ODT treatment. In addition, a comparison of supine and standing vital sign measures in subjects receiving Escitalopram Oxalate ODT indicated that Escitalopram Oxalate ODT treatment is not associated with orthostatic changes.

Weight Changes

Patients treated with Escitalopram Oxalate ODT in controlled trials did not differ from placebo-treated patients with regard to clinically important change in body weight.

Laboratory Changes

Escitalopram Oxalate ODT and placebo groups were compared with respect to (1) mean change from baseline in various serum chemistry, hematology, and urinalysis variables, and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses revealed no clinically important changes in laboratory test parameters associated with Escitalopram Oxalate ODT treatment.

ECG Changes

Electrocardiograms from Escitalopram Oxalate ODT (N=625) and placebo (N=527) groups were compared with respect to outliers defined as subjects with QTc changes over 60 msec from baseline or absolute values over 500 msec post-dose, and subjects with heart rate increases to over 100 bpm or decreases to less than 50 bpm with a 25% change from baseline (tachycardic or bradycardic outliers, respectively). None of the patients in the Escitalopram Oxalate ODT group had a QTcF interval >500 msec or a prolongation >60 msec compared to 0.2% of patients in the placebo group. The incidence of tachycardic outliers was 0.2% in the Escitalopram Oxalate ODT and the placebo group. The incidence of bradycardic outliers was 0.5% in the Escitalopram Oxalate ODT group and 0.2% in the placebo group.

QTcF interval was evaluated in a randomized, placebo and active (moxifloxacin 400 mg) controlled cross-over, escalating multiple dose study in 113 healthy subjects. The maximum mean (95% upper confidence bound) difference from placebo arm were 4.5 (6.4) and 10.7 (12.7) msec for 10 mg and suprathapeutic 30 mg escitalopram given once daily, respectively. Based on the established exposure-response relationship, the predicted QTcF change from placebo arm (95% confidence interval) under the C_{max} for the dose of 20 mg is 6.6 (7.9) msec. Escitalopram 30 mg given once daily resulted in mean C_{max} of 1.7-fold higher than the mean C_{max} for the maximum recommended therapeutic dose at steady state (20 mg). The exposure under suprathapeutic 30 mg dose is similar to the steady state concentrations expected in CYP2C19 poor metabolizers following a therapeutic dose of 20 mg.

Other Reactions Observed During the Premarketing Evaluation of Escitalopram Oxalate ODT

Following is a list of treatment-emergent adverse reactions, as defined in the introduction to the **ADVERSE REACTIONS** section, reported by the 1428 patients treated with Escitalopram Oxalate ODT for periods of up to one year in double-blind or open-label clinical trials during its premarketing evaluation. The listing does not include those reactions already listed in **Tables 2 & 3**, those reactions for which a drug cause was remote and at a rate less than 1% or lower than placebo, those reactions which were so general as to be uninformative, and those reactions reported only once which did not have a substantial probability of being acutely life threatening. Reactions are categorized by body system. Reactions of major clinical importance are described in the Warnings and Precautions section (5).

Cardiovascular: hypertension, palpitation.

Central and Peripheral Nervous System Disorders: light-headed feeling, migraine.

Gastrointestinal Disorders: abdominal cramp, heartburn, gastroenteritis.

General: allergy, chest pain, fever, hot flushes, pain in limb.

Metabolic and Nutritional Disorders: increased weight.

Musculoskeletal System Disorders: arthralgia, myalgia jaw stiffness.

Psychiatric Disorders: appetite increased, concentration impaired, irritability.

Reproductive Disorders/Female: menstrual cramps, menstrual disorder.

Respiratory System Disorders: bronchitis, coughing, nasal congestion, sinus congestion, sinus headache.

Skin and Appendages Disorders: rash.

Special Senses: vision blurred, tinnitus.

Urinary System Disorders: urinary frequency, urinary tract infection.

6.2 Post-Marketing Experience

Adverse Reactions Reported Subsequent to the Marketing of Escitalopram

The following adverse reactions have been identified during post-approval use of Escitalopram Oxalate ODT. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Blood and Lymphatic System Disorders: anemia, agranulocytis, aplastic anemia, hemolytic anemia, idiopathic thrombocytopenia purpura, leukopenia, thrombocytopenia.

Cardiac Disorders: atrial fibrillation, bradycardia, cardiac failure, myocardial infarction, tachycardia, torsade de pointes, ventricular arrhythmia, ventricular tachycardia.

Ear and labyrinth disorders: vertigo

Endocrine Disorders: diabetes mellitus, hyperprolactinemia, SIADH.

Eye Disorders: angle closure glaucoma, diplopia, mydriasis, visual disturbance.

Gastrointestinal Disorder: dysphagia, gastrointestinal hemorrhage, gastroesophageal reflux, pancreatitis, rectal hemorrhage.

General Disorders and Administration Site Conditions: abnormal gait, asthenia, edema, fall, feeling abnormal, malaise.

Hepatobiliary Disorders: fulminant hepatitis, hepatic failure, hepatic necrosis, hepatitis.

Immune System Disorders: allergic reaction, anaphylaxis.

Investigations: bilirubin increased, decreased weight, electrocardiogram QT prolongation, hepatic enzymes increased, hypercholesterolemia, INR increased, prothrombin decreased.

Metabolism and Nutrition Disorders: hyperglycemia, hypoglycemia, hypokalemia, hyponatremia.

Musculoskeletal and Connective Tissue Disorders: muscle cramp, muscle stiffness, muscle weakness, rhabdomyolysis.

Nervous System Disorders: akathisia, amnesia, ataxia, choreoathetosis, cerebrovascular accident, dysarthria, dyskinesia, dystonia, extrapyramidal disorders, grand mal seizures (or convulsions), hypoaesthesia, myoclonus, nystagmus, Parkinsonism, restless legs, seizures, syncope, tardive dyskinesia, tremor.

Pregnancy, Puerperium and Perinatal Conditions: spontaneous abortion.

Psychiatric Disorders: acute psychosis, aggression, agitation, anger, anxiety, apathy, completed suicide, confusion, depersonalization, depression aggravated, delirium, delusion, disorientation, feeling unreal, hallucinations (visual and auditory), mood swings, nervousness, nightmare, panic reaction, paranoia, restlessness, self-harm or thoughts of self-harm, suicide attempt, suicidal ideation, suicidal tendency.

Renal and Urinary Disorders: acute renal failure, dysuria, urinary retention.

Reproductive System and Breast Disorders: menorrhagia, priapism.

Respiratory, Thoracic and Mediastinal Disorders: anosmia, dyspnea, epistaxis, pulmonary embolism, hyposmia, pulmonary hypertension of the newborn.

Skin and Subcutaneous Tissue Disorders: alopecia, angioedema, dermatitis, drug reaction with eosinophilia and systemic symptoms (DRESS), ecchymosis, erythema multiforme, photosensitivity reaction, Stevens Johnson Syndrome, toxic epidermal necrolysis, urticaria.

Vascular Disorders: deep vein thrombosis, flushing, hypertensive crisis, hypotension, orthostatic hypotension, phlebitis, thrombosis.

7 DRUG INTERACTIONS

Table 6 presents clinically important drug interactions with Escitalopram Oxalate ODT.

TABLE 6 Clinically Important Drug Interactions with Escitalopram Oxalate ODT

Monoamine Oxidase Inhibitors (MAOIs)	
<i>Clinical Impact:</i>	Concomitant use of SSRIs, including Escitalopram Oxalate ODT, and MAOIs increases the risk of serotonin syndrome.
<i>Intervention:</i>	Escitalopram Oxalate ODT is contraindicated in patients taking MAOIs, including MAOIs such as linezolid or intravenous methylene blue [see <i>Dosage and Administration (2.7)</i> , <i>Contraindications (4)</i> , and <i>Warnings and Precautions (5.2)</i>].
Pimozide	
<i>Clinical Impact:</i>	Concomitant use of racemic citalopram with pimozide increases plasma concentrations of pimozide, a drug with a narrow therapeutic index, and may increase the risk of QT prolongation and/or ventricular arrhythmias compared to use of racemic citalopram alone [see <i>Clinical Pharmacology (12.3)</i>].
<i>Intervention:</i>	Escitalopram Oxalate ODT is contraindicated in patients taking pimozide [see <i>Contraindications (4)</i>].
Other Serotonergic Drugs	
<i>Clinical Impact:</i>	Concomitant use of Escitalopram Oxalate ODT and other serotonergic drugs (including other SSRIs, SNRIs, triptans, tricyclic antidepressants, opioids, lithium, buspirone, amphetamines, tryptophan, and St. John's Wort) increases the risk of serotonin syndrome.
<i>Intervention:</i>	Monitor patients for signs and symptoms of serotonin syndrome, particularly during Escitalopram Oxalate ODT initiation and dosage increases. If serotonin syndrome occurs, consider discontinuation of Escitalopram Oxalate ODT and/or concomitant serotonergic drugs [see <i>Warning and Precautions (5.2)</i>].
Drugs That Interfere With Hemostasis (NSAIDs, Aspirin, Warfarin, etc.)	
<i>Clinical Impact:</i>	Concomitant use of Escitalopram Oxalate ODT and an antiplatelet or anticoagulant may potentiate the risk of bleeding.
<i>Intervention:</i>	Inform patients of the increased risk of bleeding associated with the concomitant use of Escitalopram Oxalate ODT and antiplatelet agents and anticoagulants. For patients taking warfarin, carefully monitor the international normalized ratio [see <i>Warning and Precautions (5.7)</i>].
Sumatriptan	
<i>Clinical Impact:</i>	There have been post marketing reports describing patients with weakness, hyperreflexia, and incoordination following the use of an SSRI and sumatriptan.
<i>Intervention:</i>	If concomitant treatment with sumatriptan and an SSRI is clinically warranted, appropriate observation of the patient is advised [see <i>Warning and Precautions (5.2)</i>].
Carbamazepine	
<i>Clinical Impact:</i>	Combined administration of racemic citalopram (40 mg/day for 14 days) and carbamazepine (titrated to 400 mg/day for 35 days) did not significantly affect the pharmacokinetics of carbamazepine, a CYP3A4 substrate.
<i>Intervention:</i>	Although trough citalopram plasma levels were unaffected, given the enzyme-inducing properties of carbamazepine, the possibility that carbamazepine might increase the clearance of escitalopram should be considered if the two drugs are co-administered.
Drugs Metabolized by CYP2D6	

<i>Clinical Impact:</i>	Coadministration of escitalopram (20 mg/day for 21 days) with the tricyclic antidepressant desipramine (single dose of 50 mg), a substrate for CYP2D6, resulted in a 40% increase in C _{max} and a 100% increase in AUC of desipramine.
<i>Intervention:</i>	The clinical significance of this finding is unknown. Exercise caution during coadministration of escitalopram and drugs metabolized by CYP2D6.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antidepressants during pregnancy. Healthcare providers are encouraged to register patients by calling the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or visiting online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.

Risk Summary

Based on data from published observational studies, exposure to SSRIs, particularly in the month before delivery, has been associated with a less than 2-fold increase in the risk of postpartum hemorrhage [see *Warnings and Precautions (5.7) and Clinical Considerations*].

Available data from published epidemiologic studies and post marketing reports have not established an increased risk of major birth defects or miscarriage. There are risks of persistent pulmonary hypertension of the newborn (PPHN) (see *Data*) and poor neonatal adaptation (see *Clinical Considerations*) with exposure to selective serotonin reuptake inhibitors (SSRIs), including ESCITALOPRAM OXALATE ODT, during pregnancy. There are risks associated with untreated depression in pregnancy (see *Clinical Considerations*).

In animal reproduction studies, both escitalopram and racemic citalopram have been shown to have adverse effects on embryo/fetal and postnatal development, including fetal structural abnormalities, when administered at doses greater than human therapeutic doses (see *Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in the clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-associated maternal risk and/or embryo/fetal risk

Women who discontinue antidepressants are more likely to experience a relapse of major depression than women who continue antidepressants. This finding is from a prospective longitudinal study of 201 pregnant women with a history of major depression, who were euthymic and taking antidepressants at the beginning of pregnancy. Consider the risk of untreated depression when discontinuing or changing treatment with antidepressant medication during pregnancy and postpartum.

Maternal Adverse Reactions

Use of Escitalopram Oxalate ODT in the month before delivery may be associated with an increased risk of postpartum hemorrhage [see *Warnings and Precautions (5.7)*].

Fetal/Neonatal adverse reactions

Neonates exposed to SSRIs or SNRIs, including Escitalopram Oxalate ODT, late in third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Such complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulty, vomiting, hypoglycemia, hypotonia, hypertonia, hyperreflexia, tremor, jitteriness, irritability, and constant crying. These features are consistent with either a direct toxic effect of SSRIs and SNRIs or, possibly, a drug discontinuation syndrome. It should be noted that, in some cases, the clinical picture is consistent with serotonin syndrome [see *Warnings and Precautions (5.2)*].

Data

Human Data

Exposure to SSRIs, particularly later in pregnancy, may increase the risk for PPHN. PPHN occurs in 1-2 per 1000 live births in the general populations and is associated with substantial neonatal morbidity and mortality.

Animal Data

In a rat embryo/fetal development study, oral administration of escitalopram (56, 112, or 150 mg/kg/day) to pregnant animals during the period of organogenesis resulted in decreased fetal body weight and associated delays in ossification at the two higher doses [approximately ≥ 55 times the maximum recommended human dose (MRHD) of 20 mg/day on a mg/m² basis]. Maternal toxicity (clinical signs and decreased body weight gain and food consumption), mild at 56 mg/kg/day, was present at all dose levels. The developmental no-effect dose of 56 mg/kg/day is approximately 27 times the MRHD of 20 mg on a mg/m² basis. No malformations were observed at any of the doses tested (as high as 73 times the MRHD on a mg/m² basis).

When female rats were treated with escitalopram (6, 12, 24, or 48 mg/kg/day) during pregnancy and through weaning, slightly increased offspring mortality and growth retardation were noted at 48 mg/kg/day which is approximately 23 times the MRHD of 20 mg on a mg/m² basis. Slight maternal toxicity (clinical signs and decreased body weight gain and food consumption) was seen at this dose. Slightly increased offspring mortality was also seen at 24 mg/kg/day. The no-effect dose was 12 mg/kg/day which is approximately 6 times the MRHD of 20 mg on a mg/m² basis.

In two rat embryo/fetal development studies, oral administration of racemic citalopram (32, 56, or 112 mg/kg/day) to pregnant animals during the period of organogenesis resulted in decreased embryo/fetal growth and survival and an increased incidence of fetal abnormalities (including cardiovascular and skeletal defects) at the high dose, which is approximately 18 times the MRHD of 60 mg/day on a mg/m² basis. This dose was also associated with maternal toxicity (clinical signs, decreased body weight gain). The developmental no-effect dose was 56 mg/kg/day is approximately 9 times the MRHD on a mg/m² basis. In a rabbit study, no adverse effects on embryo/fetal development were observed at doses of racemic citalopram of up to 16 mg/kg/day, or approximately 5 times the MRHD on a mg/m² basis. Thus, developmental effects of racemic citalopram were observed at a maternally toxic dose in the rat and were not observed in the rabbit.

When female rats were treated with racemic citalopram (4.8, 12.8, or 32 mg/kg/day) from late gestation through weaning, increased offspring mortality during the first 4 days after birth and persistent offspring growth retardation were observed at the highest dose, which is approximately 5 times the MRHD of 60 mg on a mg/m² basis. The no-effect dose was 12.8 mg/kg/day is approximately 2 times the MRHD on a mg/m² basis. Similar effects on offspring mortality and growth were seen when dams were treated throughout gestation and early lactation at doses ≥ 24 mg/kg/day, approximately 4 times the MRHD on a mg/m² basis. A no-effect dose was not determined in that study.

8.2 Lactation

Risk Summary

Data from the published literature report the presence of escitalopram and desmethylescitalopram in human milk (*see Data*). There are reports of excessive sedation, restlessness, agitation, poor feeding and poor weight gain in infants exposed to escitalopram, through breast milk (*see Clinical Considerations*). There are no data on the effects of escitalopram or its metabolites on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ESCITALOPRAM OXALATE ODT and any potential adverse effects on the breastfed child from ESCITALOPRAM OXALATE ODT or from the underlying maternal condition.

Clinical Considerations

Infants exposed to ESCITALOPRAM OXALATE ODT should be monitored for excess sedation, restlessness, agitation, poor feeding and poor weight gain.

Data

A study of 8 nursing mothers on escitalopram with daily doses of 10-20 mg/day showed that exclusively breast-fed infants receive approximately 3.9% of the maternal weight-adjusted dose of escitalopram and 1.7% of the maternal weight-adjusted dose of desmethylcitalopram.

8.4 Pediatric Use

Major Depressive Disorder

The safety and effectiveness of Escitalopram Oxalate ODT for the treatment of major depressive disorder have been established in pediatric patients 12 years of age and older. Use of Escitalopram Oxalate ODT for this indication is supported by evidence from adequate and well-controlled studies in adults with additional evidence from an 8-week, flexible-dose, placebo-controlled study that compared Escitalopram Oxalate ODT 10 mg to 20 mg once daily to placebo in pediatric patients 12 to 17 years of age with major depressive disorder [see *Clinical Studies (14.1)*]. The safety of Escitalopram Oxalate ODT was similar to adult patients with MDD [see *Adverse Reactions (6.1)*].

The safety and effectiveness of Escitalopram Oxalate ODT for the treatment of major depressive disorder have not been established in pediatric patients younger than 12 years of age. In a 24-week, open-label safety study in 118 pediatric patients aged 7 to 11 years who had major depressive disorder, the safety findings were consistent with the known safety and tolerability profile for Escitalopram Oxalate ODT.

Generalized Anxiety Disorder

The safety and effectiveness of Escitalopram Oxalate ODT for the treatment of generalized anxiety disorder have been established in pediatric patients 7 years of age and older. Use of Escitalopram Oxalate ODT for this indication is supported by evidence from adequate and well-controlled studies in adults with additional evidence from an 8-week, flexible-dose, placebo-controlled study that compared Escitalopram Oxalate ODT 10 mg to 20 mg once daily to placebo in pediatric patients 7 to 17 years of age with GAD [see *Clinical Studies (14.2)*]. The safety of Escitalopram Oxalate ODT was similar to adult patients with GAD [see *Adverse Reactions (6.1)*].

The safety and effectiveness of Escitalopram Oxalate ODT for the treatment of generalized anxiety disorder have not been established in pediatric patients younger than 7 years of age.

Antidepressants increase the risk of suicidal thoughts and behaviors in pediatric patients [see *Warnings and Precautions (5.1)*]. Decreased appetite and weight loss have been observed in association with the use of SSRIs. Consequently, regular monitoring of weight and growth should be performed in children and adolescents treated with an SSRI such as Escitalopram Oxalate ODT.

Juvenile Animal Toxicity Data

In a juvenile animal study, male and female rats were administered escitalopram at 5, 40, or 80 mg/kg/day by oral gavage from postnatal day (PND) 21 to PND 69. A delay in sexual maturation was observed in both males and females at ≥ 40 mg/kg/day with a No Observed Adverse Effect Level (NOAEL) of 5 mg/kg/day. This NOAEL was associated with plasma AUC levels less than those measured at the maximum recommended dose (MRHD) in pediatrics (20 mg). However, there was no effect on reproductive function. Increased motor activity (both ambulatory and fine movements) was observed in females prior to daily dosing at ≥ 40 mg/kg/day (3.5 times the MRHD based on AUC levels). A reversible disruption of learning and memory function was observed in males at 80 mg/kg/day with a NOAEL of 40 mg/kg/day, which was associated with an AUC level 3.5 times those measured at the MRHD in pediatrics. There was no effect on learning and memory function in treated female rats.

8.5 Geriatric Use

Approximately 69 patients (6%) of the 1,144 patients receiving escitalopram in controlled trials of Escitalopram Oxalate ODT in major depressive disorder and GAD were 60 years of age or older [see *Clinical Studies (14.1, 14.2)*]. The number of elderly patients in these trials

was insufficient to adequately assess for possible differential efficacy and safety measures on the basis of age. Nevertheless, greater sensitivity of some elderly individuals to effects of Escitalopram Oxalate ODT cannot be ruled out.

In two pharmacokinetic studies, escitalopram half-life was increased by approximately 50% in subjects 65 years and older as compared to young subjects and C_{max} was unchanged [see *Clinical Pharmacology (12.3)*]. The recommended dosage of ESCITALOPRAM OXALATE ODT for elderly patients is 10 mg daily [see *Dosage and Administration (2.5)*].

SSRIs, including Escitalopram Oxalate ODT, have been associated with cases of clinically significant hyponatremia in elderly patients, who may be at greater risk for this adverse reaction [see *Warnings and Precautions (5.6)*].

Of 4,422 patients in clinical studies of racemic citalopram, 1,357 were 60 and over, 1,034 were 65 and over, and 457 were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the geriatric and younger patients, but again, greater sensitivity of some elderly individuals cannot be ruled out.

8.6 Hepatic Impairment

Increased citalopram exposure occurs in patients with hepatic impairment [see *Clinical Pharmacology (12.3)*]. The recommended dosage of ESCITALOPRAM OXALATE ODT in patients with hepatic impairment is 10 mg daily [see *Dosage and Administration (2.5)*].

8.7 Renal Impairment

Pharmacokinetics of Escitalopram Oxalate ODT in patients with a creatinine clearance less than 20 mL/minute has not been evaluated. No dosage adjustment is necessary for patients with mild or moderate renal impairment [see *Dosage and Administration (2.5)*, *Clinical Pharmacology (12.3)*].

9 DRUG ABUSE AND DEPENDENCE

9.2 Abuse and Dependence

Physical and Psychological Dependence

Animal studies suggest that the abuse liability of racemic citalopram is low. Escitalopram Oxalate ODT has not been systematically studied in humans for its potential for abuse, tolerance, or physical dependence. The premarketing clinical experience with Escitalopram Oxalate ODT did not reveal any drug-seeking behavior. However, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate Escitalopram Oxalate ODT patients for history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse (e.g., development of tolerance, incrementations of dose, drug-seeking behavior).

10 OVERDOSAGE

The following have been reported with Escitalopram Oxalate ODT tablet overdose:

- Seizures, which may be delayed, and altered mental status including coma.
- Cardiovascular toxicity, which may be delayed, including QRS and QTc interval prolongation, wide complex tachyarrhythmias, and torsade de pointes. Hypertension most commonly seen, but rarely can see hypotension alone or with co-ingestants including alcohol.
- Serotonin syndrome (patients with a multiple drug overdose with other proserotonergic drugs may have a higher risk).


Prolonged cardiac monitoring is recommended in Escitalopram Oxalate ODT overdose ingestions due to the arrhythmia risk.

Gastrointestinal decontamination with activated charcoal should be considered in patients who present early after an Escitalopram Oxalate ODT overdose.

Consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdose management recommendations.

11 DESCRIPTION

Escitalopram Oxalate ODT contains escitalopram aselective serotonin reuptake inhibitor (SSRI), present as escitalopram oxalate salt. Escitalopram is the pure S- enantiomer (single isomer) of the racemic bicyclic phthalane derivative citalopram. Escitalopram oxalate is designated S-(+)-1-[3(dimethyl-amino)propyl]-1-(p-fluorophenyl)-5-phthalanecarbonitrile oxalate with the following structural formula:



•C₂H₂O₄

The molecular formula is C₂₀H₂₁FN₂O • C₂H₂O₄ and the molecular weight is 414.40.

Escitalopram oxalate occurs as a fine, white to slightly-yellow powder and is freely soluble in methanol and dimethyl sulfoxide (DMSO), soluble in isotonic saline solution, sparingly soluble in water and ethanol, slightly soluble in ethyl acetate, and insoluble in heptane. Escitalopram Oxalate ODT, for oral use, are, *shape TBD* tablets containing 6.38 mg, 12.75 mg, 25.5 mg escitalopram oxalate in strengths equivalent to 5 mg, 10 mg, 15mg and 20 mg, respectively, of escitalopram base. The tablets also contain the following inactive ingredients: *inactive ingredients TBD, will be disclosed in the ANDA filing.*

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of antidepressant action of escitalopram, the S-enantiomer of racemic citalopram, is presumed to be linked to potentiation of serotonergic activity in the central nervous system (CNS) resulting from its inhibition of CNS neuronal reuptake of serotonin (5-HT).

12.2 Pharmacodynamics

In vitro and *in vivo* studies in animals suggest that escitalopram is a highly selective serotonin reuptake inhibitor (SSRI) with minimal effects on norepinephrine and dopamine neuronal reuptake. Escitalopram is at least 100-fold more potent than the R-enantiomer with respect to inhibition of 5-HT reuptake and inhibition of 5-HT neuronal firing rate. Tolerance to a model of antidepressant effect in rats was not induced by long-term (up to 5 weeks) treatment with escitalopram. Escitalopram has no or very low affinity for serotonergic (5-HT₁₋₇) or other receptors including alpha- and beta-adrenergic, dopamine (D₁₋₅), histamine (H₁₋₃), muscarinic (M₁₋₅), and benzodiazepine receptors. Escitalopram also does not bind to, or has low affinity for, various ion channels including Na⁺, K⁺, Cl⁻, and Ca⁺⁺ channels. Antagonism of muscarinic, histaminergic, and adrenergic receptors has been hypothesized to be associated with various anticholinergic, sedative, and cardiovascular side effects of other psychotropic drugs.

12.3 Pharmacokinetics

The single- and multiple-dose pharmacokinetics of escitalopram are linear and dose-proportional in a dose range of 10 to 30 mg/day.

With once-daily dosing, steady state plasma concentrations are achieved within approximately one week. At steady state, the extent of accumulation of escitalopram in plasma in young healthy subjects was 2.2-2.5 times the plasma concentrations observed after a single dose.

Absorption

The absolute bioavailability of citalopram is about 80% relative to an intravenous dose.

Following a single oral dose (20 mg tablet or solution) of escitalopram, peak blood levels occur at about 5 hours. Absorption of escitalopram is not affected by food.

Distribution

The binding of escitalopram to human plasma proteins is approximately 56%. The volume of distribution of citalopram is about 12 L/kg. Data specific on escitalopram are unavailable.

Elimination

Biotransformation of escitalopram is mainly hepatic, with a mean terminal half-life of about 27-32 hours. The oral clearance of escitalopram is 600 mL/min, with approximately 7% of that due to renal clearance.

Metabolism

Escitalopram is metabolized to S-DCT and S-didemethylcitalopram (S-DDCT). In humans, unchanged escitalopram is the predominant compound in plasma. At steady state, the concentration of the escitalopram metabolite S-DCT in plasma is approximately one-third that of escitalopram. The level of S-DDCT was not detectable in most subjects. *In vitro* studies show that escitalopram is at least 7 and 27 times more potent than S-DCT and S-DDCT, respectively, in the inhibition of serotonin reuptake, suggesting that the metabolites of escitalopram do not contribute significantly to the antidepressant actions of escitalopram. S-DCT and S-DDCT also have no or very low affinity for serotonergic (5-HT₁₋₇) or other receptors including alpha- and beta-adrenergic, dopamine (D₁₋₅), histamine (H₁₋₃), muscarinic (M₁₋₅), and benzodiazepine receptors. S-DCT and S-DDCT also do not bind to various ion channels including Na⁺, K⁺, Cl⁻, and Ca⁺⁺ channels. *In vitro* studies using human liver microsomes indicated that CYP3A4 and CYP2C19 are the primary isozymes involved in the N-demethylation of escitalopram.

Excretion

Following oral administrations of escitalopram, the fraction of drug recovered in the urine as escitalopram and S-demethylcitalopram (S-DCT) is about 8% and 10%, respectively.

Specific Populations

Pediatric Patients

Pediatric patients 7 to 11 years of age: Based on population PK simulations, following multiple dosing of 20 mg/day escitalopram, steady-state C_{max} and AUC_{tau} of escitalopram were increased by 93% and 86%, respectively in pediatric patients with GAD 7 to 11 years of age compared to adults.

Pediatric patients 12 to 17 years of age: In a single dose study of 10 mg escitalopram, AUC of escitalopram decreased by 19%, and C_{max} increased by 26% in healthy pediatric subjects 12 to 17 years of age compared to adults. Following multiple dosing of 40 mg/day citalopram, escitalopram elimination half-life, steady-state C_{max} and AUC were similar in pediatric patients 12 to 17 years of age with MDD compared to adults [see *Use in Specific Populations (8.4)*].

Geriatric Patients

Escitalopram pharmacokinetics in subjects ≥ 65 years of age were compared to adults in a single-dose and a multiple-dose study. Escitalopram AUC and half-life were increased by approximately 50% in elderly subjects, and C_{\max} was unchanged [see *Dosage and Administration (2.5)*, *Use in Specific Populations (8.5)*].

Male and Female Patients

Based on data from single- and multiple-dose studies measuring escitalopram in elderly, young adults, and adolescents, no dosage adjustment on the basis of gender is needed.

Patients with Hepatic Impairment

Citalopram oral clearance was reduced by 37% and half-life was doubled in patients with reduced hepatic function compared to normal subjects [see *Dosage and Administration (2.5)*, *Use in Specific Populations (8.6)*].

Patients with Renal Impairment

In patients with mild to moderate renal function impairment, oral clearance of citalopram was reduced by 17% compared to normal subjects. No information is available about the pharmacokinetics of escitalopram in patients with severely reduced renal function (creatinine clearance < 20 mL/min) [see *Use in Specific Populations (8.7)*].

Drug Interaction Studies

In vitro enzyme inhibition data did not reveal an inhibitory effect of escitalopram on CYP3A4, -1A2, -2C9, -2C19, and -2E1. Based on *in vitro* data, escitalopram would be expected to have little inhibitory effect on *in vivo* metabolism mediated by these cytochromes. While *in vivo* data to address this question are limited, results from drug interaction studies suggest that escitalopram, at a dose of 20 mg, has no 3A4 inhibitory effect and a modest 2D6 inhibitory effect [see *Drug Interactions (7)*].

CYP3A4 and CYP2C19 Inhibitors

In vitro studies indicated that CYP3A4 and -2C19 are the primary enzymes involved in the metabolism of escitalopram. However, coadministration of escitalopram (20 mg) and ritonavir (600 mg), a potent inhibitor of CYP3A4, did not significantly affect the pharmacokinetics of escitalopram. Because escitalopram is metabolized by multiple enzyme systems, inhibition of a single enzyme may not appreciably decrease escitalopram clearance.

Cimetidine

In subjects who had received 21 days of 40 mg/day racemic citalopram, combined administration of 400 mg twice a day cimetidine for 8 days resulted in an increase in citalopram AUC and C_{\max} of 43% and 39%, respectively. The clinical significance of these findings is unknown.

Digoxin

In subjects who had received 21 days of 40 mg/day racemic citalopram, combined administration of citalopram and digoxin (single dose of 1 mg) did not significantly affect the pharmacokinetics of either citalopram or digoxin.

Lithium

Coadministration of racemic citalopram (40 mg/day for 10 days) and lithium (30 mmol/day for 5 days) had no significant effect on the pharmacokinetics of citalopram or lithium. Plasma lithium levels should be monitored with appropriate adjustment to the lithium dose in accordance with standard clinical practice. Because lithium may enhance the serotonergic effects of escitalopram, caution should be exercised when Escitalopram Oxalate ODT and lithium are co-administered.

Theophylline

Combined administration of racemic citalopram (40 mg/day for 21 days) and the CYP1A2 substrate theophylline (single dose of 300 mg) did not affect the pharmacokinetics of theophylline. The effect of theophylline on the pharmacokinetics of citalopram was not evaluated.

Ketoconazole

Combined administration of racemic citalopram (40 mg) and ketoconazole (200 mg), a potent CYP3A4 inhibitor, decreased the C_{max} and AUC of ketoconazole by 21% and 10%, respectively, and did not significantly affect the pharmacokinetics of citalopram.

Ritonavir

Combined administration of a single dose of ritonavir (600 mg), both a CYP3A4 substrate and a potent inhibitor of CYP3A4, and escitalopram (20 mg) did not affect the pharmacokinetics of either ritonavir or escitalopram.

Triazolam

Combined administration of racemic citalopram (titrated to 40 mg/day for 28 days) and the CYP3A4 substrate triazolam (single dose of 0.25 mg) did not significantly affect the pharmacokinetics of either citalopram or triazolam.

Metoprolol

Administration of 20 mg/day Escitalopram Oxalate ODT for 21 days in healthy volunteers resulted in a 50% increase in C_{max} and 82% increase in AUC of the beta-adrenergic blocker metoprolol (given in a single dose of 100 mg). Increased metoprolol plasma levels have been associated with decreased cardioselectivity. Coadministration of Escitalopram Oxalate ODT and metoprolol had no clinically significant effects on blood pressure or heart rate.

Alcohol

Escitalopram Oxalate ODT did not potentiate the cognitive and motor effects of alcohol in a clinical trial. As with other psychotropic medications, the use of alcohol by patients taking Escitalopram Oxalate ODT is not recommended.

Warfarin

Administration of 40 mg/day racemic citalopram for 21 days did not affect the pharmacokinetics of warfarin, a CYP3A4 substrate. Prothrombin time was increased by 5%. The clinical significance of these findings is unknown.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Racemic citalopram was administered in the diet to NMRI/BOM strain mice and COBS WI strain rats for 18 and 24 months, respectively. There was no evidence for carcinogenicity of racemic citalopram in mice receiving up to 240 mg/kg/day. There was an increased incidence of small intestine carcinoma in rats receiving 8 or 24 mg/kg/day racemic citalopram. A no-effect dose for this finding was not established. The relevance of these findings to humans is unknown.

Mutagenesis

Racemic citalopram was mutagenic in the *in vitro* bacterial reverse mutation assay (Ames test) in 2 of 5 bacterial strains (Salmonella TA98 and TA1537) in the absence of metabolic activation. It was clastogenic in the *in vitro* Chinese hamster lung cell assay for chromosomal aberrations in the presence and absence of metabolic activation. Racemic citalopram was not mutagenic in the *in vitro* mammalian forward gene mutation assay (HPRT) in mouse lymphoma cells or in a coupled *in vitro/in vivo* unscheduled DNA synthesis (UDS) assay in rat liver. It was not clastogenic in the *in vitro* chromosomal aberration assay in human lymphocytes or in two *in vivo* mouse micronucleus assays.

Impairment of Fertility

When racemic citalopram was administered orally to 16 male and 24 female rats prior to and throughout mating and gestation at doses of 32, 48, and 72 mg/kg/day, mating was decreased at all doses, and fertility was decreased at doses \geq 32 mg/kg/day. Gestation duration was increased at 48 mg/kg/day.

13.2 Animal Toxicology and/or Pharmacology

Retinal Changes in Rats

Pathologic changes (degeneration/atrophy) were observed in the retinas of albino rats in the 2-year carcinogenicity study with racemic citalopram. There was an increase in both incidence and severity of retinal pathology in both male and female rats receiving 80 mg/kg/day. Similar findings were not present in rats receiving 24 mg/kg/day of racemic citalopram for two years, in mice receiving up to 240 mg/kg/day of racemic citalopram for 18 months, or in dogs receiving up to 20 mg/kg/day of racemic citalopram for one year.

Additional studies to investigate the mechanism for this pathology have not been performed, and the potential significance of this effect in humans has not been established.

Cardiovascular Changes in Dogs

In a one-year toxicology study, 5 of 10 beagle dogs receiving oral racemic citalopram doses of 8 mg/kg/day died suddenly between weeks 17 and 31 following initiation of treatment. Sudden deaths were not observed in rats at doses of racemic citalopram up to 120 mg/kg/day, which produced plasma levels of citalopram and its metabolites demethylcitalopram and didemethylcitalopram (DDCT) similar to those observed in dogs at 8 mg/kg/day. A subsequent intravenous dosing study demonstrated that in beagle dogs, racemic DDCT caused QT prolongation, a known risk factor for the observed outcome in dogs.

14 CLINICAL STUDIES

14.1 Major Depressive Disorder

Adults

The efficacy of Escitalopram Oxalate ODT as a treatment for major depressive disorder was established in three, 8-week, placebo-controlled studies conducted in outpatients between 18 and 65 years of age who met DSM-IV criteria for major depressive disorder. The primary outcome in all three studies was change from baseline to endpoint in the Montgomery Asberg Depression Rating Scale (MADRS).

A fixed-dose study compared 10 mg daily Escitalopram Oxalate ODT and 20 mg daily Escitalopram Oxalate ODT to placebo and 40 mg daily citalopram. The 10 mg daily and 20 mg daily Escitalopram Oxalate ODT treatment groups showed statistically significant greater mean improvement compared to placebo on the MADRS. The 10 mg and 20 mg Escitalopram Oxalate ODT groups were similar on this outcome measure.

In a second fixed-dose study of 10 mg daily Escitalopram Oxalate ODT and placebo, the 10 mg daily Escitalopram Oxalate ODT treatment group showed statistically significant greater mean improvement compared to placebo on the MADRS.

In a flexible-dose study, comparing Escitalopram Oxalate ODT, titrated between 10 mg and 20 mg daily, to placebo and citalopram, titrated between 20 mg and 40 mg daily, the Escitalopram Oxalate ODT treatment group showed statistically significant greater mean improvement compared to placebo on the MADRS.

Analyses of the relationship between treatment outcome and age, gender, and race did not suggest any differential responsiveness on the basis of these patient characteristics.

In a longer-term trial, 274 patients meeting (DSM-IV) criteria for major depressive disorder, who had responded during an initial 8 week, open-label treatment phase with Escitalopram Oxalate ODT 10 mg or 20 mg daily, were randomized to continuation of Escitalopram Oxalate ODT at their same dose, or to placebo, for up to 36 weeks of observation for relapse. Response during the open-label phase was defined by having a decrease of the MADRS total score to \leq 12. Relapse during the double-blind phase was defined as an increase of the MADRS total

score to ≥ 22 , or discontinuation due to insufficient clinical response. Patients receiving continued Escitalopram Oxalate ODT experienced a statistically significant longer time to relapse compared to those receiving placebo.

Pediatric Patients 12 years of age and older

The efficacy of Escitalopram Oxalate ODT as a treatment for major depressive disorder in pediatric patients 12 to 17 years was established in an 8-week, flexible-dose, placebo-controlled study that compared Escitalopram Oxalate ODT (10 mg to 20 mg daily) to placebo in outpatients 12 to 17 years of age inclusive who met DSM-IV criteria for major depressive disorder (MDD). The primary outcome was change from baseline to endpoint in the Children's Depression Rating Scale - Revised (CDRS-R). In this study, Escitalopram Oxalate ODT showed statistically significant greater mean improvement compared to placebo on the CDRS-R.

The efficacy of Escitalopram Oxalate ODT in the treatment of major depressive disorder in pediatric patients 12 to 17 years was established, in part, on the basis of extrapolation from the 8-week, flexible-dose, placebo-controlled study with racemic citalopram 20 mg to 40 mg daily. In this outpatient study in pediatric patients 7 to 17 years of age who met DSM-IV criteria for major depressive disorder, citalopram treatment showed statistically significant greater mean improvement from baseline, compared to placebo, on the CDRS-R; the positive results for this trial largely came from the 12 to 17 year subgroup.

Two additional flexible-dose, placebo-controlled MDD studies (one Escitalopram Oxalate ODT study in patients ages 7 to 17 years and one citalopram study patients 13 to 18 years) did not demonstrate efficacy. The safety and effectiveness of Escitalopram Oxalate ODT have not been established in pediatric patients less than 12 years of age with MDD.

14.2 Generalized Anxiety Disorder

Adults

The efficacy of Escitalopram Oxalate ODT in the treatment of generalized anxiety disorder (GAD) in adults was demonstrated in three, 8-week, multicenter, flexible-dose, placebo-controlled studies that compared Escitalopram Oxalate ODT (10 mg to 20 mg daily) to placebo in outpatients between 18 and 80 years of age who met DSM-IV criteria for GAD. In all three studies, Escitalopram Oxalate ODT showed statistically significant greater mean improvement compared to placebo on the Hamilton Anxiety Scale (HAM-A).

There were too few patients in differing ethnic and age groups to adequately assess whether or not Escitalopram Oxalate ODT has differential effects in these groups. There was no difference in response to Escitalopram Oxalate ODT between men and women.

Pediatric Patients 7 years of age and older

The efficacy of Escitalopram Oxalate ODT in the treatment for generalized anxiety disorder (GAD) in pediatric patients 7 to 17 years was established in an 8-week, flexible-dose, placebo-controlled study that compared Escitalopram Oxalate ODT (10 mg to 20 mg daily) to placebo in outpatients 7 to 17 years of age who met DSM-V criteria for GAD. The primary outcome was change from baseline to Week 8 in the Pediatric Anxiety Rating Scale (PARS) severity score for GAD. In this study, Escitalopram Oxalate ODT showed a statistically significant treatment difference when compared to placebo on the PARS severity score for GAD (Least squares mean difference: -1.42 , 95% confidence interval [-2.69 , -0.15]).

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Orally Disintegrating Tablets

5 mg Orally Disintegrating Tablets:

Bottle of *TBD* NDC # XXXX-XXXX-XX

TBD to be disclosed at ANDA filing.

10 mg Orally Disintegrating Tablets:

Bottle of *TBD* NDC # XXXX-XXXX-XX

TBD to be disclosed at ANDA filing.

15mg Orally Disintegrating Tablets:

Bottle of *TBD* NDC # XXXX-XXXX-XX

TBD to be disclosed at ANDA filing.

20 mg Orally Disintegrating Tablets:
Bottle of TBD NDC # XXXX-XXXX-XX
TBD to be disclosed at ANDA filing.

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15 to 30°C (59° to 86°F).

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Suicidal Thoughts and Behaviors

Advise patients, their families and caregivers to look for the emergence of suicidal ideation and behavior, especially during treatment and when the dose is adjusted up or down and instruct them to report such symptoms to their healthcare provider [see *Boxed Warning and Warnings and Precautions (5.1)*].

Serotonin Syndrome

Caution patients about the risk of serotonin syndrome, particularly with the with the concomitant use of Escitalopram Oxalate ODT with other serotonergic drugs including triptans, tricyclic antidepressants, opioids, lithium, tryptophan, buspirone, amphetamines, and St. John’s Wort, and with drugs that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat psychiatric disorders and also others, such as linezolid). Instruct patients to contact their health care provider or report to the emergency room if they experience signs or symptoms of serotonin syndrome [see *Warnings and Precautions (5.2), Drug Interactions (7)*].

Discontinuation Syndrome

Advise patients not to abruptly discontinue Escitalopram Oxalate ODT and to discuss any tapering regimen with their healthcare provider. Inform patients that adverse reactions can occur when Escitalopram Oxalate ODT is discontinued [see *Warnings and Precautions (5.3)*].

Activation of Mania or Hypomania

Advise patients and their caregivers to observe for signs of activation of mania/hypomania and instruct them to report such symptoms to the healthcare provider [see *Warnings and Precautions (5.5)*].

Increased Risk of Bleeding

Inform patients about the concomitant use of Escitalopram Oxalate ODT with NSAIDs, aspirin, warfarin, other antiplatelet drugs, or other anticoagulants because the combined use has been associated with an increased risk of bleeding. Advise patients to inform their healthcare providers if they are taking or planning to take any prescription or over-the-counter medications that increase the risk of bleeding [see *Warnings and Precautions (5.7)*].

Angle Closure Glaucoma

Advise patients that taking Escitalopram Oxalate ODT can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle closure glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible [see *Warnings and Precautions (5.9)*].

Sexual Dysfunction

Advise patients that use of Escitalopram Oxalate ODT may cause symptoms of sexual dysfunction in both male and female patients. Inform patients that they should discuss any changes in sexual function and potential management strategies with their healthcare provider [see *Warnings and Precautions (5.11)*].

Concomitant Medications

Since escitalopram is the active isomer of racemic citalopram (Celexa), the two agents should not be co-administered. Patients should be advised to inform their physician if they are taking, or plan to take, any prescription or over-the-counter drugs, as there is a potential for interactions.

Interference with Psychomotor Performance

Because psychoactive drugs may impair judgment, thinking, or motor skills, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that Escitalopram Oxalate ODT therapy does not affect their ability to engage in such activities.

Alcohol

Patients should be told that, although Escitalopram Oxalate ODT has not been shown in experiments with normal subjects to increase the mental and motor skill impairments caused by alcohol, the concomitant use of Escitalopram Oxalate ODT and alcohol in depressed patients is not advised.

Pregnancy

Advise pregnant women to notify their healthcare providers if they become pregnant or intend to become pregnant during treatment with ESCITALOPRAM OXALATE ODT.

Advise patients that ESCITALOPRAM OXALATE ODT use later in pregnancy may lead to increased risk for neonatal complications requiring prolonged hospitalization, respiratory support, tube feeding, and/or persistent pulmonary hypertension (PPHN) of the newborn [see *Use in Specific Populations (8.1)*].

Advise women that there is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to ESCITALOPRAM OXALATE ODT during pregnancy [see *Use in Specific Populations (8.1)*].

Lactation

Advise breastfeeding women using ESCITALOPRAM OXALATE ODT to monitor infants for excess sedation, restlessness, agitation, poor feeding and poor weight gain and to seek medical care if they notice these signs [see *Use in Specific Populations (8.2)*].

Distributed by:

TBD to be disclosed at ANDA filing.

MEDICATION GUIDE
Escitalopram Oxalate
ODT (escitalopram
ODT)

What is the most important information I should know about Escitalopram Oxalate ODT? Escitalopram Oxalate ODT may cause serious side effects, including:

- **Increased risk of suicidal thoughts or actions.** Escitalopram Oxalate ODT and other antidepressant medicines increase the risk of suicidal thoughts and actions in people 24 years of age and younger, **especially within the first few months of treatment or when the dose is changed.**
 - **Depression or other mental illnesses are the most important causes of suicidal thoughts or actions.**
- How can I watch for and try to prevent suicidal thoughts and actions?**
- Pay close attention to any changes, especially sudden changes in mood, behavior, thoughts, or feelings, or if you or your child develop suicidal thoughts or actions. This is very important when an antidepressant medicine is started or when the dose is changed.
 - Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings or if you or your child develop suicidal thoughts or actions.
 - Keep all follow-up visits with your healthcare provider as scheduled and call your healthcare provider between visits if you are worried about symptoms.

Call your healthcare provider or get emergency medical help right away if you or your child have any of the following symptoms, especially if they are new, worse, or worry you:

- | | |
|--|---|
| • attempts to commit suicide | • acting on dangerous impulses |
| • acting aggressive, being angry or violent | • thoughts about suicide or dying |
| • new or worse depression | • new or worsening anxiety |
| • panic attacks | • feeling very agitated or restless |
| • new or worse irritability | • trouble sleeping |
| • an extreme increase in activity or talking (mania) | • other unusual changes in behavior or mood |

What is Escitalopram Oxalate ODT?

Escitalopram Oxalate ODT is a prescription medicine used to treat:

- a certain type of depression called Major Depressive Disorder (MDD) in adults and children 12 years of age and older
- Generalized Anxiety Disorder (GAD) in adults and children 7 years of age and older

It is not known if Escitalopram Oxalate ODT is safe and effective for use in children under 12 years of age with MDD or children under 7 years of age with GAD.

Do not take Escitalopram Oxalate ODT if you or your child:

- are taking, or have stopped taking within the last 14 days, a medicine called a monoamine oxidase inhibitor (MAOI), including the antibiotic linezolid or intravenous methylene blue
- are taking the antipsychotic medicine pimozide
- are allergic to escitalopram or citalopram or any of the ingredients in Escitalopram Oxalate ODT. See the end of this Medication Guide for a complete list of ingredients in Escitalopram Oxalate ODT.

Ask your healthcare provider or pharmacist if you are not sure if you or your child take an MAOI, including the antibiotic linezolid or intravenous methylene blue.

Do not start taking an MAOI for at least 14 days after you or your child have stopped treatment with Escitalopram Oxalate ODT.

Before taking Escitalopram Oxalate ODT, tell your healthcare provider about all your medical conditions, including if you or your child:

- have or had seizures or convulsions
- have, or have a family history of bipolar disorder, mania, or hypomania
- have low blood sodium levels
- have or had bleeding problems
- have high pressure in the eye (glaucoma)
- have heart, liver, or kidney problems
- are pregnant or plan to become pregnant. Escitalopram Oxalate ODT may harm the unborn baby. Taking Escitalopram Oxalate ODT during the third trimester of pregnancy may cause the baby to have withdrawal symptoms, or breathing, temperature control, feeding, or other problems after birth. Talk to your healthcare provider about the risks to the baby if you or your child take Escitalopram Oxalate ODT during pregnancy.
 - Tell your healthcare provider right away if you or your child become pregnant or think you may be pregnant during treatment with Escitalopram Oxalate ODT.
 - There is a pregnancy registry for females who are exposed to Escitalopram Oxalate ODT during pregnancy. The purpose of the registry is to collect information about the health of females exposed to Escitalopram Oxalate ODT and their baby. If you or your child become pregnant during treatment with Escitalopram Oxalate ODT, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or visit online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.
- are breastfeeding or plan to breastfeed. Escitalopram Oxalate ODT passes into breast milk and may harm the baby. Talk to your healthcare provider about the best way to feed the baby during treatment with Escitalopram Oxalate ODT.
 - If you or your child breastfeed during treatment with Escitalopram Oxalate ODT, call your healthcare provider if the baby develops sleepiness or fussiness, or is not feeding or gaining weight well.

Tell your healthcare provider about all the medicines you or your child take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Escitalopram Oxalate ODT and some medicines may affect each other and may cause serious side effects.

Escitalopram Oxalate ODT may affect the way other medicines work and other medicines may affect the way Escitalopram Oxalate ODT works.

Especially tell your healthcare provider if you take:

- medicines used to treat migraine headache known as triptans
- tricyclic antidepressants
- lithium
- tramadol, fentanyl, meperidine, methadone, or other opioids
- tryptophan
- buspirone
- amphetamines
- St. John's Wort
- medicines used to treat mood, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs)
- diuretics
- medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs) and warfarin

Ask your healthcare provider if you are not sure if you or your child are taking any of these medicines. Your healthcare provider can tell you if it is safe to take Escitalopram Oxalate ODT with your other medicines.

Do not start or stop any other medicines during treatment with Escitalopram Oxalate ODT without talking to your healthcare provider first. Stopping Escitalopram Oxalate ODT suddenly may cause you or your child to have serious side effects. See, **“What are the possible side effects of Escitalopram Oxalate ODT?”**

Know the medicines you or your child take. Keep a list of them to show your healthcare provider and pharmacist when you get new medicine.

How should I take Escitalopram Oxalate ODT?

- Take Escitalopram Oxalate ODT exactly as prescribed. Your healthcare provider may need to change the dose of Escitalopram Oxalate ODT until it is the right dose for you or your child.
- Take Escitalopram Oxalate ODT 1 time each day, in the morning or the evening.
- Take Escitalopram Oxalate ODT with or without food.
- If you or your child take too much Escitalopram Oxalate ODT, call your healthcare provider or Poison Help Line at 1-800-222-1222, or go to the nearest hospital emergency room right away.

What should I avoid while taking Escitalopram Oxalate ODT?

- **Do not** drive, operate heavy machinery, or do other dangerous activities until you know how Escitalopram Oxalate ODT affects you. Escitalopram Oxalate ODT can cause sleepiness or may affect your ability to make decisions, think clearly, or react quickly.
- **Do not** drink alcohol during treatment with Escitalopram Oxalate ODT.

What are the possible side effects of Escitalopram Oxalate ODT? Escitalopram Oxalate ODT may cause serious side effects, including:

- See “**What is the most important information I should know about Escitalopram Oxalate ODT?**”
- **Serotonin syndrome.** A potentially life-threatening problem called serotonin syndrome can happen when Escitalopram Oxalate ODT is taken with certain other medicines. See “**Do not take Escitalopram Oxalate ODT if you?**” **Call your healthcare provider or go to the nearest hospital emergency room right away** if you or your child have any of the following signs and symptoms of serotonin syndrome:
 - agitation
 - confusion
 - fast heartbeat
 - sweating
 - flushing
 - seizures
 - nausea, vomiting, diarrhea
 - seeing or hearing things that are not real (hallucinations)
 - coma
 - blood pressure changes
 - shaking (tremors), stiff muscles, or muscle twitching
 - dizziness
 - high body temperature (hyperthermia)
 - loss of coordination
- **Discontinuation syndrome.** Suddenly stopping Escitalopram Oxalate ODT may cause you or your child to have serious side effects. Your healthcare provider may want to decrease the dose slowly. Symptoms may include:
 - changes in mood
 - irritability and agitation
 - dizziness
 - electric shock sensation (paresthesia)
 - anxiety
 - confusion
 - headache
 - tiredness
 - problems sleeping
 - hypomania
 - ringing in your ears (tinnitus)
 - seizures
- **Seizures (convulsions).**
- **Manic episodes.** Manic episodes may happen in people with bipolar disorder who take Escitalopram Oxalate ODT. Symptoms may include:
 - greatly increased energy
 - racing thoughts
 - unusually grand ideas
 - talking more or faster than usual
 - severe trouble sleeping
 - reckless behavior
 - excessive happiness or irritability

- **Low sodium levels in the blood (hyponatremia).** Low sodium levels in the blood that may be serious and may cause death can happen during treatment with Escitalopram Oxalate ODT. Elderly people and people who take certain medicines may be at greater risk for developing low sodium levels in the blood. Signs and symptoms may include:

- headache
- weakness or feeling unsteady which can lead to falls
- memory problems
- problems concentrating or thinking
- confusion

In more severe or more sudden cases, signs and symptoms include:

- seeing or hearing things that are not real (hallucinations)
- seizures
- stopping breathing (respiratory arrest)
- fainting
- coma

- **Increased risk of bleeding:** Taking Escitalopram Oxalate ODT with aspirin, NSAIDS, warfarin, or other blood thinners may add to this risk. Tell your healthcare provider if you have any unusual bleeding or bruising.
- **Visual problems (angle-closure glaucoma).** Escitalopram Oxalate ODT may cause a type of eye problem called angle-closure glaucoma in people with certain eye problems. You or your child may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your healthcare provider if you or your child have:

- eye pain
- changes in vision
- swelling or redness in or around the eye

- **Sexual problems (dysfunction).** Taking Escitalopram Oxalate ODT may cause sexual problems. Symptoms in males may include:

- delayed ejaculation or inability to have an ejaculation
- problems getting or keeping an erection
- decreased sex drive

Symptoms in females may include:

- decreased sex drive
- delayed orgasm or inability to have an orgasm

Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with Escitalopram Oxalate ODT. There may be treatments your healthcare provider can suggest.

The most common side effects of Escitalopram Oxalate ODT include:

- trouble sleeping
- delayed ejaculation
- nausea
- sweating
- tiredness
- sleepiness
- decreased sex drive
- delayed orgasm or inability to have an orgasm

Height and weight changes in children may happen during treatment with Escitalopram Oxalate ODT. Your child's height and weight should be monitored during treatment with ESCITALOPRAM OXALATE ODT.

These are not all the possible side effects of Escitalopram Oxalate ODT.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Escitalopram Oxalate ODT?

- Store Escitalopram Oxalate ODT at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep Escitalopram Oxalate ODT and all medicines out of the reach of children.**

General information about the safe and effective use of Escitalopram Oxalate ODT.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Escitalopram Oxalate ODT for a condition for which it was not prescribed. Do not give Escitalopram Oxalate ODT to other people, even if they have the same symptoms that you have. It may harm them. You may ask your pharmacist or healthcare provider for information about Escitalopram Oxalate ODT that is written for health professionals.

What are the ingredients in Escitalopram Oxalate ODT?

Active ingredient: escitalopram oxalate

Inactive ingredients:

Orally Disintegrating Tablets: *TBD to be disclosed at ANDA filing.*