



New Data from Verzenio® (abemaciclib) monarchE Study to Be Featured in ESMO Virtual Plenary

September 27, 2021

INDIANAPOLIS, Sept. 27, 2021 /PRNewswire/ -- New data from Eli Lilly and Company's (NYSE: LLY) monarchE study for an investigational use of Verzenio® (abemaciclib), in combination with endocrine therapy, in patients with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2) high risk early breast cancer will be presented at the October 14 European Society for Medical Oncology (ESMO) Virtual Plenary.

Details on this presentation are shared below.

<i>Presentation Date and Time:</i>	Thursday, October 14 at 19:30 CEST
<i>Title:</i>	Adjuvant abemaciclib combined with endocrine therapy (ET): Updated results from monarchE
<i>Authors:</i>	J. O'Shaughnessy et al.
<i>Publication Number:</i>	VP8_2021

The presentation will utilize an April 2021 data cutoff date, allowing for more follow-up relative to the analysis last presented at the San Antonio Breast Cancer Symposium in December 2020.

The abstract is embargoed until the start of the Virtual Plenary session. For more information, please visit: <https://www.esmo.org/meetings/october-virtual-plenary-2021>.

About the monarchE Study

[monarchE](#) is a global randomized, open-label, Phase 3 study in women and men with hormone receptor-positive (HR+), human epidermal growth factor receptor 2-negative (HER2-), node-positive, early breast cancer at high risk of recurrence. High risk of recurrence was defined by disease characteristics: either ≥ 4 positive axillary lymph nodes (pALN) or 1-3 pALN and at least one of the following criteria: tumor size ≥ 5 cm, histologic Grade 3, or Ki-67 index $\geq 20\%$. Patients had completed definitive locoregional therapy.

A total of 5,637 patients were randomized in a 1:1 ratio to receive two years of Verzenio 150 mg twice daily plus physician's choice of standard endocrine therapy, or standard endocrine therapy alone. After the treatment period, all patients will continue on endocrine therapy for five to 10 years, as clinically indicated.

The study's primary endpoint is invasive disease-free survival (IDFS). Secondary endpoints include distant relapse-free survival (DRFS), IDFS for patients with Ki-67 index $\geq 20\%$, and overall survival (OS).

Notes to Editors

About Verzenio® (abemaciclib)

Verzenio® (abemaciclib) is an inhibitor of cyclin-dependent kinases (CDK)4/6, which are activated by binding to D-cyclins. In estrogen receptor-positive (ER+) breast cancer cell lines, cyclin D1 and CDK4/6 promote phosphorylation of the retinoblastoma protein (Rb), cell cycle progression, and cell proliferation.

In vitro, continuous exposure to Verzenio inhibited Rb phosphorylation and blocked progression from G1 to S phase of the cell cycle, resulting in senescence and apoptosis (cell death). Preclinically, Verzenio dosed daily without interruption resulted in reduction of tumor size. Inhibiting CDK4/6 in healthy cells can result in side effects, some of which may be serious. Clinical evidence also suggests that Verzenio crosses the blood-brain barrier. In patients with advanced cancer, including breast cancer, concentrations of Verzenio and its active metabolites (M2 and M20) in cerebrospinal fluid are comparable to unbound plasma concentrations.

Verzenio is Lilly's first solid oral dosage form to be made using a faster, more efficient process known as continuous manufacturing. Continuous manufacturing is a new and advanced type of manufacturing within the pharmaceutical industry, and Lilly is one of the first companies to use this technology.

INDICATION FOR VERZENIO

Verzenio is indicated for the treatment of HR+, HER2- advanced or metastatic breast cancer:

- in combination with an aromatase inhibitor for postmenopausal women as initial endocrine-based therapy
- in combination with fulvestrant for women with disease progression following endocrine therapy
- as a single agent for adult patients with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting

IMPORTANT SAFETY INFORMATION FOR VERZENIO (abemaciclib)

Diarrhea occurred in 81% of patients receiving Verzenio plus an aromatase inhibitor in MONARCH 3, 86% of patients receiving Verzenio plus fulvestrant in MONARCH 2 and 90% of patients receiving Verzenio alone in MONARCH 1. Grade 3 diarrhea occurred in 9% of patients receiving Verzenio plus an aromatase inhibitor in MONARCH 3, 13% of patients receiving Verzenio plus fulvestrant in MONARCH 2 and in 20% of patients

receiving Verzenio alone in MONARCH 1. Episodes of diarrhea have been associated with dehydration and infection.

Diarrhea incidence was greatest during the first month of Verzenio dosing. In MONARCH 3, the median time to onset of the first diarrhea event was 8 days, and the median duration of diarrhea for Grades 2 and 3 were 11 and 8 days, respectively. In MONARCH 2, the median time to onset of the first diarrhea event was 6 days, and the median duration of diarrhea for Grades 2 and 3 were 9 days and 6 days, respectively. In MONARCH 3, 19% of patients with diarrhea required a dose omission and 13% required a dose reduction. In MONARCH 2, 22% of patients with diarrhea required a dose omission and 22% required a dose reduction. The time to onset and resolution for diarrhea were similar across MONARCH 3, MONARCH 2, and MONARCH 1.

Instruct patients that at the first sign of loose stools, they should start antidiarrheal therapy such as loperamide, increase oral fluids, and notify their healthcare provider for further instructions and appropriate follow-up. For Grade 3 or 4 diarrhea, or diarrhea that requires hospitalization, discontinue Verzenio until toxicity resolves to \leq Grade 1, and then resume Verzenio at the next lower dose.

Neutropenia occurred in 41% of patients receiving Verzenio plus an aromatase inhibitor in MONARCH 3, 46% of patients receiving Verzenio plus fulvestrant in MONARCH 2 and 37% of patients receiving Verzenio alone in MONARCH 1. A Grade \geq 3 decrease in neutrophil count (based on laboratory findings) occurred in 22% of patients receiving Verzenio plus an aromatase inhibitor in MONARCH 3, 32% of patients receiving Verzenio plus fulvestrant in MONARCH 2 and in 27% of patients receiving Verzenio alone in MONARCH 1. In MONARCH 3, the median time to first episode of Grade \geq 3 neutropenia was 33 days, and in MONARCH 2 and MONARCH 1, was 29 days. In MONARCH 3, median duration of Grade \geq 3 neutropenia was 11 days, and for MONARCH 2 and MONARCH 1 was 15 days.

Monitor complete blood counts prior to the start of Verzenio therapy, every 2 weeks for the first 2 months, monthly for the next 2 months, and as clinically indicated. Dose interruption, dose reduction, or delay in starting treatment cycles is recommended for patients who develop Grade 3 or 4 neutropenia.

Febrile neutropenia has been reported in <1% of patients exposed to Verzenio in the MONARCH studies. Two deaths due to neutropenic sepsis were observed in MONARCH 2. Inform patients to promptly report any episodes of fever to their healthcare provider.

Severe, life-threatening, or fatal **interstitial lung disease (ILD) and/or pneumonitis** can occur in patients treated with Verzenio and other CDK4/6 inhibitors. Across clinical trials (MONARCH 1, MONARCH 2, MONARCH 3), 3.3% of Verzenio-treated patients had ILD/pneumonitis of any grade, 0.6% had Grade 3 or 4, and 0.4% had fatal outcomes. Additional cases of ILD/pneumonitis have been observed in the post-marketing setting, with fatalities reported.

Monitor patients for pulmonary symptoms indicative of ILD/pneumonitis. Symptoms may include hypoxia, cough, dyspnea, or interstitial infiltrates on radiologic exams. Infectious, neoplastic, and other causes for such symptoms should be excluded by means of appropriate investigations.

Dose interruption or dose reduction is recommended in patients who develop persistent or recurrent Grade 2 ILD/pneumonitis. Permanently discontinue Verzenio in all patients with grade 3 or 4 ILD/pneumonitis.

Grade \geq 3 increases in **alanine aminotransferase (ALT)** (6% versus 2%) and **aspartate aminotransferase (AST)** (3% versus 1%) were reported in the Verzenio and placebo arms, respectively, in MONARCH 3. Grade \geq 3 increases in ALT (4% versus 2%) and AST (2% versus 3%) were reported in the Verzenio and placebo arms respectively, in MONARCH 2.

In MONARCH 3, for patients receiving Verzenio plus an aromatase inhibitor with Grade \geq 3 increases in ALT or AST, median time to onset was 61 and 71 days, respectively, and median time to resolution to Grade <3 was 14 and 15 days, respectively. In MONARCH 2, for patients receiving Verzenio plus fulvestrant with Grade \geq 3 increases in ALT or AST, median time to onset was 57 and 185 days, respectively, and median time to resolution to Grade <3 was 14 and 13 days, respectively.

For assessment of potential **hepatotoxicity**, monitor liver function tests (LFTs) prior to the start of Verzenio therapy, every 2 weeks for the first 2 months, monthly for the next 2 months, and as clinically indicated. Dose interruption, dose reduction, dose discontinuation, or delay in starting treatment cycles is recommended for patients who develop persistent or recurrent Grade 2, or Grade 3 or 4, hepatic transaminase elevation.

Venous thromboembolic events were reported in 5% of patients treated with Verzenio plus an aromatase inhibitor as compared to 0.6% of patients treated with an aromatase inhibitor plus placebo in MONARCH 3. Venous thromboembolic events were reported in 5% of patients treated with Verzenio plus fulvestrant in MONARCH 2 as compared to 0.9% of patients treated with fulvestrant plus placebo. Venous thromboembolic events included deep vein thrombosis, pulmonary embolism, pelvic venous thrombosis, cerebral venous sinus thrombosis, subclavian and axillary vein thrombosis, and inferior vena cava thrombosis. Across the clinical development program, deaths due to venous thromboembolism have been reported. Monitor patients for signs and symptoms of venous thrombosis and pulmonary embolism and treat as medically appropriate.

Verzenio can cause **fetal harm** when administered to a pregnant woman based on findings from animal studies and the mechanism of action. In animal reproduction studies, administration of abemaciclib to pregnant rats during the period of organogenesis caused teratogenicity and decreased fetal weight at maternal exposures that were similar to the human clinical exposure based on area under the curve (AUC) at the maximum recommended human dose. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with Verzenio and for at least 3 weeks after the last dose. There are no data on the presence of Verzenio in human milk or its effects on the breastfed child or on milk production. Advise lactating women not to breastfeed during Verzenio treatment and for at least 3 weeks after the last dose because of the potential for serious adverse reactions in breastfed infants. Based on findings in animals, Verzenio may impair fertility in males of reproductive potential.

The **most common adverse reactions (all grades, \geq 10%)** observed in **MONARCH 3 for Verzenio plus anastrozole or letrozole** and **\geq 2% higher than placebo plus anastrozole or letrozole vs placebo plus anastrozole or letrozole** were diarrhea (81% vs 30%), neutropenia (41% vs 2%), fatigue (40% vs 32%), infections (39% vs 29%), nausea (39% vs 20%), abdominal pain (29% vs 12%), vomiting (28% vs 12%), anemia (28% vs 5%), alopecia (27% vs 11%), decreased appetite (24% vs 9%), leukopenia (21% vs 2%), creatinine increased (19% vs 4%), constipation (16% vs 12%), ALT increased (16% vs 7%), AST increased (15% vs 7%), rash (14% vs 5%), pruritus (13% vs 9%), cough (13% vs 9%), dyspnea (12% vs 6%), dizziness (11% vs 9%), weight decreased (10% vs 3%), influenza-like illness (10% vs 8%), and thrombocytopenia (10% vs 2%).

The **most common adverse reactions (all grades, \geq 10%)** observed in **MONARCH 2 for Verzenio plus fulvestrant** and **\geq 2% higher than placebo**

plus fulvestrant vs placebo plus fulvestrant were diarrhea (86% vs 25%), neutropenia (46% vs 4%), fatigue (46% vs 32%), nausea (45% vs 23%), infections (43% vs 25%), abdominal pain (35% vs 16%), anemia (29% vs 4%), leukopenia (28% vs 2%), decreased appetite (27% vs 12%), vomiting (26% vs 10%), headache (20% vs 15%), dysgeusia (18% vs 3%), thrombocytopenia (16% vs 3%), alopecia (16% vs 2%), stomatitis (15% vs 10%), ALT increased (13% vs 5%), pruritus (13% vs 6%), cough (13% vs 11%), dizziness (12% vs 6%), AST increased (12% vs 7%), peripheral edema (12% vs 7%), creatinine increased (12% vs <1%), rash (11% vs 4%), pyrexia (11% vs 6%), and weight decreased (10% vs 2%).

The **most common adverse reactions (all grades, ≥10%)** observed in **MONARCH 1** with Verzenio were diarrhea (90%), fatigue (65%), nausea (64%), decreased appetite (45%), abdominal pain (39%), neutropenia (37%), vomiting (35%), infections (31%), anemia (25%), thrombocytopenia (20%), headache (20%), cough (19%), leukopenia (17%), constipation (17%), arthralgia (15%), dry mouth (14%), weight decreased (14%), stomatitis (14%), creatinine increased (13%), alopecia (12%), dysgeusia (12%), pyrexia (11%), dizziness (11%), and dehydration (10%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** that occurred in the Verzenio arm vs the placebo arm of **MONARCH 3** were neutropenia (22% vs 2%), diarrhea (9% vs 1%), leukopenia (8% vs <1%), ALT increased (7% vs 2%), and anemia (6% vs 1%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** that occurred in the Verzenio arm vs the placebo arm of **MONARCH 2** were neutropenia (27% vs 2%), diarrhea (13% vs <1%), leukopenia (9% vs 0%), anemia (7% vs 1%), and infections (6% vs 3%).

The **most frequently reported ≥5% Grade 3 or 4 adverse reactions** from **MONARCH 1** with Verzenio were neutropenia (24%), diarrhea (20%), fatigue (13%), infections (7%), leukopenia (6%), anemia (5%), and nausea (5%).

Lab abnormalities (all grades; Grade 3 or 4) for **MONARCH 3** in **≥10%** for **Verzenio plus anastrozole or letrozole** and **≥2% higher than placebo plus anastrozole or letrozole vs placebo plus anastrozole or letrozole** were increased serum creatinine (98% vs 84%; 2% vs 0%), decreased white blood cells (82% vs 27%; 13% vs <1%), anemia (82% vs 28%; 2% vs 0%), decreased neutrophil count (80% vs 21%; 22% vs 3%), decreased lymphocyte count (53% vs 26%; 8% vs 2%), decreased platelet count (36% vs 12%; 2% vs <1%), increased ALT (48% vs 25%; 7% vs 2%), and increased AST (37% vs 23%; 4% vs <1%).

Lab abnormalities (all grades; Grade 3 or 4) for **MONARCH 2** in **≥10%** for **Verzenio plus fulvestrant** and **≥2% higher than placebo plus fulvestrant vs placebo plus fulvestrant** were increased serum creatinine (98% vs 74%; 1% vs 0%), decreased white blood cells (90% vs 33%; 23% vs 1%), decreased neutrophil count (87% vs 30%; 33% vs 4%), anemia (84% vs 33%; 3% vs <1%), decreased lymphocyte count (63% vs 32%; 12% vs 2%), decreased platelet count (53% vs 15%; 2% vs 0%), increased ALT (41% vs 32%; 5% vs 1%), and increased AST (37% vs 25%; 4% vs 4%).

Lab abnormalities (all grades; Grade 3 or 4) for **MONARCH 1** were increased serum creatinine (98%; <1%), decreased white blood cells (91%; 28%), decreased neutrophil count (88%; 27%), anemia (68%; 0%), decreased lymphocyte count (42%; 14%), decreased platelet count (41%; 2%), increased ALT (31%; 3%), and increased AST (30%; 4%).

Strong and moderate CYP3A inhibitors increased the exposure of abemaciclib plus its active metabolites to a clinically meaningful extent and may lead to increased toxicity. Avoid concomitant use of the strong CYP3A inhibitor ketoconazole. Ketoconazole is predicted to increase the AUC of abemaciclib by up to 16-fold. In patients with recommended starting doses of 200 mg twice daily or 150 mg twice daily, reduce the Verzenio dose to 100 mg twice daily with concomitant use of strong CYP3A inhibitors other than ketoconazole. In patients who have had a dose reduction to 100 mg twice daily due to adverse reactions, further reduce the Verzenio dose to 50 mg twice daily with concomitant use of strong CYP3A inhibitors. If a patient taking Verzenio discontinues a strong CYP3A inhibitor, increase the Verzenio dose (after 3 to 5 half-lives of the inhibitor) to the dose that was used before starting the inhibitor. With concomitant use of moderate CYP3A inhibitors, monitor for adverse reactions and consider reducing the Verzenio dose in 50 mg decrements. Patients should avoid grapefruit products.

Avoid concomitant use of strong or moderate CYP3A inducers and consider alternative agents. Coadministration of strong or moderate CYP3A inducers decreased the plasma concentrations of abemaciclib plus its active metabolites and may lead to reduced activity.

With severe hepatic impairment (Child-Pugh Class C), reduce the Verzenio dosing frequency to once daily. The pharmacokinetics of Verzenio in patients with **severe renal impairment** (CL_{Cr} <30 mL/min), end stage renal disease, or in patients on dialysis **is unknown**. No dosage adjustments are necessary in patients with mild or moderate hepatic (Child-Pugh A or B) and/or renal impairment (CL_{Cr} ≥30-89 mL/min).

AL HCP ISI 17SEP2019

Please see full [Prescribing Information](#) for Verzenio.

About Lilly Oncology

For more than 50 years, Lilly has been dedicated to delivering life-changing medicines and support to people living with cancer and those who care for them. Lilly is determined to build on this heritage and continue making life better for all those affected by cancer around the world. To learn more about Lilly's commitment to people with cancer, please visit www.LillyOncology.com.

About Eli Lilly and Company

Lilly is a global health care leader that unites caring with discovery to create medicines that make life better for people around the world. We were founded more than a century ago by a man committed to creating high-quality medicines that meet real needs, and today we remain true to that mission in all our work. Across the globe, Lilly employees work to discover and bring life-changing medicines to those who need them, improve the understanding and management of disease, and give back to communities through philanthropy and volunteerism. To learn more about Lilly, please visit us at lilly.com and lilly.com/newsroom. P-LLY

© Lilly USA, LLC 2021. ALL RIGHTS RESERVED.

Verzenio® is a trademark owned by or licensed to Eli Lilly and Company, its subsidiaries, or affiliates.

Lilly Forward-Looking Statement

This press release contains forward-looking statements (as that term is defined in the Private Securities Litigation Reform Act of 1995) about Verzenio and reflects Lilly's current beliefs and expectations. However, as with any pharmaceutical product, there are substantial risks and uncertainties in the process of research, development, regulatory approval, and commercialization. Among other things, there can be no guarantee that future studies will be completed as planned, that future study results will be consistent with the results to date, or that Verzenio will receive additional regulatory

approvals or be commercially successful. For further discussion of these and other risks and uncertainties, see Lilly's most recent Form 10-K and Form 10-Q filings with the United States Securities and Exchange Commission. Except as required by law, Lilly undertakes no duty to update forward-looking statements to reflect events after the date of this release.

Refer to: Tracy Henrikson; tracy.henrikson@lilly.com; 609-454-7116 (Media)
Kevin Hern; hern_kevin_r@lilly.com; 317-277-1838 (Investors)

The Lilly logo is rendered in a vibrant red, cursive script. The letters are thick and fluid, with the 'L' and 'y' having prominent loops and tails. The overall style is elegant and classic, characteristic of the pharmaceutical company's branding.

 View original content to download multimedia: <https://www.prnewswire.com/news-releases/new-data-from-verzenio-abemaciclib-monarche-study-to-be-featured-in-esmo-virtual-plenary-301384959.html>

SOURCE Eli Lilly and Company