


5  
6  
7  
8 **Risk Evaluation for**  
9 **Tris(2-chloroethyl) Phosphate (TCEP)**

10  
11 **Supplemental Information File:**

12  
13  
14 **Exposure Monitoring Tornado Figures, Supplemental Tables and Data**  
15 **Integration Methods and Approach for TCEP**  
16 **CASRN: 115-96-8**  
17



18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

*September 2024*

## TABLE OF CONTENTS

---

30	<b>1 ENVIRONMENTAL MONITORING CONCENTRATIONS REPORTED BY MEDIA</b>	
31	<b>TYPE.....</b>	<b>9</b>
32	1.1 Ambient Air.....	9
33	1.1.1 Ambient Air (ng/g) – Particulate Fraction.....	9
34	1.1.2 Ambient Air (ng/m <sup>3</sup> ) – All Fractions.....	9
35	1.2 Aquatic Organisms – Fish .....	12
36	1.2.1 Aquatic Organisms – Fish (ng/g) – All Fractions.....	12
37	1.3 Aquatic Organisms – Mammal.....	14
38	1.3.1 Aquatic Organisms – Mammal (ng/g) – Lipid Fraction .....	14
39	1.4 Aquatic Organisms – Mollusk.....	15
40	1.4.1 Aquatic Organisms – Mollusk (ng/g) – All Fractions .....	15
41	1.5 Aquatic Organisms – Other .....	16
42	1.5.1 Aquatic Organisms – Other (ng/g) – Wet Fraction .....	16
43	1.6 Dietary .....	17
44	1.6.1 Dietary (ng/g) – Wet Fraction.....	17
45	1.6.2 Dietary (ng/g) – Wet Fraction.....	20
46	1.7 Drinking Water.....	21
47	1.7.1 Drinking Water (ng/L) – Not Specified Fraction .....	21
48	1.8 Dust (Indoor) .....	23
49	1.8.1 Dust (Indoor) (ng/g) – Dry Fraction .....	23
50	1.8.2 Dust (Indoor) (ng/g) – Dry Fraction .....	29
51	1.8.3 Dust (Indoor) (ng/m <sup>2</sup> ) – Dry Fraction.....	30
52	1.9 Groundwater .....	31
53	1.9.1 Groundwater (ng/L) – Not Specified Fraction.....	31
54	1.10 Human Biomonitoring – Breastmilk .....	33
55	1.10.1 Human Biomonitoring – Breastmilk (ng/L) – wet Fraction .....	33
56	1.10.2 Human Biomonitoring – Breastmilk (ng/g) – Lipid Fraction.....	33
57	1.11 Human Biomonitoring – Hair.....	34
58	1.11.1 Human Biomonitoring – Hair (ng/g) – Dry Fraction.....	34
59	1.12 Human Biomonitoring – Nails .....	35
60	1.12.1 Human Biomonitoring – Nails (ng/g) – Dry Fraction .....	35
61	1.13 Human Biomonitoring – Other.....	35
62	1.13.1 Human Biomonitoring – Other (ng/g) – Dry Fraction.....	35
63	1.13.2 Human Biomonitoring – Other (ng/g) – Dry Fraction.....	36
64	1.14 Human Biomonitoring – Plasma .....	36
65	1.14.1 Human Biomonitoring – Plasma (ng/L) – Wet Fraction .....	36
66	1.15 Human Biomonitoring – Serum .....	37
67	1.15.1 Human Biomonitoring – Serum (ng/g) – Lipid Fraction.....	37
68	1.16 Human Biomonitoring – Skin_Dermal Wipe.....	37
69	1.16.1 Human Biomonitoring – Skin_Dermal Wipe (ng/g) – Dry Fraction.....	37
70	1.16.2 Human Biomonitoring – Skin_Dermal Wipe (ng/wipe) – Dry Fraction.....	38
71	1.17 Human Biomonitoring – Urine.....	39
72	1.17.1 Human Biomonitoring – Urine (ng/g) – Creatinine Adjusted Fraction.....	39
73	1.17.2 Human Biomonitoring – Urine (ng/L) – Unadjusted Fraction .....	39
74	1.17.3 Human Biomonitoring – Urine (ng/L) – All Fractions.....	40
75	1.18 Human Biomonitoring – Silicone Wristbands .....	41
76	1.18.1 Human Biomonitoring – Silicone Wristbands (ng/g) – Not Specified Fraction .....	41

77	1.19 Indoor Air .....	42
78	1.19.1 Indoor Air (ng/m <sup>3</sup> ) – All Fractions .....	42
79	1.20 Leachate.....	46
80	1.20.1 Leachate (ng/L) – Not Specified Fraction .....	46
81	1.21 Other .....	47
82	1.21.1 Other (ng/g) – Dry Fraction.....	47
83	1.21.2 Other (ng/g) – All Fractions .....	48
84	1.21.3 Other (ng/L) – Not Specified Fraction.....	49
85	1.22 Personal Inhalation .....	49
86	1.22.1 Personal Inhalation (ng/m <sup>3</sup> ) – All Fractions .....	49
87	1.23 Precipitation.....	50
88	1.23.1 Precipitation (ng/L) – Wet Fraction.....	50
89	1.24 Sediment .....	52
90	1.24.1 Sediment (ng/g) – All Fractions .....	52
91	1.25 Soil.....	54
92	1.25.1 Soil (ng/g) – Dry Fraction.....	54
93	1.26 Surface Water .....	55
94	1.26.1 Surface Water (ng/L) – Not Specified Fraction.....	55
95	1.27 Terrestrial Organisms – Bird.....	58
96	1.27.1 Terrestrial Organisms – Bird (ng/g) – All Fractions .....	58
97	1.27.2 Terrestrial Organisms – Bird (ng/g) – Wet Fraction .....	60
98	1.28 Terrestrial Organisms – Mammal.....	61
99	1.28.1 Terrestrial Organisms – Mammal (ng/g) – All Fractions .....	61
100	1.29 Terrestrial Organisms – Plant.....	62
101	1.29.1 Terrestrial Organisms – Plant (ng/g) – Wet Fraction .....	62
102	1.30 Wastewater .....	62
103	1.30.1 Wastewater (ng/g) – Wet Fraction.....	62
104	1.30.2 Wastewater (ng/L) – Wet Fraction .....	63
105	<b>2 METHODS AND APPROACH.....</b>	<b>67</b>
106	2.1 Data Integration Methods and Approach .....	67
107	2.2 Statistical Approach of Exposure Estimates Derived from Measured Concentrations .....	68
108	2.2.1 Aggregation of Statistical Estimates.....	68
109	2.2.2 Fitting Lognormal Distributions .....	70
110	2.2.3 Fitting Normal Distributions.....	70
111	2.2.4 Quality Control of Derived Exposure Estimates .....	70
112	2.2.5 Final Exposure Estimates by Media and Pollution Source Receptor Type .....	70
113	<b>REFERENCES.....</b>	<b>71</b>
114		
115	<b>LIST OF TABLES</b>	
116	Table 1-1. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Particulate	
117	Fraction of Ambient Air.....	9
118	Table 1-2. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m <sup>3</sup> ) Levels in Ambient Air	
119	.....	11
120	Table 1-3. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Aquatic	
121	Organisms – Fish .....	13
122	Table 1-4. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Lipid	
123	Fraction of Aquatic Organisms – Mammal .....	15

124	Table 1-5. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Aquatic	
125	Organisms – Mollusk.....	16
126	Table 1-6. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet	
127	Fraction of Aquatic Organisms – Other.....	17
128	Table 1-7. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet	
129	Fraction of Dietary.....	18
130	Table 1-8. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Wet	
131	Fraction of Dietary.....	21
132	Table 1-9. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not	
133	Specified Fraction of Drinking Water.....	22
134	Table 1-10. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
135	Fraction of Dust (Indoor).....	26
136	Table 1-11. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Dry	
137	Fraction of Dust (Indoor).....	30
138	Table 1-12. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m <sup>2</sup> ) Levels in the Dry	
139	Fraction of Dust (Indoor).....	30
140	Table 1-13. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not	
141	Specified Fraction of Groundwater.....	32
142	Table 1-14. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the wet	
143	Fraction of Human Biomonitoring – Breastmilk.....	33
144	Table 1-15. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Lipid	
145	Fraction of Human Biomonitoring – Breastmilk.....	34
146	Table 1-16. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
147	Fraction of Human Biomonitoring – Hair.....	34
148	Table 1-17. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
149	Fraction of Human Biomonitoring – Nails.....	35
150	Table 1-18. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
151	Fraction of Human Biomonitoring – Other.....	36
152	Table 1-19. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Dry	
153	Fraction of Human Biomonitoring – Other.....	36
154	Table 1-20. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet	
155	Fraction of Human Biomonitoring – Plasma.....	37
156	Table 1-21. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Lipid	
157	Fraction of Human Biomonitoring – Serum.....	37
158	Table 1-22. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
159	Fraction of Human Biomonitoring – Skin_Dermal Wipe.....	38
160	Table 1-23. Summary of Peer-Reviewed Literature that Measured TCEP (ng/wipe) Levels in the Dry	
161	Fraction of Human Biomonitoring – Skin_Dermal Wipe.....	38
162	Table 1-24. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Creatinine	
163	Adjusted Fraction of Human Biomonitoring – Urine.....	39
164	Table 1-25. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the	
165	Unadjusted Fraction of Human Biomonitoring – Urine.....	40
166	Table 1-26. Summary of Peer-Reviewed Literature that Measured BCEP (ng/L) Levels in Human	
167	Biomonitoring – Urine.....	41
168	Table 1-27. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Not	
169	Specified Fraction of Human Biomonitoring – Silicone Wristbands.....	42
170	Table 1-28. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m <sup>3</sup> ) Levels in Indoor Air	44
171	Table 1-29. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not	
172	Specified Fraction of Leachate.....	47

173	Table 1-30. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
174	Fraction of Other .....	48
175	Table 1-31. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Other .....	48
176	Table 1-32. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not	
177	Specified Fraction of Other .....	49
178	Table 1-33. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m <sup>3</sup> ) Levels in Personal	
179	Inhalation .....	50
180	Table 1-34. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet	
181	Fraction of Precipitation .....	51
182	Table 1-35. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Sediment....	53
183	Table 1-36. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry	
184	Fraction of Soil .....	54
185	Table 1-37. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not	
186	Specified Fraction of Surface Water.....	56
187	Table 1-38. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Terrestrial	
188	Organisms – Bird .....	59
189	Table 1-39. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Wet	
190	Fraction of Terrestrial Organisms – Bird.....	61
191	Table 1-40. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Terrestrial	
192	Organisms – Mammal.....	62
193	Table 1-41. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet	
194	Fraction of Terrestrial Organisms – Plant.....	62
195	Table 1-42. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet	
196	Fraction of Wastewater .....	63
197	Table 1-43. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet	
198	Fraction of Wastewater .....	64
199	Table 2-1. Statistics and Methods for Data Aggregation.....	68
200	Table 2-2. Distributions Preferred Depending on Available Reported Statistics .....	69
201	Table 2-3. Assumed Percentile for Calculating Error by Statistical Estimate Type.....	70
202		

## 203 LIST OF FIGURES

---

204	Figure 1-1. Concentrations of TCEP (ng/g) in the Particulate Fraction of Ambient Air in General	
205	Population (Background) Locations in 2018 .....	9
206	Figure 1-2. Concentrations of TCEP (ng/m <sup>3</sup> ) in Ambient Air from 2000 to 2019.....	10
207	Figure 1-3. Concentrations of TCEP (ng/g) in Aquatic Organisms – Fish from 2003 to 2016.....	13
208	Figure 1-4. Concentrations of TCEP (ng/g) in the Lipid Fraction of Aquatic Organisms – Mammal from	
209	2004 to 2010 .....	15
210	Figure 1-5. Concentrations of TCEP (ng/g) in Aquatic Organisms – Mollusk in Near Facility (Highly	
211	Exposed) Locations from 2008 to 2017 .....	16
212	Figure 1-6. Concentrations of TCEP (ng/g) in the Wet Fraction of Aquatic Organisms – Other from	
213	2008 to 2018 .....	17
214	Figure 1-7. Concentrations of TCEP (ng/g) in the Wet Fraction of Dietary from 1982 to 2018 .....	18
215	Figure 1-8. Concentrations of BCEP (ng/g) in the Wet Fraction of Dietary in 2018.....	20
216	Figure 1-9. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Drinking Water from 1982	
217	to 2014 .....	22
218	Figure 1-10. Concentrations of TCEP (ng/g) in the Dry Fraction of Dust (Indoor) from 2000 to 2019..	25
219	Figure 1-11. Concentrations of BCEP (ng/g) in the Dry Fraction of Dust (Indoor) in Residential	
220	Locations in 2019.....	30
221	Figure 1-12. Concentrations of TCEP (ng/m <sup>2</sup> ) in the Dry Fraction of Dust (Indoor) from 2000 to 2016	30

222	Figure 1-13. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Groundwater from 1978 to	
223	2017.....	31
224	Figure 1-14. Concentrations of TCEP (ng/L) in the wet Fraction of Human Biomonitoring – Breastmilk	
225	in General Population (Background) Locations from 2014 to 2015.....	33
226	Figure 1-15. Concentrations of TCEP (ng/g) in the Lipid Fraction of Human Biomonitoring –	
227	Breastmilk from 1997 to 2011 .....	33
228	Figure 1-16. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Hair in	
229	General Population (Background) Locations from 2014 to 2015.....	34
230	Figure 1-17. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Nails in	
231	General Population (Background) Locations from 2014 to 2015.....	35
232	Figure 1-18. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Other in	
233	General Population (Background) Locations from 2014 to 2016.....	35
234	Figure 1-19. Concentrations of BCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Other in	
235	General Population (Background) Locations from 2014 to 2016.....	36
236	Figure 1-20. Concentrations of TCEP (ng/L) in the Wet Fraction of Human Biomonitoring – Plasma in	
237	General Population (Background) Locations from 2014 to 2016.....	36
238	Figure 1-21. Concentrations of TCEP (ng/g) in the Lipid Fraction of Human Biomonitoring – Serum in	
239	General Population (Background) Locations in 2016 .....	37
240	Figure 1-22. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring –	
241	Skin_Dermal Wipe in General Population (Background) Locations in 2012 .....	38
242	Figure 1-23. Concentrations of TCEP (ng/wipe) in the Dry Fraction of Human Biomonitoring –	
243	Skin_Dermal Wipe in General Population (Background) Locations from 2012 to 2016. 38	
244	Figure 1-24. Concentrations of BCEP (ng/g) in the Creatinine Adjusted Fraction of Human	
245	Biomonitoring – Urine in General Population (Background) Locations in 2018.....	39
246	Figure 1-25. Concentrations of TCEP (ng/L) in the Unadjusted Fraction of Human Biomonitoring –	
247	Urine in General Population (Background) Locations from 2010 to 2015 .....	40
248	Figure 1-26. Concentrations of BCEP (ng/L) in Human Biomonitoring – Urine in General Population	
249	(Background) Locations from 2011 to 2018.....	41
250	Figure 1-27. Concentrations of TCEP (ng/g) in the Not Specified Fraction of Human Biomonitoring –	
251	Silicone Wristbands in General Population (Background) Locations from 2012 to 2015 42	
252	Figure 1-28. Concentrations of TCEP (ng/m <sup>3</sup> ) in Indoor Air from 2000 to 2016 .....	44
253	Figure 1-29. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Leachate from 1994 to	
254	1995.....	47
255	Figure 1-30. Concentrations of TCEP (ng/g) in the Dry Fraction of Other in Unknown/Not Specified	
256	Locations in 2003.....	47
257	Figure 1-31. Concentrations of TCEP (ng/g) in Other from 2001 to 2008.....	48
258	Figure 1-32. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Other in General	
259	Population (Background) Locations in 2016 .....	49
260	Figure 1-33. Concentrations of TCEP (ng/m <sup>3</sup> ) in Personal Inhalation in General Population	
261	(Background) Locations from 2013 to 2016.....	50
262	Figure 1-34. Concentrations of TCEP (ng/L) in the Wet Fraction of Precipitation from 1994 to 2014... 51	
263	Figure 1-35. Concentrations of TCEP (ng/g) in Sediment from 1980 to 2017.....	53
264	Figure 1-36. Concentrations of TCEP (ng/g) in the Dry Fraction of Soil in General Population	
265	(Background) Locations from 2010 to 2014.....	54
266	Figure 1-37. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Surface Water from 1980	
267	to 2017 .....	56
268	Figure 1-38. Concentrations of TCEP (ng/g) in Terrestrial Organisms – Bird from 2000 to 2016.....	59
269	Figure 1-39. Concentrations of BCEP (ng/g) in the Wet Fraction of Terrestrial Organisms – Bird in	
270	General Population (Background) Locations from 2000 to 2012.....	61

271	Figure 1-40. Concentrations of TCEP (ng/g) in Terrestrial Organisms – Mammal from 2008 to 2018 ..	61
272	Figure 1-41. Concentrations of TCEP (ng/g) in the Wet Fraction of Terrestrial Organisms – Plant in	
273	Remote (Not Near Source) Locations from 1993 to 1994 .....	62
274	Figure 1-42. Concentrations of TCEP (ng/g) in the Wet Fraction of Wastewater from 2013 to 2018.....	63
275	Figure 1-43. Concentrations of TCEP (ng/L) in the Wet Fraction of Wastewater from 2001 to 2018 ....	64

## 276 **ABBREVIATIONS AND ACRONYMS**

---

277	AQ	Antarctica
278	AR	Argentina
279	AT	Austria
280	AU	Australia
281	BE	Belgium
282	BO	Bolivia
283	BR	Brazil
284	CA	Canada
285	CH	Switzerland
286	CL	Chile
287	CN	China
288	CO	Colombia
289	CR	Costa Rica
290	CZ	Czech Republic
291	DE	Germany
292	DK	Denmark
293	ES	Spain
294	FI	Finland
295	FOD	Frequency of detection
296	FR	France
297	GB	United Kingdom
298	GL	Greenland
299	GR	Greece
300	GSD	Geometric standard deviation
301	JP	Japan
302	KR	South Korea
303	LOD	Limits of detection
304	LOQ	Limits of quantitation
305	MX	Mexico
306	NL	Netherlands
307	NO	Norway
308	NZ	New Zealand
309	PH	Philippines
310	PL	Poland
311	PR	Puerto Rico
312	PT	Portugal
313	RO	Romania
314	SD	Standard deviation
315	SE	Sweden
316	TR	Turkey
317	US	United States
318	VN	Vietnam
319	ZA	South Africa



# 1 ENVIRONMENTAL MONITORING CONCENTRATIONS REPORTED BY MEDIA TYPE

## 1.1 Ambient Air

### 1.1.1 Ambient Air (ng/g) – Particulate Fraction

Measured concentrations of TCEP in Ambient Air with unit of ng/g, extracted from one source, are summarized in Figure 1-1 and supplemental information is provided in Table 1-1. Overall, concentrations were 300 ng/g from 18 samples collected in 2018 in one country, PL. Location types were categorized as General Population (Background). Reported detection frequency was 0.11.



**Figure 1-1. Concentrations of TCEP (ng/g) in the Particulate Fraction of Ambient Air in General Population (Background) Locations in 2018**

**Table 1-1. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Particulate Fraction of Ambient Air**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Fabiańska et al. (2019)</a>	PL	General Population (Background)	2018	18 (0.11)	N/R	Medium

N/R = Not reported

### 1.1.2 Ambient Air (ng/m<sup>3</sup>) – All Fractions


Measured concentrations of TCEP in Ambient Air with unit of ng/m<sup>3</sup>, extracted from 17 sources, are summarized in Figure 1-2 and supplemental information is provided in Table 1-2. More than one weight fraction was reported and summarized separately below:

Overall, concentrations for Combined Vapor/Gas and Particulate ranged from not detected to 58.4 ng/m<sup>3</sup> from 152 samples collected between 2000 and 2018 in 11 countries, AR, BO, BR, CA, CL, CO, CR, JP, MX, NO and US. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection frequency ranged from 0.55 to 0.94.

Overall, concentrations for Particulate ranged from not detected to 3.532 ng/m<sup>3</sup> from 855 samples collected between 2002 and 2019 in seven countries, AQ, CA, ES, FI, JP, SE and US. Location types were categorized as Unknown/Not Specified, General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection frequency ranged from 0.0 to 1.0.

Overall, concentrations for Vapor/Gas ranged from not detected to 0.143 ng/m<sup>3</sup> from 49 samples collected in 2014 in two countries, AQ and TR. Location types were categorized as General Population

353 (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection  
 354 frequency ranged from 0.8 to 1.0.  
 355



356  
 357 **Figure 1-2. Concentrations of TCEP (ng/m<sup>3</sup>) in Ambient Air from 2000 to 2019**  
 358

**Table 1-2. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m<sup>3</sup>) Levels in Ambient Air**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
Combined Vapor/Gas and Particulate						
<a href="#">Guo et al. (2017)</a>	CA, US	General Population (Background)	2013	20 (0.55)	0.0602	High
<a href="#">Bohlin-Nizzetto et al. (2019)</a>	NO	Remote (Not Near Source)	2017–2018	36 (0.56)	0.045	Medium
<a href="#">Rauert et al. (2018)</a>	AR, BR, CL, MX	General Population (Background)	2014–2016	14 (0.93)	0.08	High
<a href="#">Rauert et al. (2018)</a>	AR, BO, BR, CL, CO, CR, MX	Remote (Not Near Source)	2014–2016	36 (0.94)	0.05	High
<a href="#">Ohura et al. (2006)</a>	JP	Near Facility (Highly Exposed)	2000–2001	46 (0.91)	N/R	Medium
Particulate						
<a href="#">Peverly et al. (2015)</a>	US	General Population (Background)	2012–2014	161 (0.87)	N/R	High
<a href="#">Salamova et al. (2016)</a>	US	General Population (Background)	2012–2014	359 (0.60)	N/R	Medium
<a href="#">Clark et al. (2017)</a>	US	General Population (Background)	2013	45 (0.93)	N/R	High
<a href="#">Salamova et al. (2014)</a>	US	General Population (Background)	2012	81 (0.74)	N/R	Medium
<a href="#">Salamova et al. (2014)</a>	US	General Population (Background)	2012	16 (0.62)	N/R	Medium
<a href="#">Bradman et al. (2014)</a>	US	General Population (Background)	2010-2011	14 (0.50)	0.3	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
<a href="#">Maceira et al. (2020)</a>	ES	Near Facility (Highly Exposed)	2018–2019	24 (0.62)	0.0014	High
<a href="#">Wong et al. (2018)</a>	SE	General Population (Background)	2014-2015	24 (0.96)	0.044	Medium
<a href="#">Li et al. (2017)</a>	AQ	Remote (Not Near Source)	2014	9 (1.00)	0.0038	High
<a href="#">Sühring et al. (2016)</a>	CA	Unknown/Not Specified	2007–2013	92 (0.87)	N/R	Medium
<a href="#">Abdollahi et al. (2017)</a>	CA	General Population (Background)	2010	21 (N/R)	0.0003	High
<a href="#">Marklund et al. (2005b)</a>	FI	Remote (Not Near Source)	2003	1 (1.00)	N/R	Medium
<a href="#">Saito et al. (2007)</a>	JP	Unknown/Not Specified	2002	8 (0.00)	0.67	Medium
Vapor/Gas						
<a href="#">Li et al. (2017)</a>	AQ	Remote (Not Near Source)	2014	9 (1.00)	0.0012	High
<a href="#">Kurt-Karakus et al. (2018)</a>	TR	General Population (Background)	2014	30 (0.80)	0.073	High
<a href="#">Kurt-Karakus et al. (2018)</a>	TR	Near Facility (Highly Exposed)	2014	10 (0.80)	0.073	High

N/R = Not reported

361

## 1.2 Aquatic Organisms – Fish

362

### 1.2.1 Aquatic Organisms – Fish (ng/g) – All Fractions

363

Measured concentrations of TCEP in Aquatic Organisms – Fish with unit of ng/g, extracted from eight sources, are summarized in Figure 1-3 and supplemental information is provided in Table 1-3. More than one weight fraction was reported and summarized separately below:

366

367


368

369

Overall, concentrations for Lipid ranged from not detected to 187.0 ng/g from 55 samples collected between 2003 and 2016 in five countries, CA, ES, NO, SE and US. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source).

370 Reported detection frequency ranged from 0.21 to 1.0.

371  
 372 Overall, concentrations for Wet ranged from not detected to 26.0 ng/g from 186 samples collected  
 373 between 2004 and 2015 in four countries, CA, KR, NL and NO. Location types were categorized as  
 374 General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source).  
 375 Reported detection frequency ranged from 0.12 to 1.0.  
 376



377  
 378 **Figure 1-3. Concentrations of TCEP (ng/g) in Aquatic Organisms – Fish from 2003 to 2016**

379  
 380 **Table 1-3. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Aquatic**  
 381 **Organisms – Fish**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Lipid						
<a href="#">Guo et al. (2017)</a>	CA, US	General Population (Background)	2010	14 (0.21)	20.9	High
<a href="#">Santín et al. (2016)</a>	ES	General Population (Background)	2016	12 (0.25)	1.39	High
<a href="#">Hallanger et al. (2015)</a>	NO	Remote (Not Near Source)	2009	10 (0.70)	N/R	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Sundkvist et al. (2010)</a>	SE	General Population (Background)	2007	7 (0.57)	2.8	High
<a href="#">Sundkvist et al. (2010)</a>	SE	Near Facility (Highly Exposed)	2003–2007	4 (1.00)	2.8	High
<a href="#">Sundkvist et al. (2010)</a>	SE	Remote (Not Near Source)	2005–2007	8 (1.00)	2.8	High
Wet						
<a href="#">Choo et al. (2018)</a>	KR	General Population (Background)	2015	20 (1.00)	0.22	High
<a href="#">Choo et al. (2018)</a>	KR	General Population (Background)	2015	30 (1.00)	0.06	High
<a href="#">Choo et al. (2018)</a>	KR	General Population (Background)	2015	20 (1.00)	0.09	High
<a href="#">McGoldrick et al. (2014)</a>	CA	General Population (Background)	2009–2010	72 (0.12)	0.03	High
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	19 (0.42)	0.21	High
<a href="#">Evenset et al. (2009)</a>	NO	Remote (Not Near Source)	2004–2008	3 (1.00)	N/R	Medium
<a href="#">Evenset et al. (2009)</a>	NO	Remote (Not Near Source)	2004–2008	5 (1.00)	0.47	Medium
<a href="#">Evenset et al. (2009)</a>	NO	Remote (Not Near Source)	2008	17 (0.94)	N/R	Medium

N/R = Not reported

382

### **1.3 Aquatic Organisms – Mammal**

383

#### **1.3.1 Aquatic Organisms – Mammal (ng/g) – Lipid Fraction**

384


Measured concentrations of TCEP in Aquatic Organisms – Mammal with unit of ng/g, extracted from two sources, are summarized in Figure 1-4 and supplemental information is provided in Table 1-4.

385

386

Overall, concentrations ranged from not detected to 115.0 ng/g from 63 samples collected between 2004

387 and 2010 in two countries, ES and NO. Location types were categorized as General Population  
 388 (Background) and Remote (Not Near Source). Reported detection frequency ranged from 0.0 to 0.44.  
 389



390  
 391 **Figure 1-4. Concentrations of TCEP (ng/g) in the Lipid Fraction of Aquatic Organisms – Mammal**  
 392 **from 2004 to 2010**

393  
 394 **Table 1-4. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Lipid**  
 395 **Fraction of Aquatic Organisms – Mammal**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Hallanger et al. (2015)</a>	NO	Remote (Not Near Source)	2010	10 (0.00)	4.5	High
<a href="#">Sala et al. (2019)</a>	ES	General Population (Background)	2004–2010	9 (0.11)	1.39	Medium
<a href="#">Hallanger et al. (2015)</a>	NO	Remote (Not Near Source)	2009	10 (0.10)	N/R	High
<a href="#">Sala et al. (2019)</a>	ES	General Population (Background)	2004–2010	9 (0.44)	1.39	Medium
<a href="#">Sala et al. (2019)</a>	ES	General Population (Background)	2004–2010	10 (0.10)	1.39	Medium
<a href="#">Sala et al. (2019)</a>	ES	General Population (Background)	2004–2010	15 (0.13)	1.39	Medium

N/R = Not reported

396 **1.4 Aquatic Organisms – Mollusk**

397 **1.4.1 Aquatic Organisms – Mollusk (ng/g) – All Fractions**


398 Measured concentrations of TCEP in Aquatic Organisms – Mollusk with unit of ng/g, extracted from  
 399 two sources, are summarized in Figure 1-5 and supplemental information is provided in Table 1-5. More

400 than one weight fraction was reported and summarized separately below:

401  
402 Overall, concentrations for Lipid were not detected ng/g from 80 samples collected between 2016 and  
403 2017 in one country, PT. Location types were categorized as Near Facility (Highly Exposed). Reported  
404 detection frequency was 0.25.

405  
406 Overall, concentrations for Wet ranged from not detected to 0.82 ng/g from five samples collected in  
407 2008 in one country, NL. Location types were categorized as Near Facility (Highly Exposed). Reported  
408 detection frequency was 0.4.

409



410

411 **Figure 1-5. Concentrations of TCEP (ng/g) in Aquatic Organisms – Mollusk in Near Facility**  
412 **(Highly Exposed) Locations from 2008 to 2017**

413

414 **Table 1-5. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Aquatic**  
415 **Organisms – Mollusk**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Lipid						
<a href="#">Gadelha et al. (2019)</a>	PT	Near Facility (Highly Exposed)	2016–2017	80 (0.25)	1.2	High
Wet						
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	5 (0.40)	0.2	High

416

## 1.5 Aquatic Organisms – Other


417

### 1.5.1 Aquatic Organisms – Other (ng/g) – Wet Fraction

418 Measured concentrations of TCEP in Aquatic Organisms – Other with unit of ng/g, extracted from two  
419 sources, are summarized in Figure 1-6 and supplemental information is provided in Table 1-6. Overall,  
420 concentrations ranged from not detected to 0.33 ng/g from 61 samples collected between 2008 and 2018  
421 in two countries, NL and NO. Location types were categorized as General Population (Background) and  
422 Near Facility (Highly Exposed). Reported detection frequency ranged from 0.0 to 0.2.

423





424

425 **Figure 1-6. Concentrations of TCEP (ng/g) in the Wet Fraction of Aquatic Organisms – Other**  
 426 **from 2008 to 2018**

427

428 **Table 1-6. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet**  
 429 **Fraction of Aquatic Organisms – Other**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	5 (0.20)	0.2	High
<a href="#">Norwegian Environment (2019b)</a>	NO	General Population (Background)	2018	51 (0.00)	0.5	High
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	5 (0.20)	0.42	High

430

## 1.6 Dietary

431

### 1.6.1 Dietary (ng/g) – Wet Fraction

432

433

434


435

436

437

438

Measured concentrations of TCEP in Dietary with unit of ng/g, extracted from four sources, are summarized in Figure 1-7 and supplemental information is provided in Table 1-7. Overall, concentrations ranged from not detected to 113.0 ng/g from 363 samples collected between 1982 and 2018 in four countries, AU, BE, SE and US. Location types were categorized as fruit, dairy, grain, baby food-infant formula, vegetables, other, non-dairy beverages, meat, fish and shellfish and fats and oils. Reported detection frequency ranged from 0.0 to 0.67.



439

440 **Figure 1-7. Concentrations of TCEP (ng/g) in the Wet Fraction of Dietary from 1982 to 2018**

441

442 **Table 1-7. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet**  
 443 **Fraction of Dietary**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">FDA (1995)</a>	US	fruit	1982–1991	74 (0.04)	N/R	Medium
<a href="#">He et al. (2018b)</a>	AU	dairy	2018	9 (0.56)	0.06	Medium

<b>Citation</b>	<b>Country</b>	<b>Location Type</b>	<b>Sampling Year</b>	<b>Sample Size (Frequency of Detection)</b>	<b>Detection Limit (ng/g)</b>	<b>Overall Quality Level</b>
<a href="#">He et al. (2018b)</a>	AU	fish and shellfish	2018	9 (0.22)	0.06	Medium
<a href="#">He et al. (2018b)</a>	AU	grain	2018	12 (0.67)	0.06	Medium
<a href="#">He et al. (2018b)</a>	AU	meat	2018	12 (0.25)	0.06	Medium
<a href="#">He et al. (2018b)</a>	AU	non-dairy beverages	2018	12 (0.08)	0.021	Medium
<a href="#">He et al. (2018b)</a>	AU	other	2018	3 (0.33)	0.06	Medium
<a href="#">He et al. (2018b)</a>	AU	vegetables	2018	15 (0.60)	0.06	Medium
<a href="#">Poma et al. (2018)</a>	BE	baby food-infant formula	2015–2016	17 (N/R)	0.34	High
<a href="#">Poma et al. (2018)</a>	BE	dairy	2015–2016	27 (N/R)	0.45	High
<a href="#">Poma et al. (2018)</a>	BE	fats and oils	2015–2016	10 (0.40)	2.55	High
<a href="#">Poma et al. (2018)</a>	BE	fish and shellfish	2015–2016	53 (N/R)	0.07	High
<a href="#">Poma et al. (2018)</a>	BE	grain	2015–2016	7 (N/R)	0.09	High
<a href="#">Poma et al. (2018)</a>	BE	meat	2015–2016	38 (N/R)	0.14	High
<a href="#">Poma et al. (2018)</a>	BE	other	2015–2016	11 (N/R)	0.44	High
<a href="#">Poma et al. (2018)</a>	BE	vegetables	2015–2016	2 (0.00)	0.01	High
<a href="#">Poma et al. (2017)</a>	SE	dairy	2015	9 (0.22)	0.3	High
<a href="#">Poma et al. (2017)</a>	SE	fats and oils	2015	4 (0.00)	2.0	High
<a href="#">Poma et al. (2017)</a>	SE	fish and shellfish	2015	5 (0.00)	0.2	High


Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Poma et al. (2017)</a>	SE	fruit	2015	5 (0.20)	0.15	High
<a href="#">Poma et al. (2017)</a>	SE	grain	2015	5 (0.00)	0.5	High
<a href="#">Poma et al. (2017)</a>	SE	meat	2015	5 (0.00)	0.2	High
<a href="#">Poma et al. (2017)</a>	SE	non-dairy beverages	2015	2 (0.00)	0.45	High
<a href="#">Poma et al. (2017)</a>	SE	other	2015	8 (0.00)	0.5	High
<a href="#">Poma et al. (2017)</a>	SE	vegetables	2015	9 (0.67)	0.3	High

N/R = Not reported

444  
445  
446  
447  
448  
449  
450

### 1.6.2 Dietary (ng/g) – Wet Fraction

Measured concentrations of BCEP in Dietary with unit of ng/g, extracted from one source, are summarized in Figure 1-8 and supplemental information is provided in Table 1-8. Overall, concentrations ranged from not detected to 10.0 ng/g from 85 samples collected in 2018 in one country, AU. Location types were categorized as fruit, dairy, grain, vegetables, other, non-dairy beverages, meat and fish and shellfish. Reported detection frequency ranged from 0.0 to 0.33.



451  
452  
453

**Figure 1-8. Concentrations of BCEP (ng/g) in the Wet Fraction of Dietary in 2018**

454  
455

**Table 1-8. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Wet Fraction of Dietary**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">He et al. (2018b)</a>	AU	dairy	2018	9 (0.33)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	fish and shellfish	2018	9 (0.00)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	fruit	2018	15 (0.00)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	grain	2018	12 (0.00)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	meat	2018	12 (0.00)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	non-dairy beverages	2018	10 (0.00)	0.0013	Medium
<a href="#">He et al. (2018b)</a>	AU	other	2018	3 (0.00)	0.004	Medium
<a href="#">He et al. (2018b)</a>	AU	vegetables	2018	15 (0.00)	0.004	Medium

456

## **1.7 Drinking Water**

457

### **1.7.1 Drinking Water (ng/L) – Not Specified Fraction**

458

Measured concentrations of TCEP in Drinking Water with unit of ng/L, extracted from nine sources, are summarized in Figure 1-9 and supplemental information is provided in Table 1-9. Overall, concentrations ranged from not detected to 1,400.0 ng/L from 675 samples collected between 1982 and 2014 in six countries, CA, ES, JP, KR, PR and US. Location types were categorized as General Population (Background) and Unknown/Not Specified. Reported detection frequency ranged from 0.0 to 0.88.

459


460

461

462

463

464



465

466

467

468

469

470

**Figure 1-9. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Drinking Water from 1982 to 2014**

**Table 1-9. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not Specified Fraction of Drinking Water**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Padhye et al. (2014)</a>	US	General Population (Background)	2009–2010	8 (0.88)	N/R	Medium
<a href="#">Hopple et al. (2009)</a>	US	General Population (Background)	2004–2005	57 (0.02)	500.0	High
<a href="#">Kingsbury et al. (2008)</a>	US	General Population (Background)	2002–2004	337 (0.33)	500.0	High
<a href="#">Focazio et al. (2008)</a>	PR, US	Unknown/Not Specified	2001	73 (0.21)	100.0	Medium
<a href="#">Lebel et al. (1987)</a>	CA, US	General Population (Background)	1982–1983	20 (0.55)	N/R	Medium
<a href="#">Lee et al. (2016)</a>	KR	General Population (Background)	2014	127 (0.75)	0.7	Medium
<a href="#">Valcarcel et al. (2018)</a>	ES	General Population (Background)	2013	28 (0.75)	0.03	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Rodil et al. (2012)</a>	ES	General Population (Background)	2007–2008	24 (0.71)	4.0	Medium
<a href="#">Yasuhara (1994)</a>	JP	General Population (Background)	1994	1 (0.00)	67.5	Medium

N/R = Not reported

471

## **1.8 Dust (Indoor)**

---

472

### **1.8.1 Dust (Indoor) (ng/g) – Dry Fraction**

---

473

474

475


476

477


478

479

Measured concentrations of TCEP in Dust (Indoor) with unit of ng/g, extracted from 45 sources, are summarized in Figure 1-10 and supplemental information is provided in Table 1-10. Overall, concentrations ranged from not detected to 1,800,000.0 ng/g from 4,578 samples collected between 2000 and 2019 in 20 countries, AT, AU, BE, CA, CN, DE, DK, ES, FI, GB, GR, JP, KR, NL, NO, NZ, PT, RO, SE and US. Location types were categorized as Vehicle, Other, Public Space and Residential. Reported detection frequency ranged from 0.17 to 1.0.







481

482

483

484

485

**Figure 1-10. Concentrations of TCEP (ng/g) in the Dry Fraction of Dust (Indoor) from 2000 to 2019**

**Table 1-10. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Dust (Indoor)**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Hoffman et al. (2017)</a>	US	Residential	2014–2016	140 (N/R)	N/R	Medium
<a href="#">Phillips et al. (2018)</a>	US	Residential	2014–2016	188 (0.98)	18.7	High
<a href="#">Shin et al. (2019)</a>	US	Residential	2015–2016	38 (0.97)	25.0	Medium
<a href="#">La Guardia and Hale (2015)</a>	US	Public Space	2013	4 (1.00)	100.0	Medium
<a href="#">La Guardia and Hale (2015)</a>	US	Residential	2013	4 (1.00)	100.0	Medium
<a href="#">Stapleton et al. (2014)</a>	US	Residential	2012	30 (1.00)	N/R	High
<a href="#">Schreder and La Guardia (2014)</a>	US	Residential	2011–2012	20 (0.95)	1.0	High
<a href="#">Bradman et al. (2014)</a>	US	Public Space	2010–2011	39 (1.00)	1.0	High
<a href="#">Shin et al. (2014)</a>	US	Residential	2009–2010	30 (1.00)	1.0	High
<a href="#">Fang et al. (2013)</a>	US	Residential	2009	20 (0.50)	20.0	Medium
<a href="#">Fang et al. (2013)</a>	US	Vehicle	2009	20 (0.95)	20.0	Medium
<a href="#">Castorina et al. (2017)</a>	US	Residential	2000–2001	125 (1.00)	27.9	High
<a href="#">Tan et al. (2019)</a>	CN, US	Residential	2019	47 (1.00)	10.0	High
<a href="#">Velázquez-Gómez et al. (2019)</a>	ES	Public Space	2019	33 (1.00)	N/R	Medium
<a href="#">Velázquez-Gómez et al. (2019)</a>	ES	Residential	2019	11 (1.00)	N/R	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Velázquez-Gómez et al. (2019)</a>	ES	Vehicle	2019	14 (1.00)	N/R	Medium
<a href="#">Rantakokko et al. (2019)</a>	FI	Residential	2019	40 (1.00)	3.0	Medium
<a href="#">Liu and Mabury (2019)</a>	CA	Public Space	2018	85 (1.00)	0.4	High
<a href="#">Giovanoulis et al. (2019)</a>	SE	Public Space	2018	20 (1.00)	34.0	High
<a href="#">Langer et al. (2016)</a>	DK	Public Space	2016	151 (0.78)	600.0	High
<a href="#">Langer et al. (2016)</a>	DK	Residential	2016	497 (0.69)	600.0	High
<a href="#">Christia et al. (2018)</a>	GR	Vehicle	2016	25 (0.80)	N/R	High
<a href="#">Deng et al. (2018)</a>	CN	Public Space	2015–2016	22 (1.00)	N/R	Medium
<a href="#">Persson et al. (2018)</a>	SE	Public Space	2015–2016	31 (0.58)	6.9	High
<a href="#">Zhou et al. (2017)</a>	DE	Public Space	2015	48 (0.83)	115.0	High
<a href="#">Zhou et al. (2017)</a>	DE	Residential	2015	15 (0.80)	115.0	High
<a href="#">Zhou et al. (2017)</a>	DE	Vehicle	2015	11 (0.82)	115.0	High
<a href="#">He et al. (2018c)</a>	AU	Public Space	2015	30 (1.00)	10.0	High
<a href="#">He et al. (2018c)</a>	AU	Residential	2015	40 (1.00)	10.0	High
<a href="#">He et al. (2018c)</a>	AU	Vehicle	2015	15 (1.00)	10.0	High
<a href="#">Larsson et al. (2018)</a>	SE	Public Space	2015	100 (0.61)	1200.0	High
<a href="#">Takeuchi et al. (2015)</a>	JP	Residential	2013–2014	19 (0.95)	N/R	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Xu et al. (2016)</a>	NO	Residential	2013–2014	122 (0.76)	170.0	Medium
<a href="#">Kim and Tanabe (2017)</a>	KR	Public Space	2014	6 (0.17)	N/R	High
<a href="#">Kim and Tanabe (2017)</a>	KR	Residential	2013–2014	14 (1.00)	N/R	High
<a href="#">Kademoglou et al. (2017)</a>	GB,NO	Residential	2013–2014	20 (1.00)	44.1	Medium
<a href="#">Wallner et al. (2012)</a>	AT	Public Space	2012–2013	36 (1.00)	N/R	Medium
<a href="#">Tokumura et al. (2017)</a>	JP	Vehicle	2013	37 (1.00)	180.0	High
<a href="#">Sugeng et al. (2017)</a>	NL	Residential	2013	28 (0.82)	N/R	Medium
<a href="#">Kademoglou et al. (2017)</a>	GB	Public Space	2013	12 (1.00)	44.1	Medium
<a href="#">Ait Bamai et al. (2018)</a>	JP	Residential	2013	296 (0.84)	N/R	Medium
<a href="#">Ali et al. (2012)</a>	NZ	Residential	2012	50 (0.98)	20.0	Medium
<a href="#">Fromme et al. (2014)</a>	DE	Public Space	2011–2012	63 (1.00)	200.0	Medium
<a href="#">Brandsma et al. (2014)</a>	NL	Residential	2012	16 (1.00)	70.0	High
<a href="#">Brandsma et al. (2014)</a>	NL	Vehicle	2012	16 (1.00)	70.0	High
<a href="#">Coelho et al. (2016)</a>	PT	Residential	2010–2011	28 (0.82)	4.0	Medium
<a href="#">Brommer et al. (2012)</a>	DE	Residential	2010–2011	6 (N/R)	80.0	Medium
<a href="#">Bergh et al. (2011b)</a>	SE	Public Space	2010	20 (N/R)	N/R	Medium
<a href="#">Bergh et al. (2011b)</a>	SE	Residential	2010	10 (N/R)	N/R	Medium


Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Van den Eede et al. (2012)</a>	BE,ES,RO	Residential	2006–2010	12 (1.00)	110.0	Medium
<a href="#">Tajima et al. (2014)</a>	JP	Residential	2009–2010	256 (0.51)	1000.0	High
<a href="#">Fan et al. (2014)</a>	CA	Residential	2010	268 (0.96)	70.0	High
<a href="#">Mizouchi et al. (2015)</a>	JP	Residential	2009–2010	10 (1.00)	10.0	High
<a href="#">Bastiaensen et al. (2019a)</a>	JP	Residential	2009–2010	196 (0.59)	N/R	High
<a href="#">Luongo and Oestman (2016)</a>	SE	Residential	2008	62 (0.97)	190.0	Medium
<a href="#">Kanazawa et al. (2010)</a>	JP	Residential	2006	82 (0.95)	1300.0	Medium
<a href="#">Marklund et al. (2003)</a>	SE	Other	2003	5 (1.00)	N/R	Medium
<a href="#">Marklund et al. (2003)</a>	SE	Public Space	2003	9 (1.00)	N/R	Medium
<a href="#">Marklund et al. (2003)</a>	SE	Residential	2003	2 (1.00)	N/R	Medium
<a href="#">Marklund et al. (2003)</a>	SE	Vehicle	2003	1 (1.00)	N/R	Medium
<a href="#">Ingerowski et al. (2001)</a>	DE	Residential	2001	983 (N/R)	400.0	Medium

N/R = Not reported

488  
489  
490  
491  
492  
493

### **1.8.2 Dust (Indoor) (ng/g) – Dry Fraction**

Measured concentrations of BCEP in Dust (Indoor) with unit of ng/g, extracted from one source, are summarized in Figure 1-11 and supplemental information is provided in Table 1-11. Overall, concentrations were not detected ng/g from 47 samples collected in 2019 in two countries, CN and US. Location types were categorized as Residential. Reported detection frequency was 0.0.



494

495 **Figure 1-11. Concentrations of BCEP (ng/g) in the Dry Fraction of Dust (Indoor) in Residential**  
 496 **Locations in 2019**

497

498 **Table 1-11. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Dry**  
 499 **Fraction of Dust (Indoor)**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Tan et al. (2019)</a>	CN,US	Residential	2019	47 (0.00)	16	High


500

### 1.8.3 Dust (Indoor) (ng/m<sup>2</sup>) – Dry Fraction

501

501 Measured concentrations of TCEP in Dust (Indoor) with unit of ng/m<sup>2</sup>, extracted from four sources, are  
 502 summarized in Figure 1-12 and supplemental information is provided in Table 1-12. Overall,  
 503 concentrations ranged from not detected to 1,243,900.0 ng/m<sup>2</sup> from 180 samples collected between 2000  
 504 and 2016 in two countries, SE and US. Location types were categorized as Public Space, Unknown and  
 505 Residential. Reported detection frequency ranged from 0.0 to 1.0.  
 506

506



507

508 **Figure 1-12. Concentrations of TCEP (ng/m<sup>2</sup>) in the Dry Fraction of Dust (Indoor) from 2000 to**  
 509 **2016**

510

511 **Table 1-12. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m<sup>2</sup>) Levels in the**  
 512 **Dry Fraction of Dust (Indoor)**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>2</sup> )	Overall Quality Level
<a href="#">Dodson et al. (2017)</a>	US	Residential	2013–2014	37 (0.00)	10,763.91042	High
<a href="#">Castorina et al. (2017)</a>	US	Residential	2000–2001	125 (1.00)	27.9	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>2</sup> )	Overall Quality Level
<a href="#">Persson et al. (2018)</a>	SE	Public Space	2015–2016	16 (0.44)	0.07	High
<a href="#">Marklund et al. (2003)</a>	SE	Unknown	2003	2 (1.00)	N/R	Medium

N/R = Not reported

513

## 1.9 Groundwater

514

### 1.9.1 Groundwater (ng/L) – Not Specified Fraction

515

516


517

518

519

520

Measured concentrations of TCEP in Groundwater with unit of ng/L, extracted from 11 sources, are summarized in Figure 1-13 and supplemental information is provided in Table 1-13. Overall, concentrations ranged from not detected to 810.0 ng/L from 582 samples collected between 1978 and 2017 in four countries, DE, JP, SE and US. Location types were categorized as General Population (Background) and Near Facility (Highly Exposed). Reported detection frequency ranged from 0.0 to 1.0.



521

522

523

524

**Figure 1-13. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Groundwater from 1978 to 2017**

**Table 1-13. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not Specified Fraction of Groundwater**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Laws et al. (2011)</a>	US	Near Facility (Highly Exposed)	2009	11 (1.00)	10.0	Medium
<a href="#">Hopple et al. (2009)</a>	US	General Population (Background)	2002–2005	276 (0.02)	500.0	High
<a href="#">Buszka et al. (2009)</a>	US	Near Facility (Highly Exposed)	2000–2002	6 (0.33)	500.0	Medium
<a href="#">Barnes et al. (2008)</a>	US	Near Facility (Highly Exposed)	2000	47 (0.30)	500.0	Medium
<a href="#">Barnes et al. (2004)</a>	US	Near Facility (Highly Exposed)	2000	5 (1.00)	40.0	Medium
<a href="#">Hutchins et al. (1984)</a>	US	Near Facility (Highly Exposed)	1978	4 (N/R)	N/R	Medium
<a href="#">Gao et al. (2019)</a>	SE	General Population (Background)	2016–2017	30 (0.83)	7.2	High
<a href="#">Regnery et al. (2011)</a>	DE	General Population (Background)	2009	25 (0.56)	1.0	High
<a href="#">Regnery et al. (2011)</a>	DE	Near Facility (Highly Exposed)	2009	11 (0.91)	1.0	High
<a href="#">Fries and Puttmann (2003)</a>	DE	General Population (Background)	2000–2001	76 (N/R)	1.0	Medium
<a href="#">Fries and Puttmann (2001)</a>	DE	General Population (Background)	2000	90 (N/R)	1.0	Medium
<a href="#">Yasuhara (1994)</a>	JP	General Population (Background)	1994	1 (0.00)	67.5	Medium

N/R = Not reported



## 1.10 Human Biomonitoring – Breastmilk

528

### 1.10.1 Human Biomonitoring – Breastmilk (ng/L) – wet Fraction

529

530

531

532

533

534

Measured concentrations of TCEP in Human Biomonitoring – Breastmilk with unit of ng/L, extracted from one source, are summarized in Figure 1-14 and supplemental information is provided in Table 1-14. Overall, concentrations ranged from not detected to 470 ng/L from three samples collected between 2014 and 2015 in one country, AU. Location types were categorized as General Population (Background). Reported detection frequency was 0.67.



535

536

537

538

**Figure 1-14. Concentrations of TCEP (ng/L) in the wet Fraction of Human Biomonitoring – Breastmilk in General Population (Background) Locations from 2014 to 2015**

539

540

**Table 1-14. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the wet Fraction of Human Biomonitoring – Breastmilk**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">He et al. (2018a)</a>	AU	General Population (Background)	2014–2015	3 (0.67)	260	High

541

### 1.10.2 Human Biomonitoring – Breastmilk (ng/g) – Lipid Fraction

542

543


544

545

546

547

Measured concentrations of TCEP in Human Biomonitoring – Breastmilk with unit of ng/g, extracted from 2 sources, are summarized in Figure 1-15 and supplemental information is provided in Table 1-15. Overall, concentrations ranged from not detected to 512.0 ng/g from 93 samples collected between 1997 and 2011 in four countries, JP, PH, SE and VN. Location types were categorized as General Population (Background) and Near Facility (Highly Exposed). Reported detection frequency was 1.0.



548

549

550

551

**Figure 1-15. Concentrations of TCEP (ng/g) in the Lipid Fraction of Human Biomonitoring – Breastmilk from 1997 to 2011**

552  
553

**Table 1-15. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Lipid Fraction of Human Biomonitoring – Breastmilk**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Kim et al. (2014)</a>	JP, PH, VN	General Population (Background)	2008–2011	46 (N/R)	0.045	Medium
<a href="#">Kim et al. (2014)</a>	PH, VN	Near Facility (Highly Exposed)	2008	41 (N/R)	0.045	Medium
<a href="#">Sundkvist et al. (2010)</a>	SE	General Population (Background)	1997–2006	6 (1.00)	0.4	High

N/R = Not reported

554

## 1.11 Human Biomonitoring – Hair

555

### 1.11.1 Human Biomonitoring – Hair (ng/g) – Dry Fraction

556

Measured concentrations of TCEP in Human Biomonitoring – Hair with unit of ng/g, extracted from two sources, are summarized in Figure 1-16 and supplemental information is provided in Table 1-16.

557

558


Overall, concentrations ranged from 37.5 to 2,740 ng/g from 55 samples collected between 2014 and 2015 in one country, US. Location types were categorized as General Population (Background).

559

560

Reported detection frequency ranged from 0.68 to 0.8.

561



562

563

**Figure 1-16. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Hair in General Population (Background) Locations from 2014 to 2015**

564

565

566

**Table 1-16. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Human Biomonitoring – Hair**

567

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Liu et al. (2015)</a>	US	General Population (Background)	2015	5 (0.80)	75.0	Medium
<a href="#">Liu et al. (2016)</a>	US	General Population (Background)	2014	50 (0.68)	N/R	Medium

N/R = Not reported

568

## 1.12 Human Biomonitoring – Nails

569

### 1.12.1 Human Biomonitoring – Nails (ng/g) – Dry Fraction

570

Measured concentrations of TCEP in Human Biomonitoring – Nails with unit of ng/g, extracted from two sources, are summarized in Figure 1-17 and supplemental information is provided in Table 1-17.

571

Overall, concentrations ranged from not detected to 1860.0 ng/g from 105 samples collected between 2014 and 2015 in one country, US. Location types were categorized as General Population

572

(Background). Reported detection frequency ranged from 0.0 to 0.14.

573

(Background). Reported detection frequency ranged from 0.0 to 0.14.

574

(Background). Reported detection frequency ranged from 0.0 to 0.14.

575



576

**Figure 1-17. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Nails in General Population (Background) Locations from 2014 to 2015**

577

**Table 1-17. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Human Biomonitoring – Nails**

578

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Liu et al. (2015)</a>	US	General Population (Background)	2015	5 (0.00)	150.0	Medium
<a href="#">Liu et al. (2016)</a>	US	General Population (Background)	2014	100 (0.14)	N/R	Medium

N/R = Not reported

582

## 1.13 Human Biomonitoring – Other

583

### 1.13.1 Human Biomonitoring – Other (ng/g) – Dry Fraction

584

Measured concentrations of TCEP in Human Biomonitoring – Other with unit of ng/g, extracted from one source, are summarized in Figure 1-18 and supplemental information is provided in Table 1-18.

585

Overall, concentrations ranged from 0.055 to 41.8 ng/g from 100 samples collected between 2014 and 2016 in one country, CN. Location types were categorized as General Population (Background).

586

Reported detection frequency was 0.66.


587

Reported detection frequency was 0.66.

588

Reported detection frequency was 0.66.

589



590

**Figure 1-18. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Other in General Population (Background) Locations from 2014 to 2016**

591

(Background). Reported detection frequency was 0.66.

592

(Background). Reported detection frequency was 0.66.

593


594  
595

**Table 1-18. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Human Biomonitoring – Other**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Zhao et al. (2017)</a>	CN	General Population (Background)	2014–2016	100 (0.66)	0.11	High

596 **1.13.2 Human Biomonitoring – Other (ng/g) – Dry Fraction**

597 Measured concentrations of BCEP in Human Biomonitoring – Other with unit of ng/g, extracted from  
598 one source, are summarized in Figure 1-19 and supplemental information is provided in Table 1-19.  
599 Overall, concentrations ranged from 0.44 to 1,180 ng/g from 50 samples collected between 2014 and  
600 2016 in one country, CN. Location types were categorized as General Population (Background).  
601 Reported detection frequency was 0.88.  
602



603  
604 **Figure 1-19. Concentrations of BCEP (ng/g) in the Dry Fraction of Human Biomonitoring – Other**  
605 **in General Population (Background) Locations from 2014 to 2016**

606  
607 **Table 1-19. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Dry**  
608 **Fraction of Human Biomonitoring – Other**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Zhao et al. (2017)</a>	CN	General Population (Background)	2014–2016	50 (0.88)	0.88	High

609 **1.14 Human Biomonitoring – Plasma**

610 **1.14.1 Human Biomonitoring – Plasma (ng/L) – Wet Fraction**

611 Measured concentrations of TCEP in Human Biomonitoring – Plasma with unit of ng/L, extracted from  
612 one source, are summarized in Figure 1-20 and supplemental information is provided in Table 1-20.  
613 Overall, concentrations ranged from not detected to 230 ng/L from 25 samples collected between 2014  
614 and 2016 in one country, CN. Location types were categorized as General Population (Background).  
615 Reported detection frequency was 0.48.  
616



617  
618 **Figure 1-20. Concentrations of TCEP (ng/L) in the Wet Fraction of Human Biomonitoring –**  
619 **Plasma in General Population (Background) Locations from 2014 to 2016**

620

621 **Table 1-20. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet**  
 622 **Fraction of Human Biomonitoring – Plasma**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Zhao et al. (2017)</a>	CN	General Population (Background)	2014–2016	25 (0.48)	90	High

## 623 1.15 Human Biomonitoring – Serum

### 624 1.15.1 Human Biomonitoring – Serum (ng/g) – Lipid Fraction

625 Measured concentrations of TCEP in Human Biomonitoring – Serum with unit of ng/g, extracted from  
 626 one source, are summarized in Figure 1-21 and supplemental information is provided in Table 1-21.  
 627 Overall, concentrations ranged from 3.12 to 3.69 ng/g from 20 samples collected in 2016 in one country,  
 628 ES. Location types were categorized as General Population (Background). Reported detection frequency  
 629 was 1.0.  
 630



631  
 632 **Figure 1-21. Concentrations of TCEP (ng/g) in the Lipid Fraction of Human Biomonitoring –**  
 633 **Serum in General Population (Background) Locations in 2016**

634  
 635 **Table 1-21. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the**  
 636 **Lipid Fraction of Human Biomonitoring – Serum**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Henríquez-Hernández et al. (2017)</a>	ES	General Population (Background)	2016	20 (1.00)	N/R	High

N/R = Not reported


## 637 1.16 Human Biomonitoring – Skin\_Dermal Wipe

### 638 1.16.1 Human Biomonitoring – Skin\_Dermal Wipe (ng/g) – Dry Fraction

639 Measured concentrations of TCEP in Human Biomonitoring – Skin\_Dermal Wipe with unit of ng/g,  
 640 extracted from one source, are summarized in Figure 1-22 and supplemental information is provided in  
 641 Table 1-22. Overall, concentrations ranged from 20 to 6,920 ng/g from 30 samples collected in 2012 in  
 642 one country, US. Location types were categorized as General Population (Background). Reported

643 detection frequency was 1.0.

644



645

646 **Figure 1-22. Concentrations of TCEP (ng/g) in the Dry Fraction of Human Biomonitoring –**  
 647 **Skin\_Dermal Wipe in General Population (Background) Locations in 2012**

648

649 **Table 1-22. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry**  
 650 **Fraction of Human Biomonitoring – Skin\_Dermal Wipe**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Stapleton et al. (2014)</a>	US	General Population (Background)	2012	30 (1.00)	N/R	High

N/R = Not reported


651

**1.16.2 Human Biomonitoring – Skin\_Dermal Wipe (ng/wipe) – Dry Fraction**

652

653 Measured concentrations of TCEP in Human Biomonitoring – Skin\_Dermal Wipe with unit of ng/wipe,  
 654 extracted from four sources, are summarized in Figure 1-22 and supplemental information is provided in  
 655 Table 1-23. Overall, concentrations ranged from not detected to 3,216 ng/wipe from 400 samples  
 656 collected between 2012 and 2016 in three countries, NO, SE and US. Location types were categorized as  
 657 General Population (Background). Reported detection frequency ranged from 0.47 to 0.87.

657



658

659 **Figure 1-23. Concentrations of TCEP (ng/wipe) in the Dry Fraction of Human Biomonitoring –**  
 660 **Skin\_Dermal Wipe in General Population (Background) Locations from 2012 to 2016**

661

662 **Table 1-23. Summary of Peer-Reviewed Literature that Measured TCEP (ng/wipe) Levels in the**  
 663 **Dry Fraction of Human Biomonitoring – Skin\_Dermal Wipe**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/wipe)	Overall Quality Level
<a href="#">Phillips et al. (2018)</a>	US	General Population (Background)	2014–2016	202 (0.87)	2.7	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/wipe)	Overall Quality Level
<a href="#">Stapleton et al. (2014)</a>	US	General Population (Background)	2012	43 (0.47)	24.0	High
<a href="#">Larsson et al. (2018)</a>	SE	General Population (Background)	2015	100 (0.51)	4.5	High
<a href="#">Xu et al. (2016)</a>	NO	General Population (Background)	2013–2014	55 (0.49)	N/R	Medium

N/R = Not reported

664

## 1.17 Human Biomonitoring – Urine

665

### 1.17.1 Human Biomonitoring – Urine (ng/g) – Creatinine Adjusted Fraction

666

Measured concentrations of BCEP in Human Biomonitoring – Urine with unit of ng/g, extracted from one source, are summarized in Figure 1-23 and supplemental information is provided in Table 1-24.

667

Overall, concentrations ranged from not detected to 1900 ng/g from 213 samples collected in 2018 in

668


one country, US. Location types were categorized as General Population (Background). Reported

669

detection frequency was 0.87.

670

671



672

673

**Figure 1-24. Concentrations of BCEP (ng/g) in the Creatinine Adjusted Fraction of Human Biomonitoring – Urine in General Population (Background) Locations in 2018**

674

675

676

**Table 1-24. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Creatinine Adjusted Fraction of Human Biomonitoring – Urine**

677

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Wang et al. (2019)</a>	US	General Population (Background)	2018	213 (0.87)	2.7	High

678

### 1.17.2 Human Biomonitoring – Urine (ng/L) – Unadjusted Fraction

679

Measured concentrations of TCEP in Human Biomonitoring – Urine with unit of ng/L, extracted from three sources, are summarized in Figure 1-25 and supplemental information is provided in Table 1-25.

680

Overall, concentrations ranged from not detected to 24500 ng/L from 594 samples collected between

681


2010 and 2015 in two countries, AU and BE. Location types were categorized as General Population

682



683 (Background). Reported detection frequency ranged from 0.11 to 0.55.

684



685

686 **Figure 1-25. Concentrations of TCEP (ng/L) in the Unadjusted Fraction of Human Biomonitoring**  
 687 **– Urine in General Population (Background) Locations from 2010 to 2015**

688

689 **Table 1-25. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the**  
 690 **Unadjusted Fraction of Human Biomonitoring – Urine**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">He et al. (2018a)</a>	AU	General Population (Background)	2014–2015	400 (0.45)	22.0	High
<a href="#">Bastiaensen et al. (2019b)</a>	BE	General Population (Background)	2015	99 (0.55)	32.0	Medium
<a href="#">Van Den Eede et al. (2015)</a>	AU	General Population (Background)	2010–2013	95 (0.11)	350.0	Medium

691 **1.17.3 Human Biomonitoring – Urine (ng/L) – All Fractions**

692 Measured concentrations of BCEP in Human Biomonitoring – Urine with unit of ng/L, extracted from  
 693 four sources, are summarized in Figure 1-24 and supplemental information is provided in Table 1-26.  
 694 More than one weight fraction was reported and summarized separately below:

695


696 Overall, concentrations for Creatinine Adjusted ranged from not detected to 13.5 ng/L from 213 samples  
 697 collected in 2018 in one country, US. Location types were categorized as General Population  
 698 (Background). Reported detection frequency was 0.87.

699

700 Overall, concentrations for Unadjusted ranged from not detected to 13100.0 ng/L from 728 samples  
 701 collected between 2011 and 2015 in three countries, AU, DE and US. Location types were categorized  
 702 as General Population (Background). Reported detection frequency ranged from 0.15 to 0.75.

703





704

705 **Figure 1-26. Concentrations of BCEP (ng/L) in Human Biomonitoring – Urine in General**  
 706 **Population (Background) Locations from 2011 to 2018**

707

708 **Table 1-26. Summary of Peer-Reviewed Literature that Measured BCEP (ng/L) Levels in Human**  
 709 **Biomonitoring – Urine**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
Creatinine Adjusted						
<a href="#">Wang et al. (2019)</a>	US	General Population (Background)	2018	213 (0.87)	2.7	High
Unadjusted						
<a href="#">Dodson et al. (2014)</a>	US	General Population (Background)	2011	16 (0.75)	100.0	High
<a href="#">He et al. (2018a)</a>	AU	General Population (Background)	2014–2015	400 (0.15)	14.0	High
<a href="#">Fromme et al. (2014)</a>	DE	General Population (Background)	2011–2012	312 (0.65)	200.0	Medium

710

## 1.18 Human Biomonitoring – Silicone Wristbands


711

### 1.18.1 Human Biomonitoring – Silicone Wristbands (ng/g) – Not Specified Fraction

712

713 Measured concentrations of TCEP in Human Biomonitoring – Silicone Wristbands with unit of ng/g,  
 714 extracted from two sources, are summarized in Figure 1-27 and supplemental information is provided in  
 715 Table 1-27. Overall, concentrations ranged from not detected to 719.0 ng/g from 140 samples collected  
 716 between 2012 and 2015 in one country, US. Location types were categorized as General Population  
 717 (Background). Reported detection frequency ranged from 0.83 to 0.89.

717



718  
719  
720  
721  
722  
723  
724

**Figure 1-27. Concentrations of TCEP (ng/g) in the Not Specified Fraction of Human Biomonitoring – Silicone Wristbands in General Population (Background) Locations from 2012 to 2015**

**Table 1-27. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Not Specified Fraction of Human Biomonitoring – Silicone Wristbands**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Gibson et al. (2019)</a>	US	General Population (Background)	2015	76 (0.83)	3.27	High
<a href="#">Kile et al. (2016)</a>	US	General Population (Background)	2012–2013	64 (0.89)	3.4	Medium

725

## 1.19 Indoor Air

726

### 1.19.1 Indoor Air (ng/m<sup>3</sup>) – All Fractions

727

Measured concentrations of TCEP in Indoor Air with unit of ng/m<sup>3</sup>, extracted from 27 sources, are summarized in Figure 1-25 and supplemental information is provided in Table 1-28. More than one weight fraction was reported and summarized separately below:

728

729

730

731

Overall, concentrations for Combined Vapor/Gas and Particulate ranged from not detected to 6,000.0 ng/m<sup>3</sup> from 440 samples collected between 2000 and 2016 in seven countries, AU, BE, CA, DE, FI, JP and US. Location types were categorized as Public Space and Residential. Reported detection frequency ranged from 0.32 to 1.0.

732

733

734

735

Overall, concentrations for Particulate ranged from not detected to 136.0 ng/m<sup>3</sup> from 133 samples collected between 2002 and 2016 in four countries, CN, JP, SE and US. Location types were categorized as Public Space and Residential. Reported detection frequency ranged from 0.0 to 1.0.

736

737

738

739

Overall, concentrations for Vapor/Gas ranged from not detected to 7,100.0 ng/m<sup>3</sup> from 677 samples collected between 2000 and 2016 in six countries, CH, DE, JP, NO, SE and US. Location types were categorized as Vehicle, Public Space and Residential. Reported detection frequency ranged from 0.0 to 1.0.


740


741

742

743

744





746

747 **Figure 1-28. Concentrations of TCEP (ng/m<sup>3</sup>) in Indoor Air from 2000 to 2016**

748

749 **Table 1-28. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m<sup>3</sup>) Levels in Indoor**  
 750 **Air**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
Combined Vapor/Gas and Particulate						
<a href="#">Dodson et al. (2019)</a>	US	Public Space	2013–2015	37 (0.32)	5.6	High
<a href="#">Okeme et al. (2018b)</a>	CA	Public Space	2016	51 (0.80)	N/R	Medium
<a href="#">He et al. (2018c)</a>	AU	Public Space	2015	40 (1.00)	0.06	High
<a href="#">He et al. (2018c)</a>	AU	Residential	2015	40 (1.00)	0.06	High
<a href="#">Okeme et al. (2018b)</a>	CA	Residential	2015	102 (0.77)	N/R	Medium
<a href="#">Lazarov et al. (2015)</a>	BE	Residential	2015	6 (N/R)	0.171	Medium
<a href="#">Takeuchi et al. (2015)</a>	JP	Residential	2013–2014	21 (0.90)	0.07	High
<a href="#">Mäkinen et al. (2009)</a>	FI	Public Space	2008	3 (1.00)	N/R	Medium
<a href="#">Mäkinen et al. (2009)</a>	FI	Public Space	2008	4 (0.50)	3.0	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
<a href="#">Kanazawa et al. (2010)</a>	JP	Residential	2006	40 (0.60)	12.6	Medium
<a href="#">Ohura et al. (2006)</a>	JP	Residential	2000–2001	46 (0.89)	N/R	Medium
<a href="#">Ingerowski et al. (2001)</a>	DE	Residential	2001	50 (1.00)	N/R	Medium
Particulate						
<a href="#">La Guardia and Hale (2015)</a>	US	Public Space	2013	8 (0.00)	0.1	Medium
<a href="#">La Guardia and Hale (2015)</a>	US	Residential	2013	8 (0.00)	0.1	Medium
<a href="#">Bradman et al. (2014)</a>	US	Public Space	2010–2011	40 (0.65)	0.3	High
<a href="#">Deng et al. (2018)</a>	CN	Public Space	2015–2016	22 (1.00)	N/R	Medium
<a href="#">Wong et al. (2018)</a>	SE	Public Space	2014–2015	23 (1.00)	0.022	Medium
<a href="#">Saito et al. (2007)</a>	JP	Residential	2002	18 (N/R)	0.67	Medium
<a href="#">Saito et al. (2007)</a>	JP	Public Space	2002	14 (N/R)	0.67	Medium
Vapor/Gas						
<a href="#">Dodson et al. (2017)</a>	US	Residential	2013–2014	35 (0.17)	7.3	High
<a href="#">Persson et al. (2018)</a>	SE	Public Space	2015–2016	56 (0.00)	2.2	High
<a href="#">Sha et al. (2018)</a>	SE	Public Space	2016	36 (N/R)	0.0094	Low
<a href="#">Sha et al. (2018)</a>	SE	Residential	2016	9 (N/R)	0.0094	Low
<a href="#">Xu et al. (2016)</a>	NO	Residential	2013–2014	58 (0.93)	0.9	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
<a href="#">Tokumura et al. (2017)</a>	JP	Vehicle	2013	9 (0.00)	0.68	High
<a href="#">Fromme et al. (2014)</a>	DE	Public Space	2011–2012	63 (0.17)	4.0	Medium
<a href="#">Bergh et al. (2011b)</a>	SE	Public Space	2010	20 (N/R)	N/R	Medium
<a href="#">Bergh et al. (2011b)</a>	SE	Residential	2010	10 (N/R)	N/R	Medium
<a href="#">Luongo and Oestman (2016)</a>	SE	Residential	2008	62 (0.65)	N/R	Medium
<a href="#">Bergh et al. (2011a)</a>	SE	Residential	2006–2007	169 (N/R)	1.0	Medium
<a href="#">Hartmann et al. (2004)</a>	CH	Public Space	2004	12 (1.00)	0.15	Medium
<a href="#">Hartmann et al. (2004)</a>	CH	Vehicle	2004	4 (0.75)	0.15	Medium
<a href="#">Yoshida et al. (2006)</a>	JP	Vehicle	2004	101 (0.80)	N/R	Medium
<a href="#">Otake et al. (2004)</a>	JP	Residential	2000	27 (N/R)	N/R	Medium
<a href="#">Otake et al. (2001)</a>	JP	Residential	2000	6 (1.00)	N/R	Medium

N/R = Not reported

751

## 1.20 Leachate

752

### 1.20.1 Leachate (ng/L) – Not Specified Fraction

753

754


755

756

757

758

Measured concentrations of TCEP in leachate with unit of ng/L, extracted from three sources, are summarized in Figure 1-26 and supplemental information is provided in Table 1-29. Overall, concentrations ranged from 6 to 5,430,000,000,000.0 ng/L from 20 samples collected between 1994 and 1995 in one country, JP. Location types were categorized as Unknown/Not Specified and Near Facility (Highly Exposed). Reported detection frequency was 1.0.



759

760 **Figure 1-29. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Leachate from 1994**  
 761 **to 1995**

762

763 **Table 1-29. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not**  
 764 **Specified Fraction of Leachate**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Yasuhara et al. (1999)</a>	JP	Unknown/Not Specified	1995	11 (1.00)	N/R	Medium
<a href="#">Yasuhara (1995)</a>	JP	Near Facility (Highly Exposed)	1995	5 (1.00)	N/R	Low
<a href="#">Yasuhara (1994)</a>	JP	Near Facility (Highly Exposed)	1994	4 (1.00)	67.5	Medium

N/R = Not reported

765


## 1.21 Other

766

### 1.21.1 Other (ng/g) – Dry Fraction

767

768 Measured concentrations of TCEP in Other with unit of ng/g, extracted from one source, are  
 769 summarized in Figure 1-27 and supplemental information is provided in Table 1-30. Overall,  
 770 concentrations ranged from 0.007 to 0.039 ng/g from six samples collected in 2003 in one country, SE.  
 771 Location types were categorized as Unknown/Not Specified. Reported detection frequency was 1.0.



772

773 **Figure 1-30. Concentrations of TCEP (ng/g) in the Dry Fraction of Other in Unknown/Not**  
 774 **Specified Locations in 2003**

775

776  
777

**Table 1-30. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Other**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Marklund et al. (2005b)</a>	SE	Unknown/Not Specified	2003	6 (1.00)	N/R	Medium

N/R = Not reported


778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790

**1.21.2 Other (ng/g) – All Fractions**

Measured concentrations of TCEP in Other with unit of ng/g, extracted from three sources, are summarized in Figure 1-28 and supplemental information is provided in Table 1-31. More than one weight fraction was reported and summarized separately below:

Overall, concentrations for Particulate ranged from 0.007 to 68,000,000.0 ng/g from 12 samples collected between 2001 and 2003 in two countries, DE and SE. Location types were categorized as General Population (Background) and Unknown/Not Specified. Reported detection frequency was 1.0.

Overall, concentrations for Wet ranged from not detected to 0.55 ng/g from three samples collected in 2008 in one country, NL. Location types were categorized as Near Facility (Highly Exposed). Reported detection frequency was 0.67.



791  
792  
793  
794

**Figure 1-31. Concentrations of TCEP (ng/g) in Other from 2001 to 2008**

**Table 1-31. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Other**


Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Particulate						
<a href="#">Ingerowski et al. (2001)</a>	DE	General Population (Background)	2001	6 (1.00)	400.0	Medium
<a href="#">Marklund et al. (2005b)</a>	SE	Unknown/Not Specified	2003	6 (1.00)	N/R	Medium
Wet						



Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	3 (0.67)	0.2	High
N/R = Not reported						

795 **1.21.3 Other (ng/L) – Not Specified Fraction**

796 Measured concentrations of TCEP in Other with unit of ng/L, extracted from one source, are  
 797 summarized in Figure 1-29 and supplemental information is provided in Table 1-32. Overall,  
 798 concentrations ranged from 2.5 to 293 ng/L from 42 samples collected in 2016 in one country, AU.  
 799 Location types were categorized as General Population (Background). Reported detection frequency was  
 800 not reported.  
 801



802  
 803 **Figure 1-32. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Other in General**  
 804 **Population (Background) Locations in 2016**

805  
 806 **Table 1-32. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not**  
 807 **Specified Fraction of Other**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Teo et al. (2016)</a>	AU	General Population (Background)	2016	42 (N/R)	5	High
N/R = Not reported						

808 **1.22 Personal Inhalation**

809 **1.22.1 Personal Inhalation (ng/m<sup>3</sup>) – All Fractions**


810 Measured concentrations of TCEP in Personal Inhalation with unit of ng/m<sup>3</sup>, extracted from three  
 811 sources, are summarized in Figure 1-30 and supplemental information is provided in Table 1-33. More  
 812 than one weight fraction was reported and summarized separately below:  
 813

814 Overall, concentrations for Particulate ranged from not detected to 77.8 ng/m<sup>3</sup> from 21 samples collected  
 815 between 2015 and 2016 in two countries, CA and US. Location types were categorized as General  
 816 Population (Background). Reported detection frequency ranged from 0.44 to 1.0.  
 817

818 Overall, concentrations for Vapor/Gas ranged from 0.5 to 8.1 ng/m<sup>3</sup> from 31 samples collected between  
 819 2013 and 2014 in one country, NO. Location types were categorized as General Population

820 (Background). Reported detection frequency was 0.77.

821



822

823 **Figure 1-33. Concentrations of TCEP (ng/m<sup>3</sup>) in Personal Inhalation in General Population**  
 824 **(Background) Locations from 2013 to 2016**

825

826 **Table 1-33. Summary of Peer-Reviewed Literature that Measured TCEP (ng/m<sup>3</sup>) Levels in**  
 827 **Personal Inhalation**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/m <sup>3</sup> )	Overall Quality Level
Particulate						
<a href="#">Schreder et al. (2016)</a>	US	General Population (Background)	2015	18 (0.44)	1.5	High
<a href="#">Okeme et al. (2018a)</a>	CA	General Population (Background)	2016	3 (1.00)	0.012	Medium
Vapor/Gas						
<a href="#">Xu et al. (2016)</a>	NO	General Population (Background)	2013–2014	31 (0.77)	1.0	Medium

828

## 1.23 Precipitation

829

### 1.23.1 Precipitation (ng/L) – Wet Fraction

830

831


832

833

834

835

Measured concentrations of TCEP in Precipitation with unit of ng/L, extracted from six sources, are summarized in Figure 1-31 and supplemental information is provided in Table 1-34. Overall, concentrations ranged from not detected to 488.0 ng/L from 313 samples collected between 1994 and 2014 in three countries, AQ, DE and US. Location types were categorized as General Population (Background) and Remote (Not Near Source). Reported detection frequency ranged from 0.6 to 1.0.



836

837 **Figure 1-34. Concentrations of TCEP (ng/L) in the Wet Fraction of Precipitation from 1994 to**  
 838 **2014**

839

840 **Table 1-34. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet**  
 841 **Fraction of Precipitation**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Scott et al. (1996)</a>	US	General Population (Background)	1994	5 (0.60)	N/R	Low
<a href="#">Li et al. (2017)</a>	AQ	Remote (Not Near Source)	2014	6 (1.00)	0.21	High
<a href="#">Mihajlovic and Fries (2012)</a>	DE	General Population (Background)	2011	4 (N/R)	N/R	High
<a href="#">Mihajlovic and Fries (2012)</a>	DE	General Population (Background)	2010	4 (N/R)	N/R	High
<a href="#">Regnery and Püttmann (2010b)</a>	DE	General Population (Background)	2007–2009	167 (N/R)	2.0	High
<a href="#">Regnery and Püttmann (2010b)</a>	DE	General Population (Background)	2007–2009	29 (1.00)	2.0	High
<a href="#">Regnery and Püttmann (2009)</a>	DE	General Population (Background)	2007–2008	30 (N/R)	2.0	High

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Regnery and Puettmann (2009)</a>	DE	Remote (Not Near Source)	2007-2008	23 (N/R)	2.0	High
<a href="#">Regnery and Puettmann (2009)</a>	DE	General Population (Background)	2007-2008	8 (N/R)	2.0	High
<a href="#">Regnery and Puettmann (2009)</a>	DE	Remote (Not Near Source)	2007-2008	34 (N/R)	2.0	High
<a href="#">Fries and Puttmann (2003)</a>	DE	General Population (Background)	2001	3 (1.00)	1.0	Medium

N/R = Not reported

842

## **1.24 Sediment**

---

843

### **1.24.1 Sediment (ng/g) – All Fractions**

---

844

Measured concentrations of TCEP in Sediment with unit of ng/g, extracted from seven sources, are summarized in Figure 1-32 and supplemental information is provided in Table 1-35. More than one weight fraction was reported and summarized separately below:

845

846

847

848

Overall, concentrations for Dry ranged from not detected to 41.0 ng/g from 91 samples collected between 1980 and 2017 in seven countries, CZ, DE, JP, KR, PT, US and ZA. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Unknown/Not Specified. Reported detection frequency ranged from 0.75 to 1.0.

849

850

851

852


853

Overall, concentrations for Wet ranged from not detected to 0.35 ng/g from three samples collected in 2008 in one country, NL. Location types were categorized as Near Facility (Highly Exposed). Reported detection frequency was 0.67.

854

855

856



857

858

859

860

861

**Figure 1-35. Concentrations of TCEP (ng/g) in Sediment from 1980 to 2017**

**Table 1-35. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Sediment**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Dry						
<a href="#">Maruya et al. (2016)</a>	US	General Population (Background)	2013	16 (0.75)	N/R	High
<a href="#">Gadelha et al. (2019)</a>	PT	Near Facility (Highly Exposed)	2016–2017	12 (N/R)	0.07	High
<a href="#">Chokwe and Okonkwo (2019)</a>	ZA	Unknown/Not Specified	2017	16 (0.88)	0.24	High
<a href="#">Choo et al. (2018)</a>	KR	General Population (Background)	2015	4 (1.00)	0.01	High
<a href="#">Stachel et al. (2005)</a>	CZ,DE	Near Facility (Highly Exposed)	2002	37 (N/R)	1.0	Medium
<a href="#">Ishikawa et al. (1985)</a>	JP	General Population (Background)	1980	6 (0.83)	5.0	Medium
Wet						

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	3 (0.67)	0.2	High
N/R = Not reported						

862

## 1.25 Soil


863

### 1.25.1 Soil (ng/g) – Dry Fraction

864

865 Measured concentrations of TCEP in Soil with unit of ng/g, extracted from three sources, are  
866 summarized in Figure 1-33 and supplemental information is provided in Table 1-36. Overall,  
867 concentrations ranged from not detected to 23.48 ng/g from 18 samples collected between 2010 and  
868 2014 in two countries, DE and TR. Location types were categorized as General Population  
869 (Background). Reported detection frequency was not reported.

869



870

871

**Figure 1-36. Concentrations of TCEP (ng/g) in the Dry Fraction of Soil in General Population (Background) Locations from 2010 to 2014**

872

873

874

**Table 1-36. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Dry Fraction of Soil**

875

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Kurt-Karakus et al. (2018)</a>	TR	General Population (Background)	2014	8 (N/R)	3.4	High
<a href="#">Mihajlović et al. (2011)</a>	DE	General Population (Background)	2011	6 (N/R)	0.2	Medium
<a href="#">Mihajlovic and Fries (2012)</a>	DE	General Population (Background)	2010–2011	4 (N/R)	0.2	High
N/R = Not reported						

## 1.26 Surface Water

877

### 1.26.1 Surface Water (ng/L) – Not Specified Fraction

878

Measured concentrations of TCEP in Surface Water with unit of ng/L, extracted from 29 sources, are summarized in Figure 1-34 and supplemental information is provided in Table 1-37. Overall, concentrations ranged from not detected to 2,019.0 ng/L from 3,283 samples collected between 1980 and 2017 in 14 countries, AQ, AU, CA, DE, DK, ES, FR, GB, GL, JP, KR, PT, SE and US. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection frequency ranged from 0.0 to 1.0.

879


880

881

882


883

884



885

886  
887  
888  
889  
890  
891



**Figure 1-37. Concentrations of TCEP (ng/L) in the Not Specified Fraction of Surface Water from 1980 to 2017**

**Table 1-37. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Not Specified Fraction of Surface Water**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Maruya et al. (2016)</a>	US	General Population (Background)	2013	17 (0.65)	5.0	High
<a href="#">Sengupta et al. (2014)</a>	US	General Population (Background)	2011	30 (1.00)	N/R	Medium
<a href="#">Padhye et al. (2014)</a>	US	General Population (Background)	2009–2010	8 (N/R)	N/R	Medium
<a href="#">Giorgino et al. (2007)</a>	US	General Population (Background)	2002–2005	14 (0.36)	500.0	High
<a href="#">Kolpin et al. (2002)</a>	US	Near Facility (Highly Exposed)	1999–2000	85 (0.58)	40.0	High
<a href="#">Scott et al. (1996)</a>	CA, US	General Population (Background)	1994	43 (1.00)	N/R	Low
<a href="#">Gadelha et al. (2019)</a>	PT	Near Facility (Highly Exposed)	2016–2017	12 (N/R)	0.13	High
<a href="#">Gao et al. (2019)</a>	SE	General Population (Background)	2016–2017	8 (0.25)	7.2	High
<a href="#">McDonough et al. (2018)</a>	CA, GL	Remote (Not Near Source)	2014–2016	13 (0.46)	0.22	High
<a href="#">Blum et al. (2018a)</a>	SE	General Population (Background)	2014–2015	16 (0.88)	0.15	High



Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Blum et al. (2018b)</a>	SE	Near Facility (Highly Exposed)	2014–2015	20 (0.60)	N/R	High
<a href="#">Choo et al. (2018)</a>	KR	General Population (Background)	2015	4 (1.00)	0.24	High
<a href="#">Li et al. (2017)</a>	AQ	Remote (Not Near Source)	2014	25 (0.88)	0.21	High
<a href="#">Loos et al. (2017)</a>	DE	General Population (Background)	2013	71 (1.00)	0.29	High
<a href="#">Gustavsson et al. (2018)</a>	SE	General Population (Background)	2013	28 (0.57)	0.68	High
<a href="#">Scott et al. (2014)</a>	AU	General Population (Background)	2011–2012	285 (0.44)	10.0	High
<a href="#">Cristale et al. (2013)</a>	GB	General Population (Background)	2011	13 (1.00)	2.4	Medium
<a href="#">Matamoros et al. (2012)</a>	DK	Near Facility (Highly Exposed)	2010	29 (1.00)	N/R	High
<a href="#">Regnery and Püttmann (2010b)</a>	DE	General Population (Background)	2008–2009	52 (1.00)	2.0	High
<a href="#">Calderón-Preciado et al. (2011)</a>	ES	General Population (Background)	2008–2009	8 (0.00)	55.0	Medium
<a href="#">Regnery and Püttmann (2010a)</a>	DE	General Population (Background)	2007–2009	151 (N/R)	1.0	High
<a href="#">Gourmelon et al. (2010)</a>	FR	Near Facility (Highly Exposed)	2009	20 (0.25)	40.0	Medium
<a href="#">Rodil et al. (2012)</a>	ES	General Population (Background)	2007–2008	28 (0.64)	0.004	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Quednow and Püttmann (2009)</a>	DE	General Population (Background)	2003–2006	1,650 (0.91)	5.0	High
<a href="#">Andresen et al. (2007)</a>	DE	General Population (Background)	2005	14 (N/R)	0.3	High
<a href="#">Andresen et al. (2004)</a>	DE	General Population (Background)	2002	44 (N/R)	N/R	Medium
<a href="#">Fries and Puttmann (2003)</a>	DE	General Population (Background)	2000–2001	9 (0.89)	1.0	Medium
<a href="#">Fries and Puttmann (2001)</a>	DE	General Population (Background)	2000	561 (N/R)	1.0	Medium
<a href="#">Ishikawa et al. (1985)</a>	JP	General Population (Background)	1980	9 (1.00)	10.0	Medium
<a href="#">Ishikawa et al. (1985)</a>	JP	General Population (Background)	1980	16 (0.88)	10.0	Medium

N/R = Not reported

892

## **1.27 Terrestrial Organisms – Bird**

893

### **1.27.1 Terrestrial Organisms – Bird (ng/g) – All Fractions**

894

Measured concentrations of TCEP in Terrestrial Organisms – Bird with unit of ng/g, extracted from seven sources, are summarized in Figure 1-35 and supplemental information is provided in Table 1-38. More than one weight fraction was reported and summarized separately below:

895

896

897

898

899

900

901

902

Overall, concentrations for Wet ranged from not detected to 39.0 ng/g from 160 samples collected between 2000 and 2012 in four countries, CA, NL, NO and US. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection frequency ranged from 0.0 to 1.0.

903


904

905

906

907

Overall, concentrations for Dry ranged from not detected to 3,000.0 ng/g from 40 samples collected between 2008 and 2016 in three countries, ES, NL and NO. Location types were categorized as General Population (Background), Near Facility (Highly Exposed) and Remote (Not Near Source). Reported detection frequency ranged from 0.0 to 1.0.



908

909 **Figure 1-38. Concentrations of TCEP (ng/g) in Terrestrial Organisms – Bird from 2000 to 2016**

910

911 **Table 1-38. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in**

912 **Terrestrial Organisms – Bird**


Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Wet						
<a href="#">Guo et al. (2018)</a>	US	General Population (Background)	2000–2012	24 (0.00)	N/R	High
<a href="#">Guo et al. (2018)</a>	US	General Population (Background)	2000–2012	22 (0.55)	1.74	High
<a href="#">Huber et al. (2015)</a>	NO	Remote (Not Near Source)	2012	16 (1.00)	N/R	High
<a href="#">Chen et al. (2012)</a>	CA	Remote (Not Near Source)	2010	13 (0.77)	0.1	Medium
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	16 (0.00)	0.03	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	8 (0.00)	0.03	Medium
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	24 (N/R)	0.03	Medium
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	8 (N/R)	0.03	Medium
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	16 (N/R)	0.03	Medium
<a href="#">Greaves and Letcher (2014)</a>	CA	Remote (Not Near Source)	2010	8 (0.38)	0.03	Medium
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	5 (N/R)	0.26	High
Dry						
<a href="#">Monclús et al. (2018)</a>	ES	General Population (Background)	2016	14 (0.43)	1.0	High
<a href="#">Eulaers et al. (2014)</a>	NO	Remote (Not Near Source)	2011	21 (1.00)	1.0	High
<a href="#">Brandsma et al. (2015)</a>	NL	Near Facility (Highly Exposed)	2008	5 (0.00)	0.26	High
N/R = Not reported						

913 **1.27.2 Terrestrial Organisms – Bird (ng/g) – Wet Fraction**

914 Measured concentrations of BCEP in Terrestrial Organisms – Bird with unit of ng/g, extracted from one  
915 source, are summarized in Figure 1-36 and supplemental information is provided in Table 1-39. Overall,  
916 concentrations ranged from 0.38 to 26 ng/g from 21 samples collected between 2000 and 2012 in one  
917 country, US. Location types were categorized as General Population (Background). Reported detection  
918 frequency was 1.0.  
919

920



921

922 **Figure 1-39. Concentrations of BCEP (ng/g) in the Wet Fraction of Terrestrial Organisms – Bird**  
 923 **in General Population (Background) Locations from 2000 to 2012**

924

925 **Table 1-39. Summary of Peer-Reviewed Literature that Measured BCEP (ng/g) Levels in the Wet**  
 926 **Fraction of Terrestrial Organisms – Bird**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Stubbings et al. (2018)</a>	US	General Population (Background)	2000–2012	21 (1.00)	N/R	High

N/R = Not reported

927

## 1.28 Terrestrial Organisms – Mammal

928

### 1.28.1 Terrestrial Organisms – Mammal (ng/g) – All Fractions

929

929 Measured concentrations of TCEP in Terrestrial Organisms – Mammal with unit of ng/g, extracted from  
 930 two sources, are summarized in Figure 1-37 and supplemental information is provided in Table 1-40.  
 931 More than one weight fraction was reported and summarized separately below:


932

933 Overall, concentrations for Lipid ranged from 1.91 to 52.5 ng/g from 20 samples collected between 2008  
 934 and 2010 in one country, NO. Location types were categorized as Remote (Not Near Source). Reported  
 935 detection frequency was 0.1.

936

937 Overall, concentrations for Wet ranged from not detected to 0.115 ng/g from 21 samples collected  
 938 between 2017 and 2018 in one country, NO. Location types were categorized as General Population  
 939 (Background). Reported detection frequency was 0.0.

940



941

942 **Figure 1-40. Concentrations of TCEP (ng/g) in Terrestrial Organisms – Mammal from 2008 to**  
 943 **2018**

944

945  
946

**Table 1-40. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in Terrestrial Organisms – Mammal**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
Lipid						
<a href="#">Hallanger et al. (2015)</a>	NO	Remote (Not Near Source)	2008–2010	20 (0.10)	N/R	High
Wet						
<a href="#">Heimstad et al. (2019)</a>	NO	General Population (Background)	2017–2018	21 (0.00)	0.23	High
N/R = Not reported						

947

## 1.29 Terrestrial Organisms – Plant

948

### 1.29.1 Terrestrial Organisms – Plant (ng/g) – Wet Fraction

949

Measured concentrations of TCEP in Terrestrial Organisms – Plant with unit of ng/g, extracted from one source, are summarized in Figure 1-38 and supplemental information is provided in Table 1-41. Overall, concentrations ranged from 1.25 to 1950 ng/g from nine samples collected between 1993 and 1994 in one country, US. Location types were categorized as Remote (Not Near Source). Reported detection frequency was 0.67.


950

951

952

953

954



955

956

**Figure 1-41. Concentrations of TCEP (ng/g) in the Wet Fraction of Terrestrial Organisms – Plant in Remote (Not Near Source) Locations from 1993 to 1994**

957

958

959

960

**Table 1-41. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet Fraction of Terrestrial Organisms – Plant**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Aston et al. (1996)</a>	US	Remote (Not Near Source)	1993–1994	9 (0.67)	2.5	Medium

961

## 1.30 Wastewater

962

### 1.30.1 Wastewater (ng/g) – Wet Fraction


963

Measured concentrations of TCEP in Wastewater with unit of ng/g, extracted from three sources, are summarized in Figure 1-39 and supplemental information is provided in Table 1-42. Overall, concentrations ranged from 0.5 to 198.0 ng/g from 74 samples collected between 2013 and 2018 in three

964

965

966 countries, CA, NO and US. Location types were categorized as Raw Influent and Treated Effluent.  
 967 Reported detection frequency ranged from 0.5 to 1.0.  
 968



969  
 970 **Figure 1-42. Concentrations of TCEP (ng/g) in the Wet Fraction of Wastewater from 2013 to 2018**  
 971


972 **Table 1-42. Summary of Peer-Reviewed Literature that Measured TCEP (ng/g) Levels in the Wet**  
 973 **Fraction of Wastewater**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/g)	Overall Quality Level
<a href="#">Kim et al. (2017)</a>	US	Raw Influent	2013–2015	16 (1.00)	1.0	High
<a href="#">Kim et al. (2017)</a>	US	Treated Effluent	2013–2015	38 (0.50)	1.0	High
<a href="#">Kim et al. (2017)</a>	US	Treated Effluent	2013–2015	16 (1.00)	1.0	High
<a href="#">Norwegian Environment (2019a)</a>	NO	Treated Effluent	2018	2 (N/R)	N/R	Medium
<a href="#">Woudneh et al. (2015)</a>	CA	Raw Influent	2015	1 (1.00)	0.1	Medium
<a href="#">Woudneh et al. (2015)</a>	CA	Treated Effluent	2015	1 (1.00)	0.1	Medium

N/R = Not reported

974 **1.30.2 Wastewater (ng/L) – Wet Fraction**

975 Measured concentrations of TCEP in Wastewater with unit of ng/L, extracted from 16 sources, are  
 976 summarized in Figure 1-40 and supplemental information is provided in Table 1-43. Overall,  
 977 concentrations ranged from not detected to 42800.0 ng/L from 305 samples collected between 2001 and  
 978 2018 in eight countries, AU, BE, DE, ES, FR, NO, SE and US. Location types were categorized as  
 979 Untreated Combined Sewer Overflow, Raw Influent, Treated Effluent and Untreated Effluent at  
 980 Discharge Origin. Reported detection frequency ranged from 0.0 to 1.0.  
 981



982

983 **Figure 1-43. Concentrations of TCEP (ng/L) in the Wet Fraction of Wastewater from 2001 to 2018**

984

985 **Table 1-43. Summary of Peer-Reviewed Literature that Measured TCEP (ng/L) Levels in the Wet**  
 986 **Fraction of Wastewater**

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Kim et al. (2017)</a>	US	Raw Influent	2013–2015	16 (1.00)	50.0	High



Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">Kim et al. (2017)</a>	US	Treated Effluent	2013–2015	16 (1.00)	50.0	High
<a href="#">Schreder and La Guardia (2014)</a>	US	Treated Effluent	2011–2012	2 (1.00)	1.0	High
<a href="#">Schreder and La Guardia (2014)</a>	US	Untreated Effluent at Discharge Origin	2011–2012	21 (1.00)	1.0	High
<a href="#">Laws et al. (2011)</a>	US	Treated Effluent	2009	1 (1.00)	200.0	Medium
<a href="#">Jackson and Sutton (2008)</a>	US	Raw Influent	2006	10 (0.20)	6250.0	Medium
<a href="#">Jackson and Sutton (2008)</a>	US	Treated Effluent	2006	3 (0.67)	N/R	Medium
<a href="#">Loraine and Pettigrov (2006)</a>	US	Treated Effluent	2001–2002	6 (0.50)	760.0	Medium
<a href="#">Norwegian Environment (2019a)</a>	NO	Treated Effluent	2018	2 (N/R)	N/R	Medium
<a href="#">Gao et al. (2019)</a>	SE	Raw Influent	2017	4 (1.00)	7.2	High
<a href="#">Gao et al. (2019)</a>	SE	Treated Effluent	2016–2017	8 (0.88)	7.2	High
<a href="#">Been et al. (2017)</a>	BE	Raw Influent	2015–2016	8 (1.00)	1.1	Medium
<a href="#">Launay et al. (2016)</a>	DE	Untreated Combined Sewer Overflow	2014	9 (N/R)	50.0	High
<a href="#">Launay et al. (2016)</a>	DE	Untreated Effluent at Discharge Origin	2014	7 (N/R)	50.0	High
<a href="#">Blum et al. (2017)</a>	SE	Treated Effluent	2013	10 (0.80)	N/R	Medium

Citation	Country	Location Type	Sampling Year	Sample Size (Frequency of Detection)	Detection Limit (ng/L)	Overall Quality Level
<a href="#">O'Brien et al. (2015)</a>	AU	Raw Influent	2011	15 (0.93)	200.0	High
<a href="#">Gourmelon et al. (2010)</a>	FR	Treated Effluent	2009	14 (1.00)	40.0	Medium
<a href="#">Rodil et al. (2012)</a>	ES	Raw Influent	2007–2008	11 (1.00)	10.0	Medium
<a href="#">Rodil et al. (2012)</a>	ES	Treated Effluent	2007–2008	11 (1.00)	10.0	Medium
<a href="#">Meyer and Bester (2004)</a>	DE	Raw Influent	2003	0 (N/R)	6.1	Medium
<a href="#">Meyer and Bester (2004)</a>	DE	Treated Effluent	2003	18 (0.00)	6.1	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Raw Influent	2002–2003	18 (N/R)	N/R	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Treated Effluent	2002–2003	17 (N/R)	N/R	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Raw Influent	2002–2003	9 (N/R)	N/R	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Treated Effluent	2002–2003	34 (N/R)	N/R	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Treated Effluent	2002–2003	18 (N/R)	N/R	Medium
<a href="#">Marklund et al. (2005a)</a>	SE	Treated Effluent	2002–2003	9 (N/R)	N/R	Medium
<a href="#">Fries and Puttmann (2003)</a>	DE	Raw Influent	2001	4 (1.00)	1.0	Medium
<a href="#">Fries and Puttmann (2003)</a>	DE	Treated Effluent	2001	4 (0.75)	1.0	Medium

N/R = Not reported

**2.1 Data Integration Methods and Approach**

---

989 Extracted study data required further processing to allow for the standardization and integration of  
990 TCEP data across all studies. Where studies reported data values for metabolites of TCEP, including  
991 BCEP (bis(2-chloroethyl) phosphate (CASRN 4050-56-0), these values were extracted separately in  
992 DistillerSR and data summaries are reported separately in this report for TCEP and its individual  
993 metabolites.  
994

995  
996 To enable comparison of data across studies, all extracted environmental monitoring and biomonitoring  
997 concentrations were converted to common unit by medium (*i.e.*, ng/L for aqueous media, ng/g for solid  
998 phase media, ng/m<sup>3</sup> for air media). Study-reported summary statistics were used, as available, to  
999 characterize the concentrations for all unique scenarios including minimums and maximum  
1000 concentrations, measures of central tendency, percentiles, measures of variance, frequencies of  
1001 detection, and reported limits of detection (LOD) and/or limits of quantitation (LOQ). In cases where  
1002 point data were available, summary statistics were calculated for each unique scenario depending on the  
1003 number of point values. If only one point value was reported per unique scenario, it was treated as an  
1004 arithmetic mean. For unique scenarios with 2–9 point values, arithmetic means, medians, standard  
1005 deviations, and minimum and maximums were calculated. For unique scenarios with 10 or more point  
1006 values, the 25th, 50th, and 90th percentiles also were calculated.  
1007

1008 A left-censoring protocol was applied to impute the lower bound of concentration ranges in cases where  
1009 the reported frequency of detection (FOD) was less than 100 percent, meaning that TCEP, or metabolite,  
1010 was not detected in at least one sample. Specifically, a value of one-half the highest reported LOD or  
1011 LOQ (if no LOD available) was imputed as the minimum value for each unique scenario. In cases where  
1012 authors reported values as “not detected” (*e.g.*, “ND”, “< LOD”, “BL0D”) without providing a value,  
1013 the same left-censoring protocol was applied. In the case where values were reported with an indicator  
1014 that the values were estimated (*e.g.*, typically above LOD and below LOQ), those values were used in  
1015 the data aggregation directly. Where no LOD or LOQ were provided, no substitution was possible. If the  
1016 FOD was zero, and no limits were reported, the study aggregate was dropped from consideration. Other  
1017 issues in study reported detection limits included when a range of detection limits were reported across  
1018 all chemicals in the analytical method. These limits were dropped since no concentration could be  
1019 attributed to the TSCA chemical specifically.  
1020

1021 Data were first aggregated by like media (*e.g.*, surface water, ambient air) and then generally by unit and  
1022 sampling phase (*e.g.* particulate or vapor phases in air) or weight fraction type (*e.g.* wet versus dry  
1023 weights). Media-specific aggregations were employed as appropriate (*e.g.*, microenvironments for  
1024 inhalation of indoor air, taxa and tissue type for terrestrial and aquatic organisms), and further  
1025 aggregation was performed to group data by pollution source receptor type (*i.e.*, General Population  
1026 (Background), Near Facility (Highly Exposed), Remote (Not Near Point Source)).  
1027

1028 All data aggregation, unit conversion, range and central tendency standardization, and estimation of  
1029 derived exposures were performed computationally with a workflow, data management system, and  
1030 computational pipeline developed specifically to support EPA risk evaluations. All data and statistical  
1031 analyses were performed on DistillerSR reports of quality control reviewed data. The data computational  
1032 pipeline was prepared using scripts in Python 3.9 using the pandas, scipy and xlrd libraries and  
1033 visualized with services developed in NodeJS and D3.  
1034

1035 Section 1 of this supplement provides a data summary plot for each media by unit. Each plot presents  
 1036 summary statistics for each study aggregated by pollution source receptor type and setting or  
 1037 microenvironment (*i.e.*, General Population (Background), Near Facility (Highly Exposed), Remote  
 1038 (Not Near Point Source)). Because individual studies often present multiple unique scenarios that can be  
 1039 grouped into a single representative aggregate for the study, available statistics were combined and the  
 1040 ranges observations (*e.g.*, minimum, maximum, and percentiles) and central tendencies (*e.g.* arithmetic  
 1041 mean, geometric mean, and median), and overall FOD where possible were calculated.

1043 Within each plot, data are separated by unit basis of sampling fraction, then monitoring data from the  
 1044 U.S. are presented first, followed by studies with data from mixed locations (*i.e.*, U.S. and other  
 1045 countries), finally by studies with data from non-U.S. sources. For each grouping, data are presented  
 1046 from newest to oldest, based on latest year of sampling. Differentiation by tissue type for ecological  
 1047 monitoring media is indicated in the tick label. The lighter region of each bar represents the overall  
 1048 range of data and the darker region represents the range of central tendency reported in each study.  
 1049 Triangles indicate the arithmetic mean and 90th percentile estimates are plotted over the bars for study  
 1050 aggregates that reported enough statistical results to reconstruct a lognormal or normal distribution. The  
 1051 statistical methods used to calculate the central and high-end estimates are described in the following  
 1052 section. The tables that follow each plot provide summary information for each study aggregate such as  
 1053 the sampling location and dates, sample size and FOD, maximum LOD or LOQ (if no LOD was  
 1054 reported), and overall study quality judgement from data evaluation.

## 1055 **2.2 Statistical Approach of Exposure Estimates Derived from Measured** 1056 **Concentrations**

1057 Following the aggregation and standardization of reported study data from DistillerSR, the statistical  
 1058 methods described were applied to enhance the comparability and informative value of the available  
 1059 information. All statistical calculations were performed with Python scripts included as steps within the  
 1060 computational pipeline of the methodology.

### 1061 **2.2.1 Aggregation of Statistical Estimates**

1062 Studies were aggregated as described in the previous section. Based on this aggregation and study-  
 1063 reported statistics, normal and lognormal distributions were estimated based on available data. In cases  
 1064 where more than one statistic type (*i.e.*, mean, median, minimum, maximum, percentile, and variability  
 1065 measures) each type was handled as described in Table 2-1 below.

1066 **Table 2-1. Statistics and Methods for Data Aggregation**

Statistic Type	Description of Calculation Method for Aggregate Estimate
Arithmetic means	$\sum_{j=1}^K w_j \bar{x}_j$ , where $\bar{x}_j = \sum_{i=1}^{N_j} x_{j,i}$
Medians	$\sum_{j=1}^K w_j \cdot med_j$ , where $med_j$ is the median of dataset $J$
Percentiles	$\sum_{j=1}^K w_j \cdot perc_j$ , where $perc_j$ is the percentile of dataset $J$
Minimums	$\min\{m_1, \dots, m_K\}$ , where $m_j = \min\{x_{j,1}, \dots, x_{j,N_j}\}$
Maximums	$\max\{M_1, \dots, M_n\}$ , where $M_j = \max\{x_{j,1}, \dots, x_{j,N_j}\}$
Geometric means	$\exp\left(\sum_{j=1}^K w_j \cdot \ln(GM_j)\right)$ , where $GM_j = \exp\left(\frac{1}{n} \sum_{i=1}^{N_j} \ln(x_{j,i})\right)$
Geometric standard deviations	$\exp\left(\sqrt{\frac{1}{K-1} \left(\sum_{j=1}^K \ln(GSD_j)\right)}\right)$ , where $GSD_j = \exp\left(\sqrt{\sum_{i=1}^{N_j} \left(\ln\left(\frac{x_{j,i}}{GM_j}\right)\right)^2 / N_j}\right)$

Variances	$\frac{1}{K-1} \sum_{j=1}^K V_j$ , where $V_j = \frac{1}{N_j-1} \sum_{i=1}^{N_j} (x_{j,i} - \bar{x}_j)^2$
Standard deviations	$\sqrt{\frac{1}{K-1} \sum_{j=1}^K \sigma_j^2}$ , where $\sigma_j = \sqrt{\frac{1}{N_j-1} \sum_{i=1}^{N_j} (x_{j,i} - \bar{x}_j)^2}$

1068

1069

In cases where measures of variability were provided, no fitting was required to build a distribution. If geometric means and geometric standard deviations (GSDs) were provided they were used directly to construct a lognormal distribution by using the mean of geometric means ( $\exp(\mu)$ ) and the sample weighted mean of GSD ( $\sigma$ ). Using this distribution, the central tendency was estimated by calculating the arithmetic mean and 90th percentile using the equations below.

1070

1071

1072

1073

1074

- Equation for arithmetic mean estimates from lognormal distribution:  $e^{(\mu + \frac{\sigma^2}{2})}$

1075

1076

- Equation for estimating 90th percentile from lognormal distribution:  $e^{(\mu + \sigma * 1.282)}$

1077

1078

1079

1080

1081

1082

If arithmetic means and standard deviations (SDs) or variance were provided and no other statistics indicate that the data are not normally distributed, then a normal distribution was derived using the available statistics. If arithmetic means, medians, and SDs were provided and means and medians were within 5 percent relative percent difference, then a normal distribution was assumed and derived using the provided arithmetic mean and measure of variation. When a normal distribution was assumed the arithmetic mean (assumed to be median) and 90th percentile was calculated using the equations below.

1083

- Equation for arithmetic mean for normal distribution:  $\mu$

1084

1085

- Equation for 90th percentile from normal distribution:  $\mu + 1.282\sigma$

1086

1087

1088

1089

If a variation was not provided or a normal distribution was not assumed, Table 2-2 describes the preferred distributions used based on the available statistics in the study aggregate. In some cases, the preferred distribution was not used, see the Quality Control section (Section 2.2.4) for this justification.

1090

**Table 2-2. Distributions Preferred Depending on Available Reported Statistics**

Case Type	Description of Available Statistics Per Study Aggregate	Distribution Type Preferred
Case 0A	Geometric mean and GSD	Lognormal
Case 0B	Median and GSD	Lognormal
Case 1A	(Mean == Median) and SD	Normal
Case 1B	Mean and SD (no Median provided)	Normal
Case 2A	Median and (min or max or percentile)	Lognormal
Case 2B	Median and (FOD < 1 and LOD/LOQ)	Lognormal
Case 3A	Mean only and (min or max or percentile)	Lognormal
Case 3B	Mean only and (FOD < 1 and LOD/LOQ)	Lognormal
Case 4	Median and mean only	Lognormal
All other cases	Not enough data to build distribution	N/A

GSD = geometric standard deviation; SD = standard deviation; FOD = frequency of detection; LOD = limit of detection; LOQ = limit of quantitation

1091 **2.2.2 Fitting Lognormal Distributions**

---

1092 In cases where the study data provided median values, the average median was substituted for geometric  
1093 mean, and the remaining statistics were used to estimate the GSD by minimizing the sum of squared  
1094 errors for all provided statistical estimates. Sum of squared errors was calculated by comparing the mean  
1095 of the residual statistic to the estimated value produced by the fitted distribution, based on the  
1096 assumptions in Table 2-3 that defined the percentiles assumed for each statistic type.  
1097

1098 **Table 2-3. Assumed Percentile for Calculating Error by Statistical Estimate Type**

Mean of Statistical Estimate by Type	Assumed Percentile for Calculating Error
Maximum	0.99
Minimum	0.01
nth percentile ( <i>e.g.</i> , 25th percentile)	n/100 ( <i>e.g.</i> , 0.25)
Half limit of quantitation substituted minimum	0.005
Half limit of detection substituted minimum	0.0025

1099 This methodology requires a central tendency estimate and at least one data point on the distribution in  
1100 order to fit a lognormal distribution. Thus, lognormal distributions were fitted for studies that provided  
1101 an arithmetic mean and at least one data point on the curve. In these cases, both the geometric mean and  
1102 the GSD were derived by minimizing the sum of the squared errors for all estimates.  
1103

1104 **2.2.3 Fitting Normal Distributions**

---

1105 Normal distributions also were constructed for all study aggregates using an approach similar to the  
1106 approach for geometric distributions described in Section 2.2.1. Study-reported means were assumed to  
1107 be medians, and standard deviations were calculated by minimizing the sum of squares error of all  
1108 available estimates.

1109 **2.2.4 Quality Control of Derived Exposure Estimates**

---

1110 As a quality control measure, the estimated medians and arithmetic means were evaluated to verify that  
1111 the estimated values fell within the range of the reported data. Estimates were not used if they fell  
1112 outside of the range of the reported data, typically an indicator of anomalous data. In addition, derived  
1113 GSDs were not used if they exceeded 10 for the lognormal distributions, mean estimates were not used  
1114 if they exceeded 100 percent relative percent difference from residual means. In these cases, the  
1115 estimates from the normal distributions were used when normal distributions could be derived.

1116 **2.2.5 Final Exposure Estimates by Media and Pollution Source Receptor Type**

---

1117 Central tendency exposure values that carried forward to risk evaluation after passing the QC process  
1118 were summarized for each media aggregate by taking the sample weighted mean of the arithmetic mean  
1119 estimates from the selected distribution (*i.e.*, lognormal or normal). Similarly, the 90th percentile  
1120 estimates carried forward to risk evaluation were calculated as the sample weighted mean of 90th  
1121 percentile estimates.

## 1122 REFERENCES

---

- 1123 [Abdollahi, A; Eng, A; Jantunen, LM; Ahrens, L; Shoeib, M; Parnis, JM; Harner, T.](#) (2017).  
1124 Characterization of polyurethane foam (PUF) and sorbent impregnated PUF (SIP) disk passive  
1125 air samplers for measuring organophosphate flame retardants. *Chemosphere* 167: 212-219.  
1126 <http://dx.doi.org/10.1016/j.chemosphere.2016.09.111>
- 1127 [Ait Bamai, Y; Araki, A; Nomura, T; Kawai, T; Tsuboi, T; Kobayashi, S; Miyashita, C; Takeda, M;](#)  
1128 [Shimizu, H; Kishi, R.](#) (2018). Association of filaggrin gene mutations and childhood eczema and  
1129 wheeze with phthalates and phosphorus flame retardants in house dust: The Hokkaido study on  
1130 Environment and Children's Health. *Environ Int* 121: 102-110.  
1131 <http://dx.doi.org/10.1016/j.envint.2018.08.046>
- 1132 [Ali, N; Dirtu, AC; van Den Eede, N; Goosey, E; Harrad, S; Neels, H; T Mannetje, A; Coakley, J;](#)  
1133 [Douwes, J; Covaci, A.](#) (2012). Occurrence of alternative flame retardants in indoor dust from  
1134 New Zealand: Indoor sources and human exposure assessment. *Chemosphere* 88: 1276-1282.  
1135 <http://dx.doi.org/10.1016/j.chemosphere.2012.03.100>
- 1136 [Andresen, JA; Grundmann, A; Bester, K.](#) (2004). Organophosphorus flame retardants and plasticisers in  
1137 surface waters. *Sci Total Environ* 332: 155-166.  
1138 <http://dx.doi.org/10.1016/j.scitotenv.2004.04.021>
- 1139 [Andresen, JA; Muir, D; Ueno, D; Darling, C; Theobald, N; Bester, K.](#) (2007). Emerging pollutants in the  
1140 North Sea in comparison to Lake Ontario, Canada, data. *Environ Toxicol Chem* 26: 1081-1089.  
1141 <http://dx.doi.org/10.1897/06-416R.1>
- 1142 [Aston, LS; Noda, J; Seiber, JN; Reece, CA.](#) (1996). Organophosphate flame retardants in needles of  
1143 *Pinus ponderosa* in the Sierra Nevada foothills. *Bull Environ Contam Toxicol* 57: 859-866.  
1144 <http://dx.doi.org/10.1007/s001289900269>
- 1145 [Barnes, KK; Christenson, SC; Kolpin, DW; Focazio, M; Furlong, ET; Zaugg, SD; Meyer, MT; Barber,](#)  
1146 [LB.](#) (2004). Pharmaceuticals and other organic waste water contaminants within a leachate  
1147 plume downgradient of a municipal landfill. *Ground Water Monit Remediat* 24: 119-126.  
1148 <http://dx.doi.org/10.1111/j.1745-6592.2004.tb00720.x>
- 1149 [Barnes, KK; Kolpin, DW; Furlong, ET; Zaugg, SD; Meyer, MT; Barber, LB.](#) (2008). A national  
1150 reconnaissance of pharmaceuticals and other organic wastewater contaminants in the United  
1151 States--I) groundwater. *Sci Total Environ* 402: 192-200.  
1152 <http://dx.doi.org/10.1016/j.scitotenv.2008.04.028>
- 1153 [Bastiaensen, M; Ait Bamai, Y; Araki, A; Van Den Eede, N; Kawai, T; Tsuboi, T; Kishi, R; Covaci, A.](#)  
1154 (2019a). Biomonitoring of organophosphate flame retardants and plasticizers in children:  
1155 Associations with house dust and housing characteristics in Japan. *Environ Res* 172: 543-551.  
1156 <http://dx.doi.org/10.1016/j.envres.2019.02.045>
- 1157 [Bastiaensen, M; Malarvannan, G; Been, F; Yin, S; Yao, Y; Huygh, J; Clotman, K; Schepens, T; Jorens,](#)  
1158 [PG; Covaci, A.](#) (2019b). Metabolites of phosphate flame retardants and alternative plasticizers in  
1159 urine from intensive care patients. *Chemosphere* 233: 590-596.  
1160 <http://dx.doi.org/10.1016/j.chemosphere.2019.05.280>
- 1161 [Been, F; Bastiaensen, M; Lai, FY; van Nuijs, ALN; Covaci, A.](#) (2017). Liquid chromatography-tandem  
1162 mass spectrometry analysis of biomarkers of exposure to phosphorus flame retardants in  
1163 wastewater to monitor community-wide exposure. *Ind Eng Chem Anal Ed* 89: 10045–10053.  
1164 <http://dx.doi.org/10.1021/acs.analchem.7b02705>
- 1165 [Bergh, C; Aberg, KM; Svartengren, M; Emenius, G; Oestman, C.](#) (2011a). Organophosphate and  
1166 phthalate esters in indoor air: a comparison between multi-storey buildings with high and low  
1167 prevalence of sick building symptoms. *J Environ Monit* 13: 2001-2009.  
1168 <http://dx.doi.org/10.1039/c1em10152h>



1169 [Bergh, C; Torgrip, R; Emenius, G; Ostman, C.](#) (2011b). Organophosphate and phthalate esters in air and  
1170 settled dust - a multi-location indoor study. *Indoor Air* 21: 67-76.  
1171 <http://dx.doi.org/10.1111/j.1600-0668.2010.00684.x>  
1172 [Blum, KM; Andersson, PL; Ahrens, L; Wiberg, K; Haglund, P.](#) (2018a). Persistence, mobility and  
1173 bioavailability of emerging organic contaminants discharged from sewage treatment plants. *Sci*  
1174 *Total Environ* 612: 1532-1542. <http://dx.doi.org/10.1016/j.scitotenv.2017.09.006>  
1175 [Blum, KM; Andersson, PL; Renman, G; Ahrens, L; Gros, M; Wiberg, K; Haglund, P.](#) (2017). Non-  
1176 target screening and prioritization of potentially persistent, bioaccumulating and toxic domestic  
1177 wastewater contaminants and their removal in on-site and large-scale sewage treatment plants.  
1178 *Sci Total Environ* 575: 265-275. <http://dx.doi.org/10.1016/j.scitotenv.2016.09.135>  
1179 [Blum, KM; Haglund, P; Gao, Q; Ahrens, L; Gros, M; Wiberg, K; Andersson, PL.](#) (2018b). Mass fluxes  
1180 per capita of organic contaminants from on-site sewage treatment facilities. *Chemosphere* 201:  
1181 864-873. <http://dx.doi.org/10.1016/j.chemosphere.2018.03.058>  
1182 [Bohlin-Nizzetto, P; Aas, W; Nikiforov, V.](#) (2019). Monitoring of Environmental Contaminants in Air  
1183 and Precipitation, 2018. (Report M-1419). Norwegian Environment Agency.  
1184 <https://www.miljodirektoratet.no/globalassets/publikasjoner/m1419/m1419.pdf>  
1185 [Bradman, A; Castorina, R; Gaspar, F; Nishioka, M; Colón, M; Weathers, W; Egeghy, PP; Maddalena,](#)  
1186 [R; Williams, J; Jenkins, PL; McKone, TE.](#) (2014). Flame retardant exposures in California early  
1187 childhood education environments. *Chemosphere* 116: 61-66.  
1188 <http://dx.doi.org/10.1016/j.chemosphere.2014.02.072>  
1189 [Brandsma, SH; de Boer, J; van Velzen, MJ; Leonards, PE.](#) (2014). Organophosphorus flame retardants  
1190 (PFRs) and plasticizers in house and car dust and the influence of electronic equipment.  
1191 *Chemosphere* 116: 3-9. <http://dx.doi.org/10.1016/j.chemosphere.2014.02.036>  
1192 [Brandsma, SH; Leonards, P; Leslie, HA; de Boer, J.](#) (2015). Tracing organophosphorus and brominated  
1193 flame retardants and plasticizers in an estuarine food web. *Sci Total Environ* 505: 22-31.  
1194 <http://dx.doi.org/10.1016/j.scitotenv.2014.08.072>  
1195 [Brommer, S; Harrad, S; Van den Eede, N; Covaci, A.](#) (2012). Concentrations of organophosphate esters  
1196 and brominated flame retardants in German indoor dust samples. *J Environ Monit* 14: 2482-  
1197 2487. <http://dx.doi.org/10.1039/c2em30303e>  
1198 [Buszka, PM; Yeskis, DJ; Kolpin, DW; Furlong, ET; Zaugg, SD; Meyer, MT.](#) (2009). Waste-indicator  
1199 and pharmaceutical compounds in landfill-leachate-affected ground water near Elkhart, Indiana,  
1200 2000-2002. *Bull Environ Contam Toxicol* 82: 653-659. [http://dx.doi.org/10.1007/s00128-009-](http://dx.doi.org/10.1007/s00128-009-9702-z)  
1201 [9702-z](#)  
1202 [Calderón-Preciado, D; Matamoros, V; Bayona, JM.](#) (2011). Occurrence and potential crop uptake of  
1203 emerging contaminants and related compounds in an agricultural irrigation network. *Sci Total*  
1204 *Environ* 412-413: 14-19. <http://dx.doi.org/10.1016/j.scitotenv.2011.09.057>  
1205 [Castorina, R; Butt, C; Stapleton, HM; Avery, D; Harley, KG; Holland, N; Eskenazi, B; Bradman, A.](#)  
1206 (2017). Flame retardants and their metabolites in the homes and urine of pregnant women  
1207 residing in California (the CHAMACOS cohort). *Chemosphere* 179: 159-166.  
1208 <http://dx.doi.org/10.1016/j.chemosphere.2017.03.076>  
1209 [Chen, D; Letcher, RJ; Chu, S.](#) (2012). Determination of non-halogenated, chlorinated and brominated  
1210 organophosphate flame retardants in herring gull eggs based on liquid chromatography-tandem  
1211 quadrupole mass spectrometry. *J Chromatogr A* 1220: 169-174.  
1212 <http://dx.doi.org/10.1016/j.chroma.2011.11.046>  
1213 [Chokwe, TB; Okonkwo, JO.](#) (2019). Occurrence, distribution and ecological risk assessment of  
1214 organophosphorus flame retardants and plasticizers in sediment samples along the Vaal River  
1215 catchment, South Africa. *Emerging Contaminants* 5: 173-178.  
1216 <http://dx.doi.org/10.1016/j.emcon.2019.05.003>



1217 [Choo, G; Cho, HS; Park, K; Lee, JW; Kim, P; Oh, JE.](#) (2018). Tissue-specific distribution and  
1218 bioaccumulation potential of organophosphate flame retardants in crucian carp. *Environ Pollut*  
1219 239: 161-168. <http://dx.doi.org/10.1016/j.envpol.2018.03.104>  
1220 [Christia, C; Poma, G; Besis, A; Samara, C; Covaci, A.](#) (2018). Legacy and emerging organophosphorus  
1221 flame retardants in car dust from Greece: Implications for human exposure. *Chemosphere* 196:  
1222 231-239. <http://dx.doi.org/10.1016/j.chemosphere.2017.12.132>  
1223 [Clark, AE; Yoon, S; Sheesley, RJ; Usenko, S.](#) (2017). Spatial and Temporal Distributions of  
1224 Organophosphate Ester Concentrations from Atmospheric Particulate Matter Samples Collected  
1225 across Houston, TX. *Environ Sci Technol* 51: 4239-4247.  
1226 <http://dx.doi.org/10.1021/acs.est.7b00115>  
1227 [Coelho, SD; Sousa, AC; Isobe, T; Kim, JW; Kunisue, T; Nogueira, AJ; Tanabe, S.](#) (2016). Brominated,  
1228 chlorinated and phosphate organic contaminants in house dust from Portugal. *Sci Total Environ*  
1229 569-570: 442-449. <http://dx.doi.org/10.1016/j.scitotenv.2016.06.137>  
1230 [Cristale, J; Katsoyiannis, A; Sweetman, AJ; Jones, KC; Lacorte, S.](#) (2013). Occurrence and risk  
1231 assessment of organophosphorus and brominated flame retardants in the River Aire (UK).  
1232 *Environ Pollut* 179: 194-200. <http://dx.doi.org/10.1016/j.envpol.2013.04.001>  
1233 [Deng, WJ; Li, N; Wu, R; Richard, WKS; Wang, Z; Ho, W.](#) (2018). Phosphorus flame retardants and  
1234 Bisphenol A in indoor dust and PM2.5 in kindergartens and primary schools in Hong Kong.  
1235 *Environ Pollut* 235: 365-371. <http://dx.doi.org/10.1016/j.envpol.2017.12.093>  
1236 [Dodson, RE; Bessonneau, V; Udesky, JO; Nishioka, M; McCauley, M; Rudel, RA.](#) (2019). Passive  
1237 indoor air sampling for consumer product chemicals: A field evaluation study. *J Expo Sci*  
1238 *Environ Epidemiol* 29: 95-108. <http://dx.doi.org/10.1038/s41370-018-0070-9>  
1239 [Dodson, RE; Udesky, JO; Colton, MD; McCauley, M; Camann, DE; Yau, AY; Adamkiewicz, G; Rudel,](#)  
1240 [RA.](#) (2017). Chemical exposures in recently renovated low-income housing: Influence of  
1241 building materials and occupant activities. *Environ Int* 109: 114-127.  
1242 <http://dx.doi.org/10.1016/j.envint.2017.07.007>  
1243 [Dodson, RE; Van den Eede, N; Covaci, A; Perovich, LJ; Brody, JG; Rudel, RA.](#) (2014). Urinary  
1244 Biomonitoring of Phosphate Flame Retardants: Levels in California Adults and  
1245 Recommendations for Future Studies. *Environ Sci Technol* 48: 13625-13633.  
1246 <http://dx.doi.org/10.1021/es503445c>  
1247 [Eulaers, I; Jaspers, VL; Halley, DJ; Lepoint, G; Nygård, T; Pinxten, R; Covaci, A; Eens, M.](#) (2014).  
1248 Brominated and phosphorus flame retardants in White-tailed Eagle *Haliaeetus albicilla* nestlings:  
1249 bioaccumulation and associations with dietary proxies ( $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and  $\delta^{34}\text{S}$ ). *Sci Total Environ*  
1250 478: 48-57. <http://dx.doi.org/10.1016/j.scitotenv.2014.01.051>  
1251 [Evenset, A; Leknes, H; Christensen, GN; Warner, N; Remberger, M; Gabrielsen, GW.](#) (2009).  
1252 Screening of new contaminants in samples from the Norwegian Arctic: Silver, platinum,  
1253 sucralose, bisphenol A, tetrabromobisphenol A, siloxanes, phtalates (DEHP), phosphororganic  
1254 flame retardants. In *Akvaplan-niva rapport, no 4351-1.* (TA report no. 2510/2009; SPFO report  
1255 no. 1049/2009). Oslo, Norway: Norwegian Pollution Control Authority.  
1256 [https://brage.npolar.no/npolar-](https://brage.npolar.no/npolar-xmlui/bitstream/handle/11250/173176/ScreeningContaminantsArctic.pdf)  
1257 [xmlui/bitstream/handle/11250/173176/ScreeningContaminantsArctic.pdf](https://brage.npolar.no/npolar-xmlui/bitstream/handle/11250/173176/ScreeningContaminantsArctic.pdf)  
1258 [Fabińska, MJ; Kozielska, B; Koniecznyński, J; Bielaczyc, P.](#) (2019). Occurrence of organic phosphates  
1259 in particulate matter of the vehicle exhausts and outdoor environment - A case study. *Environ*  
1260 *Pollut* 244: 351-360. <http://dx.doi.org/10.1016/j.envpol.2018.10.060>  
1261 [Fan, X; Kubwabo, C; Rasmussen, PE; Wu, F.](#) (2014). Simultaneous determination of thirteen  
1262 organophosphate esters in settled indoor house dust and a comparison between two sampling  
1263 techniques. *Sci Total Environ* 491-492: 80-86. <http://dx.doi.org/10.1016/j.scitotenv.2013.12.127>  
1264 [Fang, M; Webster, TF; Gooden, D; Cooper, EM; McClean, MD; Carignan, C; Makey, C; Stapleton,](#)  
1265 [HM.](#) (2013). Investigating a novel flame retardant known as V6: measurements in baby products,

1266 house dust, and car dust. *Environ Sci Technol* 47: 4449-4454.  
1267 <http://dx.doi.org/10.1021/es400032v>

1268 **FDA**. (1995). Accumulated pesticide and industrial chemical findings from a ten-year study of ready-to-  
1269 eat foods. *J AOAC Int* 78: 614-630. <http://dx.doi.org/10.1093/jaoac/78.3.614>

1270 **Focazio, MJ; Kolpin, DW; Barnes, KK; Furlong, ET; Meyer, MT; Zaugg, SD; Barber, LB; Thurman,**  
1271 **ME**. (2008). A national reconnaissance for pharmaceuticals and other organic wastewater  
1272 contaminants in the United States--II) untreated drinking water sources. *Sci Total Environ* 402:  
1273 201-216. <http://dx.doi.org/10.1016/j.scitotenv.2008.02.021>

1274 **Fries, E; Puttmann, W**. (2001). Occurrence of organophosphate esters in surface water and ground water  
1275 in Germany. *J Environ Monit* 3: 621-626. <http://dx.doi.org/10.1039/b105072a>

1276 **Fries, E; Puttmann, W**. (2003). Monitoring of the three organophosphate esters TBP, TCEP and TBEP in  
1277 river water and ground water (Oder, Germany). *J Environ Monit* 5: 346-352.  
1278 <http://dx.doi.org/10.1039/b210342g>

1279 **Fromme, H; Lahrz, T; Kraft, M; Fembacher, L; Mach, C; Dietrich, S; Burkardt, R; Völkel, W; Göen, T.**  
1280 (2014). Organophosphate flame retardants and plasticizers in the air and dust in German daycare  
1281 centers and human biomonitoring in visiting children (LUPE 3). *Environ Int* 71: 158-163.  
1282 <http://dx.doi.org/10.1016/j.envint.2014.06.016>

1283 **Gadelha, JR; Rocha, AC; Camacho, C; Eljarrat, E; Peris, A; Aminot, Y; Readman, JW; Boti, V;**  
1284 **Nannou, C; Kapsi, M; Albanis, T; Rocha, F; Machado, A; Bordalo, A; Valente, LMP; Nunes,**  
1285 **ML; Marques, A; Almeida, CMR**. (2019). Persistent and emerging pollutants assessment on  
1286 aquaculture oysters (*Crassostrea gigas*) from NW Portuguese coast (Ria De Aveiro). *Sci Total*  
1287 *Environ* 666: 731-742. <http://dx.doi.org/10.1016/j.scitotenv.2019.02.280>

1288 **Gao, Q; Blum, KM; Gago-Ferrero, P; Wiberg, K; Ahrens, L; Andersson, PL**. (2019). Impact of on-site  
1289 wastewater infiltration systems on organic contaminants in groundwater and recipient waters. *Sci*  
1290 *Total Environ* 651: 1670-1679. <http://dx.doi.org/10.1016/j.scitotenv.2018.10.016>

1291 **Gibson, EA; Stapleton, HM; Calero, L; Holmes, D; Burke, K; Martinez, R; Cortes, B; Nematollahi, A;**  
1292 **Evans, D; Anderson, KA; Herbstman, JB**. (2019). Differential exposure to organophosphate  
1293 flame retardants in mother-child pairs. *Chemosphere* 219: 567-573.  
1294 <http://dx.doi.org/10.1016/j.chemosphere.2018.12.008>

1295 **Giorgino, MJ; Rasmussen, RB; Pfeifle, CM**. (2007). Occurrence of organic wastewater compounds in  
1296 selected surface-water supplies, Triangle Area of North Carolina, 2002-2005 (pp. 29). (Scientific  
1297 Investigations Report 2007-5054). U.S. Geological Survey.  
1298 <http://dx.doi.org/10.3133/sir20075054>

1299 **Giovanoulis, G; Nguyen, MA; Arwidsson, M; Langer, S; Vestergren, R; Lagerqvist, A**. (2019).  
1300 Reduction of hazardous chemicals in Swedish preschool dust through article substitution actions.  
1301 *Environ Int* 130: 104921. <http://dx.doi.org/10.1016/j.envint.2019.104921>

1302 **Gourmelon, M; Caprais, MP; Mieszkin, S; Marti, R; Wéry, N; Jardé, E; Derrien, M; Jadas-Hécart, A;**  
1303 **Communal, PY; Jaffrezic, A; Pourcher, AM**. (2010). Development of microbial and chemical  
1304 MST tools to identify the origin of the faecal pollution in bathing and shellfish harvesting waters  
1305 in France. *Water Res* 44: 4812-4824. <http://dx.doi.org/10.1016/j.watres.2010.07.061>

1306 **Greaves, AK; Letcher, RJ**. (2014). Comparative body compartment composition and in ovo transfer of  
1307 organophosphate flame retardants in North American Great Lakes herring gulls. *Environ Sci*  
1308 *Technol* 48: 7942-7950. <http://dx.doi.org/10.1021/es501334w>

1309 **Guo, J; Venier, M; Salamova, A; Hites, RA**. (2017). Bioaccumulation of Dechloranes, organophosphate  
1310 esters, and other flame retardants in Great Lakes fish. *Sci Total Environ* 583: 1-9.  
1311 <http://dx.doi.org/10.1016/j.scitotenv.2016.11.063>

1312 **Guo, JH; Simon, K; Romanak, K; Bowerman, W; Venier, M**. (2018). Accumulation of flame retardants  
1313 in paired eggs and plasma of bald eagles. *Environ Pollut* 237: 499-507.  
1314 <http://dx.doi.org/10.1016/j.envpol.2018.02.056>

1315 [Gustavsson, J; Wiberg, K; Ribeli, E; Nguyen, MA; Josefsson, S; Ahrens, L.](#) (2018). Screening of  
1316 organic flame retardants in Swedish river water. *Sci Total Environ* 625: 1046-1055.  
1317 <http://dx.doi.org/10.1016/j.scitotenv.2017.12.281>

1318 [Hallanger, IG; Sagerup, K; Evenset, A; Kovacs, KM; Leonards, P; Fuglei, E; Routti, H; Aars, J; Strøm,](#)  
1319 [H; Lydersen, C; Gabrielsen, GW.](#) (2015). Organophosphorous flame retardants in biota from  
1320 Svalbard, Norway. *Mar Pollut Bull* 101: 442-447.  
1321 <http://dx.doi.org/10.1016/j.marpolbul.2015.09.049>

1322 [Hartmann, PC; Bürgi, D; Giger, W.](#) (2004). Organophosphate flame retardants and plasticizers in indoor  
1323 air. *Chemosphere* 57: 781-787. <http://dx.doi.org/10.1016/j.chemosphere.2004.08.051>

1324 [He, C; Covaci, A; Heffernan, AL; Baduel, C; Harden, FA; Mueller, JF; Toms, LML; Nele Van Den, E;](#)  
1325 [Hobson, P; Thai, P; Wang, X; Li, Y.](#) (2018a). Urinary metabolites of organophosphate esters:  
1326 Concentrations and age trends in Australian children. *Environ Int* 111: 124-130.  
1327 <http://dx.doi.org/10.1016/j.envint.2017.11.019>

1328 [He, C; Wang, X; Tang, S; Phong, T; Li, Z; Baduel, C; Mueller, JF.](#) (2018b). Concentrations of  
1329 Organophosphate Esters and Their Specific Metabolites in Food in Southeast Queensland,  
1330 Australia: Is Dietary Exposure an Important Pathway of Organophosphate Esters and Their  
1331 Metabolites? *Environ Sci Technol* 52: 12765-12773. <http://dx.doi.org/10.1021/acs.est.8b03043>

1332 [He, C; Wang, X; Thai, P; Baduel, C; Gallen, C; Banks, A; Bainton, P; English, K; Mueller, JF.](#) (2018c).  
1333 Organophosphate and brominated flame retardants in Australian indoor environments: Levels,  
1334 sources, and preliminary assessment of human exposure. *Environ Pollut* 235: 670-679.  
1335 <http://dx.doi.org/10.1016/j.envpol.2017.12.017>

1336 [Heimstad, ES; Nygård, T; Herzke, D; Bohlin-Nizzetto, P.](#) (2019). Environmental pollutants in the  
1337 terrestrial and urban environment, 2018. (Report M-1402). Oslo, Norway: Norwegian  
1338 Environment Agency.  
1339 <https://www.miljodirektoratet.no/globalassets/publikasjoner/m1402/m1402.pdf>

1340 [Henríquez-Hernández, LA; Carretón, E; Camacho, M; Montoya-Alonso, JA; Boada, LD; Bernal Martín,](#)  
1341 [V; Falcón Cordón, Y; Falcón Cordón, S; Zumbado, M; Luzardo, OP.](#) (2017). Potential Role of  
1342 Pet Cats As a Sentinel Species for Human Exposure to Flame Retardants. *Front Vet Sci* 4: 79.  
1343 <http://dx.doi.org/10.3389/fvets.2017.00079>

1344 [Hoffman, K; Lorenzo, A; Butt, CM; Hammel, SC; Henderson, BB; Roman, SA; Scheri, RP; Stapleton,](#)  
1345 [HM; Sosa, JA.](#) (2017). Exposure to flame retardant chemicals and occurrence and severity of  
1346 papillary thyroid cancer: A case-control study. *Environ Int* 107: 235-242.  
1347 <http://dx.doi.org/10.1016/j.envint.2017.06.021>

1348 [Hopple, JA; Delzer, GC; Kingsbury, JA.](#) (2009). Anthropogenic organic compounds in source water of  
1349 selected community water systems that use groundwater, 2002-05 (pp. 76). (SIR 2009-5200).  
1350 Reston, VA: U.S. Geological Survey. <http://dx.doi.org/10.3133/sir20095200>

1351 [Huber, S; Warner, NA; Nygård, T; Remberger, M; Harju, M; Uggerud, HT; Kaj, L; Hanssen, L.](#) (2015).  
1352 A broad cocktail of environmental pollutants found in eggs of three seabird species from remote  
1353 colonies in Norway. *Environ Toxicol Chem* 34: 1296-1308. <http://dx.doi.org/10.1002/etc.2956>

1354 [Hutchins, SR; Tomson, MB; Wilson, JT; Ward, CH.](#) (1984). Fate of trace organics during rapid  
1355 infiltration of primary waste water at Fort Devens, Massachusetts (USA). *Water Res* 18: 1025-  
1356 1036. [http://dx.doi.org/10.1016/0043-1354\(84\)90255-0](http://dx.doi.org/10.1016/0043-1354(84)90255-0)

1357 [Ingerowski, G; Friedle, A; Thumulla, J.](#) (2001). Chlorinated ethyl and isopropyl phosphoric acid triesters  
1358 in the indoor environment--an inter-laboratory exposure study. *Indoor Air* 11: 145-149.  
1359 <http://dx.doi.org/10.1034/j.1600-0668.2001.011003145.x>

1360 [Ishikawa, S; Taketomi, M; Shinohara, R.](#) (1985). Determination of trialkyl phosphates and triaryl  
1361 phosphates in environmental samples. *Water Res* 19: 119-126. [http://dx.doi.org/10.1016/0043-1354\(85\)90332-X](http://dx.doi.org/10.1016/0043-1354(85)90332-X)

1362

1363 [Jackson, J; Sutton, R.](#) (2008). Sources of endocrine-disrupting chemicals in urban wastewater, Oakland,  
1364 CA. *Sci Total Environ* 405: 153-160. <http://dx.doi.org/10.1016/j.scitotenv.2008.06.033>  
1365 [Kademoglou, K; Xu, F; Padilla-Sanchez, JA; Haug, LS; Covaci, A; Collins, CD.](#) (2017). Legacy and  
1366 alternative flame retardants in Norwegian and UK indoor environment: Implications of human  
1367 exposure via dust ingestion. *Environ Int* 102: 48-56.  
1368 <http://dx.doi.org/10.1016/j.envint.2016.12.012>  
1369 [Kanazawa, A; Saito, I; Araki, A; Takeda, M; Ma, M; Saijo, Y; Kishi, R.](#) (2010). Association between  
1370 indoor exposure to semi-volatile organic compounds and building-related symptoms among the  
1371 occupants of residential dwellings. *Indoor Air* 20: 72-84. [http://dx.doi.org/10.1111/j.1600-](http://dx.doi.org/10.1111/j.1600-0668.2009.00629.x)  
1372 [0668.2009.00629.x](http://dx.doi.org/10.1111/j.1600-0668.2009.00629.x)  
1373 [Kile, ML; Scott, RP; O'Connell, SG; Lipscomb, S; Macdonald, M; McClelland, M; Anderson, KA.](#)  
1374 (2016). Using silicone wristbands to evaluate preschool children's exposure to flame retardants.  
1375 *Environ Res* 147: 365-372. <http://dx.doi.org/10.1016/j.envres.2016.02.034>  
1376 [Kim, H; Tanabe, SI.](#) (2017). Measuring Degree of Contamination by Semi-volatile Organic Compounds  
1377 (SVOC) in Interiors of Korean Homes and Kindergartens. *Journal of Asian Architecture and*  
1378 *Building Engineering* 16: 661-668. <http://dx.doi.org/10.3130/jaabe.16.661>  
1379 [Kim, J; Isobe, T; Muto, M; Nguyen Minh, T; Katsura, K; Malarvannan, G; Sudaryanto, A; Chang, KH;](#)  
1380 [Prudente, M; Pham Hung, V; Takahashi, S; Tanabe, S.](#) (2014). Organophosphorus flame  
1381 retardants (PFRs) in human breast milk from several Asian countries. *Chemosphere* 116: 91-97.  
1382 <http://dx.doi.org/10.1016/j.chemosphere.2014.02.033>  
1383 [Kim, UJ; Oh, JK; Kannan, K.](#) (2017). Occurrence, removal, and environmental emission of  
1384 organophosphate flame retardants/plasticizers in a wastewater treatment plant in New York  
1385 State. *Environ Sci Technol* 51: 7872-7880. <http://dx.doi.org/10.1021/acs.est.7b02035>  
1386 [Kingsbury, JA; Delzer, GC; Hopple, JA.](#) (2008). Anthropogenic organic compounds in source water of  
1387 nine community water systems that withdraw from streams, 2002–05 (pp. 68). (Scientific  
1388 Investigations Report 2008–5208). Reston, VA: U.S. Geological Survey.  
1389 <http://pubs.usgs.gov/sir/2008/5208/>  
1390 [Kolpin, DW; Furlong, ET; Meyer, MT; Thurman, EM; Zaugg, SD; Barber, LB; Buxton, HT.](#) (2002).  
1391 Pharmaceuticals, hormones, and other organic wastewater contaminants in US streams, 1999-  
1392 2000: A national reconnaissance. *Environ Sci Technol* 36: 1202-1211.  
1393 <http://dx.doi.org/10.1021/es011055j>  
1394 [Kurt-Karakus, P; Alegria, H; Birgul, A; Gungormus, E; Jantunen, L.](#) (2018). Organophosphate ester  
1395 (OPEs) flame retardants and plasticizers in air and soil from a highly industrialized city in  
1396 Turkey. *Sci Total Environ* 625: 555-565. <http://dx.doi.org/10.1016/j.scitotenv.2017.12.307>  
1397 [La Guardia, MJ; Hale, RC.](#) (2015). Halogenated flame-retardant concentrations in settled dust,  
1398 respirable and inhalable particulates and polyurethane foam at gymnastic training facilities and  
1399 residences. *Environ Int* 79: 106-114. <http://dx.doi.org/10.1016/j.envint.2015.02.014>  
1400 [Langer, S; Fredricsson, M; Weschler, CJ; Bekö, G; Strandberg, B; Remberger, M; Toftum, J; Clausen,](#)  
1401 [G.](#) (2016). Organophosphate esters in dust samples collected from Danish homes and daycare  
1402 centers. *Chemosphere* 154: 559-566. <http://dx.doi.org/10.1016/j.chemosphere.2016.04.016>  
1403 [Larsson, K; de Wit, CA; Sellström, U; Sahlström, L; Lindh, CH; Berglund, M.](#) (2018). Brominated  
1404 flame retardants and organophosphate esters in preschool dust and children's hand wipes.  
1405 *Environ Sci Technol* 52: 4878-4888. <http://dx.doi.org/10.1021/acs.est.8b00184>  
1406 [Launay, MA; Dittmer, U; Steinmetz, H.](#) (2016). Organic micropollutants discharged by combined sewer  
1407 overflows - Characterisation of pollutant sources and stormwater-related processes. *Water Res*  
1408 104: 82-92. <http://dx.doi.org/10.1016/j.watres.2016.07.068>  
1409 [Laws, BV; Dickenson, ER; Johnson, TA; Snyder, SA; Drewes, JE.](#) (2011). Attenuation of contaminants  
1410 of emerging concern during surface-spreading aquifer recharge. *Sci Total Environ* 409: 1087-  
1411 1094. <http://dx.doi.org/10.1016/j.scitotenv.2010.11.021>



1412 [Lazarov, B; Swinnen, R; Spruyt, M; Maes, F; Van Campenhout, K; Goelen, E; Covaci, A; Stranger, M.](#)  
1413 (2015). Air sampling of flame retardants based on the use of mixed-bed sorption tubes-a  
1414 validation study. *Environ Sci Pollut Res Int* 22: 18221-18229. [http://dx.doi.org/10.1007/s11356-](http://dx.doi.org/10.1007/s11356-015-5028-z)  
1415 [015-5028-z](http://dx.doi.org/10.1007/s11356-015-5028-z)

1416 [Lebel, GL; Williams, DT; Benoit, FM.](#) (1987). Use of large-volume resin cartridges for the  
1417 determination of organic contaminants in drinking water derived from the great lakes. In IH  
1418 Suffet; M Malaiyandi (Eds.), *Advances in Chemistry* (pp. 309-326). Washington, DC: American  
1419 Chemical Society. <http://dx.doi.org/10.1021/ba-1987-0214.ch014>

1420 [Lee, S; Jeong, W; Kannan, K; Moon, HB.](#) (2016). Occurrence and exposure assessment of  
1421 organophosphate flame retardants (OPFRs) through the consumption of drinking water in Korea.  
1422 *Water Res* 103: 182-188. <http://dx.doi.org/10.1016/j.watres.2016.07.034>

1423 [Li, J; Xie, Z; Mi, W; Lai, S; Tian, C; Emeis, KC; Ebinghaus, R.](#) (2017). Organophosphate esters in air,  
1424 snow, and seawater in the North Atlantic and the arctic. *Environ Sci Technol* 51: 6887-6896.  
1425 <http://dx.doi.org/10.1021/acs.est.7b01289>

1426 [Liu, LY; He, K; Hites, RA; Salamova, A.](#) (2016). Hair and nails as noninvasive biomarkers of human  
1427 exposure to brominated and organophosphate flame retardants. *Environ Sci Technol* 50: 3065-  
1428 3073. <http://dx.doi.org/10.1021/acs.est.5b05073>

1429 [Liu, LY; Salamova, A; He, K; Hites, RA.](#) (2015). Analysis of polybrominated diphenyl ethers and  
1430 emerging halogenated and organophosphate flame retardants in human hair and nails. *J*  
1431 *Chromatogr A* 1406: 251-257. <http://dx.doi.org/10.1016/j.chroma.2015.06.003>

1432 [Liu, R; Mabury, SA.](#) (2019). Organophosphite antioxidants in indoor dust represent an indirect source of  
1433 organophosphate esters. *Environ Sci Technol* 53: 1805-1811.  
1434 <http://dx.doi.org/10.1021/acs.est.8b05545>

1435 [Loos, R; Tavazzi, S; Mariani, G; Suurkuusk, G; Paracchini, B; Umlauf, G.](#) (2017). Analysis of emerging  
1436 organic contaminants in water, fish and suspended particulate matter (SPM) in the Joint Danube  
1437 Survey using solid-phase extraction followed by UHPLC-MS-MS and GC-MS analysis. *Sci*  
1438 *Total Environ* 607-608: 1201-1212. <http://dx.doi.org/10.1016/j.scitotenv.2017.07.039>

1439 [Loraine, GA; Pettigrov, ME.](#) (2006). Seasonal Variations in Concentrations of Pharmaceuticals and  
1440 Personal Care Products in Drinking Water and Reclaimed Wastewater in Southern California.  
1441 *Environ Sci Technol* 40: 687-695. <http://dx.doi.org/10.1021/es051380x>

1442 [Luongo, G; Oestman, C.](#) (2016). Organophosphate and phthalate esters in settled dust from apartment  
1443 buildings in Stockholm. *Indoor Air* 26: 414-425. <http://dx.doi.org/10.1111/ina.12217>

1444 [Maceira, A; Pecikoza, I; Marcé, RM; Borrull, F.](#) (2020). Multi-residue analysis of several high-  
1445 production-volume chemicals present in the particulate matter from outdoor air. A preliminary  
1446 human exposure estimation. *Chemosphere* 252: 126514.  
1447 <http://dx.doi.org/10.1016/j.chemosphere.2020.126514>

1448 [Mäkinen, MSE; Mäkinen, MRA; Koistinen, JTB; Pasanen, AL; Pasanen, PO; Kalliokoski, PJ; Korpi,](#)  
1449 [AM.](#) (2009). Respiratory and dermal exposure to organophosphorus flame retardants and  
1450 tetrabromobisphenol A at five work environments. *Environ Sci Technol* 43: 941-947.  
1451 <http://dx.doi.org/10.1021/es802593t>

1452 [Marklund, A; Andersson, B; Haglund, P.](#) (2003). Screening of organophosphorus compounds and their  
1453 distribution in various indoor environments. *Chemosphere* 53: 1137-1146.  
1454 [http://dx.doi.org/10.1016/S0045-6535\(03\)00666-0](http://dx.doi.org/10.1016/S0045-6535(03)00666-0)

1455 [Marklund, A; Andersson, B; Haglund, P.](#) (2005a). Organophosphorus flame retardants and plasticizers  
1456 in Swedish sewage treatment plants. *Environ Sci Technol* 39: 7423-7429.  
1457 <http://dx.doi.org/10.1021/es051013l>

1458 [Marklund, A; Andersson, B; Haglund, P.](#) (2005b). Traffic as a source of organophosphorus flame  
1459 retardants and plasticizers in snow. *Environ Sci Technol* 39: 3555-3562.  
1460 <http://dx.doi.org/10.1021/es0482177>

- 1461 [Maruya, KA; Dodder, NG; Sengupta, A; Smith, DJ; Lyons, JM; Heil, AT; Drewes, JE.](#) (2016).  
1462 Multimedia screening of contaminants of emerging concern (CECS) in coastal urban watersheds  
1463 in southern California (USA). *Environ Toxicol Chem* 35: 1986-1994.  
1464 <http://dx.doi.org/10.1002/etc.3348>
- 1465 [Matamoros, V; Arias, CA; Nguyen, LX; Salvadó, V; Brix, H.](#) (2012). Occurrence and behavior of  
1466 emerging contaminants in surface water and a restored wetland. *Chemosphere* 88: 1083-1089.  
1467 <http://dx.doi.org/10.1016/j.chemosphere.2012.04.048>
- 1468 [McDonough, CA; De Silva, AO; Sun, C; Cabrerizo, A; Adelman, D; Soltwedel, T; Bauerfeind, E; Muir,](#)  
1469 [DCG; Lohmann, R.](#) (2018). Dissolved organophosphate esters and polybrominated diphenyl  
1470 ethers in remote marine environments: Arctic surface water distributions and net transport  
1471 through Fram Strait. *Environ Sci Technol* 52: 6208-6216.  
1472 <http://dx.doi.org/10.1021/acs.est.8b01127>
- 1473 [McGoldrick, DJ; Letcher, RJ; Barresi, E; Keir, MJ; Small, J; Clark, MG; Sverko, E; Backus, SM.](#)  
1474 (2014). Organophosphate flame retardants and organosiloxanes in predatory freshwater fish from  
1475 locations across Canada. *Environ Pollut* 193: 254-261.  
1476 <http://dx.doi.org/10.1016/j.envpol.2014.06.024>
- 1477 [Meyer, J; Bester, K.](#) (2004). Organophosphate flame retardants and plasticisers in wastewater treatment  
1478 plants. *J Environ Monit* 6: 599-605. <http://dx.doi.org/10.1039/b403206c>
- 1479 [Mihajlovic, I; Fries, E.](#) (2012). Atmospheric deposition of chlorinated organophosphate flame retardants  
1480 (OFR) onto soils. *Atmos Environ* 56: 177-183. <http://dx.doi.org/10.1016/j.atmosenv.2012.03.054>
- 1481 [Mihajlović, I; Miloradov, MV; Fries, E.](#) (2011). Application of Twisselmann extraction, SPME, and  
1482 GC-MS to assess input sources for organophosphate esters into soil. *Environ Sci Technol* 45:  
1483 2264-2269. <http://dx.doi.org/10.1021/es103870f>
- 1484 [Mizouchi, S; Ichiba, M; Takigami, H; Kajiwara, N; Takamuku, T; Miyajima, T; Kodama, H; Someya,](#)  
1485 [T; Ueno, D.](#) (2015). Exposure assessment of organophosphorus and organobromine flame  
1486 retardants via indoor dust from elementary schools and domestic houses. *Chemosphere* 123: 17-  
1487 25. <http://dx.doi.org/10.1016/j.chemosphere.2014.11.028>
- 1488 [Monclús, L; Lopez-Bejar, M; De la Puente, J; Covaci, A; Jaspers, VLB.](#) (2018). First evaluation of the  
1489 use of down feathers for monitoring persistent organic pollutants and organophosphate ester  
1490 flame retardants: A pilot study using nestlings of the endangered cinereous vulture (*Aegypius*  
1491 *monachus*). *Environ Pollut* 238: 413-420. <http://dx.doi.org/10.1016/j.envpol.2018.03.065>
- 1492 [Norwegian Environment, A.](#) (2019a). Environmental contaminants in an urban fjord, 2018. (M-1441).  
1493 Trondheim, Norway. [https://www.miljodirektoratet.no/publikasjoner/2019/september-](https://www.miljodirektoratet.no/publikasjoner/2019/september-2019/envir-onmental-contaminants-in-an-urban-fjord-2018/)  
1494 [2019/envir-onmental-contaminants-in-an-urban-fjord-2018/](https://www.miljodirektoratet.no/publikasjoner/2019/september-2019/envir-onmental-contaminants-in-an-urban-fjord-2018/)
- 1495 [Norwegian Environment, A.](#) (2019b). Monitoring of environmental contaminants in freshwater  
1496 ecosystems 2018 - Occurrence and biomagnification. (Report M-1411).  
1497 [https://www.miljodirektoratet.no/publikasjoner/2019/juni-2019/monitoring-of-environmental-](https://www.miljodirektoratet.no/publikasjoner/2019/juni-2019/monitoring-of-environmental-contaminants-in-freshwater-ecosystems-2018--occurrence-and-biomagnification/)  
1498 [contaminants-in-freshwater-ecosystems-2018--occurrence-and-biomagnification/](https://www.miljodirektoratet.no/publikasjoner/2019/juni-2019/monitoring-of-environmental-contaminants-in-freshwater-ecosystems-2018--occurrence-and-biomagnification/)
- 1499 [O'Brien, JW; Thai, PK; Brandsma, SH; Leonards, PE; Ort, C; Mueller, JF.](#) (2015). Wastewater analysis  
1500 of Census day samples to investigate per capita input of organophosphorus flame retardants and  
1501 plasticizers into wastewater. *Chemosphere* 138: 328-334.  
1502 <http://dx.doi.org/10.1016/j.chemosphere.2015.06.014>
- 1503 [Ohura, T; Amagai, T; Senga, Y; Fusaya, M.](#) (2006). Organic air pollutants inside and outside residences  
1504 in Shimizu, Japan: Levels, sources and risks. *Sci Total Environ* 366: 485-499.  
1505 <http://dx.doi.org/10.1016/j.scitotenv.2005.10.005>
- 1506 [Okeme, JO; Nguyen, LV; Lorenzo, M; Dhal, S; Pico, Y; Arrandale, VH; Diamond, ML.](#) (2018a).  
1507 Polydimethylsiloxane (silicone rubber) brooch as a personal passive air sampler for semi-volatile  
1508 organic compounds. *Chemosphere* 208: 1002-1007.  
1509 <http://dx.doi.org/10.1016/j.chemosphere.2018.05.196>

1510 [Okeme, JO; Yang, C; Abdollahi, A; Dhal, S; Harris, SA; Jantunen, LM; Tsirlin, D; Diamond, ML.](#)  
1511 (2018b). Passive air sampling of flame retardants and plasticizers in Canadian homes using  
1512 PDMS, XAD-coated PDMS and PUF samplers. *Environ Pollut* 239: 109-117.  
1513 <http://dx.doi.org/10.1016/j.envpol.2018.03.103>

1514 [Otake, T; Yoshinaga, J; Yanagisawa, Y.](#) (2001). Analysis of organic esters of plasticizer in indoor air by  
1515 GC-MS and GC-FPD. *Environ Sci Technol* 35: 3099-3102. <http://dx.doi.org/10.1021/es001914o>

1516 [Otake, T; Yoshinaga, J; Yanagisawa, Y.](#) (2004). Exposure to phthalate esters from indoor environment. *J*  
1517 *Expo Sci Environ Epidemiol* 14: 524-528. <http://dx.doi.org/10.1038/sj.jea.7500352>

1518 [Padhye, LP; Yao, H; Kung'u, FT; Huang, CH.](#) (2014). Year-long evaluation on the occurrence and fate  
1519 of pharmaceuticals, personal care products, and endocrine disrupting chemicals in an urban  
1520 drinking water treatment plant. *Water Res* 51: 266-276.  
1521 <http://dx.doi.org/10.1016/j.watres.2013.10.070>

1522 [Persson, J; Wang, T; Hagberg, J.](#) (2018). Organophosphate flame retardants and plasticizers in indoor  
1523 dust, air and window wipes in newly built low-energy preschools. *Sci Total Environ* 628-629:  
1524 159-168. <http://dx.doi.org/10.1016/j.scitotenv.2018.02.053>

1525 [Peeverly, AA; Ma, Y; Venier, M; Rodenburg, Z; Spak, SN; Hornbuckle, KC; Hites, RA.](#) (2015).  
1526 Variations of flame retardant, polycyclic aromatic hydrocarbon, and pesticide concentrations in  
1527 Chicago's atmosphere measured using passive sampling. *Environ Sci Technol* 49: 5371-5379.  
1528 <http://dx.doi.org/10.1021/acs.est.5b00216>

1529 [Phillips, AL; Hammel, SC; Hoffman, K; Lorenzo, AM; Chen, A; Webster, TF; Stapleton, HM.](#) (2018).  
1530 Children's residential exposure to organophosphate ester flame retardants and plasticizers:  
1531 Investigating exposure pathways in the TESIE study. *Environ Int* 116: 176-185.  
1532 <http://dx.doi.org/10.1016/j.envint.2018.04.013>

1533 [Poma, G; Glynn, A; Malarvannan, G; Covaci, A; Darnerud, PO.](#) (2017). Dietary intake of phosphorus  
1534 flame retardants (PFRs) using Swedish food market basket estimations. *Food Chem Toxicol* 100:  
1535 1-7. <http://dx.doi.org/10.1016/j.fct.2016.12.011>

1536 [Poma, G; Sales, C; Bruyland, B; Christia, C; Gosciny, S; Van Loco, J; Covaci, A.](#) (2018). Occurrence  
1537 of organophosphorus flame retardants and plasticizers (PFRs) in Belgian foodstuffs and  
1538 estimation of the dietary exposure of the adult population. *Environ Sci Technol* 52: 2331-2338.  
1539 <http://dx.doi.org/10.1021/acs.est.7b06395>

1540 [Quednow, K; Püttmann, W.](#) (2009). Temporal concentration changes of DEET, TCEP, terbutryn, and  
1541 nonylphenols in freshwater streams of Hesse, Germany: possible influence of mandatory  
1542 regulations and voluntary environmental agreements. *Environ Sci Pollut Res Int* 16: 630-640.  
1543 <http://dx.doi.org/10.1007/s11356-009-0169-6>

1544 [Rantakokko, P; Kumar, E; Braber, J; Huang, T; Kiviranta, H; Cequier, E; Thomsen, C.](#) (2019).  
1545 Concentrations of brominated and phosphorous flame retardants in Finnish house dust and  
1546 insights into children's exposure. *Chemosphere* 223: 99-107.  
1547 <http://dx.doi.org/10.1016/j.chemosphere.2019.02.027>

1548 [Rauert, C; Harner, T; Schuster, JK; Eng, A; Fillmann, G; Castillo, LE; Fentanes, O; Villa Ibarra, MV;  
1549 Miglioranza, KSB; Rivadeneira, IM; Pozo, K; Zuluaga, BHA.](#) (2018). Atmospheric  
1550 Concentrations of New Persistent Organic Pollutants and Emerging Chemicals of Concern in the  
1551 Group of Latin America and Caribbean (GRULAC) Region. *Environ Sci Technol* 52: 7240-  
1552 7249. <http://dx.doi.org/10.1021/acs.est.8b00995>

1553 [Regnery, J; Puettmann, W.](#) (2009). Organophosphorus flame retardants and plasticizers in rain and snow  
1554 from middle Germany. *CLEAN - Soil, Air, Water* 37: 334-342.  
1555 <http://dx.doi.org/10.1002/clen.200900050>

1556 [Regnery, J; Püttmann, W.](#) (2010a). Occurrence and fate of organophosphorus flame retardants and  
1557 plasticizers in urban and remote surface waters in Germany. *Water Res* 44: 4097-4104.  
1558 <http://dx.doi.org/10.1016/j.watres.2010.05.024>

1559 [Regnery, J; Püttmann, W.](#) (2010b). Seasonal fluctuations of organophosphate concentrations in  
1560 precipitation and storm water runoff. *Chemosphere* 78: 958-964.  
1561 <http://dx.doi.org/10.1016/j.chemosphere.2009.12.027>  
1562 [Regnery, J; Püttmann, W; Merz, C; Berthold, G.](#) (2011). Occurrence and distribution of  
1563 organophosphorus flame retardants and plasticizers in anthropogenically affected groundwater. *J*  
1564 *Environ Monit* 13: 347-354. <http://dx.doi.org/10.1039/c0em00419g>  
1565 [Rodil, R; Quintana, JB; Concha-Graña, E; López-Mahía, P; Muniategui-Lorenzo, S; Prada-Rodríguez,](#)  
1566 [D.](#) (2012). Emerging pollutants in sewage, surface and drinking water in Galicia (NW Spain).  
1567 *Chemosphere* 86: 1040-1049. <http://dx.doi.org/10.1016/j.chemosphere.2011.11.053>  
1568 [Saito, I; Onuki, A; Seto, H.](#) (2007). Indoor organophosphate and polybrominated flame retardants in  
1569 Tokyo. *Indoor Air* 17: 28-36. <http://dx.doi.org/10.1111/j.1600-0668.2006.00442.x>  
1570 [Sala, B; Giménez, J; de Stephanis, R; Barceló, D; Eljarrat, E.](#) (2019). First determination of high levels  
1571 of organophosphorus flame retardants and plasticizers in dolphins from Southern European  
1572 waters. *Environ Res* 172: 289-295. <http://dx.doi.org/10.1016/j.envres.2019.02.027>  
1573 [Salamova, A; Ma, Y; Venier, M; Hites, RA.](#) (2014). High levels of organophosphate flame retardants in  
1574 the great lakes atmosphere. *Environ Sci Technol Lett* 1: 8-14.  
1575 <http://dx.doi.org/10.1021/ez400034n>  
1576 [Salamova, A; Peverly, AA; Venier, M; Hites, RA.](#) (2016). Spatial and temporal trends of particle phase  
1577 organophosphate ester concentrations in the atmosphere of the great lakes. *Environ Sci Technol*  
1578 50: 13249-13255. <http://dx.doi.org/10.1021/acs.est.6b04789>  
1579 [Santín, G; Eljarrat, E; Barceló, D.](#) (2016). Simultaneous determination of 16 organophosphorus flame  
1580 retardants and plasticizers in fish by liquid chromatography-tandem mass spectrometry. *J*  
1581 *Chromatogr A* 1441: 34-43. <http://dx.doi.org/10.1016/j.chroma.2016.02.058>  
1582 [Schreder, ED; La Guardia, MJ.](#) (2014). Flame retardant transfers from U.S. households (dust and  
1583 laundry wastewater) to the aquatic environment. *Environ Sci Technol* 48: 11575-11583.  
1584 <http://dx.doi.org/10.1021/es502227h>  
1585 [Schreder, ED; Uding, N; La Guardia, MJ.](#) (2016). Inhalation a significant exposure route for chlorinated  
1586 organophosphate flame retardants. *Chemosphere* 150: 499-504.  
1587 <http://dx.doi.org/10.1016/j.chemosphere.2015.11.084>  
1588 [Scott, BF; Sverko, E; Maguire, RJ.](#) (1996). Determination of benzothiazole and alkylphosphates in water  
1589 samples from the Great Lakes drainage basin by gas chromatography/atomic emission detection.  
1590 *Water Qual Res J Can* 31: 341-360. <http://dx.doi.org/10.2166/wqrj.1996.021>  
1591 [Scott, PD; Bartkow, M; Blockwell, SJ; Coleman, HM; Khan, SJ; Lim, R; McDonald, JA; Nice, H;](#)  
1592 [Nugegoda, D; Pettigrove, V; Tremblay, LA; Warne, MS; Leusch, FD.](#) (2014). A national survey  
1593 of trace organic contaminants in Australian rivers. *J Environ Qual* 43: 1702-1712.  
1594 <http://dx.doi.org/10.2134/jeq2014.01.0012>  
1595 [Sengupta, A; Lyons, JM; Smith, DJ; Drewes, JE; Snyder, SA; Heil, A; Maruya, KA.](#) (2014). The  
1596 occurrence and fate of chemicals of emerging concern in coastal urban rivers receiving discharge  
1597 of treated municipal wastewater effluent. *Environ Toxicol Chem* 33: 350-358.  
1598 <http://dx.doi.org/10.1002/etc.2457>  
1599 [Sha, B; Dahlberg, AK; Wiberg, K; Ahrens, L.](#) (2018). Fluorotelomer alcohols (FTOHs), brominated  
1600 flame retardants (BFRs), organophosphorus flame retardants (OPFRs) and cyclic volatile  
1601 methylsiloxanes (cVMSs) in indoor air from occupational and home environments. *Environ*  
1602 *Pollut* 241: 319-330. <http://dx.doi.org/10.1016/j.envpol.2018.04.032>  
1603 [Shin, H; Moschet, C; Young, TM; Bennett, DH.](#) (2019). Measured concentrations of consumer product  
1604 chemicals in California house dust: Implications for sources, exposure, and toxicity potential.  
1605 *Indoor Air* 30: 60-75. <http://dx.doi.org/10.1111/ina.12607>  
1606 [Shin, HM; McKone, TE; Nishioka, MG; Fallin, MD; Croen, LA; Hertz-Picciotto, I; Newschaffer, CJ;](#)  
1607 [Bennett, DH.](#) (2014). Determining source strength of semivolatile organic compounds using



1608 measured concentrations in indoor dust. *Indoor Air* 24: 260-271.  
1609 <http://dx.doi.org/10.1111/ina.12070>

1610 [Stachel, B; Jantzen, E; Knoth, W; Kruger, F; Lepom, P; Oetken, M; Reincke, H; Sawal, G; Schwartz, R;](#)  
1611 [Uhlig, S.](#) (2005). The Elbe Flood in August 2002—Organic Contaminants in Sediment Samples  
1612 Taken After the Flood Event. *J Environ Sci Health A Tox Hazard Subst Environ Eng* 40: 265-  
1613 287. <http://dx.doi.org/10.1081/ESE-200045531>

1614 [Stapleton, HM; Misenheimer, J; Hoffman, K; Webster, TF.](#) (2014). Flame retardant associations  
1615 between children's handwipes and house dust. *Chemosphere* 116: 54-60.  
1616 <http://dx.doi.org/10.1016/j.chemosphere.2013.12.100>

1617 [Stubbings, WA; Guo, JH; Simon, K; Romanak, K; Bowerman, W; Venier, M.](#) (2018). Flame retardant  
1618 metabolites in addled bald eagle eggs from the Great Lakes region. *Environ Sci Technol Lett* 5:  
1619 354-359. <http://dx.doi.org/10.1021/acs.estlett.8b00163>

1620 [Sugeng, EJ; Leonards, PEG; van de Bor, M.](#) (2017). Brominated and organophosphorus flame retardants  
1621 in body wipes and house dust, and an estimation of house dust hand-loadings in Dutch toddlers.  
1622 *Environ Res* 158: 789-797. <http://dx.doi.org/10.1016/j.envres.2017.07.035>

1623 [Sühring, R; Diamond, ML; Scheringer, M; Wong, F; Pucko, M; Stern, G; Burt, A; Hung, H; Fellin, P;](#)  
1624 [Li, H; Jantunen, LM.](#) (2016). Organophosphate esters in Canadian Arctic air: Occurrence, levels  
1625 and trends. *Environ Sci Technol* 50: 7409. <http://dx.doi.org/10.1021/acs.est.6b00365>

1626 [Sundkvist, AM; Olofsson, U; Haglund, P.](#) (2010). Organophosphorus flame retardants and plasticizers in  
1627 marine and fresh water biota and in human milk. *J Environ Monit* 12: 943-951.  
1628 <http://dx.doi.org/10.1039/b921910b>

1629 [Tajima, S; Araki, A; Kawai, T; Tsuboi, T; Ait Bamai, Y; Yoshioka, E; Kanazawa, A; Cong, S; Kishi, R.](#)  
1630 (2014). Detection and intake assessment of organophosphate flame retardants in house dust in  
1631 Japanese dwellings. *Sci Total Environ* 478: 190-199.  
1632 <http://dx.doi.org/10.1016/j.scitotenv.2013.12.121>

1633 [Takeuchi, S; Tanaka-Kagawa, T; Saito, I; Kojima, H; Jin, K; Satoh, M; Kobayashi, S; Jinno, H.](#) (2015).  
1634 Differential determination of plasticizers and organophosphorus flame retardants in residential  
1635 indoor air in Japan. *Environ Sci Pollut Res Int* 25: 7113-7120. [http://dx.doi.org/10.1007/s11356-](http://dx.doi.org/10.1007/s11356-015-4858-z)  
1636 [015-4858-z](#)

1637 [Tan, H; Yang, L; Yu, Y; Guan, Q; Liu, X; Li, L; Chen, D.](#) (2019). Co-existence of organophosphate di-  
1638 and tri-esters in house dust from South China and Midwestern United States: Implications for  
1639 human exposure. *Environ Sci Technol* 53: 4784-4793. <http://dx.doi.org/10.1021/acs.est.9b00229>

1640 [Teo, TL; Coleman, HM; Khan, SJ.](#) (2016). Presence and select determinants of organophosphate flame  
1641 retardants in public swimming pools. *Sci Total Environ* 569-570: 469-475.  
1642 <http://dx.doi.org/10.1016/j.scitotenv.2016.06.085>

1643 [Tokumura, M; Hatayama, R; Tatsu, K; Naito, T; Takeda, T; Raknuzzaman, M; Al-Mamun, MH;](#)  
1644 [Masunaga, S.](#) (2017). Organophosphate flame retardants in the indoor air and dust in cars in  
1645 Japan. *Environ Monit Assess* 189: 48. <http://dx.doi.org/10.1007/s10661-016-5725-1>

1646 [Valcarcel, Y; Valdehita, A; Becerra, E; Lopez de Alda, M; Gil, A; Gorga, M; Petrovic, M; Barcelo, D;](#)  
1647 [Navas, JM.](#) (2018). Determining the presence of chemicals with suspected endocrine activity in  
1648 drinking water from the Madrid region (Spain) and assessment of their estrogenic, androgenic  
1649 and thyroidal activities. *Chemosphere* 201: 388-398.  
1650 <http://dx.doi.org/10.1016/j.chemosphere.2018.02.099>

1651 [Van den Eede, N; Dirtu, AC; Ali, N; Neels, H; Covaci, A.](#) (2012). Multi-residue method for the  
1652 determination of brominated and organophosphate flame retardants in indoor dust. *Talanta* 89:  
1653 292-300. <http://dx.doi.org/10.1016/j.talanta.2011.12.031>

1654 [Van Den Eede, N; Heffernan, A; Aylward, LL; Hobson, P; Neels, H; Mueller, JF; Covaci, A.](#) (2015).  
1655 Age as a determinant of phosphate flame retardant exposure of the Australian population and

1656 identification of novel urinary PFR metabolites. *Environ Int* 74: 1-8.  
1657 <http://dx.doi.org/10.1016/j.envint.2014.09.005>

1658 [Velázquez-Gómez, M; Hurtado-Fernández, E; Lacorte, S.](#) (2019). Differential occurrence, profiles and  
1659 uptake of dust contaminants in the Barcelona urban area. *Sci Total Environ* 648: 1354-1370.  
1660 <http://dx.doi.org/10.1016/j.scitotenv.2018.08.058>

1661 [Wallner, P; Kundi, M; Moshhammer, H; Piegler, K; Hohenblum, P; Scharf, S; Fröhlich, M; Damberger,](#)  
1662 [B; Tapplere, P; Hutter, HP.](#) (2012). Indoor air in schools and lung function of Austrian school  
1663 children. *J Environ Monit* 14: 1976-1982. <http://dx.doi.org/10.1039/c2em30059a>

1664 [Wang, Y; Li, W; Martínez-Moral, MP; Sun, H; Kannan, K.](#) (2019). Metabolites of organophosphate  
1665 esters in urine from the United States: Concentrations, temporal variability, and exposure  
1666 assessment. *Environ Int* 122: 213-221. <http://dx.doi.org/10.1016/j.envint.2018.11.007>

1667 [Wong, F; De Wit, CA; Newton, SR.](#) (2018). Concentrations and variability of organophosphate esters,  
1668 halogenated flame retardants, and polybrominated diphenyl ethers in indoor and outdoor air in  
1669 Stockholm, Sweden. *Environ Pollut* 240: 514-522.  
1670 <http://dx.doi.org/10.1016/j.envpol.2018.04.086>

1671 [Woudneh, MB; Benskin, JP; Wang, G; Grace, R; Hamilton, MC; Cosgrove, JR.](#) (2015). Quantitative  
1672 determination of 13 organophosphorous flame retardants and plasticizers in a wastewater  
1673 treatment system by high performance liquid chromatography tandem mass spectrometry. *J*  
1674 *Chromatogr A* 1400: 149-155. <http://dx.doi.org/10.1016/j.chroma.2015.04.026>

1675 [Xu, F; Giovanoulis, G; van Waes, S; Padilla-Sanchez, JA; Papadopoulou, E; Magnér, J; Haug, LS;](#)  
1676 [Neels, H; Covaci, A.](#) (2016). Comprehensive study of human external exposure to  
1677 organophosphate flame retardants via air, dust, and hand wipes: The importance of sampling and  
1678 assessment strategy. *Environ Sci Technol* 50: 7752-7760.  
1679 <http://dx.doi.org/10.1021/acs.est.6b00246>

1680 [Yasuhara, A.](#) (1994). DETERMINATION OF TRIS(2-CHLOROETHYL) PHOSPHATE IN  
1681 LEACHATES FROM LANDFILLS BY CAPILLARY GAS-CHROMATOGRAPHY USING  
1682 FLAME PHOTOMETRIC DETECTION. *J Chromatogr A* 684: 366-369.  
1683 [http://dx.doi.org/10.1016/0021-9673\(94\)00643-1](http://dx.doi.org/10.1016/0021-9673(94)00643-1)

1684 [Yasuhara, A.](#) (1995). Chemical components in leachates from hazardous wastes landfills in Japan.  
1685 *Toxicol Environ Chem* 51: 113-120. <http://dx.doi.org/10.1080/02772249509358229>

1686 [Yasuhara, A; Shiraishi, H; Nishikawa, M; Yamamoto, T; Nakasugi, O; Okumura, T; Kenmotsu, K;](#)  
1687 [Fukui, H; Nagase, M; Kawagoshi, Y.](#) (1999). Organic components in leachates from hazardous  
1688 waste disposal sites. *Waste Manag Res* 17: 186-197. <http://dx.doi.org/10.1034/j.1399->  
1689 [3070.1999.00038.x](http://dx.doi.org/10.1034/j.1399-3070.1999.00038.x)

1690 [Yoshida, T; Matsunaga, I; Tomioka, K; Kumagai, S.](#) (2006). Interior air pollution in automotive cabins  
1691 by volatile organic compounds diffusing from interior materials: I. Survey of 101 types of  
1692 Japanese domestically produced cars for private use. *Indoor Built Environ* 15: 425-444.  
1693 <http://dx.doi.org/10.1177/1420326X06069395>

1694 [Zhao, F; Chen, M; Gao, F; Shen, H; Hu, J.](#) (2017). Organophosphorus flame retardants in pregnant  
1695 women and their transfer to chorionic villi. *Environ Sci Technol* 51: 6489-6497.  
1696 <http://dx.doi.org/10.1021/acs.est.7b01122>

1697 [Zhou, L; Hiltcher, M; Püttmann, W.](#) (2017). Occurrence and human exposure assessment of  
1698 organophosphate flame retardants in indoor dust from various microenvironments of the  
1699 Rhine/Main region, Germany. *Indoor Air* 27: 1113-1127. <http://dx.doi.org/10.1111/ina.12397>