

Air Quality Criteria for Particulate Matter

Volume I of II

Air Quality Criteria for Particulate Matter

Volume I

National Center for Environmental Assessment-RTP Office
Office of Research and Development
U.S. Environmental Protection Agency
Research Triangle Park, NC

DISCLAIMER

This document has been reviewed in accordance with U.S. Environmental Protection Agency policy and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

PREFACE

National Ambient Air Quality Standards (NAAQS) are promulgated by the United States Environmental Protection Agency (EPA) to meet requirements set forth in Sections 108 and 109 of the U.S. Clean Air Act (CAA). Sections 108 and 109 require the EPA Administrator (1) to list widespread air pollutants that reasonably may be expected to endanger public health or welfare; (2) to issue air quality criteria for them that assess the latest available scientific information on nature and effects of ambient exposure to them; (3) to set “primary” NAAQS to protect human health with adequate margin of safety; (4) to set “secondary” NAAQS to protect against welfare effects (e.g., effects on vegetation, ecosystems, visibility, climate, manmade materials, etc.); and (5) to periodically (every 5 years) review and revise, as appropriate, the criteria and NAAQS for a given listed pollutant or class of pollutants.

The original NAAQS for particulate matter (PM), issued in 1971 as “total suspended particulate” (TSP) standards, were revised in 1987 to focus on protecting against human health effects associated with exposure to ambient PM less than 10 microns ($\leq 10 \mu\text{m}$) that are capable of being deposited in thoracic (tracheobronchial and alveolar) portions of the lower respiratory tract. Later periodic reevaluation of newly available scientific information, as presented in the last previous version of this “*Air Quality Criteria for Particulate Matter*” document published in 1996, provided key scientific bases for PM NAAQS decisions published in July 1997. More specifically, the PM_{10} NAAQS set in 1987 ($150 \mu\text{g}/\text{m}^3$, 24-h; $50 \mu\text{g}/\text{m}^3$, annual average) were retained in modified form and new standards ($65 \mu\text{g}/\text{m}^3$, 24-h; $15 \mu\text{g}/\text{m}^3$, annual average) for particles $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$) were promulgated in July 1997.

This final version of revised *Air Quality Criteria for Particulate Matter* assesses new scientific information that has become available (published or accepted for publication) mainly

between early 1996 through April 2002, although a few important new studies published through 2003 are also considered. Several previous successive drafts of this document were released for public comment and review by the Clean Air Scientific Advisory Committee (CASAC), to obtain comments on the organization and structure of the document, the issues addressed, the approaches employed in assessing and interpreting the newly available information on PM exposures and effects, and the key findings and conclusions arrived at by this assessment. Public comments and CASAC review recommendations were taken into account in making revisions to this document for incorporation into this final draft. Evaluations contained in this document will be drawn on to provide inputs to associated PM Staff Paper analyses prepared by EPA's Office of Air Quality Planning and Standards (OAQPS) to pose alternatives for consideration by the EPA Administrator with regard to proposal and, ultimately, promulgation of decisions on potential retention or revision of the current PM NAAQS.

The document describes the nature, sources, distribution, measurement, and concentrations of PM in outdoor (ambient) environments. It also evaluates the latest data on human exposures to ambient PM and consequent health effects in exposed human populations, to support decision making regarding primary, health-related PM NAAQS. The document also evaluates ambient PM environmental effects on vegetation and ecosystems, visibility, and man-made materials, as well as atmospheric PM effects on climate change processes, to support decision making bearing on secondary PM NAAQS.

Preparation of this document was coordinated by EPA's National Center for Environmental Assessment in Research Triangle Park (NCEA-RTP). NCEA-RTP scientific staff, together with experts from other EPA/ORD laboratories and academia, contributed to writing of document chapters. The NCEA of EPA acknowledges the contributions provided by authors, contributors, and reviewers and the diligence of its staff and contractors in the preparation of this document.

Air Quality Criteria for Particulate Matter

VOLUME I

1.	INTRODUCTION	1-1
2.	PHYSICS, CHEMISTRY, AND MEASUREMENT OF OF PARTICULATE MATTER	2-1
	APPENDIX 2A: Techniques for Measuring of Semivolatile Organic Compounds	2A-1
	APPENDIX 2B: Analytical Techniques	2B-1
3.	CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER	3-1
	APPENDIX 3A: Composition of Particulate Matter Source Emissions	3A-1
	APPENDIX 3B: Organic Composition of Particulate Matter	3B-1
	APPENDIX 3C: Aerosol Composition Data from the Speciation Network	3C-1
	APPENDIX 3D: Spatial and Temporal Variability of the Nationwide AIRS PM _{2.5} and PM _{10-2.5} Data Sets	3D-1
	APPENDIX 3E: Characterization of PM _{2.5} , PM ₁₀ , and PM _{10-2.5} Concentrations at IMPROVE Sites	3E-1
4.	ENVIRONMENTAL EFFECTS OF AIRBORNE PARTICULATE MATTER	4-1
	APPENDIX 4A: Common and Latin Names	4A-1
5.	HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS	5-1

Air Quality Criteria for Particulate Matter
(cont'd)

VOLUME II

6.	DOSIMETRY OF PARTICULATE MATTER	6-1
7.	TOXICOLOGY OF PARTICULATE MATTER IN HUMANS AND LABORATORY ANIMALS	7-1
	APPENDIX 7A: Rat-to-Human Dose Extrapolation	7A-1
	APPENDIX 7B: Ambient Bioaerosols	7B-1
8.	EPIDEMIOLOGY OF HUMAN HEALTH EFFECTS ASSOCIATED WITH AMBIENT PARTICULATE MATTER	8-1
	APPENDIX 8A: Short-Term PM Exposure-Mortality Studies: Summary Tables	8A-1
	APPENDIX 8B: Particulate Matter-Morbidity Studies: Summary Tables	8B-1
9.	INTEGRATIVE SYNTHESIS	9-1

Table of Contents

	<u>Page</u>
List of Tables	I-xiii
List of Figures	I-xix
Authors, Contributors, and Reviewers	I-xxxii
U.S. Environmental Protection Agency Project Team for Development of Air Quality Criteria for Particulate Matter	I-xl
U.S. Environmental Protection Agency Science Advisory Board (SAB) Staff Office Clean Air Scientific Advisory Committee (CASAC) Particulate Matter Review Panel	I-xliii
Abbreviations and Acronyms	I-xlvi
1. INTRODUCTION	1-1
1.1 BACKGROUND	1-1
1.1.1 Legislative Requirements	1-1
1.1.2 Criteria and NAAQS Review Process	1-2
1.1.3 History of Earlier PM Criteria and NAAQS Reviews	1-4
1.1.4 The 1997 PM NAAQS Revision	1-5
1.1.5 Coordinated PM Research Program	1-7
1.2 CURRENT PM CRITERIA AND NAAQS REVIEW	1-12
1.2.1 Key Milestones and Procedures for Document Preparation	1-12
1.2.2 Assessment Approaches	1-16
1.3 DOCUMENT ORGANIZATION	1-18
REFERENCES	1-19
2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER	2-1
2.1 PHYSICS AND CHEMISTRY OF PARTICULATE MATTER	2-2
2.1.1 Basic Concepts	2-2
2.1.2 Physical Properties and Processes	2-3
2.1.2.1 Definitions of Particle Diameter	2-3
2.1.2.2 Aerosol Size Distributions	2-5
2.1.2.3 Ultrafine Particles	2-28
2.1.3 Chemistry of Atmospheric Particulate Matter	2-37
2.1.3.1 Chemical Composition and Its Dependence on Particle Size	2-37
2.1.3.2 Primary and Secondary Particulate Matter	2-38
2.1.3.3 Particle-Vapor Partitioning	2-39
2.1.3.4 Atmospheric Lifetimes and Removal Processes	2-49
2.1.4 Comparison of Fine and Coarse Particles	2-51
2.2 MEASUREMENT OF PARTICULATE MATTER	2-51
2.2.1 Particle Measurements of Interest	2-51
2.2.2 Issues in Measurement of Particulate Matter	2-54
2.2.2.1 Artifacts Due to Chemical Reactions	2-55

Table of Contents
(cont'd)

	<u>Page</u>
2.2.2.2	Treatment of Semivolatile Components of Particulate Matter 2-56
2.2.2.3	Upper Cut Point 2-57
2.2.2.4	Cut Point for Separation of Fine and Coarse Particulate Matter 2-58
2.2.2.5	Treatment of Pressure, Temperature, and Relative Humidity 2-58
2.2.2.6	Time Resolution 2-59
2.2.2.7	Accuracy and Precision 2-59
2.2.3	Measurement of Semivolatile Particulate Matter 2-62
2.2.3.1	Particle-Bound Water 2-62
2.2.3.2	Nitrate and Organic Species 2-68
2.2.3.3	Continuous Measurement of Semivolatile and Nonvolatile Mass 2-75
2.2.4	U.S. Environmental Protection Agency Monitoring Methods 2-79
2.2.4.1	The Federal Reference Methods for Measurement of Equilibrated Mass for PM ₁₀ , PM _{2.5} , and PM _{10-2.5} 2-79
2.2.5	Speciation Monitoring 2-93
2.2.6	Inorganic Elemental Analyses 2-96
2.2.7	Elemental and Organic Carbon in Particulate Matter 2-97
2.2.8	Ionic Species 2-99
2.2.9	Continuous Monitoring 2-99
2.2.10	Measurements of Individual Particles 2-101
2.2.11	Low Flow Filter Samples for Multiday Collection of Particulate Matter 2-103
2.3	SUMMARY AND KEY POINTS 2-104
2.3.1	Chemistry and Physics of Atmospheric Particles 2-104
2.3.2	Measurement of Atmospheric Particles 2-107
2.3.3	Key Points 2-111
REFERENCES 2-113
APPENDIX 2A:	TECHNIQUES FOR MEASURING OF SEMIVOLATILE ORGANIC COMPOUNDS 2A-1
APPENDIX 2B:	ANALYTICAL TECHNIQUES 2B-1
2B.1	INORGANIC ELEMENTS 2B-1
2B.1.1	Energy Dispersive X-Ray Fluorescence 2B-1
2B.1.2	Synchrotron Induced X-ray Fluorescence 2B-2
2B.1.3	Proton (or Particle) Induced X-ray Emission 2B-2
2B.1.4	Proton Elastic Scattering Analysis 2B-3
2B.1.5	Total Reflection X-Ray Fluorescence 2B-3
2B.1.6	Instrumental Neutron Activation Analysis 2B-4

Table of Contents
(cont'd)

		<u>Page</u>
2B.1.7	Atomic Absorption Spectrophotometry	2B-5
2B.1.8	Inductively Coupled Plasma with Atomic Emission Spectroscopy	2B-6
2B.1.9	Inductively Coupled Plasma with Mass Spectroscopy	2B-6
2B.1.10	Scanning Electron Microscopy	2B-7
2B.2	ORGANIC AND INORGANIC CARBON	2B-8
2B.3	CONTINUOUS METHODS	2B-17
2B.3.1	Continuous Measurement of Mass	2B-17
2B.3.2	Continuous Measurement of Organic and/or Elemental Carbon	2B-21
2B.3.3	Continuous Measurements of Nitrate and Sulfate	2B-24
2B.4	OTHER CONTINUOUS MEASUREMENTS	2B-26
3.	CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER	3-1
3.1	INTRODUCTION	3-1
3.2	PATTERNS AND TRENDS IN AMBIENT PM CONCENTRATIONS	3-2
3.2.1	Seasonal Variability in PM Concentrations	3-13
3.2.2	Diurnal (Circadian) Variability in PM Concentrations	3-30
3.2.3	Relationships Among Particulate Matter in Different Size Fractions	3-32
3.2.4	Relationships Between Mass and Chemical Component Concentrations	3-34
3.2.5	Spatial Variability in Particulate Matter and its Components	3-40
3.3	SOURCES OF PRIMARY AND SECONDARY PM	3-59
3.3.1	Chemistry of Secondary PM Formation	3-62
3.3.2	Source Contributions to Ambient PM Determined by Receptor Models	3-72
3.3.3	Background Concentrations of PM in the United States	3-82
3.3.4	Emissions Estimates for Primary PM, and Precursors to Secondary PM (SO ₂ , NO _x , VOCs, and NH ₃) in the United States	3-92
3.3.5	Uncertainties of Emissions Inventories	3-98
3.4	SUMMARY AND KEY CONCLUSIONS	3-101
	REFERENCES	3-106
APPENDIX 3A:	Spatial and Temporal Variability of the Nationwide AIRS PM _{2.5} and PM _{10-2.5} Data Sets	3A-1
	REFERENCES	3A-3

Table of Contents
(cont'd)

	<u>Page</u>
APPENDIX 3B: Aerosol Composition Data from the Speciation Network	3B-1
REFERENCES	3B-3
APPENDIX 3C: Organic Composition of Particulate Matter	3C-1
REFERENCES	3C-14
APPENDIX 3D: Composition of Particulate Matter Source Emissions	3D-1
REFERENCES	3D-25
APPENDIX 3E: Characterization of PM _{2.5} , PM ₁₀ , and PM _{10-2.5} Concentrations at IMPROVE Sites	3E-1
REFERENCES	3E-6
4. ENVIRONMENTAL EFFECTS OF AIRBORNE PARTICULATE MATTER	4-1
4.1 INTRODUCTION	4-1
4.2 EFFECTS OF AMBIENT AIRBORNE PARTICULATE MATTER ON VEGETATION AND NATURAL ECOSYSTEMS	4-1
4.2.1 Ecological Attributes	4-2
4.2.2 Ecosystem Exposures – Particle Deposition	4-4
4.2.2.1 Fine and Coarse Particulate Matter	4-5
4.2.2.2 Diversity of Deposition Modes	4-8
4.2.2.3 Magnitude of Deposition	4-44
4.2.3 Assessment of Atmospheric PM Deposition Effects	4-55
4.2.3.1 Effects on Vegetation and Ecosystems	4-61
4.2.3.2 Ecosystem Response to Stresses	4-79
4.2.4 Urban Ecosystems	4-148
4.3 AIRBORNE PARTICLE EFFECTS ON VISIBILITY	4-152
4.3.1 Introduction	4-152
4.3.2 Factors Affecting Atmospheric Visibility	4-154
4.3.2.1 Optical Properties of the Atmosphere and Atmospheric Particles	4-154
4.3.2.2 Relative Humidity Effects on Particle Size and Light-Scattering Properties	4-161
4.3.3 Relationships Between Particles and Visibility	4-165
4.3.4 Photographic Modeling of Visibility Impairment	4-173
4.3.5 Visibility Monitoring Methods and Networks	4-174
4.3.6 Visibility Impairment: Trends and Current Conditions	4-177
4.3.6.1 Trends in Visibility Impairment	4-177
4.3.6.2 Current Conditions	4-185
4.3.7 Societal Impacts of Particulate Matter Visibility Effects	4-186
4.3.7.1 Economic Studies	4-186
4.3.7.2 Public Perception and Attitude Studies	4-188
4.4 PARTICULATE MATTER EFFECTS ON MATERIALS	4-191
4.4.1 Corrosive Effects of Particles and Sulfur Dioxide on Man-Made Surfaces	4-191

Table of Contents
(cont'd)

	<u>Page</u>	
4.4.1.1	Metals	4-192
4.4.1.2	Painted Finishes	4-193
4.4.1.3	Stone and Concrete	4-196
4.4.2	Soiling and Discoloration of Man-Made Surfaces	4-202
4.4.2.1	Stones and Concrete	4-203
4.4.2.2	Household and Industrial Paints	4-203
4.5	ATMOSPHERIC PARTICULATE MATTER, CLIMATE CHANGE, AND EFFECTS ON SOLAR UV-B RADIATION TRANSMISSION	4-204
4.5.1	Atmospheric Particle Interactions with Solar and Terrestrial Radiation Related to Climate Change	4-206
4.5.1.1	The Projected Impacts of Global Climate Change	4-206
4.5.1.2	Airborne Particle Relationships to Global Warming and Climate Change	4-209
4.5.2	Atmospheric Particulate Matter Effects on the Transmission of Solar Ultraviolet Radiation Transmission: Impacts on Human Health and the Environment	4-220
4.5.2.1	Potential Effects of Increased Ultraviolet Radiation Transmission	4-220
4.5.2.2	Airborne Particle Effects on Atmospheric Transmission of Solar Ultraviolet Radiation	4-223
4.6	SUMMARY AND KEY CONCLUSIONS	4-227
4.6.1	Particulate Matter Effects on Vegetation and Ecosystems	4-227
4.6.2	Particulate Matter-Related Effects on Visibility	4-235
4.6.3	Particulate Matter-Related Effects on Materials	4-238
4.6.4	Effects of Atmospheric Particulate Matter on Global Warming Processes and Transmission of Solar Ultraviolet Radiation	4-239
	REFERENCES	4-241
	APPENDIX 4A: Common and Latin Names	4A-1
5.	HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS	5-1
5.1	INTRODUCTION	5-1
5.1.1	Purpose	5-1
5.1.2	Particulate Matter Mass and Constituents	5-2
5.1.3	Relationship to Past Documents	5-3
5.1.4	Chapter Structure	5-4
5.2	BASIC CONCEPTS OF EXPOSURE	5-5
5.2.1	The Concept of Exposure	5-5
5.2.2	Components of Exposure	5-6
5.2.3	Quantification of Exposure	5-9

Table of Contents
(cont'd)

		<u>Page</u>
5.2.4	Methods To Estimate Personal Exposure	5-10
5.2.4.1	Direct Measurement Methods	5-11
5.2.4.2	Indirect or Modeling Methods	5-15
5.2.4.3	Methods for Estimating Personal Exposure to Ambient Particulate Matter	5-22
5.3	SUMMARY OF PARTICULATE MATTER MASS DATA	5-29
5.3.1	Types of Particulate Matter Exposure Measurement Studies	5-29
5.3.2	Available Data	5-30
5.3.2.1	Personal Exposure Data	5-30
5.3.2.2	Microenvironmental Data	5-35
5.3.2.3	Traffic-Related Microenvironments	5-43
5.3.2.4	Reanalyses of Previously-Reported Particulate Matter Exposure Data	5-43
5.3.3	Factors Influencing and Key Findings on Particulate Matter Exposures	5-46
5.3.3.1	Relationship of Personal/Microenvironmental Particulate Matter with Ambient Particulate Matter	5-46
5.3.3.2	Factors That Affect Relationships Between Personal Exposure and Ambient PM	5-66
5.3.3.3	Effect of Ambient Sources on Exposures to Particulate Matter	5-87
5.3.3.4	Correlations of Particulate Matter with Other Pollutants	5-89
5.4	SUMMARY OF PARTICULATE MATTER CONSTITUENT DATA	5-94
5.4.1	Introduction	5-94
5.4.2	Monitoring Studies That Address Particulate Matter Constituents	5-94
5.4.3	Key Findings	5-94
5.4.3.1	Correlations of Personal and Indoor Concentrations with Ambient Concentrations of Particulate Matter Constituents	5-94
5.4.4	Factors Affecting Correlations Between Ambient Measurements and Personal or Microenvironmental Measurements of Particulate Matter Constituents	5-105
5.4.5	Limitations of Available Data	5-106
5.5	IMPLICATIONS OF USING AMBIENT PM CONCENTRATIONS IN TOXICOLOGICAL AND EPIDEMIOLOGICAL STUDIES OF PM HEALTH EFFECTS	5-107
5.5.1	Toxicology	5-107
5.5.2	Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiological Analyses	5-107

Table of Contents
(cont'd)

	<u>Page</u>	
5.5.2.1	Associations Between Personal Exposures and Ambient Particulate Matter Concentrations	5-110
5.5.2.2	Role of Compositional Differences in Exposure Characterization for Epidemiology	5-114
5.5.2.3	Role of Spatial Variability in Exposure Characterization for Epidemiology	5-115
5.5.3	Analysis of Exposure Measurement Error Issues in Particulate Matter Epidemiology	5-116
5.5.3.1	Time-Series Analyses	5-117
5.5.3.2	Studies of Chronic Effects	5-117
5.5.4	Conclusions from Analysis of Exposure Measurement Errors on Particulate Matter Epidemiology	5-120
5.6	SUMMARY OF OBSERVATIONS AND LIMITATIONS	5-121
	REFERENCES	5-129

List of Tables

<u>Number</u>		<u>Page</u>
1-1	Key Scientific Uncertainties Related to the Source-To-Response Framework	1-10
1-2	Research Topics and Questions Recommended by National Research Council to be Addressed by Expanded M Research Program	1-11
1-3	Key Milestones in the Development of This Document	1-13
2-1	Correlations Between TEOM Measurements in Phoenix	2-26
2-2	Comparison of Ambient Particles, Fine Particles (ultrafine plus accumulation-mode) and Coarse Particles	2-52
2-3	Particulate Matter Components/Parameters of Interest for Health, Ecological, or Radiative Effects; for Source Category Apportionment Studies; or for Air Quality Modeling Evaluation Studies	2-53
2-4	Summary of Sensitivity Studies of WINS Impactor Performance	2-84
2-5	PM _{2.5} Samplers Currently Designated as FRMs for PM _{2.5} Mass Concentrations	2-85
2-6	Measurement Methods for Inorganic Elements	2-97
2-7	Methods for Continuous Measurement of PM Mass, PM Components, etc.	2-100
3-1	Distribution of Ratios of PM _{2.5} to PM ₁₀ and Correlations Between PM _{2.5} and PM ₁₀ , PM _{2.5} and PM _{10-2.5} , and PM _{10-2.5} and PM ₁₀ Found at Collocated Monitoring Sites in Seven Aerosol Characteristic (EPA/HEI) Regions in 1999	3-33
3-2	Concentrations (ng/m ³) of PM _{2.5} , PM _{10-2.5} , and Selected Elements (ng/m ³) in the PM _{2.5} and PM _{10-2.5} Size Ranges with Standard Deviations (SD) and Correlations Between Elements and PM _{2.5} Mass in Philadelphia, PA	3-35
3-3	Concentrations (in ng/m ³) of PM _{2.5} , PM _{10-2.5} , and Selected Elements in the PM _{2.5} and PM _{10-2.5} Size Range with Standard Deviations (SD) and Correlations (r) Between Elements and PM _{2.5} and PM _{10-2.5} Mass in Phoenix, AZ	3-36
3-4a-d	Measures of the Spatial Variability of PM _{2.5} Concentrations Within Selected Metropolitan Statistical Areas	3-41

List of Tables
(Cont'd)

<u>Number</u>		<u>Page</u>
3-5	Measures of the Spatial Variability of PM _{10-2.5} Concentrations Within Selected Metropolitan Statistical Areas	3-49
3-6	Correlation Coefficients for Spatial Variation of PM _{2.5} Mass and Different Sources for Pairs of Sampling Sites in the South Coast Air Basin (1986)	3-57
3-7	Correlation Coefficients for Spatial Variation of PM _{2.5} Mass and Different Components for Pairs of Sampling Sites in Philadelphia (1994)	3-57
3-8	Constituents of Atmospheric Particles and their Major Sources	3-60
3-9	Receptor Model Source Contributions to PM _{2.5}	3-76
3-10	Receptor Model Source Contributions to PM ₁₀	3-77
3-11	Emissions of Primary PM _{2.5} by Various Sources in 1999	3-94
3-12	Emissions of Precursors to Secondary PM _{2.5} Formation by Various Sources in 1999	3-95
3-13	Nationwide Changes in Ambient Concentrations and Emissions of PM ₁₀ and Gaseous Precursors to Secondary Particulate Matter from 1992 to 2001	3-97
3A-1	Performance Metrics for PM _{2.5} from Collocated Samplers	3A-4
3B-1	PM _{2.5} Speciation Samplers by Location: Sites Selected for Summary in 2004 PM AQCD	3B-4
3B-2a	Burlington, VT Summary Data (October 2001 to September 2002)	3B-5
3B-2b	Philadelphia, PA Summary Data (October 2001 to September 2002)	3B-6
3B-2c	Atlanta, GA Summary Data (October 2001 to September 2002)	3B-7
3B-2d	Detroit, MI Summary Data (October 2001 to September 2002)	3B-8
3B-2e	Chicago, IL Summary Data (October 2001 to September 2002)	3B-9
3B-2f	St. Louis, MI Summary Data (October 2001 to September 2002)	3B-10
3B-2g	Houston, TX Summary Data (October 2001 to September 2002)	3B-11

List of Tables
(Cont'd)

<u>Number</u>		<u>Page</u>
3B-2h	Minneapolis, MN Summary Data (October 2001 to September 2002)	3B-12
3B-2i	Boulder, CO Summary Data (October 2001 to September 2002)	3B-13
3B-2j	Phoenix, AZ Summary Data (October 2001 to September 2002)	3B-14
3B-2k	Seattle, WA Summary Data (October 2001 to September 2002)	3B-15
3B-2l	Sacramento, CA Summary Data (October 2001 to September 2002)	3B-16
3B-2m	Riverside-Rubidoux, CA Summary Data (October 2001 to September 2002)	3B-17
3B-3	Blank Corrections for Elemental Carbon, Organic Carbon, and Total Carbon in the Speciation Network	3B-18
3C-1	Particulate Organic and Elemental Carbon Concentrations (in $\mu\text{g C/m}^3$) Based on Studies Published After 1995	3C-2
3C-2	Particulate Organic Compound Concentrations (in ng C/m^3) Based on Studies Published After 1990 at Selected Sites	3C-5
3D-1	Average Abundances of Major Elements in Soil and Crustal Rock	3D-2
3D-2	Summary of Particle-Phase Organic Compounds Present in Fine Particle Road Dust Sample	3D-5
3D-3	Composition of Fine Particles Released by Various Stationary Sources in the Philadelphia Area	3D-7
3D-4a	Organic and Elemental Carbon Fractions of Diesel and Gasoline Engine Particulate Matter Exhaust	3D-12
3D-4b	Contribution of Organic Carbon to Particulate Matter Carbon Emissions in Motor Vehicle Exhaust Collected from Vehicles Operated on Chassis Dynamometers	3D-13
3D-5	Emission Rates (mg/mi) for Constituents of Particulate Matter from Gasoline and Diesel Vehicles	3D-14
3D-6	Summary of Particle-Phase Organic Compounds Emitted from Motor Vehicles	3D-17

List of Tables
(Cont'd)

<u>Number</u>		<u>Page</u>
3D-7	Mass Emissions, Organic Carbon, and Elemental Carbon Emissions from Residential Combustion of Wood	3D-19
3D-8	Summary of Particle-Phase Organic Compounds Emitted from the Combustion of Wood in Fireplaces	3D-21
3D-9	Mean Aerosol Composition at Tropical Site (Sriwijaya University, Sumatra, Indonesia) Affected Heavily by Biomass Burning Emissions	3D-22
3E-1	Ranges of Annual Mean PM Concentrations at IMPROVE Monitoring Sites ($\mu\text{g}/\text{m}^3$)	3E-3
3E-2	Ranges of Annual 90th Percentile Values of Daily PM Concentrations at IMPROVE Monitoring Sites ($\mu\text{g}/\text{m}^3$)	3E-4
4-1	Essential Ecological Attributes and Reporting Categories	4-3
4-2	Types and Determinants of Particulate Deposition and Impact to Vegetation	4-9
4-3	Relative Importance of Wet, Dry, Particulate, and Total Deposition to Three Forest Sites	4-12
4-4	Reported Mean Deposition Velocities for Sulfate, Chlorine, Nitrate, and Ammonium-Ion-Containing Particles	4-37
4-5	Representative Empirical Measurements of Deposition Velocity for Particulate Deposition	4-38
4-6	Reported Mean Deposition Velocities for Potassium, Sodium, Calcium, and Magnesium Base Cation Containing Particles	4-38
4-7	Mean Annual Nitrogen Deposition (equivalents/ha/year) from Fine and Coarse Particles Compared to Total Nitrogen Deposition from all Sources to a Variety of Forest Ecosystems	4-46
4-8	Mean Annual Sulfate Deposition (equivalents/ha/year) from Fine and Coarse Particles Compared to Total Sulfur Deposition from all Sources to a Variety of Forest Ecosystems	4-47

List of Tables
(Cont'd)

<u>Number</u>		<u>Page</u>
4-9	Mean Annual Base Cation Deposition (equivalents/ha/year) from Fine and Coarse Particles Compared to Total Base Cation Deposition from all Sources to a Variety of Forest Ecosystems	4-48
4-10	Mean Particle Size, Deposition Rates, and Derived Deposition Velocities for Heavy Metal Deposition to the Upper Canopy (inert plates or leaves) of an Upland Oak Forest	4-52
4-11	Total Heavy Metal Deposition to Temperate Latitude Forests	4-52
4-12	Primary Goods and Services Provided by Ecosystems	4-58
4-13	Ecosystem Functions Impacted by Air Pollution Effects on Temperate Forest Ecosystems	4-84
4-14	Nitrogen-Saturated Forests in North America, Including Estimated N Inputs and Outputs	4-102
4-15	Visibility Measurement Techniques	4-171
4-16	Residential Visibility Contingent Valuation Study Results	4-189
4-17	Residential Visibility Valuation Study Results for Los Angeles and San Francisco	4-190
4-18	Corrosive Effects of Particulate Matter and Sulfur Dioxide on Metals	4-194
4-19	Corrosive Effects of Particulate Matter and Sulfur Dioxide on Stone	4-197
4-20	Examples of Impacts Resulting from Projected Changes in Extreme Climate Events	4-208
4-21	Effects of Reactive Nitrogen	4-230
5-1	Types of Particulate Matter Used in Exposure and Concentration Variables ...	5-7
5-2	Activity Pattern Studies Included in the Consolidated Human Activity Database	5-18
5-3	Personal Exposure Models for Particulate Matter	5-19
5-4	Summary of Recent PM Personal Exposure Studies	5-31

List of Tables
(cont'd)

<u>Number</u>	<u>Page</u>
5-5	Summary of Recent Microenvironmental PM Measurement Studies 5-36
5-6	Papers Reporting Reanalyses of Particulate Matter Exposure Studies 5-44
5-7	Personal Monitoring Studies for Particulate Matter: Measured Concentrations and Correlation Coefficients 5-49
5-8	Mean Concentration for PM Mass Reported for the Baltimore (Williams et al., 2000a,b,c) and the Fresno (Evans et al., 2000) Studies 5-58
5-9	Daily-average Correlation Coefficients Reported for the Baltimore (Williams et al., 2000a,b,c) and the Fresno (Evans et al., 2000) Studies 5-58
5-10	Regression Analysis Reported for Indoor/Outdoor Relationships for PM _{2.5} in the Baltimore (Williams et al., 2000a,b,c) and Fresno (Evans et al., 2000) Studies 5-58
5-11	Volume Mean Diameter and Maximum PM _{2.5} Concentrations of Indoor Particle Sources 5-84
5-12	Correlations Between Personal PM _{2.5} and Ambient Pollutant Concentrations 5-91
5-13	Correlations Between Hourly Personal PM _{2.5} and Gaseous Pollutants 5-93
5-14	Studies That have Measured Particulate Matter Constituents in Personal Exposure Samples 5-96
5-15	Studies That Have Measured Particulate Matter Constituents in Microenvironmental Samples 5-97
5-16	Summary Statistics for Personal, Indoor, and Outdoor Concentrations of Selected Aerosol Components in Two Pennsylvania Communities 5-99
5-17	Regression Analysis of Indoor Versus Outdoor Concentrations 5-101
5-18	Mixed Model Analysis of Personal Versus Outdoor Concentrations 5-103
5-19	Regression Analysis of Indoor Versus Outdoor Concentrations 5-103
5-20	Regression Analysis of Indoor Versus Outdoor Concentrations 5-104

List of Figures

<u>Number</u>		<u>Page</u>
1-1	A general framework for integrating PM research	1-9
2-1	Distribution of coarse (c), accumulation (a), and nuclei (n) mode particles by three characteristics: (a) number, N; (b) surface area, S; and (c) volume, V for the grand average continental size distribution	2-8
2-2	Particle size distributions by number: (a) number concentrations are shown on a logarithmic scale to display the wide range by site and size; (b) number concentrations for the average urban distribution are shown on a linear scale	2-9
2-3	Size distribution by volume for the averaged (a) rural and urban-influenced rural number distribution shown in Figure 2-2a and a distribution from south central New Mexico, and (b) for the averaged urban and freeway-influenced urban number distributions shown in Figure 2-2a	2-10
2-4	Volume size distribution, measured in traffic, showing fine and coarse particles and the nuclei and accumulation modes of fine particles	2-11
2-5	Sub-micrometer number size distribution observed in a boreal forest in Finland showing the tri-modal structure of fine particles	2-12
2-6	An idealized size distribution, that might be observed in traffic, showing fine and coarse particles and the nucleation, Aitken, and accumulation modes that comprise fine particles	2-13
2-7	Specified particle penetration (size-cut curves) through an ideal (no-particle-loss) inlet for five different size-selective sampling criteria	2-16
2-8	An idealized distribution of ambient particulate matter showing fine-mode particles and coarse-mode particles and the fractions collected by size-selective samplers	2-18
2-9	Comparison of penetration curves for two PM ₁₀ beta gauge samplers using cyclone inlets	2-19
2-10	Three examples of impactor size distributions and distributions resulting from fitting several log normal distributions to the impactor data	2-24
2-11	Particulate matter concentrations in Spokane, WA, during the August 30, 1996 dust storm	2-27
2-12	Typical engine exhaust size distribution	2-32

List of Figures

(cont'd)

<u>Number</u>		<u>Page</u>
2-13	Number size distributions showing measurement of a nucleation burst mode in a boreal forest in Finland	2-33
2-14	Examples of the measured 1-h average particle number size distributions and the log normal fits to the modes of the data	2-34
2-15a-f	Fitted multi-model particle size distribution at different sampling distances from Freeway 405: (a) 30 m downwind, (b) 60 m downwind, (c) 90 m downwind, (d) 150 m downwind, (e) 300 m downwind, (f) 300 m upwind ...	2-35
2-15g	Combination of Figures 2-15(a-e), with $dN/d \log D_p$ scale	2-36
2-16a	Particle growth curves showing fully reversible hygroscopic growth of sulfuric acid (H_2SO_4) particles, deliquescent growth of ammonium sulfate $[(NH_4)_2 SO_4]$ particles at the deliquescent point (A, about 80% relative humidity [RH]), reversible hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached	2-42
2-16b	Theoretical predictions and experimental measurements of growth of NH_4HSO_4 particles at relative humidities between 95 and 100%	2-43
2-17	Concentration of the dissolved gas in the particle normalized by its initial concentration as a function of airway generation for $H = 10^4 M atm^{-1}$ for particle diameters of 0.1, 0.3, 0.7, and 1 μm	2-46
2-18	Schematic showing major nonvolatile and semivolatile components of $PM_{2.5}$.	2-57
2-19	Aerosol water content expressed as a mass percentage, as a function of relative humidity	2-65
2-20	Amount of ammonium nitrate volatilized from Teflon filters, expressed as a percentage of the measured $PM_{2.5}$ mass, for the SCAQS and CalTech studies, for spring and fall sampling periods	2-69
2-21	Average concentration ($\mu g/m^3$) of nonvolatile and semivolatile PM components in three cities	2-76
2-22	Comparison of mass measurements with collocated RAMS (real-time data), PC-BOSS (1-h data), FRM $PM_{2.5}$ sampler (average of 24-h data), and a conventional TEOM monitor (real-time data)	2-77

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
2-23	Schematic diagram of the sample collection portion of the PM _{2.5} FRM sampler	2-81
2-24	Schematic view of the final design of the WINS	2-82
2-25	Evaluation of the final version of the WINS	2-83
2-26	Schematic diagram showing the principle of virtual impaction	2-91
2-27	Size distribution of particles divided by chemical classification into organic, marine, and soil (or crustal)	2-102
2B-1	Thermogram for a sample containing rock dust (a carbonate source) and diesel exhaust, shows three traces that correspond to temperature, filter transmittance, and Flame Ionization Detector (FID) detector response	2B-10
2B-2	Examples of thermograms obtained by (a) the IMPROVE protocol, and by (b) the NIOSH protocol	2B-12
3-1a	1999-2001 county-wide average annual mean PM ₁₀ concentrations (µg/m ³) for counties with PM ₁₀ monitors	3-7
3-1b	1999-2001 highest county-wide 98th percentile 24-h average PM ₁₀ concentrations (µg/m ³) for counties with PM ₁₀ monitors	3-7
3-2	Nationwide trend in ambient PM ₁₀ concentration from 1992 through 2001	3-8
3-3	Trend in PM ₁₀ annual mean concentrations by EPA region, 1992 through 2001 (µg/m ³)	3-9
3-4a	1999-2001 county-wide average annual mean PM _{2.5} concentrations (µg/m ³) for counties with PM _{2.5} monitors	3-10
3-4b	1999-2001 highest county-wide 98th percentile 24-h average PM _{2.5} concentrations (µg/m ³) for counties with PM _{2.5} monitors	3-10
3-5	Collection of annual distribution of 24-h average PM _{2.5} concentrations observed in U.S. and Canadian health studies conducted during the 1980's and early 1990's	3-11

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
3-6a	1999-2000 estimated county-wide average annual mean PM _{10-2.5} concentrations (µg/m ³) for counties with collocated PM _{2.5} and PM ₁₀ monitors	3-14
3-6b	1999-2000 estimated county-wide highest 98th percentile 24-h PM _{10-2.5} concentrations (µg/m ³) for counties with collocated PM _{2.5} and PM ₁₀ monitors	3-15
3-7a,b	Quarterly distribution of 24-h average PM _{2.5} concentrations for selected monitors in the (a) Philadelphia, PA and (b) Cleveland, OH MSAs	3-16
3-7c,d	Quarterly distribution of 24-h average PM _{2.5} concentrations for selected monitors in the (c) Dallas, TX and (d) Los Angeles, CA MSAs	3-17
3-8	Seasonal concentrations of PM _{2.5} and PM ₁₀ measured in the four MAACS cities	3-20
3-9a,b	Quarterly distribution of 24-h average PM _{10-2.5} concentrations for selected sites in the (a) Cleveland, OH; (b) Dallas, TX MSAs	3-22
3-9c	Quarterly distribution of 24-h average PM _{10-2.5} concentrations for selected sites in the Los Angeles, CA MSAs	3-23
3-10	Frequency distribution of 24-h average PM _{2.5} concentrations measured at the Presbyterian home (PBY) monitoring site in southwestern Philadelphia from 1992 to 1995	3-25
3-11	Concentrations of 24-h average PM _{2.5} measured at the EPA site in Phoenix, AZ from 1995 to 1997	3-26
3-12	Frequency distribution of 24-h average PM _{2.5} concentrations measured at the EPA site in Phoenix, AZ from 1995 to 1997	3-27
3-13	Frequency distribution of 24-h average PM _{2.5} measurements obtained from all California Air Resources Board dichotomous sampler sites from 1989 to 1998	3-27
3-14	Frequency distribution of 24-h average PM _{10-2.5} concentrations obtained from all California Air Resource Board Dichotomous sampler sites from 1989 to 1998	3-28

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
3-15	Concentrations of 24-h average PM _{2.5} measured at the Riverside-Rubidoux site from 1989 to 1998	3-28
3-16	Frequency distribution of 24-h average PM _{2.5} concentrations measured at the Riverside-Rubidoux site from 1989 to 1994	3-29
3-17	Intraday variability of hourly average PM _{2.5} concentrations across the United States	3-31
3-18	Occurrence of differences between pairs of sites in three MSAs	3-47
3-19	Intersite correlation coefficients for PM _{2.5} , PM ₁₀ , and PM _{10-2.5}	3-51
3-20	PM _{2.5} chemical components in downtown Los Angeles and Burbank (1986) have similar characteristics	3-54
3-21	Concentrations of PM _{2.5} chemical components in Rubidoux and downtown Los Angeles (1986)	3-55
3-22	Monthly average Saharan dust components in all size fractions of the aerosol sampled in Miami, FL (from 1974 to 1996)	3-85
3-23	PM _{2.5} and PM ₁₀ concentrations measured at Chilliwack Airport, located in southwestern British Columbia, just before and during the Asian desert dust episode of April and May 1998	3-86
3-24	Time series of 24-h average PM ₁₀ concentrations observed in the Rio Grande Valley during May 1998	3-88
3-25	PM ₁₀ concentrations observed in St. Louis, MO, during May 1998	3-89
3A-1	Philadelphia, PA-NJ metropolitan statistical area	3A-5
3A-2	Washington, DC metropolitan statistical area	3A-6
3A-3	Norfolk, VA metropolitan statistical area	3A-7
3A-4	Columbia, SC metropolitan statistical area	3A-8
3A-5	Atlanta, GA metropolitan statistical area	3A-9

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
3A-6	Birmingham, AL metropolitan statistical area	3A-10
3A-7	Tampa, FL metropolitan statistical area	3A-11
3A-8	Cleveland, OH metropolitan statistical area	3A-12
3A-9	Pittsburgh, PA metropolitan statistical area	3A-13
3A-10	Steubenville, OH-Weirton, WV metropolitan statistical area	3A-14
3A-11	Detroit MI metropolitan statistical area	3A-15
3A-12	Grand Rapids, MI metropolitan statistical area	3A-16
3A-13	Milwaukee, WI metropolitan statistical area	3A-17
3A-14	Chicago, IL metropolitan statistical area	3A-18
3A-15	Gary, IN metropolitan statistical area	3A-19
3A-16	Louisville, KY metropolitan statistical area	3A-20
3A-17	St. Louis, MO metropolitan statistical area	3A-21
3A-18	Baton Rouge, LA metropolitan statistical area	3A-22
3A-19	Kansas City, KS-MO metropolitan statistical area	3A-23
3A-20	Dallas, TX metropolitan statistical area	3A-24
3A-21	Boise, ID metropolitan statistical area	3A-25
3A-22	Salt Lake City, UT metropolitan statistical area	3A-26
3A-23	Seattle, WA metropolitan statistical area	3A-27
3A-24	Portland, OR metropolitan statistical area	3A-28
3A-25	Los Angeles-Long Beach, CA metropolitan statistical area	3A-29
3A-26	Riverside-San Bernadino, CA metropolitan statistical area	3A-30

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
3A-27	San Diego, CA metropolitan statistical area	3A-31
3A-28	Columbia, SC metropolitan statistical area	3A-32
3A-29	Tampa, FL metropolitan statistical area	3A-33
3A-30	Cleveland, OH metropolitan statistical area	3A-34
3A-31	Steubenville, OH metropolitan statistical area	3A-35
3A-32	Detroit, MI metropolitan statistical area	3A-36
3A-33	Milwaukee, WI metropolitan statistical area	3A-37
3A-34	Chicago, IL metropolitan statistical area	3A-38
3A-35	Gary, IN metropolitan statistical area	3A-39
3A-36	Louisville, KY metropolitan statistical area	3A-40
3A-37	St. Louis, MO metropolitan statistical area	3A-41
3A-38	Baton Rouge, LA metropolitan statistical area	3A-42
3A-39	Dallas, TX metropolitan statistical area	3A-43
3A-40	Salt Lake City, UT metropolitan statistical area	3A-44
3A-41	Portland, OR metropolitan statistical area	3A-45
3A-42	Los Angeles, CA metropolitan statistical area	3A-46
3A-43	Riverside, CA metropolitan statistical area	3A-47
3A-44	San Diego, CA metropolitan statistical area	3A-48
3D-1	Size distribution of particles generated in a laboratory resuspension chamber	3D-2
3D-2	Size distribution of California source emissions, 1986	3D-4

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
3D-3	Chemical abundances for PM _{2.5} emissions from paved road dust in Denver, CO	3D-5
3D-4	Chemical abundances for PM _{2.5} emissions from wood burning in Denver, CO	3D-18
3E-1a,b	Interannual variability in 24-h average PM _{2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY	3E-7
3E-2a,b	Seasonal variability in 24-h average PM _{2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY	3E-8
3E-3a,b	Seasonal variability in 24-h average PM _{10-2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY	3E-9
4-1	The diversity of fine PM from sites in the western and eastern United States	4-6
4-2	Relative importance of three modes of deposition of nitrate (A) and sulfate (B) at high elevation sites (Unsworth and Wilshaw, 1989; Fowler et al., 1989; Mueller, 1988; Aneja and Murthy, 1994)	4-11
4-3	A simplified resistance catena representing the factors controlling deposition of particles to the surface	4-16
4-4	The relationship between deposition velocity of selected particulate materials and the distribution of the material between the coarse- and fine-aerosol fractions	4-17
4-5	The relationship between particle diameter and deposition velocity for particles	4-18
4-6	Vertical stratification of diverse, chemically speciated particles in a mixed oak forest	4-33
4-7	The relationship between particle size and concentration below a spruce canopy with wind velocity at a height of 16.8 m equaling 5 m/s	4-34

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
4-8	Range of percent of total deposition delivered in the dry particulate fraction, across the sites of the Integrated Forest Study	4-40
4-9	Contribution of particulate deposition to total deposition of nitrogen, sulfur, and base cations	4-40
4-10	Mean percent of total nitrogen, sulfur, or base cation deposition contributed by fine plus coarse particles	4-49
4-11	Annual total deposition of heavy metals to Hubbard Brook Experimental Forest, NH	4-53
4-12	Sample stressors and the essential ecological attributes they affect (after Science Advisory Board, 2002)	4-56
4-13	Linkages among various ecosystem goods and services (food, water, biodiversity, forest products) and other driving forces (climate change)	4-60
4-14	Effects of environmental stress on forest trees are presented on a hierarchial scale for the leaf, branch, tree, and stand levels of organization . . .	4-80
4-15	Illustration of the nitrogen cascade showing the movement of the human-produced reactive nitrogen (Nr) as it cycles through the various environmental reservoirs in the atmosphere, terrestrial ecosystems, and aquatic ecosystems	4-97
4-16	Nitrogen cycle (dotted lines indicate processes altered by nitrogen saturation)	4-99
4-17	Diagrammatic overview of excess nitrogen in North America	4-103
4-18	Schematic of sources and sinks of hydrogen ions in a forest	4-117
4-19	Key elements of proposed framework for determining critical loads for nitrogen and sulfur in the United States	4-124
4-20	Calcium deposition in > 2- μ m particles, < 2- μ m particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites	4-128

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
4-21	Magnesium deposition in > 2- μ m particles, < 2- μ m particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites	4-129
4-22	Potassium deposition in > 2- μ m particles, < 2- μ m particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites	4-129
4-23	Base cation deposition in > 2- μ m particles, < 2- μ m particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites	4-130
4-24	Total cation leaching (total height of bar) balanced by sulfate and nitrate estimated from particulate deposition (assuming no ecosystem retention, particulate sulfur and nitrogen) and by other sources (both deposition and internal) of sulfate and nitrate (other sulfur and nitrogen sources) and by other anions in the Integrated Forest Study sites	4-131
4-25	Soil exchangeable Ca ²⁺ pools and net annual export of Ca ²⁺ (deposition minus leaching times 25 years) in the Integrated Forest Study sites	4-133
4-26	Soil exchangeable Mg ²⁺ pools and net annual export of Mg ²⁺ (deposition minus leaching times 25 years) in the Integrated Forest Study sites	4-133
4-27	Soil exchangeable K ⁺ pools and net annual export of K ⁺ (deposition minus leaching times 25 years) in the Integrated Forest Study sites	4-134
4-28a	Simulated soil solution mineral acid anions in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition	4-138
4-28b	Simulated soil solution base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition	4-139
4-29a	Simulated soil solution Al in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition	4-140
4-29b	Simulated soil solution soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition	4-141
4-30a	Simulated soil solution mineral acid anions in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition	4-142

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
4-30b	Simulated soil solution base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition	4-143
4-31	Relationship of plant nutrients and trace metals with vegetation	4-147
4-32	Light reflected from a target toward an observer	4-156
4-33	Light-scattering efficiency factor (per cross-sectional area), Q , for a homogeneous sphere with an index of refraction of 1.50 as a function of the size parameter, $\alpha = \pi D/\lambda$	4-159
4-34	Volume-specific light-scattering efficiency as a function of particle diameter D_p	4-161
4-35	Particle growth curve as a function of relative humidity (RH) showing deliquescent growth of ammonium sulfate $[(NH_4)_2 SO_4]$ particles at the deliquescent point (A, about 80% RH), reversible hygroscopic growth of ammonium sulfate solution droplets at $RH > 80\%$, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached	4-162
4-36	Comparison of extinction (Mm^{-1}) and visual range (km)	4-167
4-37	Proportionality of observed daytime haziness to fine particle mass concentration in Los Angeles	4-168
4-38	Relative humidity adjustment factor, $f(RH)$, for ammonium sulfate as a function of relative humidity	4-170
4-39a	Aggregate visibility trends (in deciviews) for 10 eastern Class 1 areas	4-180
4-39b	Aggregate visibility trends (in deciviews) for 26 western Class 1 areas	4-180
4-40	(a) Eastern class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component; (b) Western class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component	4-182
4-41a	Light extinction trends in Tucson, Arizona from 1993 to 2002	4-184
4-41b	Light extinction trends in Phoenix, Arizona from 1994 to 2001	4-184

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
4-42	Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750	4-213
5-1	Hypothetical exposure time profile: pollutant exposure as a function of time showing how the averaged exposure, integrated exposure, and peak exposure relate to the instantaneous exposure	5-9
5-2	Distribution of F_{INF} (a) and α (b) for daytime as estimated from PTEAM study data	5-25
5-3	Comparison of correlation coefficients for longitudinal analyses of personal exposure versus ambient concentrations for individual subjects for $PM_{2.5}$ and sulfate	5-61
5-4	Personal exposure versus ambient concentrations for $PM_{2.5}$ and sulfate for well, moderately, and poorly ventilated indoor environments	5-64
5-5	Regression analyses of aspects of daytime personal exposure to PM_{10} estimated using data from the PTEAM study: (a) total personal exposure to PM, T, regressed on ambient concentration, C; (b) personal exposure to ambient PM, A, regressed on C; and (c) personal exposure to nonambient PM, N, regressed on C	5-65
5-6	Air exchange rates measured in homes throughout the United States	5-72
5-7	Box plots of hourly air exchange rates stratified by season in Boston, MA during 1998	5-73
5-8	Geometric mean infiltration factor (indoor/outdoor ratio) for hourly nighttime, nonsource data for two seasons	5-74
5-9	Regression of air exchange rate on absolute indoor-outdoor temperature difference	5-76
5-10	Comparison of deposition rates from Long et al. (2001a) with literature values (from Abt et al., 2000b)	5-78
5-11	Penetration efficiencies and deposition rates from models of nightly average data	5-79
5-12	Mean hourly indoor/outdoor particle concentration ratio from an unoccupied residence in Fresno, CA during spring 1999	5-85

List of Figures
(cont'd)

<u>Number</u>		<u>Page</u>
5-13	Personal versus outdoor SO_4^{2-} in State College, PA	5-100
5-14	Plots of nonambient exposure to PM_{10} showing (a) daily individual daytime values from PTEAM data and (b) daily-average values from THEES data	5-113

Authors, Contributors, and Reviewers

CHAPTER 1. INTRODUCTION

Principal Authors

Dr. Lester D. Grant—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Dennis J. Kotchmar—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

CHAPTER 2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER

Principal Authors

Dr. William Wilson—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Candis S. Claiborn—Washington State University, Laboratory for Atmospheric Research,
Department of Civil and Environmental Engineering, P.O. Box 642910, Pullman, WA 99164

Dr. Brooke L. Hemming—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributing Authors

Dr. Tom Cahill, University of California, Davis, One Shields Ave., Davis, CA 95616

Dr. Judith C. Chow, Desert Research Institute, 2215 Raggio Parkway, Reno, NV 89512

Max R. Peterson, Research Triangle Institute, P.O. Box 12194, Research Triangle Park, NC
27709

Dr. James J. Schauer, University of Wisconsin, College of Engineering, Department of Civil and
Environmental Engineering, Madison, WI 53706

Dr. Barbara J. Turpin, The State University of New Jersey, Rutgers, Department of
Environmental Sciences and Rutgers Cooperative Extension, New Brunswick, NJ 08901-8551

Dr. John G. Watson, Desert Research Institute, 2215 Raggio Parkway, Reno, NV 89512

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers

Dr. Edward O. Edney—National Exposure Research Laboratory (E205-02)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. R.R. Eldred, Crocker Nuclear Laboratory, University of California, Davis, Davis, CA 95616

Mr. William Ewald—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

James B. Flanagan—Research Triangle Institute, P.O. Box 12194, Research Triangle Park, NC
27709

Dr. Judith Graham—American Chemistry Council, 1300 Wilson Boulevard, Arlington, VA
22207

Dr. Lester D. Grant—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Jim Homolya—Office of Air Quality Planning and Standards (C339-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

R.K.M. Jayanty—Research Triangle Institute, P.O. Box 12194, Research Triangle Park, NC
27709

Mr. Scott Mathias—Office of Air Quality Planning and Standards (C539-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Frank McElroy—National Exposure Research Laboratory (MD-46),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Joseph Pinto—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Joann Rice—Office of Air Quality Planning and Standards (C339-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27709

Ed E. Rickman—Research Triangle Institute, P.O. Box 12194, Research Triangle Park, NC
27709

Dr. Bret Schichtel—Colorado State University, Cooperative Institute for Research in the
Atmosphere, Foothills Campus, Fort Collins, CO 80523-1375

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. John J. Vandenberg—National Center for Environmental Assessment (8601D), U.S. Environmental Protection Agency, 1200 Pennsylvania Avenue, NW, Washington, DC 20460

Dr. Russell Wiener—National Exposure Research Laboratory (D205-03)
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

***CHAPTER 3. CONCENTRATIONS, SOURCES, AND
EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER***

Principal Author

Dr. Joseph P. Pinto—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributing Authors

Dr. Allen S. Lefohn—A.S.L. & Associates, 111 North Last Chance Gulch, Helena, MT 59601

Dr. Barbara J. Turpin—The State University of New Jersey, Rutgers, Department of
Environmental Sciences and Rutgers Cooperative Extension, New Brunswick, NJ 08901-8551

Dr. James J. Schauer—University of Wisconsin, College of Engineering, Department of Civil
and Environmental Engineering, Madison, WI 53706

Dr. Robert D. Willis, ManTech Environmental, Inc., Research Triangle Park, NC 27711

Contributors and Reviewers

Dr. JoEllen Bandemeyer—Research Triangle Institute, PO Box 12194, Research Triangle Park,
NC 27709

Dr. Lyle Chinkin—Sonoma Technology, 1360 Redwood Way, Suite C, Petaluma, CA 94549

Dr. Candis S. Claiborn—Washington State University, Laboratory for Atmospheric Research,
Department of Civil and Environmental Engineering, P.O. Box 642910, Pullman, WA 99164

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. Steven Colome—Integrated Environmental Services, 5319 University Drive, #430, Irvine, CA 92612

Mr. Tom Coulter—National Exposure Research Laboratory (D234-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Edward O. Edney—National Exposure Research Laboratory (E205-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Terence Fitz-Simons—Office of Air Quality Planning and Standards (C304-01),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Sheldon Friedlander—University of California at Los Angeles, Department of Chemical
Engineering, 5531 Boelter Hall, Los Angeles, CA 90095

Dr. Lester D. Grant—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lynn Hildemann—Civil and Environmental Engineering Department, Stanford University,
Stanford, CA 94305

Mr. Phil Lorang—Office of Air Quality Planning and Standards (D205-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Karen Magliano—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

Mr. Scott Mathias—Office of Air Quality Planning and Standards (C539-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Nehzat Mottallebi—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

Mr. Tom Pace—Office of Air Quality Planning and Standards (D205-01), U.S. Environmental
Protection Agency, Research Triangle Park, NC 27711

Dr. Richard Poirot—VT Air Program, Building 3 South, 103 South Main Street, Waterbury, VT
05671

Mr. Win Setiawan—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers
(cont'd)

Dr. John J. Vandenberg—National Center for Environmental Assessment (8601D), U.S. Environmental Protection Agency, 1200 Pennsylvania Avenue, NW, Washington, DC 20460

Mr. Dane Westerdahl—California Air Resources Board, 2020 L Street, Sacramento, CA 95814

CHAPTER 4. ENVIRONMENTAL EFFECTS OF PARTICULATE MATTER

Principal Authors

Ms. Beverly Comfort—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. J.H.B. Garner—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lester D. Grant—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. David A. Grantz—University of California/Riverside, Kearney Agricultural Center, 9240 South Riverbend Avenue, Parlier, CA 93648

Dr. Brooke L. Hemming—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Dale W. Johnson—Environmental and Resource Science, 1000 Valley Road, University of Nevada, Reno, NV 89512

Dr. William Malm—National Park Service, Air Resources Division, CIRA, Colorado State University, Fort Collins, CO

Dr. Joseph P. Pinto—National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers

(cont'd)

Contributing Authors

Dr. Paul J. Hanson—Environmental Sciences Division, Oak Ridge National Laboratory,
P.O. Box 2008, Bethel Valley Road, Building 1059, Oak Ridge, TN 37831-6422

Dr. William H. Smith—Professor Emeritus, Yale University School of Forestry and
Environmental Studies, 370 Prospect Street, New Haven, CT 06511

Contributors and Reviewers

Dr. Larry T. Cupitt—National Exposure Research Laboratory (D305-01),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Rich Damberg—Office of Air Quality Planning and Standards (C539-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Russell R. Dickerson—University of Maryland, Department of Meteorology,
Stadium Drive, College Park, MD 20742

Dr. Anne Grambsch—National Center for Environmental Assessment (8601D),
U. S. Environmental Protection Agency, Washington, DC 20036

Dr. Sagar V. Krupa—University of Minnesota, Department of Plant Pathology,
495 Borlaug Hall, 1991 Upper Buford Circle, St. Paul, MN 55108

Dr. Alan J. Krupnick—Quality of the Environment Division, Resources for the Future,
1616 P Street, NW, Washington, DC 20036

Mr. Paul T. Roberts—Sonoma Technology, Inc., 1360 Redwood Way - Suite C,
Petaluma, CA 94954

Mr. John Spence—Cary, NC 27511

Dr. John J. Vandenberg—National Center for Environmental Assessment (8601D), U.S.
Environmental Protection Agency, 1200 Pennsylvania Avenue, NW, Washington, DC 20460

Ms. Debra Meyer Wefering—Duckterather Weg 61, Bergisch Gladbach, Germany 54169

Ms. Kay Whitfield—Office of Air Quality Planning and Standards (C243-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Richard Zepp—National Exposure Research Laboratory (IOD),
U. S. Environmental Protection Agency, Athens, GA

Authors, Contributors, and Reviewers
(cont'd)

***CHAPTER 5. HUMAN EXPOSURE TO PARTICULATE MATTER
AND ITS CONSTITUENTS***

Principal Authors

Dr. William E. Wilson—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Linda S. Sheldon—National Exposure Research Laboratory (E205-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Halûk Özkaynak—National Exposure Research Laboratory (E205-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Contributing Authors

Dr. Janet Burke—National Exposure Research Laboratory (E205-02),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Roy Fortmann—National Exposure Research Laboratory (E205-02),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. David T. Mage—Institute for Survey Research, Temple University,
Philadelphia, PA 19122-6099

Mr. Thomas McCurdy—National Exposure Research Laboratory (E205-02),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Gary Norris—National Exposure Research Laboratory (E205-03),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Anne Rea—National Exposure Research Laboratory (E205-04),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Alan Vette—National Exposure Research Laboratory (E205-04),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Authors, Contributors, and Reviewers
(cont'd)

Contributors and Reviewers

Dr. Lester D. Grant—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. John Langstaff—Office of Air Quality Planning and Standards (C539-01)
U.S. Environmental Protection Agency, Research Triangle park, NC 27711

Dr. Joseph Pinto—National Center for Environmental Assessment (B243-01),
U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Harvey Richmond—Office of Air Quality Planning and Standards (C539-01),
U. S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. John J. Vandenberg—National Center for Environmental Assessment (8601D), U.S.
Environmental Protection Agency, 1200 Pennsylvania Avenue, NW, Washington, DC 20460

**U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA
FOR PARTICULATE MATTER**

Executive Direction

Dr. Lester D. Grant—Director, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Scientific Staff

Dr. Robert W. Elias—PM Team Leader, Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. William E. Wilson—Air Quality Coordinator, Physical Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. James Brown—Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Beverly Comfort—Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. William Ewald—Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. J.H.B. Garner—Ecological Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Brooke Hemming—Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Dennis J. Kotchmar—Medical Officer, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Joseph P. Pinto—Physical Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Dr. Lori White—Health Scientist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

**U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA
FOR PARTICULATE MATTER**

(cont'd)

Technical Support Staff

Ms. Nancy Broom—Information Technology Manager, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Douglas B. Fennell—Technical Information Specialist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Emily R. Lee—Management Analyst, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Diane H. Ray—Program Specialist, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Ms. Donna Wicker—Administrative Officer, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Mr. Richard Wilson—Clerk, National Center for Environmental Assessment (B243-01), U.S. Environmental Protection Agency, Research Triangle Park, NC 27711

Document Production Staff

Ms. Carolyn T. Perry—Manager/Word Processor, Computer Sciences Corporation, 2803 Slater Road, Suite 220, Morrisville, NC 27560

Dr. Barbara Liljequist—Technical Editor, Computer Sciences Corporation, 2803 Slater Road, Suite 220, Morrisville, NC 27560

Dr. Carol A. Seagle—Technical Editor, Computer Sciences Corporation, 2803 Slater Road, Suite 220, Morrisville, NC 27560

Ms. Jessica Long—Graphic Artist, Computer Sciences Corporation, 2803 Slater Road, Suite 220, Morrisville, NC 27560

Mr. Matthew Kirk—Graphic Artist, Computer Sciences Corporation, 2803 Slater Road, Suite 220, Morrisville, NC 27560

**U.S. ENVIRONMENTAL PROTECTION AGENCY
PROJECT TEAM FOR DEVELOPMENT OF AIR QUALITY CRITERIA
FOR PARTICULATE MATTER**

(cont'd)

Document Production Staff

(cont'd)

Mr. John A. Bennett—Technical Information Specialist, Library Associates of Maryland,
11820 Parklawn Drive, Suite 400, Rockville, MD 20852

Ms. Sandra L. Hughey—Technical Information Specialist, Library Associates of Maryland,
11820 Parklawn Drive, Suite 400, Rockville, MD 20852

Ms. Rebecca Caffey—Records Management Technician, Reference Retrieval and Database
Entry Clerk, InfoPro, Inc., 8200 Greensboro Drive, Suite 1450, McLean, VA 22102

**U.S. ENVIRONMENTAL PROTECTION AGENCY
SCIENCE ADVISORY BOARD (SAB) STAFF OFFICE
CLEAN AIR SCIENTIFIC ADVISORY COMMITTEE (CASAC)
PARTICULATE MATTER REVIEW PANEL***

CASAC Chair

Dr. Philip Hopke—Bayard D. Clarkson Distinguished Professor, Department of Chemical Engineering, Clarkson University, Box 5708, Potsdam, NY 13699-5708

CASAC Members

Dr. Ellis Cowling—University Distinguished Professor At-Large, North Carolina State University, Colleges of Natural Resources and Agriculture and Life Sciences, North Carolina State University, 1509 Varsity Drive, Raleigh, NC 27695-7632

Dr. James D. Crapo—Chairman, Department of Medicine, National Jewish Medical and Research Center, 1400 Jackson Street, Denver, CO, 80206, and Chief Executive Officer (CEO) of Aeolus Pharmaceuticals, Inc.

Dr. Frederick J. Miller—Vice President for Research, CIIT Centers for Health Research, 6 Davis Drive, P.O. Box 12137, Research Triangle Park, NC 27709

Mr. Richard L. Poirot—Environmental Analyst, Air Pollution Control Division, Department of Environmental Conservation, Vermont Agency of Natural Resources, Bldg. 3 South, 103 South Main Street, Waterbury, VT 05671-0402

Dr. Frank Speizer—Edward Kass Professor of Medicine, Channing Laboratory, Harvard Medical School, 181 Longwood Avenue, Boston, MA 02115-5804

Dr. Barbara Zielinska—Research Professor, Division of Atmospheric Science, Desert Research Institute, 2215 Raggio Parkway, Reno, NV 89512-1095

* Members of this CASAC Panel consist of:

a. CASAC Members: Experts appointed to the statutory Clean Air Scientific Advisory Committee by the EPA Administrator; and

b. CASAC Consultants: Experts appointed by the SAB Staff Director to serve on one of the CASAC's National Ambient Air Quality Standards (NAAQS) Panels for a particular criteria air pollutant.

**U.S. ENVIRONMENTAL PROTECTION AGENCY
SCIENCE ADVISORY BOARD (SAB) STAFF OFFICE
CLEAN AIR SCIENTIFIC ADVISORY COMMITTEE (CASAC)
PARTICULATE MATTER REVIEW PANEL***
(cont'd)

CASAC Consultants

Dr. Jane Q. Koenig—Professor, Department of Environmental Health, School of Public Health and Community Medicine, University of Washington, Box 357234, Seattle, WA 98195-7234

Dr. Petros Koutrakis—Professor of Environmental Science, Environmental Health, School of Public Health, Harvard University, HSPH, 401 Park Dr., Room 410 West, Boston, MA 02215

Dr. Allan Legge—President, Biosphere Solutions, 1601 11th Avenue NW, Calgary, Alberta, CANADA T2N 1H1

Dr. Paul J. Liroy—Associate Director and Professor, Environmental and Occupational Health Sciences Institute, UMDNJ - Robert Wood Johnson Medical School, 170 Frelinghuysen Road, Rm 301, Piscataway, NJ 08854

Dr. Morton Lippmann—Professor, Nelson Institute of Environmental Medicine, New York University School of Medicine, 57 Old Forge Road, Tuxedo, NY 10987

Dr. Joe Mauderly—Vice President, Senior Scientist, and Director, National Environmental Respiratory Center, Lovelace Respiratory Research Institute, 2425 Ridgecrest Drive SE, Albuquerque, NM, 87108

Dr. Roger O. McClellan—Consultant, 1370 Quaking Aspen Pine, Albuquerque, NM, 87111

Dr. Gunter Oberdorster—Professor of Toxicology, Department of Environmental Medicine, School of Medicine and Dentistry, University of Rochester, 575 Elmwood Avenue, Box 850, Rochester, NY 14642

Dr. Robert D. Rowe—President, Stratus Consulting, Inc., , PO Box 4059, Boulder, CO 80306-4059

Dr. Jonathan M. Samet—Professor and Chair, Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, 615 N. Wolfe Street, Suite W6041, Baltimore, MD 21205-2179

Dr. Sverre Vedal—Professor of Medicine, School of Public Health and Community Medicine Department of Environmental and Occupational Health Sciences, University of Washington, 4225 Roosevelt Way NE, Suite 100, Seattle, WA 98105-6099

**U.S. ENVIRONMENTAL PROTECTION AGENCY
SCIENCE ADVISORY BOARD (SAB) STAFF OFFICE
CLEAN AIR SCIENTIFIC ADVISORY COMMITTEE (CASAC)
PARTICULATE MATTER REVIEW PANEL***
(cont'd)

CASAC Consultants
(cont'd)

Mr. Ronald White—Research Scientist, Epidemiology, Bloomberg School of Public Health, Room W6035, The Johns Hopkins University, 615 N. Wolfe St., Rm W6035, Baltimore, MD 21205

Dr. Warren H. White—Visiting Professor, Crocker Nuclear Laboratory, University of California-Davis, , Davis, CA 95616-8569

Dr. George T. Wolff—Principal Scientist, General Motors Corporation, 300 Renaissance Center (MC482-C27-B76), Detroit, MI 48265-3000

EPA Science Advisory Board Staff

Dr. Vanessa Vu—SAB Staff Office Director, EPA Science Advisory Board Staff Office (Mail Code 1400F), 1200 Pennsylvania Avenue, N.W., Washington DC, 20460

Mr. Fred Butterfield—CASAC Designated Federal Officer, EPA Science Advisory Board Staff Office (Mail Code 1400F), 1200 Pennsylvania Avenue, N.W., Washington, DC 20460

Abbreviations and Acronyms

AAS	atomic absorption spectrophotometry
AC	air conditioning
ACGIH	American Conference of Governmental Industrial Hygienists
ACS	American Cancer Society
ADS	annular denuder system
AES	atomic emission spectroscopy
AIRS	Aerometric Information Retrieval System
AM	alveolar macrophage
ANC	acid neutralizing capacity
APS	aerodynamic particle sizer
AQCD	Air Quality Criteria Document
AQI	Air Quality Index
ARIES	Aerosol Research and Inhalation Epidemiology Study
ASOS	Automated Surface Observing System
ATOFMS	aerosol time-of-flight mass spectrometry
AWOS	Automated Weather Observing System
BaP	benzo(a)pyrene
BASE	Building Assessment and Survey Evaluation
BC	black carbon
BNF	bacterial nitrogen fertilization
BOSS	Brigham Young University Organic Sampling System
BYU	Brigham Young University
CAA	Clean Air Act
CAAM	continuous ambient mass monitor
CAMNET	Coordinated Air Monitoring Network
CARB	California Air Resources Board

CASAC	Clean Air Scientific Advisory Committee
CASTNet	Clean Air Status and Trends Network
CC	carbonate carbon
CCPM	continuous coarse particle monitor
CCSEM	computer-controlled scanning electron microscopy
CEN	European Standardization Committee
CFCs	chlorofluorocarbons
CFR	Code of Federal Regulations
CHAD	Consolidated Human Activity Database
CI	confidence interval
CIF	charcoal-impregnated cellulose fiber; charcoal-impregnated filter
CMAQ	Community Model for Air Quality
CMB	chemical mass balance
CMSA	Consolidated Metropolitan Statistical Area
CO AQCD	Air Quality Criteria Document for Carbon Monoxide
COD	coefficient of divergence
COH	coefficient of haze
CONCDEP	average of unitless regional deposition and precipitation-weighted concentrations
COPD	chronic obstructive pulmonary disease
CPC	condensation particle counter
CRP	Coordinated Research Program
CSIRO	Commonwealth Scientific and Industrial Research Organisation
CSMCS	Carbonaceous Species Methods Comparison Study
CSS	coastal sage scrub
CTM	chemistry-transport model
CV	coefficient of variation
CVM	contigent valuation method
D_a	aerodynamic diameter

DAQ	Department of Air Quality
DMPS	differential mobility particle sizer
DMS	dimethyl sulfide
D_p	Stokes particle diameter
dv	deciview
EAD	electrical aerosol detector
EC	elemental carbon
EDXRF	energy dispersive X-ray fluorescence
EEA	essential ecological attribute
ENSO	El Niño-Southern Oscillation
EPEC	Ecological Processes and Effects Committee
ESP	electrostatic precipitator
ETS	environmental tobacco smoke
EXPOLIS	Air Pollution Exposure Distributions within Adult Urban Populations in Europe
FID	flame ionization detection
FRD	flame photometric detector
FRM	Federal Reference Method
GAM	general additive model
GC	gas chromatography
GCM	General Circulation Model
GCVTC	Grand Canyon Visibility Transport Commission
GC/MSD	gas chromatography/mass-selective detection
GHG	greenhouse gas
GSD	geometric standard deviation
HAPs	hazardous air pollutants
<i>H</i>	Henry's law constant
HBEF	Hubbard Brook Experimental Forest
HDS	honeycomb denuder/filter pack sampler

HEADS	Harvard-EPA Annular Denuder System
HEI	Health Effects Institute
HI	Harvard Impactor
hivol	high volume sampler
HPEM	Harvard personal environmental monitor
HTGC/MS	high temperature gas chromatography/mass spectrometry
HVAC	heating, ventilation, or air conditioning
IC	ion chromatography
ICP	inductively coupled plasma
IFS	Integrated Forest Study
IMPROVE	Interagency Monitoring of Protected Visual Environments
INAA	instrumental neutron activation analysis
IOVPS	integrated organic vapor/particle sampler
IP	inhalable particle
IPCC	Intergovernmental Panel on Climate Change
IPM	inhalable particulate matter
IPN	Inhalable Particulate Network
ISO	International Standards Organization
LAI	leaf area indices
LOD	level of detection
LWC	liquid water content
LWCA	liquid water content analyzer
MAA	mineral acid anion
MAACS	Metropolitan Acid Aerosol Characterization Study
MADPro	Mountain Acid Deposition Program
MAQSIP	Multiscale Air Quality Simulation Platform
MCM	mass concentration monitor
MHVOC	more highly volatile organic compound

MDL	minimum detection level
MOUDI	microorifice uniform deposit impactor; multiple orifice uniform deposit impactor
MS	mass spectroscopy
MSA	methane sulfonic acid
MSA	metropolitan statistical area
mv	motor vehicle
NAAQS	National Ambient Air Quality Standards
NAMS	National Ambient Monitoring Stations
NAPAP	National Acid Precipitation Assessment Program
NARSTO	North American Research Strategy for Tropospheric Ozone
NCEA	National Center for Environmental Assessment
NDDN	National Dry Deposition Network
NERL	National Exposure Research Laboratory
NESCAUM	Northeast States for Coordinated Air Use Management
NFRAQS	North Frontal Range Air Quality Study
NHAPS	National Human Activity Pattern Survey
NIOSH	National Institute for Occupational Safety and Health
NIR	near infrared radiation
NIST	National Institute of Standards and Technology
NO _x	nitrogen oxides
Nr	reactive nitrogen
NRC	National Research Council
NuCM	nutrient cycling model
OAQPS	Office of Air Quality Planning and Standards
OC	organic carbon
ORD	Office of Research and Development
PAH	polycyclic aromatic hydrocarbon; polynuclear aromatic hydrocarbon
PAN	peroxyacetyl nitrate

PAR	photosynthetically active radiation
PBL	planetary boundary layer
PBPs	primary biological particles
PBW	particle-bound water
PBY	Presbyterian Home
PC	particle concentrator
PC	pyrolytic carbon
PC-BOSS	Particulate Concentrator-Brigham Young University Organic Sampling System
PCA	principal component analysis
PCBs	polychlorinated biphenyls
PCDD	polychlorinated dibenzo- <i>p</i> -dioxins
PCDF	polychlorinated dibenzofurans
PCM	particle composition monitor
pDR	personal DataRAM
PEM	personal exposure monitor
PESA	proton (or particle) elastic scattering analysis
PIXE	proton (or particle) induced X-ray emission
PM	particulate matter
PM _{2.5}	fine particulate matter
PM ₁₀	combination of coarse and fine particulate matter
PM ₁₀₋₂₅	coarse particulate matter
PM ₉₀	90th percentile difference in concentration
PM ACQD	Air Quality Criteria Document for Particulate Matter
PMF	positive matrix factorization
POP	persistent organic pollutant
PRB	policy-relevant background
PTEAM	Particle Total Exposure Assessment Methodology
PTEP	PM ₁₀ Technical Enhancement Program

PTFE	polytetrafluoroethylene (Teflon)
PUF	polyurethane foam
PWC	precipitation-weighted concentrations
r	Pearson correlation coefficient
RAMS	Regional Air Monitoring Study
RAMS	real-time total ambient mass sampler
RAPS	Regional Air Pollution Study
RAS	Roll-Around System
RCS	Random Component Superposition
RFO	residual fuels oils
RH	relative humidity
RIVM	Dutch National Institute of Public Health and the Environment
ROS	reactive oxygen species
RPM	reconstructed particulate mass
RPM	respirable particulate matter
RRMS	relatively remote monitoring sites
RSP	respirable particulate matter
RSP	respirable suspended particles
RTI	Research Triangle Institute
RTP	Research Triangle Park
RUE	radiation use efficiency
SA	Sierra Anderson
SAB	Science Advisory Board
SCAQ5	Southern California Air Quality Study
SCENES	Subregional Cooperative Electric Utility, Department of Defense, National Park Services, and Environmental Protection Agency Study
SCOS97	1997 Southern California Ozone Study
sd, SD	standard deviation
sec	secondary

SEM	scanning electron microscopy
SES	sample equilibration system
SEV	Sensor Equivalent Visibility
SHEDS	Stochastic Human Exposure and Dose Simulation
SIP	State Implementation Plan
SLAMS	State and Local Air Monitoring Stations
SMPS	scanning mobility particle sizer
SMSAs	Standard Metropolitan Statistical Areas
SoCAB	South Coast Air Basin
SOPM	secondary organic particulate matter
SO _x	sulfur oxides
SRM	standard reference method
STN	Speciation Trends Network
SUVB	solar ultraviolet B radiation
SVOC	semivolatile organic compound
SVM	semivolatile material
S-XRF	synchrotron induced X-ray fluorescence
TAR	Third Assessment Report
TC	total carbon
TDMA	Tandem Differential Mobility Analyzer
TEO	trace element oxides
TEOM	tapered element oscillating microbalance
THEES	Total Human Environmental Exposure Study
TMO	thermal manganese oxidation
TNF	tumor necrosis factor
TO	thermal-optical
TOFMS	time-of-flight mass spectroscopy
TOR	thermal-optical reflectance

TOT	thermal-optical transmission
TPM	thoracic particulate matter
TRM	traffic-related microenvironment
TRXRF	total reflection X-ray fluorescence
TSP	total suspended particulate
TVOC	total volatile organic compounds
UAM-IV	Urban Airshed Model Version IV
UCM	unresolved complex mixture
UNEP	United Nations Environment Programme
URG	University Research Glassware
VAPS	Versatile Air Pollution Sampler
VMD	volume mean diameter
VOC	volatile organic compound
WMO	World Meteorological Organization
VR	visual range
WINS	Well Impactor Ninety-Six
WRAC	Wide Range Aerosol Classifier
WTA	willing to accept
WTP	willing to pay
XAD	polystyrene-divinyl benzene
XRF	X-ray fluorescence

1. INTRODUCTION

This document is an update of “*Air Quality Criteria for Particulate Matter*” published by the U.S. Environmental Protection Agency (EPA) in 1996, and it will serve as the basis for Congressionally-mandated periodic review by EPA of the National Ambient Air Quality Standards for Particulate Matter (PM NAAQS). The present document critically assesses the latest scientific information relative to determining the health and welfare effects associated with the presence of various concentrations of PM in ambient air. It builds upon the assessment in the previous 1996 EPA Air Quality Criteria Document for Particulate Matter (1996 PM AQCD; U.S. Environmental Protection Agency, 1996a) by focusing on assessment and integration of information most relevant to PM NAAQS criteria development, based on pertinent peer-reviewed literature published or accepted for publication mainly through 2002, as well as a few more recent important studies. This introductory chapter presents (1) background information that summarizes legislative requirements, the criteria and NAAQS review process, the history of PM NAAQS reviews including the chronology of changes in key elements of the standards, and the coordinated PM research program that has guided much of the more recent research in this area; (2) an overview of the current PM criteria and NAAQS review (including key milestones), as well as assessment approaches and procedures used in preparing this document; and (3) an orientation to the general organizational structure of this document.

1.1 BACKGROUND

1.1.1 Legislative Requirements

As indicated in U.S. Code (1991), Sections 108 and 109 of the U.S. Clean Air Act (CAA) (42 U.S.C. Sections 7408 and 7409) govern the establishment, review, and revision of NAAQS. Section 108(a) directs the EPA Administrator to list pollutants, which, in the Administrator’s judgment, cause or contribute to air pollution that may reasonably be anticipated to endanger public health or welfare and to issue air quality criteria for them. The air quality criteria are to

reflect the latest scientific information useful in indicating the kind and extent of all identifiable effects on public health and welfare that may be expected from the presence of the pollutant in ambient air.

Section 109 directs the Administrator of EPA to propose and promulgate “primary” and “secondary” NAAQS for criteria air pollutants listed under Section 108. Section 109(b)(1) defines a primary standard as a level of air quality, the attainment and maintenance of which, in the judgment of the Administrator, based on the criteria and allowing for an adequate margin of safety, is requisite to protect the public health. Section 109(b)(2) defines a secondary standard as one which, in the judgment of the Administrator, based on the criteria, is requisite to protect public welfare from any known or anticipated adverse effects associated with the presence of the pollutant in ambient air. Welfare effects, defined in Section 302(h), include (but are not limited to) effects on soils, water, crops, vegetation, man-made materials, animals, wildlife, weather, visibility, and climate; damage to and deterioration of property; hazards to transportation; and effects on economic values, personal comfort, and well-being. Section 109(d)(1) requires the periodic review and, as appropriate, revision of existing criteria and standards. Also, Section 109(d)(2) requires an independent committee of non-EPA experts, the Clean Air Scientific Advisory Committee (CASAC), to provide advice and recommendations to the EPA Administrator regarding the scientific soundness and appropriateness of criteria and NAAQS for PM and other criteria air pollutants (i.e., ozone, nitrogen dioxide, sulfur oxides, carbon monoxide, lead).

1.1.2 Criteria and NAAQS Review Process

The EPA’s periodic reviews of criteria and NAAQS for a given criteria air pollutant progress through a number of steps, beginning with the preparation of an Air Quality Criteria Document (AQCD) by EPA’s National Center for Environmental Assessment Division in Research Triangle Park, NC (NCEA-RTP), which critically assesses the scientific information upon which the NAAQS are to be based. Building upon the AQCD, staff within EPA’s Office of Air Quality Planning and Standards (OAQPS) prepare a Staff Paper that evaluates the policy implications of the key studies and scientific information contained in the AQCD and presents

staff conclusions and recommendations of options for the Administrator to consider. The Staff Paper is intended to help “bridge the gap” between the scientific assessment contained in the AQCD and the judgments required of the Administrator in determining whether it is appropriate to retain or to revise the NAAQS. Iterative drafts of both of the AQCD and the Staff Paper (as well as any other analyses supporting the Staff Paper, such as exposure and/or risk assessments) are made available for public comment and are reviewed by CASAC. Following CASAC review of these documents, the Administrator proposes decisions on whether to retain or revise the NAAQS based on the information in these documents, taking into account CASAC advice and recommendations and public comments. The Administrator’s proposed decisions are published in the *Federal Register*, with a preamble that presents the rationale for the decisions and solicits public comment. After taking into consideration comment received on the proposed decisions, the Administrator’s final decisions are promulgated in a final *Federal Register* notice that addresses significant comments received on the proposal.

Decisions on the NAAQS involve consideration of the four basic elements of a standard: *indicator, averaging time, form, and level*. The indicator defines the pollutant to be measured in the ambient air for the purpose of determining compliance with the standard. The averaging time defines the time period over which air quality measurements are to be obtained and averaged, considering evidence of effects associated with various time periods of exposure. The form of a standard defines the air quality statistic that is to be compared to the level of the standard (i.e., an ambient concentration of the indicator pollutant) in determining whether an area attains the standard. The form of the standard specifies the air quality measurements that are to be used for compliance purposes (e.g., the 98th percentile of an annual distribution of daily concentrations; the annual arithmetic average), the monitors from which the measurements are to be obtained (e.g., one or more population-oriented monitors in an area), and whether the statistic is to be averaged across multiple years. These basic elements of a standard are the primary focus of the staff conclusions and recommendations in the Staff Paper and in the subsequent rulemaking, building upon the policy-relevant scientific information assessed in the AQCD and on the policy analyses contained in the Staff Paper. These four elements taken together determine the degree of public health and welfare protection afforded by the NAAQS.

1.1.3 History of Earlier PM Criteria and NAAQS Reviews

Selection of appropriate indicator(s) for the PM NAAQS has long posed a unique challenge, in that unlike the other criteria air pollutants, PM is the generic term for a broad class of physically and chemically diverse substances that exist in ambient air as discrete particles (liquid droplets or solids) over a wide range of sizes. These airborne particles originate from a variety of stationary and mobile sources. Primary particles are emitted directly into ambient air; whereas secondary particles are formed in the atmosphere by transformation of gaseous emissions such as sulfur oxides (SO_x), nitrogen oxides (NO_x), and volatile organic compounds (VOCs). The physical and chemical properties of PM vary greatly with time, region, meteorology, and source category, thus complicating assessment of ambient PM health and welfare effects. Particles in ambient air are usually distributed bimodally in two somewhat overlapping size categories: (1) fine (diameter generally < 2.5 μm) and (2) coarse (diameter generally > 1.0 μm). Particles in these two size fractions tend to differ in terms of formation mechanisms, sources of origin, composition, and behavior in the atmosphere and human respiratory tract.

EPA first promulgated primary and secondary NAAQS for PM on April 30, 1971 (Federal Register, 1971). These standards measured PM as “total suspended particulate” (TSP), which refers to ambient PM up to a nominal size of 25 to 45 μm. The primary standards for PM (measured as TSP) were 260 μg/m³ (24-h average), not to be exceeded more than once per year, and 75 μg/m³ (annual geometric mean). The secondary standard (measured as TSP) was 150 μg/m³ (24-h average), not to be exceeded more than once per year.

EPA completed the next review of PM air quality criteria and standards in July 1987, revising the 1971 standards to protect against adverse health effects of inhalable airborne particles that can be deposited in the lower (thoracic) regions of the human respiratory tract, with “PM₁₀” (i.e., those particles collected by a sampler with a specified penetration curve yielding an upper 50% cut-point of 10-μm aerodynamic diameter) as the indicator (Federal Register, 1987). EPA established identical primary and secondary PM₁₀ standards for two averaging times: 150 μg/m³ (24-h average, with no more than one expected exceedance per year) and 50 μg/m³ (expected annual arithmetic mean, averaged over three years).

1.1.4 The 1997 PM NAAQS Revision

The last previous review of the air quality criteria and standards for PM was initiated in April 1994 by EPA announcing its intention to develop a revised *Air Quality Criteria for Particulate Matter*. Several workshops were held in November 1994 and January 1995 to discuss important new health effects information useful in preparing initial PM AQCD draft materials. Also, plans for review of the PM criteria and standards under a highly accelerated, court-ordered schedule were presented by EPA at a CASAC public meeting in December 1994. A court order entered in *American Lung Association v. Browner*, CIV-93-643-TUC-ACM (U.S. District Court of Arizona, 1995), as subsequently modified, required publication of EPA's final decision on the review of the PM NAAQS by July 19, 1997.

Several external review drafts of the revised PM AQCD were prepared by NCEA-RTP and made available for public comment and CASAC review (at public meetings held in August 1995, December 1995, and February 1996). The CASAC completed its review of the PM AQCD in early 1996, advising the EPA Administrator in a March 15, 1996 letter (Wolff, 1996) that "although our understanding of the health effects of PM is far from complete, a revised Criteria Document which incorporates the Panel's latest comments will provide an adequate review of the available scientific data and relevant studies of PM." Revisions made in response to public and CASAC comments were then incorporated by NCEA-RTP, as appropriate, into the final 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). The associated PM Staff Paper, prepared by OAQPS staff, drew upon the 1996 PM AQCD and other assessments to pose options for the Administrator to consider in making PM NAAQS decisions. Drafts of the PM Staff Paper also underwent public comment and CASAC review, which informed the final 1996 version (U.S. Environmental Protection Agency, 1996b).

The 1996 PM AQCD and PM Staff Paper provide detailed information on atmospheric formation, ambient concentrations, and health effects of ambient air PM, as well as quantitative estimates of human health risks associated with exposure to ambient PM. The main focus of these documents was on recent epidemiologic evidence reporting associations between ambient concentrations of PM and a range of serious health effects. Special attention was given to several size-specific classes of particles, including PM_{10} and the principal fractions of PM_{10} ,

referred to as the fine ($PM_{2.5}$) and coarse ($PM_{10-2.5}$) fractions. $PM_{2.5}$ refers to those particles collected by samplers having penetration curves yielding an upper 50% cut point of 2.5 μm aerodynamic diameter. $PM_{10-2.5}$ refers to those particles in an aggregate sample having an upper 50% cut point of 10 μm and a lower 50% cut point of 2.5 μm aerodynamic diameter. In other words, the coarse fraction ($PM_{10-2.5}$) refers to inhalable particles that remain if fine ($PM_{2.5}$) particles are removed from a sample of PM_{10} particles. As discussed in the 1996 PM AQCD, fine and coarse fraction particles can be differentiated by their sources and formation processes, by their chemical and physical properties, and by their behavior in the atmosphere.

Taking into account information and assessments presented in the 1996 PM AQCD and Staff Paper, CASAC advice and recommendations, and public comments received on proposed revisions to the PM NAAQS published in December 1996 (Federal Register, 1996), the EPA Administrator promulgated significant revisions to the PM NAAQS in July 1997 (Federal Register, 1997). In that decision, the PM NAAQS were revised in several respects. While it was determined that the PM NAAQS should continue to focus on particles less than or equal to 10 μm in diameter, it was also determined that the fine and coarse fractions of PM_{10} should be considered separately. New standards were added, using $PM_{2.5}$ as the indicator for fine particles; and PM_{10} standards were retained for the purpose of regulating coarse-fraction particles. Two new $PM_{2.5}$ standards were set: an annual standard of 15 $\mu g/m^3$, based on the 3-year average of annual arithmetic mean $PM_{2.5}$ concentrations from single or multiple community-oriented monitors; and a 24-h standard of 65 $\mu g/m^3$, based on the 3-year average of the 98th percentile of 24-h $PM_{2.5}$ concentrations at each population-oriented monitor within an area. To continue to address coarse-fraction particles, the annual PM_{10} standard was retained, and the form, but not the level, of the 24-h PM_{10} standard was revised to be based on the 99th percentile of 24-h PM_{10} concentrations at each monitor in an area. The secondary standards were revised by making them identical in all respects to the $PM_{2.5}$ and PM_{10} primary standards.

Following promulgation of the revised PM NAAQS, legal challenges were filed by a large number of parties, addressing a broad range of issues. In May 1998, the U.S. Court of Appeals for the District of Columbia Circuit issued an initial opinion that upheld EPA's decision to establish fine particle standards, holding that such standards were amply justified by the growing

body of empirical evidence demonstrating a relationship between fine particle pollution and adverse health effects. American Trucking Associations v. Browner, 175 F. 3d 1027, 1055-56 (D.C. Cir. 1999a) (rehearing granted in part and denied in part, 195 F. 3d 4 (D.C. Cir. 1999b), affirmed in part and reversed in part, Whitman v. American Trucking Associations, 531 U.S. 457 (2001). Further, the court found “ample support” for EPA’s decision to regulate coarse fraction particles, although it vacated the revisions to the 1987 PM₁₀ standards on the basis of PM₁₀ being a “poorly matched indicator for coarse particulate pollution” because PM₁₀ includes fine particles (Id. at 1053-55). As a result of this aspect of the court’s ruling, which EPA did not appeal, the 1987 PM₁₀ standards remain in effect.

In addition, the court broadly held that EPA’s approach to establishing the level of the standards in its 1997 decisions on both the PM and Ozone NAAQS (which were promulgated on the same day and considered together by the court in this aspect of its opinion) effected “an unconstitutional delegation of legislative authority” (Id. at 1034-40). EPA appealed this aspect of the court’s ruling to the U.S. Supreme Court. In February 2001, the U.S. Supreme Court unanimously reversed the Court of Appeals’ ruling on the constitutional issue and sent the case back to the Court of Appeals for resolution of any remaining issues that had not been addressed in that court’s earlier rulings. Whitman v. American Trucking Associations, 531 U.S. 457, 475-76 (2001). In March 2002, the Court of Appeals rejected all remaining challenges to the standards, finding that the 1997 PM_{2.5} standards were reasonably supported by the record and were not “arbitrary or capricious.” American Trucking Associations v. EPA, 283 F. 3d 355, 369-72 (D.C. Cir. 2002). Thus, the 1997 PM_{2.5} standards also remain in effect.

1.1.5 Coordinated PM Research Program

Shortly after promulgation of the 1997 PM NAAQS decisions, NCEA-RTP published a PM Health Risk Research Needs Document (U.S. Environmental Protection Agency, 1998a) that identified research needed to improve scientific information supporting future reviews of the PM NAAQS. The document provided a foundation for PM research coordination among Federal agencies and other research organizations, as well as input to National Research Council (NRC) deliberations on PM research. The Office of Research and Development (ORD) of EPA also

moved quickly to broaden its ongoing PM research activities by developing, in partnership with other Federal agencies, a coordinated interagency PM research program. This interagency program has focused and continues to focus mainly on expanding scientific knowledge of ambient PM exposure and health effects, as well as on developing improved monitoring methods and cost-effective mitigation strategies. The interagency effort also promotes substantially expanded coordination with other research organizations, including the Health Effects Institute (HEI) and other state-, university-, and industry-sponsored research groups. Beginning in the fall of 1997, public participation was and continues to be encouraged through workshops and review of EPA's PM Research Program documentation.

In response to Congressional requirements in EPA's Fiscal Year 1998 Appropriation, the NRC established its Committee on Research Priorities for Airborne Particulate Matter in January 1998. This NRC PM Research Committee's charge was to identify the most important research priorities relevant to setting primary (health-based) PM NAAQS, to develop a conceptual plan for PM research, and to monitor EPA's research progress toward improved understanding of the relationship between PM and public health. The NRC PM Research Committee issued a series of reports (National Research Council, 1998, 1999, 2001) which recommended that expanded PM research efforts be planned and carried out in relation to a general conceptual framework, as shown in Figure 1-1. That framework essentially calls for research aimed at (a) identifying sources of airborne particles or gaseous precursor emissions and characterization of processes involved in atmospheric transformation, transport, and fate of ambient PM; (b) delineating temporal and spatial patterns of air quality indicators (e.g., $PM_{2.5}$, $PM_{10-2.5}$, PM_{10} mass concentrations) of ambient PM and apportionment of observed variations in such ambient PM indicators to various emission sources; (c) characterizing human exposures to ambient PM as one important component of total personal exposure to particles, as modified by time-activity patterns and varying microenvironmental exposure to particles of indoor or ambient origin; (d) characterizing resulting respiratory tract deposition, clearance, retention, and disposition of inhaled particles, as determinants of dose to target tissues (e.g., locally in the lungs or via systemic translocation to the heart or other organs); and (e) delineating of mechanisms of damage and repair plausibly leading to (f) human health responses, as extrapolated from or




Figure 1-1. A general framework for integrating PM research. Note that this figure is not intended to represent a framework for research management. Such a framework would include multiple pathways for the flow of information.

Source: National Research Council (2001).

quantified by experimental animal or human exposure (toxicology) studies and/or observational (epidemiology) studies.

Research conducted under a PM Research Program structured in relation to the conceptual framework shown in Figure 1-1 would be expected to (a) reduce key scientific uncertainties regarding interrelationships between PM sources, ambient concentrations, exposures, dose to target tissues, and resulting health effects and thereby (b) improve the scientific underpinnings for both current and future periodic PM NAAQS reviews. Table 1-1 highlights some types of key uncertainties identified by the NRC PM Research Committee in relation to elements of the source-to-response conceptual framework illustrated in Figure 1-1. The NRC Committee went on to delineate a series of 10 research topics that they recommended be addressed by an expanded PM research program aimed at answering a set of broadly stated questions, as shown in Table 1-2.

The EPA PM Research Program, structured to address topics shown in Table 1-2, has encompassed studies to improve understanding of the formation and composition of fine PM, to improve the measurement and estimation of population exposures to ambient PM, to delineate characteristics or components of PM that are responsible for its health effects, to evaluate factors that increase susceptibility to PM effects in some subpopulations, and to elucidate mechanisms

TABLE 1-1. KEY SCIENTIFIC UNCERTAINTIES RELATED TO THE SOURCE-TO-RESPONSE FRAMEWORK

Source	→	Concentration (or other indicator)
<ul style="list-style-type: none"> • Contribution of various emission sources to ambient and indoor particulate matter concentrations • Relative contribution of various sources to the most toxic components of particulate matter 		
Concentration (indicator)	→	Exposure
<ul style="list-style-type: none"> • Relationship between ambient (indoor) particulate matter and the composition of particles to which people are exposed • Contribution of ambient particulate matter to total personal exposure for: <ul style="list-style-type: none"> – Susceptible subpopulations – General population • Variation in relationship of ambient particulate matter concentrations to human exposure by place • Variation in contribution of ambient particulate matter concentrations to total human exposure over time • Covariance of particulate matter exposures with exposures to other pollutants • Relationships between outdoor ambient and personal exposures for particulate matter and co-pollutants 		
Exposure	→	Dose
<ul style="list-style-type: none"> • Relationship between inhaled concentration and dose of particulate matter and constituents at the tissue level in susceptible subjects <ul style="list-style-type: none"> – Asthma – Chronic obstructive pulmonary disease (COPD) – Heart disease – Age: infants and elderly – Others 		
Dose	→	Response
<ul style="list-style-type: none"> • Mechanisms linking morbidity and mortality to particulate matter dose to or via the lungs <ul style="list-style-type: none"> – Inflammation – Host defenses – Neural mechanisms 		

Source: National Research Council (2001).

TABLE 1-2. RESEARCH TOPICS AND QUESTIONS RECOMMENDED BY NATIONAL RESEARCH COUNCIL (NRC) TO BE ADDRESSED BY EXPANDED PM RESEARCH PROGRAM

RESEARCH TOPIC 1.	OUTDOOR MEASURES VERSUS ACTUAL HUMAN EXPOSURES
	<ul style="list-style-type: none"> • <i>What are the quantitative relationships between concentrations of particulate matter and gaseous co-pollutants measured at stationary outdoor air monitoring sites and the contributions of these concentrations to actual personal exposures, especially for subpopulations and individuals?</i>
RESEARCH TOPIC 2.	EXPOSURES OF SUSCEPTIBLE SUBPOPULATIONS TO TOXIC PARTICULATE MATTER COMPONENTS
	<ul style="list-style-type: none"> • <i>What are the exposures to biologically important constituents and specific characteristics of particulate matter that cause responses in potentially susceptible subpopulations and the general population?</i>
RESEARCH TOPIC 3.	CHARACTERIZATION OF EMISSION SOURCES
	<ul style="list-style-type: none"> • <i>What are the size distribution, chemical composition, and mass-emission rates of particulate matter emitted from the collection of primary-particle sources in the United States, and what are the emissions of reactive gases that lead to secondary particle formation through atmospheric chemical reactions?</i>
RESEARCH TOPIC 4.	AIR-QUALITY MODEL DEVELOPMENT AND TESTING
	<ul style="list-style-type: none"> • <i>What are the linkages between emission sources and ambient concentrations of the biologically important components of particulate matter?</i>
RESEARCH TOPIC 5.	ASSESSMENT OF HAZARDOUS PARTICULATE MATTER COMPONENTS
	<ul style="list-style-type: none"> • <i>What is the role of physicochemical characteristics of particulate matter in eliciting adverse health effects?</i>
RESEARCH TOPIC 6.	DOSIMETRY: DEPOSITION AND FATE OF PARTICLES IN THE RESPIRATORY TRACT
	<ul style="list-style-type: none"> • <i>What are the deposition patterns and fate of particles in the respiratory tract of individuals belonging to presumed susceptible subpopulations?</i>
RESEARCH TOPIC 7.	COMBINED EFFECTS OF PARTICULATE MATTER AND GASEOUS POLLUTANTS
	<ul style="list-style-type: none"> • <i>How can the effects of particulate matter be disentangled from the effects of other pollutants? How can the effects of long-term exposure to particulate matter and other pollutants be better understood?</i>
RESEARCH TOPIC 8.	SUSCEPTIBLE SUBPOPULATIONS
	<ul style="list-style-type: none"> • <i>What subpopulations are at increased risk of adverse health outcomes from particulate matter?</i>
RESEARCH TOPIC 9.	MECHANISMS OF INJURY
	<ul style="list-style-type: none"> • <i>What are the underlying mechanisms (local pulmonary and systemic) that can explain the epidemiologic findings of mortality/morbidity associated with exposure to ambient particulate matter?</i>
RESEARCH TOPIC 10.	ANALYSIS AND MEASUREMENT
	<ul style="list-style-type: none"> • <i>To what extent does the choice of statistical methods in the analysis of data from epidemiologic studies influence estimates of health risks from exposures to particulate matter? Can existing methods be improved? What is the effect of measurement error and misclassification on estimates of the association between air pollution and health?</i>

Source: National Research Council (2001).

by which these effects are produced. The results from these efforts, and related efforts by other Federal agencies and the general scientific community during the past several years, have substantially enhanced the scientific and technical bases for future decisions on the PM NAAQS.

1.2 CURRENT PM CRITERIA AND NAAQS REVIEW

1.2.1 Key Milestones and Procedures for Document Preparation

As with other NAAQS reviews, critical assessment of relevant scientific information is presented in this updated PM AQCD. Key milestones in the development of the present PM AQCD are shown in Table 1-3 and discussed below. It is important to note that development of the document involved substantial external peer and public review through (a) public workshops involving the general aerosol scientific community, (b) iterative reviews of successive drafts by CASAC, and (c) comments from the public on successive drafts. This final document reflects extensive external input received through these reviews and serves to ensure that the review of the PM standards is based on critical assessment of the latest available pertinent science.

The procedures for developing this updated PM AQCD built on the experience derived from the most recent previous PM, Ozone, and CO AQCD preparation efforts. Briefly, the respective responsibilities for production of the present PM AQCD are as follows. An NCEA-RTP PM Team was responsible for developing and implementing a project plan for preparing the PM AQCD, taking into account inputs from individuals in other EPA program and policy offices identified as part of the EPA PM Work Group. The resulting project plan (i.e., the PM Document Development Plan) was discussed with CASAC in May 1998 and was appropriately revised. A literature search was ongoing throughout the preparation of this document to identify, to the extent possible, all pertinent PM literature published since early 1996. Additionally, EPA published in the Federal Register (1) a request for information asking for recently available research information on PM that may not yet be published and (2) a request for individuals with the appropriate type and level of expertise to contribute to the writing of PM AQCD materials to identify themselves (U.S. Environmental Protection Agency, 1998b). The specific authors of chapters or sections of the proposed document were selected on the basis of their expertise on

TABLE 1-3. KEY MILESTONES IN THE DEVELOPMENT OF THIS DOCUMENT

<i>Key Milestones</i>	<i>Dates</i>
PM NAAQS Review Plan to CASAC	October 1997
Prepare AQCD Development Plan	November 1997 to January 1998
Begin Literature Search	February 1998
Federal Register Call for Information/Sources Sought	April 1998
CASAC Meeting on PM AQCD Development Plan	May 1998
Prepare Workshop Drafts of Chapters	May to December 1998
Peer Review Workshop	April 1999
Prepare First External Review Draft PM AQCD	March to September 1999
First External Review Draft PM AQCD	October 1999
Public Comment Period on First Draft	October 1999 to January 2000
CASAC Meeting on First Draft	December 1999
Second External Review Draft PM AQCD	March 2001
Public Comment Period on Second Draft	April to July 2001
CASAC Meeting on Second Draft	July 2001
Third External Review Draft PM AQCD	April 2002
Public Comment Period on Third Draft	May to July 2002
CASAC Meeting on Third Draft	July 2002
Fourth External Review Draft PM AQCD	June 2003
Public Comment Period on Fourth Draft	June to August 2003
CASAC Meeting on Fourth Draft	August 2003
CASAC Consultation on Proposed Revisions to Chapter 9	October 2003
CASAC Consultation on Revisions to Chapters 7 and 8	November 2003
Revised Draft Chapters 7 and 8	December 2003
Public Comment Period on Revised Draft Chapters 7 and 8	January 2004
CASAC Teleconference Review of Revised Draft Chapters 7 and 8	February 2004
Revised Draft Chapters 7, 8 and 9	June 2004
Public Comment Period on Revised Draft Chapters 7, 8 and 9	July 2004
CASAC Meeting on Revised Draft Chapters 7, 8 and 9	July 2004
Revised Draft Chapter 9	August 2004
Public Comment Period on Revised Chapter 9	September 2004
CASAC Review of Revised Chapter 9	September 2004
Final PM AQCD	October 2004

the subject areas and their familiarity with the relevant literature; and these included both EPA and non-EPA scientific experts. The project team defined critical issues and topics to be addressed by the authors and provided direction in order to emphasize evaluation of those studies most clearly identified as important for standard setting. It should be noted that materials contributed by non-EPA authors were, at times, modified by EPA PM Team staff in response to internal and/or external review comments, e.g., by the public or CASAC, and that EPA is responsible for the ultimate content of this PM AQCD.

The main focus of this document is the evaluation and interpretation of pertinent atmospheric science information, air quality data, human exposure information, and health and welfare effects information newly published since that assessed in the 1996 PM AQCD. Initial draft versions of AQCD chapters were evaluated via expert peer-review workshop discussions and/or written peer reviews that focused on the selection of pertinent studies to be included in the chapters, the potential need for additional information to be added to the chapters, and the quality of the characterization and interpretation of the literature. The authors of the draft chapters then revised them on the basis of workshop and/or written expert review comments. These and other integrative materials were incorporated into the First External Review Draft of the PM AQCD (October 1999), which was made available for public comment and was the subject of consultation with CASAC at a December 1999 public meeting.

In order to foster timely presentation and publication of newly emerging PM research findings, EPA cosponsored and helped to organize an Air and Waste Management Association International Speciality Conference, entitled “PM 2000: Particulate Matter and Health,” held in January 2000 in Charleston, SC. The conference was cosponsored in cooperation with several other government agencies and/or private organizations that also fund PM research. Topics covered included new research results concerning the latest advances in PM atmospheric sciences (e.g., PM formation, transport, transformation), PM exposure, PM dosimetry and extrapolation modeling, PM toxicology (e.g., mechanisms, laboratory animal models, human clinical responses), and PM epidemiology. The main purpose of the conference was to facilitate having the latest scientific information available in time for incorporation as quickly as possible into the Second External Review Draft of the PM AQCD. Hence, arrangements were made for

scientists to submit written manuscripts on papers or posters presented at the PM 2000 Conference for expedited peer-review by several major journals, so that decisions on acceptance for publication could be made by mid-2000. The evaluations and findings set forth in the Second External Review Draft (March 2001) included consideration of such published PM 2000 papers, as well as extensive additional information published elsewhere since that addressed in the First External Review Draft. The Second External Review Draft was made available for public comment and was reviewed by CASAC at a July 2001 public meeting.

Further revisions were then incorporated into the Third External Review Draft (April 2002) to reflect both public comment and CASAC review of the Second Draft, as well as assessment of additional extensive new information published since that addressed in the Second Draft. Shortly after EPA released the Third External Review Draft in May 2002 for public comment and CASAC review, the HEI announced that researchers at Johns Hopkins University had discovered problems with applications of statistical software used in a number of important epidemiologic studies on links between ambient air PM and mortality and morbidity effects. The Third External Review Draft was reviewed by CASAC at a July 2002 public meeting, although it was recognized that discussion of the epidemiology information would need to be further revised after the newly surfaced statistical issues were appropriately addressed. In response to the surfacing of such statistical issues, which affected numerous PM time-series studies that used General Additive Models (GAM) and were published post-1995, EPA took steps in consultation with CASAC to identify particularly policy-relevant studies and to encourage researchers to reanalyze affected studies and to submit them expeditiously for peer review by a special expert panel convened by HEI. The results of reanalyses for more than 30 studies were peer-reviewed and published, along with commentary by the HEI expert peer-review panel, in an HEI Special Report (Health Effects Institute, May 2003).

Discussion of the newly addressed statistical issues and reanalyses results was incorporated into the Fourth External Review Draft (June 2003), which was made available for public comment and was reviewed by CASAC at an August 2003 public meeting. The Fourth Draft also incorporated changes made in response to earlier public comments and CASAC reviews, including pertinent peer-reviewed literature published or accepted for publication mainly

through April 2002. CASAC completed its review of Chapters 1 through 6 of the Fourth Draft PM AQCD at the August 2003 meeting, but comments on the remaining chapters (7, 8, and 9) were judged to warrant further revision and review of those chapters. EPA subsequently consulted with CASAC and made available revised Chapters 7, 8, and 9 for public comment and CASAC review, as indicated in Table 1-3. The CASAC completed its review in September, 2004, as indicated in an October 4, 2004 letter (Hopke, 2004) to the EPA Administrator. Revisions made in response to final CASAC and public comments were then incorporated by the NCEA-RTP PM team, as appropriate, into this final 2004 PM AQCD (U.S. Environmental Protection Agency, 2004) released in October, 2004.

1.2.2 Assessment Approaches

The assessment presented in this document is framed by: (1) the selection of pertinent issues to be addressed; (2) the selection of relevant studies and an approach to the presentation of information drawn from those studies; and (3) the selection of an approach to interpreting and integrating the body of evidence evaluated in the document.

As an initial matter, the NCEA-RTP PM team focused on selecting pertinent issues to be addressed in this assessment. Preliminary issues were drawn from among those highlighted in the 1996 PM AQCD and Staff Paper and in CASAC and public reviews of those documents, the 1997 PM NAAQS promulgation process, and the 1998 PM Health Risk Research Needs Document (U.S. Environmental Protection Agency, 1998a). Further identification and refinement of issues resulted from the NRC review and reports on PM research priorities, as discussed in Section 1.1.5 above. The CASAC review of the PM AQCD Development Plan and public comments on early draft AQCD materials at various stages of their development also provided important inputs regarding issue identification. The issues selected are reflected throughout this document and are most concisely identified in the introductory section at the beginning of each of the ensuing chapters.

The selection of relevant studies to be included in this assessment was based on a detailed review of new research published in the peer-reviewed literature since early 1996, including materials accepted for publication mainly through April 2002 (and, thus, appearing mostly

during 2002). Limited coverage of some more recent studies is also included as deemed appropriate in light of their special importance. For example, information derived from the HEI Special Report (Health Effects Institute, May 2003), discussed above in Section 1.2.1, has been integrated into this assessment. Literature discussed in detail in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) generally is not discussed in depth in this document; rather key findings from the 1996 review are concisely summarized as appropriate. Limited treatment is included, however, for some earlier studies judged to be of particular importance in this review of the PM NAAQS. Key literature is summarized in tables, whereas text discussions focus mainly on evaluation and integration of the literature, including discussion of alternative points of view where scientific controversy exists. This approach to study selection and presentation reflects recommendations from CASAC aimed at development of a more concise document than the 1996 PM AQCD.

The scientific assessments presented in this document are primarily driven by the large body of available epidemiologic evidence evaluating associations between ambient PM, alone and in combination with other air pollutants, and various health endpoints. In such a case, an approach to interpreting the evidence most fundamentally focuses on characterizing the causal significance of observed associations. In so doing, it is appropriate to consider various aspects of the evidence of associations, in particular drawing from those presented earlier in Hill's classic monograph (Hill, 1965) and extensively used by the scientific community in conducting such evidence-based reviews, e.g., in the 2004 Surgeon General's Report on smoking (Centers for Disease Control and Prevention, 2004). As discussed in Chapters 8 and 9, a number of these aspects are judged to be particularly salient in evaluating the body of evidence available in this review, including the aspects described by Hill as strength, consistency, temporality, biologic gradient, experiment, plausibility, and coherence. These interrelated aspects are considered in the evaluation of epidemiologic evidence presented in Chapter 8 and are also more broadly addressed in the Chapter 9 integrative synthesis.

1.3 DOCUMENT ORGANIZATION

This document is basically organized to assess information related to topics along the same flow of issues presented in the NRC conceptual framework shown in Figure 1-1, including information related to effects on both human health and the environment. The document consists of nine chapters presented in two volumes. Volume I contains this general introduction (Chapter 1), as well as Chapters 2 through 5. Chapters 2 and 3 provide background information on the physical and chemical properties of PM and related compounds; sources and emissions; atmospheric transport, transformation, and fate of PM; methods for the collection and measurement of PM; and U.S. ambient air PM concentrations. Chapter 4 assesses welfare-related PM effects on vegetation and ecosystems, visibility, and man-made materials, as well as climate-related effects (including effects on atmospheric transmission of solar radiation), and it includes limited information on economic impacts of some welfare effects. Chapter 5 discusses factors affecting exposure of the general population to ambient PM.

The second volume contains Chapters 6 through 9. Chapter 6 assesses information concerning dosimetry of inhaled particles in the respiratory tract. Chapter 7 assesses the toxicology of specific types of PM constituents and potential mechanisms of action, based primarily on laboratory animal studies and controlled human exposure studies. Chapter 8 assesses the epidemiologic literature. Lastly, Chapter 9 integrates key information on PM-related health effects, drawing from assessments in prior chapters of the literature on exposure, dosimetry, toxicology, and epidemiology, as well as highlighting key information regarding important welfare effects associated with ambient PM.

REFERENCES

- Centers for Disease Control and Prevention. (2004) The health consequences of smoking: a report of the Surgeon General. Atlanta, GA: U.S. Department of Health and Human Services, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. Available: http://www.cdc.gov/tobacco/sgr/sgr_2004/chapters.htm (18 August, 2004).
- Federal Register. (1971) National primary and secondary ambient air quality standards. F. R. (April 30) 36: 8186-8201.
- Federal Register. (1987) Revisions to the national ambient air quality standards for particulate matter. F. R. (July 1) 52: 24,634-24,669.
- Federal Register. (1996) National ambient air quality standards for particulate matter; proposed rule. F. R. (December 13) 61: 65,638-65,713.
- Federal Register. (1997) National ambient air quality standards for particulate matter; final rule. F. R. (July 18) 62: 38,652-38,752.
- Health Effects Institute. (2003) Revised analyses of time-series studies of air pollution and health. Boston, MA: Health Effects Institute; special report.
- Hill, A. B. (1965) The environment and disease: association or causation? *Proc. R. Soc. Med.* 58: 295-300.
- Hopke, P. K. (2004) Clean Air Scientific Advisory Committee (CASAC) Particulate Matter (PM) Review Panel's ongoing peer review of the agency's draft Air Quality Criteria for Particulate Matter [letter to Michael O. Leavitt, Administrator, U.S. EPA]. Washington, DC: U.S. Environmental Protection Agency, Clean Air Scientific Advisory Committee; EPA-SAB-CASAC-04-009; October 4.
- National Research Council. (1998) Research priorities for airborne particulate matter. I. Immediate priorities and a long-range research portfolio. Washington, DC: National Academy Press. Available: <http://www.nap.edu/catalog/6131.html> (4 June 2003).
- National Research Council. (1999) Research priorities for airborne particulate matter. II. Evaluating research progress and updating the portfolio. Washington, DC: National Academy Press. Available: <http://www.nap.edu/books/0309066387/html/> (4 June 2003).
- National Research Council. (2001) Research priorities for airborne particulate matter. III. Early research progress. Washington, DC: National Academy Press. Available: <http://www.nap.edu/books/0309073375/html/> (4 June 2003).
- U.S. Code. (1991) Clean Air Act, §108, air quality criteria and control techniques, §109, national ambient air quality standards. U. S. C. 42: §§7408-7409.
- U.S. Court of Appeals for the District of Columbia. (1999a) American Trucking Associations, Inc. vs. U.S. Environmental Protection Agency. 175 F3d 1027 (D.C. Cir. 1999).
- U.S. Court of Appeals for the District of Columbia. (1999b) American Trucking Associations, Inc. vs. U.S. Environmental Protection Agency. 195 F.3d 4 (D.C. Cir. 1999), affirmed in part, reversed in part, and remanded..
- U.S. Court of Appeals for the District of Columbia. (2002) American Trucking Associations, Inc. vs. U.S. Environmental Protection Agency. 283 F.3d 355, 378-79 (D.C. Cir. 2002).
- U.S. District Court of Arizona. (1995) American Lung Association v. Browner. West's Federal Supplement 884 F.Supp. 345 (No. CIV 93-643 TUC ACM).
- U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1996b) Review of the national ambient air quality standards for particulate matter: policy assessment of scientific and technical information. OAQPS staff paper. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/452/R-96-013. Available from: NTIS, Springfield, VA; PB97-115406REB.
- U.S. Environmental Protection Agency. (1998a) Particulate matter research needs for human health risk assessment to support future reviews of the national ambient air quality standards for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/R-97/132F.
- U.S. Environmental Protection Agency. (1998b) Review of national ambient air quality standards for particulate matter. *Commer. Bus. Daily*: February 19. Available: <http://cbdnet.access.gpo.gov/index.html> [1999, November 24].

U.S. Supreme Court. (2001) *Whitman v. American Trucking Association*. 531 U.S. 457 (nos. 99-1257 and 99-1426).
Wolff, G. T. (1996) Closure by the Clean Air Scientific Advisory Committee (CASAC) on the draft Air Quality
Criteria for Particulate Matter [letter to Carol M. Browner, Administrator, U.S. EPA]. Washington, DC:
U.S. Environmental Protection Agency, Clean Air Scientific Advisory Committee.; EPA-SAB-CASAC-LTR-
96-005; March 15.

2. PHYSICS, CHEMISTRY, AND MEASUREMENT OF PARTICULATE MATTER

Chapter 3 of the 1996 EPA document Air Quality Criteria for Particulate Matter (1996 PM AQCD; U.S. Environmental Protection Agency, 1996a) contained an extensive review of the physics and chemistry of airborne particulate matter (PM). Chapter 2 of this revised version of the PM AQCD also provides background information on the physics and chemistry of atmospheric particles, information useful in understanding the subsequent chapters. The chapters in this document are organized to generally follow the sequence of key elements that make up the risk assessment framework described in Chapter 1 (Section 1.2.2) beginning with sources and continuing to effects as shown in Figure 1-1. Thus, this chapter discusses new background information useful in evaluating PM effects on human health and welfare and in preparing related risk assessments used to support PM standard-setting decisions. Information important for implementation of PM standards, but not essential to the standard setting process, is not the focus in this chapter. The reader is referred to the NARSTO (North American Research Strategy for Tropospheric Ozone) Fine Particle Assessment (NARSTO, 2003) for information relevant to air quality management for PM.

Unlike other criteria pollutants (O_3 , CO, SO_2 , NO_2 , and Pb), PM is not a specific chemical entity but is a mixture of particles from different sources and of different sizes, compositions, and properties. Emphasis is placed here on discussion of differences between fine and coarse particles and differences between ultrafine particles and accumulation-mode particles within fine particles.

Since PM is defined quantitatively by measurement techniques, it will be useful to discuss our understanding of the relationships between PM suspended in the atmosphere, PM inhaled by people, and PM measured by various sampling and analytical techniques. Chapter 4 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) contained a review of the state of the art of PM measurement technology. Since that time, considerable progress has been made in understanding problems in the measurement of PM mass, chemical composition, and physical parameters. Progress has also been made in developing new and improved measurement techniques, especially for continuous measurements. Therefore, a more extensive survey on

measurement problems and on newly developed measurement techniques is included here in Section 2.2. For more detail and older references, the reader is referred to Chapters 3 and 4 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).

2.1 PHYSICS AND CHEMISTRY OF PARTICULATE MATTER

2.1.1 Basic Concepts

Atmospheric particles originate from a variety of sources and possess a range of morphological, chemical, physical, and thermodynamic properties. Examples of atmospheric particles include combustion-generated particles, such as diesel soot or fly ash; photochemically produced particles, such as those found in urban haze; salt particles formed from sea spray; and soil-like particles from resuspended dust. Some particles are liquid; some are solid. Others may contain a solid core surrounded by liquid. Atmospheric particles contain inorganic ions, metallic compounds, elemental carbon, organic compounds, and crustal compounds. Some atmospheric particles are hygroscopic and contain particle-bound water. The organic fraction is especially complex, containing hundreds (probably thousands) of organic compounds. (See Appendix 3C for information on the composition of the organic fraction and the concentration of specific organic compounds.) Primary particles are emitted directly from sources; whereas secondary particles are formed from gases through chemical reactions in the atmosphere involving atmospheric oxygen (O_2) and water vapor (H_2O); reactive species such as ozone (O_3); radicals such as the hydroxyl ($\bullet OH$) and nitrate ($\bullet NO_3$) radicals; and pollutants such as sulfur dioxide (SO_2), nitrogen oxides (NO_x), and organic gases from natural and anthropogenic sources. The particle formation process includes nucleation of particles from low-vapor pressure gases emitted from sources or formed in the atmosphere by chemical reactions, condensation of low-vapor pressure gases on existing particles, and coagulation of particles. Thus, any given particle may contain PM from many sources. Because a particle from a given source is likely to be composed of a mixture of chemical components and because particles from different sources may coagulate to form a new particle, atmospheric particles may be considered a mixture of mixtures. The composition and behavior of particles are fundamentally linked with those of the surrounding gas. An aerosol may be defined as a suspension of solid or liquid particles in air. The term aerosol includes both the particles and all vapor or gas phase components of air.

However, the term aerosol is sometimes used to refer to the suspended particles only. In this document, “particulate” is used only as an adjective, as in particulate matter.

A complete description of the atmospheric aerosol would include an accounting of the chemical composition, morphology, and size of each particle, as well as the relative abundance of each particle type as a function of particle size (Friedlander, 1970). However, the physical and chemical characteristics of particles are usually measured separately. Size distributions by particle number used to calculate surface area and volume distributions often are determined by physical means, such as electrical mobility, aerodynamic behavior, or light scattering. Chemical composition usually is determined by analysis of collected samples, although some species can be measured in situ. The mass and average chemical composition of particles segregated according to aerodynamic diameter by cyclones or impactors can also be determined. However, recent developments in single particle analysis techniques by electron microscopy with X-ray analysis of single particles (but not agglomerates) collected on a substrate or by mass spectroscopy of individual suspended particles provide elemental composition of individual particles by particle size and, thus, are bringing the description envisioned by Friedlander closer to reality.

2.1.2 Physical Properties and Processes

2.1.2.1 Definitions of Particle Diameter

The diameter of a spherical particle may be determined by optical or electron microscopy, by light scattering and Mie theory, by its electrical mobility, or by its aerodynamic behavior. However, atmospheric particles often are not spherical. Therefore, their diameters are described by an “equivalent” diameter (i.e., the diameter of a sphere that would have the same physical behavior). An optical diameter is the diameter of a spherical particle, with the same refractive index as the particle used to calibrate the optical particle sizer, that scatters the same amount of light into the solid angle measured. Diffusion and gravitational settling are important physical behaviors for particle transport, collection, and removal processes, including deposition in the respiratory tract. Different equivalent diameters are used depending on which process is more important. For smaller particles, diffusion is more important and the Stokes diameter is often used. For larger particles, gravitational setting is more important and the aerodynamic diameter is often used.

The Stokes diameter, D_p , describes particle size based on the aerodynamic drag force imparted on a particle when its velocity differs from that of the surrounding fluid. For a smooth, spherically shaped particle, D_p exactly equals the physical diameter of the particle. For irregularly shaped particles, D_p is the diameter of an equivalent sphere that would have the same aerodynamic resistance. Electrical mobility analyzers classify particles according to their electrical mobility. Particles of equal Stokes diameters that carry the same electric charge will have the same electrical mobility. Hence, for spherical particles, the electrical mobility diameter would equal the Stokes diameter. The mobility diameter can be considered the diameter of a spherical particle that would have the same electrical mobility. The particle mobility can be related to the particle diffusion coefficient and Brownian diffusion velocity through the Stokes-Einstein equation. Thus, the Stokes diameter is the appropriate parameter for particle behavior governed by diffusion. The Stokes diameter, D_p , is used in size distributions based on light scattering and mobility analysis. The Stokes diameter is independent of density.

The aerodynamic diameter, D_a , however, depends on particle density. It is defined as the diameter of a spherical particle with an equal gravitational settling velocity but a material density of 1 g/cm^3 . Cascade impactors separate particles based on their aerodynamic diameter, and aerodynamic particle sizers measure the aerodynamic diameter. Respirable, thoracic, and inhalable sampling and $\text{PM}_{2.5}$ and PM_{10} sampling are based on particle aerodynamic diameter. For particles greater than about $0.5 \text{ }\mu\text{m}$, the aerodynamic diameter is generally the quantity of interest. For smaller particles, the Stokes diameter may be more useful. Particles with the same physical size and shape but different densities will have the same Stokes diameter but different aerodynamic diameters.

The aerodynamic diameter, D_a , is related to the Stokes diameter, D_p , by:

$$D_a = D_p \left(\frac{\rho C_p}{C_a} \right)^{1/2} \quad (2-1)$$

where ρ is the particle density, and C_p and C_a are the Cunningham slip factors evaluated for the particle diameters D_p and D_a respectively. The slip factor is a function of the ratio between particle diameter and mean free path of the suspending gas ($0.066 \text{ }\mu\text{m}$ for air at one atmosphere pressure and $20 \text{ }^\circ\text{C}$). C is an empirical factor that accounts for the reduction in the drag force on

particles due to the “slip” of the gas molecules at the particle surface. C is an important factor for particles less than $1\ \mu\text{m}$ in diameter, for which the surrounding air cannot be modeled by a continuous fluid. For large particles ($D_p > 5\ \mu\text{m}$) $C = 1$; while for smaller particles $C > 1$.

For particles with diameters greater than the mean free path, λ , the aerodynamic diameter given by equation (2-1) is approximated by:

$$D_a = D_p (\rho)^{1/2} \quad (\text{for } D_p \gg \lambda) \quad (2-2)$$

This expression, which shows that aerodynamic diameter is directly proportional to the square root of the particle density, is often used for particles as small as $0.5\ \mu\text{m}$. For particles with diameters much smaller than the mean free path, the slip factor must be taken into account. In this case, the aerodynamic diameter is directly proportional to the particle density,

$$D_a = (\rho) D_p \quad (\text{for } D_p \ll \lambda) \quad (2-3)$$

Detailed definitions of the various sizes and their relationships are given in standard aerosol textbooks (e.g., Friedlander [2000], Reist [1984, 1993], Seinfeld and Pandis [1998], Hinds [1999], Vincent [1989, 1995], Willeke and Baron [1993], Baron and Willeke [2002], and Fuchs [1964, 1989]).

2.1.2.2 Aerosol Size Distributions

Particle size, as indexed by one of the “equivalent” diameters, is an important parameter in determining the properties, effects, and fate of atmospheric particles. The atmospheric deposition rates of particles and, therefore, their residence times in the atmosphere are a strong function of their Stokes and aerodynamic diameters. Particle diameters also influence the deposition patterns of particles within the lung. Because light scattering is strongly dependent on the optical particle size, the amount of light scattering per unit PM mass will be dependent on the size distribution of atmospheric particles. Therefore, the effects of atmospheric particles on visibility, radiative balance, and climate will be influenced by the size distribution of the

particles. Studies using cascade impactors or cyclones measure the particle-size distribution directly in aerodynamic diameter. The diameters of atmospheric particles range from 1 nm to 100 μm , spanning 5 orders of magnitude. A variety of different instruments, measuring a variety of equivalent diameters, are required to cover this range.

Older particle counting studies used optical particle counters to cover the range of 0.3 to 30 μm diameter. Diameters of particles below 0.5 μm were measured as mobility diameters. The particle diameters used in size distribution graphs from these studies usually are given as physical or Stokes diameters rather than aerodynamic diameters. In recent years, aerodynamic particle sizers have been developed that give a direct measurement of the aerodynamic diameter in the range of approximately 0.7 to 10 μm diameter. These instruments have been used with electrical mobility analyzers that measure the mobility diameter of particles from 3 nm to approximately 0.5 μm (McMurry, 2000). Unfortunately, there is no agreed-upon technique for combining the various equivalent diameters. Some workers use various assumptions to combine the various measurements into one presentation; others report each instrument separately. Therefore, the user of size distribution data should be careful to determine exactly which equivalent diameter is reported.

Particle Size Distribution Functions

The distribution of particles with respect to size is an important physical parameter governing particle behavior. Because atmospheric particles cover several orders of magnitude in particle size, size distributions often are expressed in terms of the logarithm of the particle diameter on the X-axis and the measured differential concentration on the Y-axis:

$\Delta N/\Delta(\log D_p)$ = the number of particles per cm^3 of air having diameters in the size range from $\log D_p$ to $\log(D_p + \Delta D_p)$. Because logarithms do not have dimensions, it is necessary to think of the distribution as a function of $\log(D_p/D_{p0})$, where the reference diameter $D_{p0} = 1 \mu\text{m}$ is not explicitly stated. If $\Delta N/\Delta(\log D_p)$ is plotted on a linear scale, the number of particles between D_p and $D_p + \Delta D_p$ is proportional to the area under the curve of $\Delta N/\Delta(\log D_p)$ versus $\log D_p$. Similar considerations apply to distributions of surface, volume, and mass. When approximated by a function, the distributions are usually given as $dN/d(\log D_p)$ rather than $\Delta N/\Delta(\log D_p)$.

Atmospheric Aerosol Size Distributions

Whitby (1978) published an analysis of over 1,000 particle size distributions measured at various locations in the United States. Figure 2-1 shows the number, surface area, and volume distributions for the grand average continental size distribution. Volume, surface area, and number distributions are plotted on an arithmetic scale such that the volume, surface area, or number of particles in any specified size range is proportional to the corresponding area under the curve. These distributions show that most of the particles are quite small, below 0.1 μm ; whereas most of the particle volume (and therefore most of the mass) is found in particles $> 0.1 \mu\text{m}$. Other averaged atmospheric size distributions are shown in Figures 2-2 and 2-3 (Whitby, 1978; Whitby and Sverdrup, 1980). Figures 2-2a and 2-2b describe the number of particles as a function of particle diameter for rural, urban-influenced rural, urban, and freeway-influenced urban aerosols. For some of the same data, the particle volume distributions are shown in Figures 2-3a and 2-3b. Whitby (1978) observed that the size distributions typically had three peaks which he called “modes.” The entire size distribution could be characterized well by a trimodal model consisting of three additive log-normal distributions. The mode with a peak between 5 and 30 μm diameter formed by mechanical processes was called the coarse particle mode; the mode with a peak between 0.15 and 0.5 μm formed by condensation and coagulation was called the accumulation mode; and the mode with a peak between 0.015 and 0.04 μm whose size was influenced by nucleation as well as by condensation and coagulation was called the transient nuclei or Aitken nuclei range, subsequently shortened to the nuclei mode. The nuclei mode could be seen in the number and surface distribution but only in special situations was it noticeable in the mass or volume distributions. The accumulation and nuclei modes taken together were called fine particles. An experimental size distribution showing modes and formation mechanisms is given in Figure 2-4. This size distribution was measured in traffic. Therefore, the nuclei mode is clearly separated from the accumulation mode and larger than it would be in size-distributions measured farther from sources of nuclei-mode particles.

Whitby (1978) concluded

“The distinction between ‘fine particles’ and ‘coarse particles’ is a fundamental one. There is now an overwhelming amount of evidence that not only are two modes in the mass or volume distribution usually observed, but that these fine and coarse modes are usually chemically quite different. The physical separation of




Figure 2-1. Distribution of coarse (c), accumulation (a), and nuclei (n) mode particles by three characteristics: (a) number, N; (b) surface area, S; and (c) volume, V for the grand average continental size distribution. DGV = geometric mean diameter by volume; DGS = geometric mean diameter by surface area; DGN = geometric mean diameter by number; D_p = particle diameter.

Source: Whitby (1978).




Figure 2-2. Particle size distributions by number: (a) number concentrations are shown on a logarithmic scale to display the wide range by site and size; (b) number concentrations for the average urban distribution are shown on a linear scale. For the linear scale, the area under any part of the curve is proportional to particle number in that size range.

Source: Whitby (1978); Whitby and Sverdrup (1980).

the fine and coarse modes originates because condensation produces fine particles while mechanical processes produce mostly coarse particles . . . the dynamics of fine particle growth ordinarily operate to prevent the fine particles from growing larger than about 1 μm . Thus, the fine and coarse modes originate separately, are transformed separately, are removed separately, and are usually chemically different . . . practically all of the sulfur found in atmospheric aerosol is found in the fine particle fraction. Thus, the distinction between fine and coarse fractions is of fundamental importance to any discussion of aerosol physics, chemistry, measurement, or aerosol air quality standards.”

Whitby’s (1978) conclusions were based on extensive studies of size distributions in a number of western and midwestern locations during the 1970s (Whitby et al., 1974; Willeke and Whitby, 1975; Whitby, 1978; Wilson et al., 1977; Whitby and Sverdrup, 1980). No size




Figure 2-3. Size distribution by volume for the averaged (a) rural and urban-influenced rural number distribution shown in Figure 2-2a and a distribution from south-central New Mexico, and (b) urban and freeway-influenced urban number distributions shown in Figure 2-2a.

Source: Whitby and Sverdrup (1980); Kim et al. (1993) and south-central New Mexico.

distribution studies of similar scope have been published since then. Newer results from particle counting and impactor techniques, including data from Europe (U.S. Environmental Protection Agency, 1996a) and Australia (Keywood et al., 1999, 2000), show similar results for the accumulation and coarse modes. Extensive measurements of particle size distributions, as part of the EPA's Supersites program, are providing considerable new data for analysis.

Whitby's (1978) conclusions have held up remarkably well. However, ideas about the sub-0.1 μm diameter range have changed somewhat as newer instruments provided measurements extending to smaller sizes and with greater resolution in size and time (McMurry et al., 2000). Depending on the source, temperature, saturated vapor pressure of the components,




Figure 2-4. Volume size distribution, measured in traffic, showing fine and coarse particles and the nuclei and accumulation modes of fine particles. DGV (geometric mean diameter by volume, equivalent to volume median diameter) and σ_g (geometric standard deviation) are shown for each mode. Also shown are transformation and growth mechanisms (e.g., nucleation, condensation, and coagulation).

Source: Adapted from Wilson et al. (1977) and Wilson and Suh (1997).

and the age of the aerosol, size distributions have been observed with peaks (including multiple peaks) throughout the sub-0.1 μm diameter size range. Sub-0.1 μm diameter peaks have been observed in rural areas (O'Dowd, 2002) as well as for brief periods (nucleation bursts) in urban areas (Woo et al., 2001a). Based on these and other observations, discussed in detail in Section 2.1.2.3, aerosol scientists now classify particles in the sub-0.1 μm size range as *ultrafine particles* and divide this size range into a *nucleation region* (< 10 nm) and an *Aitken (nuclei) region* (10 to 100 nm), as shown in Figure 2-5. Other studies, discussed in detail in the 1996 PM




Figure 2-5. Sub-micrometer number size distribution observed in a boreal forest in Finland showing the trimodal structure of fine particles. The total particle number concentration was 1,011 particles/cm³ (10-min average).

Source: Mäkelä et al. (1997).

AQCD (U.S. Environmental Protection Agency, 1996a), have shown that in fog or clouds or at very high relative humidities the accumulation mode may split into a larger size (more hygroscopic or droplet) submode and a smaller size (less hygroscopic or condensation) submode.

Definitions of Particle Size Fractions

In the preceding discussion several subdivisions of the aerosol size distribution were identified. Aerosol scientists use several different approaches or conventions in the classification of particles by size. These include: (1) modes, based on the observed size distributions and formation mechanisms; (2) dosimetry or occupational health sizes, based on the entrance into various compartments of the respiratory system; and (3) cut point, usually based on the 50% cut point of the specific sampling device, including legally specified, regulatory cut points for air quality standards.

Modal. The modal classification as first proposed by Whitby (1978) is shown in Figures 2-1 and 2-4. The newer modes introduced since 1978 are shown in Figure 2-5. An idealized distribution showing all four modes is shown in Figure 2-6. The nucleation and Aitken modes are best observed in the number distribution. However, the Aitken mode can be seen in the volume distribution in traffic or near traffic or other sources of ultrafine particles (Figures 2-3b and 2-4). The observed modal structure is frequently approximated by several lognormal distributions. Definitions of terms used to describe size distributions in modal terms are given below.




Figure 2-6. An idealized size distribution, that might be observed in traffic, showing fine and coarse particles and the nucleation, Aitken, and accumulation modes that comprise fine particles. Also shown are the major formation and growth mechanisms of the four modes of atmospheric particles.

Nucleation Mode: Freshly formed particles with diameters below about 10 nm, observed during active nucleation events. The lower limit, where particles and molecular clusters or large molecules overlap, is uncertain. Current techniques limit measurements to particles 3 nm or greater.

Aitken Mode: Larger particles with diameters between about 10 and 100 nm. The Aitken mode may result from growth of smaller particles or nucleation from higher concentrations of precursors.

Accumulation Mode: Particles with diameters from about 0.1 μm to just above the minimum in the mass or volume distributions which usually occurs between 1 and 3 μm .

Fine Particles: Fine particles include the nucleation, Aitken, and accumulation modes, i.e., particles from the lowest measurable size, currently about 3 nm, to just above the minimum in the mass or volume distribution which generally occurs between 1 and 3 μm .

Coarse Mode or Coarse Particles: Particles with diameters mostly greater than the minimum in the particle mass or volume distributions, which generally occurs between 1 and 3 μm .

Ultrafine Particles: Ultrafine particles are not a mode. In the air pollution literature, they are generally defined by size alone, i.e., particles with diameters of 0.1 μm (100 nm) or less. They include the nucleation mode and much of the Aitken mode. They may also be defined as particles whose properties differ from those of the bulk material because of their small size.

Modes are defined primarily in terms of their formation mechanisms but also differ in sources, composition, transport and fate, as well as in size. Nucleation mode applies to newly formed particles which have had little chance to grow by condensation or coagulation. Aitken mode particles are also recently formed particles that are still actively undergoing coagulation. However, because of higher concentrations of precursors or more time for condensation and

coagulation, Aitken particles have grown to larger sizes. Fine particles grow by coagulation (two particles combining to form one) or by condensation (low-equilibrium vapor pressure gas molecules condensing on a particle). As the particle size increases, the rate of growth by coagulation and condensation decreases and particles “accumulate” in the accumulation mode size range. Thus, accumulation-mode particles normally do not grow into the coarse particle size range. However, during conditions of high relative humidity, hygroscopic accumulation mode particles grow in size, increasing the overlap of fine and coarse particles. The accumulation mode may split into a (hygroscopic) droplet mode and a (non-hygroscopic) condensation mode. In addition, gas-phase pollutants may dissolve and react in the particle-bound water of hygroscopic particles, leading to an increase in the dry size. The combination of nucleation, Aitken, and accumulation modes are called fine particles (or sometimes fine-mode particles). Fine particles are formed primarily by combustion or chemical reactions of gases yielding products with low saturated vapor pressures. Fine particles are composed of metals (and metal oxides), black or elemental carbon, primary and secondary organic compounds, and sulfate, nitrate, ammonium and hydrogen ions.

The coarse mode refers to particles formed by the mechanical breakdown of minerals, crustal material, and organic debris. In addition to primary minerals and organic material, the coarse mode may include sea salt, nitrate formed from the reaction of nitric acid with sodium chloride, and sulfate formed from the reaction of sulfur dioxide with basic particles. The accumulation mode and the coarse mode overlap in the region between 1 and 3 μm (and occasionally over an even larger range). In this region, the chemical composition of individual particles can usually, but not always, allow identification of a source or formation mechanism, permitting identification of a particle as belonging to the accumulation or coarse mode.

Occupational Health or Dosimetric Size Cuts. The occupational health community has defined size fractions in terms of their entrance into various compartments of the respiratory system. This convention classifies particles into inhalable, thoracic, and respirable particles according to their upper size cuts. Inhalable particles enter the respiratory tract, beginning with the head airways. Thoracic particles travel past the larynx and reach the lung airways and the gas-exchange regions of the lung. Respirable particles are a subset of thoracic particles that are more likely to reach the gas-exchange region of the lung. In the past, exact definitions of these

terms have varied among organizations. As of 1993, a unified set of definitions was adopted by the American Conference of Governmental Industrial Hygienists (ACGIH, 1994), the International Standards Organization (ISO), and the European Standardization Committee (CEN). The curves which define inhalable (IPM), thoracic (TPM), and respirable (RPM) particulate matter are shown in Figure 2-7. These curves should not be taken to indicate that particles $> 4 \mu\text{m } D_a$ do not reach the gas exchange regions or that particles $< 4 \mu\text{m } D_a$ do not deposit in the bronchi. See Figure 6-13 in Chapter 6 for a graphical characterization of particle deposition in regions of the respiratory system as a function of particle size.




Figure 2-7. Specified particle penetration (size-cut curves) through an ideal (no-particle-loss) inlet for five different size-selective sampling criteria. Regulatory size cuts are defined in the Code of Federal Regulations (PM_{2.5} [2001c], PM₁₀ [2001a]). PM_{2.5} is also defined in the Federal Register (1997). Size-cut curves for inhalable particulate matter (IPM), thoracic particulate matter (TPM) and respirable particulate matter (RPM) size cuts are computed from definitions given by American Conference of Governmental and Industrial Hygienists (1994).

Size-Selective Sampling. Another set of definitions of particle size fractions arises from considerations of size-selective sampling. Size-selective sampling refers to the collection of particles below or within a specified aerodynamic size range. Size fractions are usually specified by the 50% cut point size; e.g., PM_{2.5} refers to particles collected by a sampling device that collects 50% of 2.5 μm particles and rejects 50% of 2.5 μm particles. However, size fractions are defined not merely by the 50% cut point, but by the entire penetration curve. Examples of penetration curves are given in Figure 2-7. Thus, as shown by Figure 2-7, a PM_{2.5} sampler, as defined by the Federal Reference Method, rejects 94% of 3 μm particles, 50% of 2.5 μm particles, and 16% of 2 μm particles. Samplers with the same 50% cut point, but differently shaped penetration curves, would collect different fractions of PM. Size-selective sampling has arisen in an effort to measure particle size fractions with some special significance (e.g., health, visibility, source apportionment, etc.), to measure mass size distributions, or to collect size-segregated particles for chemical analysis. Dichotomous samplers split the particles into smaller and larger fractions that may be collected on separate filters. However, some fine particles (≈10%) are collected with the coarse particle fraction. Cascade impactors use multiple size cuts to obtain a distribution of size cuts for mass or chemical composition measurements. One-filter samplers with a variety of upper size cuts are also used, e.g., PM_{2.5}, PM₁₀.

An idealized particle size distribution with the normally observed division of ambient aerosols into fine and coarse particles and the size fractions collected by the WRAC, TSP, PM₁₀, PM_{2.5} and PM_{10-2.5} samplers is shown in Figure 2-8. PM₁₀ samplers, as defined in Appendix J to Title 40 Code of Federal Regulations (40 CFR), Part 50 (Code of Federal Regulations, 2001a; Federal Register, 1987), collect all of the fine-mode particles and part of the coarse-mode particles. The upper cut point is defined as having a 50% collection efficiency at 10 ± 0.5 μm aerodynamic diameter. The slope of the collection efficiency curve is defined in amendments to 40 CFR, Part 53 (Code of Federal Regulations, 2001b).

An example of a PM_{2.5} size-cut curve is also shown in Figure 2-7. The PM_{2.5} size-cut curve, however, is defined by the design of the Federal Reference Method (FRM) sampler. The basic design of the FRM sampler is given in the Federal Register (1997, 1998) and in 40 CFR, Part 50, Appendix L (Code of Federal Regulations, 2001c). Additional performance specifications are given in 40 CFR, Parts 53 and 58 (Code of Federal Regulations, 2001b,d). In order to be used for measurement of PM_{2.5} to determine compliance with the PM_{2.5} NAAQS,




Figure 2-8. An idealized distribution of ambient particulate matter showing fine-mode particles and coarse-mode particles and the fractions collected by size-selective samplers. (WRAC is the Wide Range Aerosol Classifier which collects the entire coarse mode [Lundgren and Burton, 1995].)

Source: Adapted from Wilson and Suh (1997) and Whitby (1978).

each specific sampler design and its associated manual of operational procedures must be designated as a reference method under 40 CFR, Part 53 in Section 1.2 of Appendix L (Code of Federal Regulations, 2001c). Thus $PM_{2.5}$ FRM samplers may have somewhat different designs (see Table 2-5 in Section 2.2.4.1.2).

Papers discussing PM_{10} or $PM_{2.5}$ frequently insert an explanation such as “ PM_x (particles less than $x \mu m$ diameter)” or “ PM_x (nominally, particles with aerodynamic diameter $\leq x \mu m$).” While these explanations may seem easier than “ PM_x (particles collected with an upper 50% cut point of $x \mu m$ aerodynamic diameter and a specified penetration curve),” they are not entirely correct and may be misleading, because they imply an upper 100% cut point of $x \mu m$. Some

countries use PM_{10} to refer not to samplers with a 50% cut at $10 \mu m D_a$, but to samplers with 100% rejection of all particles greater than $10 \mu m D_a$. Such samplers miss a fraction of coarse thoracic PM. An example is shown in Figure 2-9.




Figure 2-9. Comparison of penetration curves for two PM_{10} beta gauge samplers using cyclone inlets. The Wedding PM_{10} sampler uses the EPA definition of PM_x as $x = 50\%$ cut point. The Kimoto PM_{10} defines PM_x as $x =$ the 100% cut point (or zero penetration).

Source: Tsai and Cheng (1996).

PM_{10} , as defined by EPA, refers to particles collected by a sampler with an upper 50% cut point of $10 \mu m D_a$ and a specific, fairly sharp, penetration curve. $PM_{2.5}$ is analogously defined. Although there is not yet an FRM, $PM_{10-2.5}$ refers either to particles collected by a sampler with an upper 50% cut point of $10 \mu m D_a$ and a lower 50% cut point of $2.5 \mu m D_a$ or to the difference between the particle concentration measured by a PM_{10} monitor and a $PM_{2.5}$ monitor. In all

cases, the fraction of PM collected depends on the entire penetration curve (or curves); i.e., for $PM_{2.5}$ some particles $> 2.5 \mu m D_a$ are collected and not all particles $< 2.5 \mu m D_a$ are collected.

A $PM_{10-2.5}$ size fraction may be obtained from a dichotomous sampler or by subtracting the mass collected by a $PM_{2.5}$ sampler from the mass collected by a PM_{10} sampler. The resulting $PM_{10-2.5}$ mass, or $PM_{10-2.5}$, is sometimes called “coarse” particles or “thoracic coarse” particles. However, it would be more correct to call $PM_{10-2.5}$ an indicator of the thoracic component of coarse particles (because it excludes some coarse particles below $2.5 \mu m D_a$ and above $10 \mu m D_a$). Also, $PM_{2.5}$ should be considered an indicator of fine particles (because it contains some coarse particles). It would also be appropriate to call PM_{10} an indicator of thoracic particles. PM_{10} and thoracic PM, as shown in Figure 2-7, have the same 50% cut point. However, the thoracic cut is not as sharp as the PM_{10} cut; therefore, thoracic PM contains some particles between 10 and $30 \mu m$ diameter that are excluded from the PM_{10} fraction.

Over the years, the terms *fine* and *coarse*, as applied to particles, have lost the precise meaning given in Whitby’s (1978) definition. In any given article, therefore, the meaning of fine and coarse, unless defined, must be inferred from the author’s usage. In this document, “fine particles” means all particles in the nucleation, Aitken, and accumulation modes; and “coarse particles” means all particles in the coarse mode. Fine particles and $PM_{2.5}$ are not equivalent terms.

Selection of Cut Points for Regulatory Size Cuts

TSP. Regulatory size cuts are a specific example of size-selective sampling. Prior to 1987, the indicator for the NAAQS for PM was TSP. TSP is defined by the design of the High Volume sampler (hivol) that collects all of the fine particles but only part of the coarse particles (Figure 2-8). The upper cut-off size of the hivol depends on the wind speed and direction and may vary from 25 to $40 \mu m$. Newer PM samplers are usually designed to have an upper cut point and its standard deviation that are independent of wind direction and relatively independent of wind speed.

PM_{10} . In 1987, the NAAQS for PM were revised to use PM_{10} , rather than TSP, as the indicator for the NAAQS for PM (Federal Register, 1987). The use of PM_{10} as an indicator is an example of size-selective sampling based on a regulatory size cut (Federal Register, 1987). The

selection of cut point characteristics depends upon the application for the sampling device. A separation that simulates the removal of particles by the human upper respiratory system would appear to be a good choice for both health risk and regulatory monitoring (i.e., it would measure what gets into the lungs). The ACGIH-ISO-CEN penetration curve for thoracic particles (particles able to pass the larynx and penetrate into the bronchial and alveolar regions of the lung) has a 50% cut point at 10 μm aerodynamic diameter (D_a). The selection of PM_{10} as an indicator was based on health considerations and was intended to focus regulatory concern on those particles small enough to enter the thoracic region of the human respiratory tract. The PM_{10} is an indicator of thoracic particles and is a compromise between the desire to collect all particles that might enter the thoracic regions and the need to design a sampler with a collection efficiency independent of particle size or wind speed. As shown in Figure 2-7, the PM_{10} penetration efficiency curve matches the definition for thoracic PM given by the American Conference of Government and Industrial Hygienists (1994) very well except for a slight under-collection of particles between 10 and 30 μm in diameter. While the U.S. PM_{10} separation curve is sharper than the thoracic penetration curve, it has the advantage of reducing the problem of maintaining the finite collection efficiency specified by the thoracic penetration curve for particles larger than 10 μm D_a . (See Section 2.1.2.2 and Figure 2-7.)

Current PM_{10} samplers have upper cut points that are stable under normal operating conditions. However, problems may occur under unusual or adverse conditions. Ono et al. (2000) reported the results of a study in which several PM_{10} samplers were collocated and operated at various sites at Owens Lake, CA, a location with high concentrations of coarse PM. Samplers included the Partisol, the dichotomous, the Wedding high-volume, and the Graseby high-volume samplers in addition to the tapered element oscillating microbalance (TEOM) monitor. They found that the TEOM and Partisol samplers agreed to within 6%, on average. The dichotomous sampler and the Graseby and Wedding high-volume samplers, however, measured significantly lower PM_{10} concentrations than the TEOM (on average 10, 25, and 35% lower, respectively). These lower concentrations were attributed to a decrease in the cut point at higher wind speeds and to a dirty inlet. Since the 10 μm cut point is on a part of the size distribution curve where the concentration is changing rapidly, the amount of PM collected is sensitive to small changes in the cut point. Therefore, the cut point needs to be specified very

precisely and the balance between a design standard and a performance standard will need to be considered to deal with this problem.

PM_{2.5}: Cut Point for Separation of Fine and Coarse Particles. The PM_{2.5} standard set in 1997 is also an example of size-selective sampling based on a regulatory size cut (Federal Register, 1997). The PM_{2.5} standard was based primarily on epidemiologic studies using concentrations measured with PM_{2.5} samplers as an exposure index. However, the PM_{2.5} sampler was not designed to collect all respirable particles; it was designed to separate fine and coarse particles and to collect fine particles because of their different sources and properties (Whitby, 1978; Miller et al., 1979). Thus, the need to attain a PM_{2.5} standard tends to focus regulatory concern on control of the sources of fine particles.

Fine and coarse particles differ not only in size but also in formation mechanisms, sources, and chemical, physical, and biological properties. They also differ in concentration-exposure relationships, dosimetry (deposition and retention in the respiratory system), toxicity, and health effects as observed in epidemiologic studies. Thus, it is desirable to measure fine and coarse PM separately in order to properly allocate health effects to either fine or coarse PM and to correctly determine sources by receptor modeling approaches. For example, sulfates in fine particles are associated with hydrogen or ammonium ions, while sulfates in coarse particles are associated with basic metal ions. Transition metals in coarse particles are likely to be associated with soil and tend to be less soluble (and presumably less bioavailable) than transition metals in fresh combustion particles found in fine particles.

In the early 1970s, aerosol scientists were beginning to recognize the existence of a minimum between 1 and 3 μm in the distribution of particle size by volume (Whitby et al., 1974). However, the limited size distribution information available at that time (two distributions with a minimum near 1 μm and two with a minimum near 2.5 μm) did not permit an unambiguous definition of the appropriate cut point size for the separation of the two modes. A cut point of 2.5 μm was chosen for a new dichotomous sampler (Loo et al., 1976; Jaklevic et al., 1977) designed for use in the Regional Air Pollution Study in St. Louis, MO. The 2.5 μm cut point was subsequently used as an indicator of fine particles in a number of studies including the Harvard Six-City Studies of the relationships between mortality and PM concentrations (Dockery et al., 1993; Schwartz et al., 1996). In an analysis reported in 1979, EPA scientists

endorsed the need to measure fine and coarse particles separately (Miller et al., 1979). Based on the availability of a dichotomous sampler with a separation size of $2.5 \mu\text{m } D_a$, they recommended $2.5 \mu\text{m } D_a$ as the cut point between fine and coarse particles. Because of the wide use of this cut point, the $\text{PM}_{2.5}$ fraction is frequently referred to as “fine” particles. However, although the $\text{PM}_{2.5}$ sample will contain most of the fine particles, except to a lesser extent during conditions of high relative humidity, it may also collect a small fraction of the coarse particles, especially in dry areas or during dry conditions. A $2.5 \mu\text{m}$ cut point was also used in the Inhalable Particle Network (Suggs and Burton, 1983), which provided data for another major epidemiologic study of PM-mortality relationships using an American Cancer Society cohort (Pope et al., 1995). Therefore, at the time of the last previous review of the NAAQS for PM (U.S. Environmental Protection Agency, 1996a,b), a number of epidemiologic studies demonstrating a statistical relationship between $\text{PM}_{2.5}$ concentrations and mortality were available.

During the previous review of the PM standards, EPA conducted an extensive review of the cut point to be used for a fine particle standard, including consideration of PM_1 as an alternative to $\text{PM}_{2.5}$. The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) contains a review of the available information on size distributions. As shown in Figure 2-10 (adapted from the 1966 PM AQCD; U.S. Environmental Protection Agency, 1996a), published size distributions exhibit considerable variability in the intermodal region (1 to $3.0 \mu\text{m}$ diameter). In Figure 2-10a, Philadelphia, very little mass is found in the intermodal region and it is not clear whether it should be considered fine or coarse. In Figure 2-10b, Phoenix, a tail of the coarse mode extends to $1 \mu\text{m}$ or below. These two size distributions were fit with three lognormal distributions. However, the intermodal mode probably does not have physical significance. In Figure 2.10c, Claremont, South Coast Basin, a size distribution is shown with the accumulation mode split into a condensation mode and a droplet mode, which extends to $3.0 \mu\text{m}$ or above (John et al., 1990). The droplet mode occurs under high relative humidity (RH) conditions, usually with very high fine PM concentrations, and is believed to result from reactions involving gases dissolving and reacting in the particle-bound water of the particle droplets (Hering and Friedlander, 1982; McMurry and Wilson, 1983). Although Figure 2-10c shows nitrate concentrations, similar distributions have been observed for sulfate (John et al., 1990).




Figure 2-10. Three examples of impactor size distributions and distributions resulting from fitting several lognormal distributions to the impactor data. These distributions show the variation in the particle size range of the minimum in mass between the accumulation mode and the coarse mode.

Source: a and b, adapted from Lundgren and Hausknecht (1982); c, John et al. (1990).

It is now understood that the size range between 1.0 and 3.0 μm , sometimes called the intermodal region, may contain either accumulation-mode or coarse-mode material or both, i.e., the two modes may overlap in this region (Kegler et al., 2001). The experimental information on the composition and source of the intermodal mass was discussed extensively in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Depending largely on the RH, a significant amount of either accumulation- or coarse-mode material may be found in the intermodal region between 1.0 and 3.0 μm . The analysis demonstrated the important role of relative humidity in influencing the size of particles in both the accumulation and coarse modes.

As the RH increases, hygroscopic accumulation-mode particles will increase in size due to accumulation of particle-bound water. At high RH, some originally submicrometer-sized accumulation-mode PM may be found with a D_a above 1 μm . At an RH of 100%, such as is found in fog and clouds, accumulation-mode PM may extend above 2.5 μm D_a . What is not well understood is whether such particles will shrink to diameters below 1 μm as the RH decreases or whether reactions occurring in the wet particles will result in an increase in nonaqueous mass, so that, even at low RH, the diameters would exceed 1 μm . On the other hand, at very low RH, coarse particles may be fragmented into smaller sizes, and small amounts of coarse PM may be found with an $D_a < 2.5 \mu\text{m}$ (Lundgren et al., 1984; Lundgren and Burton, 1995). Thus, a $\text{PM}_{2.5}$ sample will contain all of the accumulation-mode PM except during periods of high RH. However, under low-RH conditions, it may also contain a small fraction of the coarse PM.

Considerations that led the EPA to choose to retain 2.5 μm as the cut point for the separation of fine and coarse particles included the following three points. First, epidemiologic data showing statistical relationships between fine PM and health outcomes were based largely on $\text{PM}_{2.5}$. Second, while PM_1 would exclude a tail of the coarse mode in some locations, in other locations it would miss a portion of the fine PM. Since the growth of the droplet mode is associated with very high fine PM concentrations, this would result in falsely low fine PM measurements on days with the highest fine PM concentrations. Third, only limited data on the concentration and composition of the intermodal PM mass was available. The selection of a cut point of 2.5 μm as a basis for the EPA's 1997 NAAQS for fine particles (Federal Register, 1997) and its continued use in many health effects studies reflect the importance placed on a more complete inclusion of accumulation-mode particles, while recognizing that the intrusion of coarse particles can occur under some conditions with this cut point.

Since the 1966 PM AQCD (U.S. Environmental Protection Agency, 1996a), several papers have addressed the issue of intermodal mass in terms of $PM_{2.5}$ - PM_1 . Kegler et al. (2001) analyzed data from Phoenix including TEOM measurements of PM_1 , $PM_{2.5}$, and PM_{10} as well as filter measurements of $PM_{2.5}$ and concluded that while $PM_{2.5-1}$ was dominated by soil components, it also contained some nonsoil components. Their analysis suggested that there were two sources of the coarse mass found in $PM_{2.5-1}$: (1) resuspension of soil dust by natural wind (windblown dust), which would be prominent at high wind speeds, and (2) resuspension due to roadway turbulence generated by motor vehicles (road dust), which would occur at all natural wind speeds. Correlations among the various PM size ranges measured in Phoenix are given in Table 2-1. The high correlation found between $PM_{2.5}$ and PM_1 ($r = 0.97$) suggests that the use of PM_1 instead of $PM_{2.5}$ would not significantly change epidemiologic relationships between PM mass and health outcomes in Phoenix.

TABLE 2-1. CORRELATIONS BETWEEN TEOM MEASUREMENTS IN PHOENIX

	PM_1	$PM_{2.5-1}$	$PM_{2.5}$	$PM_{10-2.5}$	PM_{10}
PM_1	1	0.69	0.97	0.65	0.81
$PM_{2.5-1}$	0.69	1	0.84	0.84	0.89
$PM_{2.5}$	0.97	0.84	1	0.75	0.89
$PM_{10-2.5}$	0.65	0.84	0.75	1	0.97
PM_{10}	0.81	0.89	0.89	0.97	1

In areas where winds cause high concentrations of windblown soil, there is evidence that a significant amount of coarse-mode PM may be found below 2.5 μm . An example, taken from data collected during the August 1996 dust storm in Spokane, WA, is shown in Figure 2-11. Note that the PM_{10} scale is 10 times that of the other size fractions. PM_1 , although high in the morning, goes down as the wind increases and PM_{10} , $PM_{2.5}$, and $PM_{2.5-1}$ go up. During the peak of the dust storm, around 9:00 p.m., $PM_{2.5-1}$ was 88% of $PM_{2.5}$. For the 24-h period, $PM_{2.5-1}$ was 54% of $PM_{2.5}$. However, PM_1 was not affected by the intrusion of coarse particles. Similar




Figure 2-11. Particulate matter concentrations in Spokane, WA, during the August 30, 1996 dust storm.

Source: Claiborn et al. (2000).

considerations probably apply to short-term intrusions of dust transported from distant sources such as the Sahara and Gobi deserts (Husar et al., 2001). In Spokane, the correlations between PM₁ and PM_{2.5-1} (Haller et al., 1999) were lower, $r = -0.37$ (spring) to 0.26 (winter), than those observed in Phoenix (Kegler et al., 2001), $r = 0.69$.

Pope et al. (1999) found that removing days with high PM₁₀ values due to windblown dust resulted in the observation of a statistically significant relationship between PM₁₀ and mortality in Salt Lake City. This relationship was not observed when days with windblown dust events were included in the PM₁₀ concentration time series. The elevated values of PM_{2.5}, but not PM₁, observed during the windblown dust events in Spokane (Claiborn et al., 2000) suggest that a PM_{2.5} time series would also be impacted by windblown dust while a PM₁ time series would not. Thus, the contribution of soil to PM_{2.5} could possibly contribute to erroneous conclusions from some epidemiologic analyses.

A cut point of 1.0 μm could reduce the misclassification of coarse-mode material as fine, especially in areas with high levels of windblown soil, but under high RH conditions could result in some accumulation-mode material being misclassified as coarse. A reduction in RH, either intentionally or inadvertently, will reduce the mass mean diameter of the accumulation-mode particles. Studies of the changes in particle size with changes in relative humidity suggest that only a small fraction of accumulation-mode particles will be above 1.0 μm in diameter at RH below 60%, but a substantial fraction will grow to above 1.0 μm at RH above 80% (Hitzenberger et al., 1997; McMurry and Stolzenburg, 1989; U.S. Environmental Protection Agency, 1996a). Studies in Europe (Berner, 1989; see Figure 3-31 in the 1966 PM AQCD) show that dehumidification of ambient PM by heating will reduce the size of accumulation mode particles, leaving little accumulation mode PM above 1 μm in aerodynamic diameter. However, heating will also remove ammonium nitrate and semivolatile organic compounds. Currently, the data are insufficient to determine how much dehumidification by diffusion drying would reduce the size of accumulation mode particles.

Under high RH circumstances, a monitor using a 1.0 μm D_a cut point can achieve better modal separation if the air stream is dehumidified to some fixed humidity that would remove all or most particle-bound water without evaporating semivolatile components. New techniques requiring the reduction of RH by diffusion drying prior to collection have been developed for measuring fine PM minus particle-bound water but including semivolatile nitrate and organic compounds. With such techniques, measurements with a 1 μm (or slightly higher) cut point, in conjunction with concurrent $\text{PM}_{2.5}$ measurements, would be useful for exposure, epidemiologic, and source-apportionment studies, especially in areas where intrusion of coarse-mode particles into the intermodal range is likely.

2.1.2.3 Ultrafine Particles

As discussed in Chapter 7 (Toxicology of Particulate Matter in Humans and Laboratory Animals) and in Chapter 8 (Epidemiology of Human Health Effects Associated with Ambient Particulate Matter), some scientists argue that ultrafine particles may pose potential health problems and that some health effects may be associated with particle number or particle surface area as well as, or more closely than, with particle mass. Some additional attention will be given

here to ultrafine particles, because they contribute the major portion of particle number and a significant portion of particle surface area.

Formation and Growth of Fine Particles

Several processes influence the formation and growth of particles. New particles may be formed by nucleation from gas phase material. Particles may grow by condensation as gas phase material condenses on existing particles; particles may also grow by coagulation as two particles combine to form one. Gas phase material condenses preferentially on smaller particles, and the rate constant for coagulation of two particles decreases as the particle size increases. Therefore, ultrafine particles grow into the accumulation mode; but accumulation-mode particles do not normally grow into the coarse mode (see Figure 2-6). More information and references on the formation and growth of fine particles can be found in the 1996 AQCD PM (U.S. Environmental Protection Agency, 1996a).

Equilibrium Vapor Pressures

An important parameter in particle nucleation and in particle growth by condensation is the saturation ratio, S , defined as the ratio of the partial pressure of a species, p , to its equilibrium vapor pressure above a flat surface at a specified temperature, p_o : $S = p/p_o$. For either condensation or nucleation to occur, the species vapor pressure must exceed its equilibrium vapor pressure. For particles, the equilibrium vapor pressure is not the same as p_o . Two effects are important: (1) the Kelvin effect, which is an increase in the equilibrium vapor pressure above the surface due to its curvature (very small particles have higher vapor pressures and will not be stable to evaporation until they attain a critical size) and (2) the solute effect, which is a decrease in the equilibrium vapor pressure of the liquid due to the presence of other compounds in solution. Organic compounds may also be adsorbed onto ultrafine carbonaceous particles.

For an aqueous solution of a nonvolatile salt, the presence of the salt decreases the equilibrium vapor pressure of the water vapor around the droplet. This effect is in the opposite direction of the Kelvin effect, which increases the equilibrium water vapor pressure around a droplet because of its curvature. The existence of an aqueous solution will also influence the vapor pressure of water-soluble species. The vapor pressure behavior of mixtures of several liquids or of liquids containing several solutes is complex.

New Particle Formation

When the vapor concentration of a species exceeds its equilibrium concentration (expressed as its equilibrium vapor pressure), it is considered condensable. Condensable species can either condense on the surface of existing particles or can nucleate to form new particles. The relative importance of nucleation versus condensation depends on the rate of formation of the condensable species and on the surface or cross-sectional area of existing particles (McMurry and Friedlander, 1979). In ambient urban environments, the available particle surface area is usually sufficient to rapidly scavenge the newly formed condensable species. Formation of new, ultrafine particles is usually not observable in mass or volume distributions except near sources of condensable species. Wilson et al. (1977) reported observations of the Aitken nuclei mode in traffic. However, bursts of new particle formation have been observed in urban areas in the number distribution (Woo et al., 2001a; McMurray et al., 2000). New particle formation also can be observed in cleaner, remote regions. Bursts of new particle formation in the atmosphere under clean conditions usually occur when aerosol surface area concentrations are low (Covert et al., 1992). High concentrations of nuclei mode particles have been observed in regions with low particle mass concentrations, indicating that new particle formation is inversely related to the available aerosol surface area (Clarke, 1992).

Sources of Ultrafine Particles

Ultrafine particles are the result of nucleation of gas phase species to form condensed phase species with a very low equilibrium vapor pressure. In the atmosphere, four major classes of substances yield PM with equilibrium vapor pressures low enough to form nuclei mode particles:

- (1) *Particles containing heavy metals.* Nuclei mode particles of metal oxides or other metal compounds are generated when metallic impurities in coal or oil are vaporized during combustion and the vapor undergoes nucleation. Metallic ultrafine particles also may be formed from metals in lubricating oil or fuel additives that are vaporized during combustion of gasoline or diesel fuels. Ultrafine metallic particles were discussed in Section 6.9 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).
- (2) *Elemental carbon (EC) or soot.* Elemental carbon particles are formed primarily by the condensation of C_2 molecules generated during the combustion process. Because EC has a very low equilibrium vapor pressure, ultrafine EC particles can nucleate even at high temperatures (Kittelson, 1998; Morawska et al., 1998).

- (3) *Organic carbon (OC)*. Recent smog chamber studies and indoor experiments show that atmospheric oxidation of certain organic compounds often found in the atmosphere can produce highly oxidized organic compounds with an equilibrium vapor pressure sufficiently low to result in nucleation (Kamens et al., 1999; Weschler and Shields, 1999).

- (4) *Sulfates*. Sulfuric acid (H_2SO_4) molecules are generated in the atmosphere by conversion of sulfur dioxide (SO_2) to H_2SO_4 . As H_2SO_4 is formed, it can either nucleate to form new ultrafine particles, or it can condense on existing ultrafine or accumulation mode particles (Clark and Whitby, 1975; Whitby, 1978). Nucleation theory allows the calculation of nucleation rates for both binary nucleation, involving water and sulfuric acid (Easter and Peters, 1994) or ternary nucleation, which requires sulfuric acid, ammonia (NH_3), and water (Korhonen et al., 1999). Kulmala et al. (2000) compared binary and ternary nucleation rates and concluded that the ternary rate is 1,000 times the binary rate. Results from an aerosol dynamics model with a ternary nucleation scheme indicate that nucleation in the troposphere should be ubiquitous and yield a reservoir of thermodynamically stable sulfate clusters 1 to 3 nm in diameter. The growth of these clusters to a detectable size (> 3 nm diameter) is limited by the availability of condensable vapor. Observations of atmospheric particle formation and growth from a continental and a coastal site suggest that a growth process including ternary nucleation is responsible for the formation of cloud condensation nuclei. Nucleation processes in the atmosphere may also involve organic compounds as well as sulfuric acid, ammonia, and water. However, current formulations of nucleation theory only include the three inorganic components. (The possible formation of ultrafine NH_4NO_3 by reaction of NH_3 and nitric acid [HNO_3] vapor has not been investigated.)

Vehicle engine exhaust may include all the substances described above. Ultrafine particles are observed in the emissions from spark, diesel, and jet engines (Kittelson, 1998). In these cases, it seems likely that EC, organic compounds, ammonia and sulfuric acid from sulfur in the fuel, as well as metal additives in the fuel or fuel oil, may contribute to the formation of ultrafine particles (Tobias et al., 2001). An example of particles from a diesel engine showing number and volume distributions is shown in Figure 2-12.

Recent Measurements of Ultrafine Particles

Instruments developed during the past decade permit measurement of size distributions of particles down to 3 nm in diameter. The use of these techniques has led to new information on the formation of new particles by nucleation. Such measurements have been carried out during intensive field measurement campaigns, during continuous measurements in urban areas in several European cities, and in the United States as a part of the Supersite program (McMurry




Figure 2-12. Typical engine exhaust size distribution.

Source: Kittelson (1998).

et al., 2000; Woo et al., 2001a). Nucleation has been observed in the free troposphere (Weber et al., 1999; Clarke, 1992; Schröder and Ström, 1997; Raes et al., 1997), in outflows of evaporating convective clouds (Clarke et al., 1998; Hegg et al., 1990, 1991; Radke and Hobbs, 1991; Perry and Hobbs, 1994), in the marine boundary layer (Covert et al., 1992; Hoppel et al., 1994; Van Dingenen et al., 1995; Weber et al., 1998; Clarke et al., 1998), downwind of coastal regions during low tide (McGovern et al., 1996; McGovern, 1999), on mountains (Weber et al., 1995, 1997; Raes et al., 1997; Wiedensohler et al., 1997), over forests (Mäkelä et al., 1997; Kulmala et al., 1998; O'Dowd et al., 2002), downwind of certain biogenic emissions (Weber et al., 1998), in urban areas (Birmili and Wiedensohler, 1998; McMurry et al., 2000; Woo et al., 2001a), near freeways (Zhu et al., 2002a,b), in engine exhaust (Kittelson, 1998; Tobias et al., 2001), and in homes (Wallace and Howard-Reed, 2002). Nucleation events in outdoor air almost always occur during daylight, indicating that photochemistry plays a role in producing the gas phase precursors of new particles.

The number size distributions observed over a boreal forest in Finland before and during the initial stages of a nucleation event are shown in Figure 2-13. The Aitken and accumulation modes can be seen clearly before the nucleation event. The nucleation mode, with a peak




Figure 2-13. Number size distributions showing measurement of a nucleation burst mode in a boreal forest in Finland.

Source: O'Dowd et al. (2002).

between 3 and 7 nm, appears during the event. Figure 2-14 shows the variety of size distributions that may be observed as nuclei are formed and grow, based on size distributions measured in the Arctic marine boundary layer (Covert et al., 1996). These distributions all show a trimodal distribution within the fine-particle size range. The changes in size distribution due to coagulation (and dilution) immediately downwind of a freeway (Zhu et al., 2002b) are shown in Figures 2-15(a-f) and 2-15(g). At 30 m downwind, the nucleation mode number concentration is larger than that of the Aitken mode, but by 60 m downwind that has reversed, because coagulation removes particles from the nucleation mode and adds particles to the Aitken mode.

Strong evidence suggests that sulfuric acid vapor often participates in nucleation. However, condensation of sulfuric acid and its associated water and ammonium ions typically accounts for only 10 to 20% of the observed growth rates for freshly nucleated particles. Therefore, organic compounds may account for much of the formation and growth of freshly nucleated particles. Evidence of nucleation of organic particles comes from smog chamber studies (Kamens et al., 1999) and from field studies over forests (Mäkelä et al., 1997; Kulmala et al., 1998; O'Dowd et al., 2002). Nucleation of organic particles may also occur indoors due to the reaction of infiltrated ozone with indoor terpenes from air fresheners or cleaning solutions (Weschler and Shields, 1999). The observation of bursts of nuclei-mode particles in Atlanta




Figure 2-14. Examples of the measured 1-h average particle number size distributions and the log normal fits to the modes of the data. Squares are measured data, solid lines are the fitted lognormal modes determined by DistFit™. These modes (nucleation between 3 and 20 nm, Aitken between 20 and 100 nm, and accumulation above 100 nm) can be observed in most examples.

Source: Covert et al. (1996).

(Woo et al., 2001a), perhaps due to unusually high rates of production of condensable species, suggests that high concentrations of ultrafine particles may be a normal occurrence in polluted urban areas.




Figure 2-15a-f. Fitted multi-model particle size distribution at different sampling distances from Freeway 405: (a) 30 m downwind, (b) 60 m downwind, (c) 90 m downwind, (d) 150 m downwind, (e) 300 m downwind, (f) 300 m upwind. Size distributions were normalized to the control CPC's reading. Note different scales for $dN/d \log D_p$ axis. Modal parameters given are: geometric mean diameter, μ_g ; and geometric standard deviation, σ_g .

Source: Zhu et al. (2002b).




Figure 2-15g. Combination of Figures 2-15(a-e), with $dN/d \log D_p$ scale. Ultrafine particle size distribution at different sampling locations near Freeway 405 in Los Angeles, CA.

Source: Zhu et al. (2002b).

Concentration of Ultrafine Particles: A Balance Between Formation and Removal

Nuclei-mode particles may be removed by dry deposition or by their growth into the accumulation mode. Such growth takes place as other low vapor pressure material condenses on the particles or as nuclei-mode particles coagulate with themselves or with accumulation mode particles. Because the rate of coagulation would vary with the concentration of accumulation-mode particles, it might be expected that the concentration of nuclei-mode particles would increase with a decrease in accumulation-mode mass. On the other hand, the concentration of particles would be expected to decrease with a decrease in the rate of generation of particles by reduction in emissions of metal and carbon particles or a decrease in the rate of generation of H_2SO_4 or condensable organic vapor. The rate of generation of H_2SO_4 depends on the concentrations of SO_2 and the hydroxyl radical ($\bullet OH$), which is generated primarily by reactions involving ozone (O_3). Thus, reductions in SO_2 and O_3 would lead to a decrease in the

rate of generation of H_2SO_4 and condensable organic vapor and to a decrease in the concentration of nuclei-mode particles. The balance between formation and removal is uncertain. However, these processes can be modeled using a general dynamic equation for particle size distribution (Friedlander, 2000) or by aerosol dynamics modules in newer air quality models (Binkowski and Shanker, 1995; Binkowski and Ching, 1995).

2.1.3 Chemistry of Atmospheric Particulate Matter

The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental carbon; a great variety of organic compounds; crustal material; and (at coastal locations) sea salt. Atmospheric PM also contains a large number of elements in various compounds and concentrations. More information and references on the composition of PM measured in a large number of studies in the United States, may be found in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Also, in this document, the concentrations and composition of ambient PM are discussed in Chapter 3 (Section 3.1), and ambient data for concentrations and composition of $\text{PM}_{2.5}$ are given in Appendices 3A, 3B, and 3C.

2.1.3.1 Chemical Composition and Its Dependence on Particle Size

Studies conducted in most parts of the United States indicate that sulfate, ammonium, and hydrogen ions; elemental carbon, secondary organic compounds, and primary organic species from cooking and combustion; and certain transition metals are found predominantly in the fine particle mode. Crustal materials such as calcium, aluminum, silicon, magnesium, and iron are found predominately in the coarse particles. Some primary organic materials such as pollen, spores, and plant and animal debris are also found predominantly in the coarse mode. Certain components such as potassium and nitrate may be found in both the fine and coarse particle modes, but they originate from different sources or mechanisms. Potassium in coarse particles comes from soil. Potassium in fine particles originates in emissions from burning wood or cooking meat. Nitrate in fine particles comes primarily from the reaction of gas-phase nitric acid with gas-phase ammonia forming particulate ammonium nitrate. Nitrate in coarse particles comes primarily from the reaction of gas-phase nitric acid with preexisting coarse particles.

2.1.3.2 Primary and Secondary Particulate Matter

Particulate matter can be primary or secondary. PM is called “primary” if it is in the same chemical form in which it was emitted into the atmosphere, but it is called “secondary” if it is formed by chemical reactions in the atmosphere. Primary coarse particles are usually formed by mechanical processes and include windblown dust, sea salt, road dust, and combustion-generated particles such as fly ash and soot. Primary fine particles are emitted from sources either directly as particles or as vapors that rapidly condense to form ultrafine or nuclei-mode particles. This includes soot from diesel engines, a great variety of organic compounds condensed from incomplete combustion or cooking, and compounds of As, Se, Zn, etc. that condense from vapor formed during combustion or smelting. The concentration of primary particles depends on their emission rate, transport and dispersion, and removal rate from the atmosphere.

Secondary PM is formed by chemical reactions of free, adsorbed, or dissolved gases. Most secondary fine PM is formed from condensable vapors generated by the chemical reactions of gas-phase precursors. Secondary formation processes can result in either the formation of new particles or the addition of PM to preexisting particles. Most of the sulfate and nitrate and a portion of the organic compounds in atmospheric particles are formed by chemical reactions that occur in the atmosphere. Secondary aerosol formation depends on numerous factors, including the concentrations of precursors; the concentrations of other gaseous reactive species such as ozone, hydroxyl radical, peroxy radicals, or hydrogen peroxide; atmospheric conditions including solar radiation and RH; and the interactions of precursors and preexisting particles within cloud or fog droplets or in the liquid film on solid particles. As a result, it is considerably more difficult to relate ambient concentrations of secondary species to sources of precursor emissions than it is to identify the sources of primary particles. A significant effort is currently being directed toward the identification and modeling of organic products of photochemical smog, including the conversion of gases to PM. More information of the transformation of precursor gases into secondary PM is given in Chapter 3 (Section 3.3.1).

Particle strong acidity is due almost entirely to the presence of H_2SO_4 or NH_4HSO_4 . Thus, the acidity of atmospheric particles depends on both the amount of SO_2 that is oxidized to SO_3 and subsequently forms H_2SO_4 as well as the amount of ammonia available to react with the sulfuric acid. Nitric acid is more volatile than sulfuric acid. Thus, if gas phase SO_3 or sulfuric acid or particles containing H_2SO_4 or NH_4HSO_4 contact particles containing NH_4NO_3 , nitric acid

gas will be released with the remaining ammonia contributing to the further neutralization of the acid. Little NH_4NO_3 is found in atmospheres containing significant particle strong acidity. However, as SO_2 emissions are reduced to the point that there is more than enough ammonia to neutralize the sulfuric acid, NH_4NO_3 particles will begin to form. Thus, ammonia emissions and concentrations relative to those of SO_2 and H_2SO_4 are important in determining the strong acidity in the atmosphere and the concentration of particulate NH_4NO_3 . Therefore, once SO_2 emissions have been reduced to the point that ammonia and sulfate are in balance to form $(\text{NH}_4)_2\text{SO}_4$, further reductions in SO_2 will not result in an equivalent reduction in airborne PM, because one $(\text{NH}_4)_2\text{SO}_4$ unit will be replaced by two NH_4NO_3 units.

2.1.3.3 Particle-Vapor Partitioning

Several atmospheric aerosol species, such as water, ammonium nitrate and certain organic compounds, exist in an equilibrium between gaseous and condensed phases and are called semivolatile. The equilibrium between water vapor and particle-bound water is well understood and can be modeled accurately. A variety of thermodynamic models have also been developed to predict the temperature and relative humidity dependence of the ammonium nitrate equilibria with gaseous nitric acid and ammonia. However, under some atmospheric conditions (such as cool, cold, or very clean air), the relative concentrations of the gas and solid phases are not accurately predicted by equilibrium considerations alone, and transport kinetics can be important. The gas-particle distribution of semivolatile organic compounds depends on the equilibrium vapor pressure of the compound, total particle surface area, particle composition, atmospheric temperature, and relative humidity. Although it generally is assumed that the gas-particle partitioning of semivolatile organics is in equilibrium in the atmosphere, neither the equilibria nor the kinetics of redistribution are well understood. Diurnal temperature fluctuations cause gas-particle partitioning to be dynamic on a time scale of a few hours and can cause semivolatile compounds to evaporate during the sampling process. The pressure drop across the filter can also contribute to the loss of semivolatile compounds. The dynamic changes in gas-particle partitioning caused by changes in temperature, pressure, and gas-phase concentration, both in the atmosphere and after collection, cause serious sampling problems that are discussed in Section 2.2.3, *Measurement of Semivolatile Particulate Matter*.

Equilibria with Water Vapor

As a result of the equilibrium of water vapor with liquid water in hygroscopic particles, many ambient particles contain liquid water (particle-bound water). Unless removed, this particle-bound water will be measured as a component of the particle mass. Particle-bound water plays an important role in atmospheric chemistry and physics. It influences the size of particles, and in turn, the light scattering and aerodynamic properties of particles. Particle aerodynamic properties are important in dry deposition to surfaces, deposition to airway surfaces during breathing, and deposition in sampling instrumentation. The aqueous solution provides a medium for reactions of dissolved gases, including reactions that do not take place in the gas phase. The aqueous solutions also may act as a carrier to convey soluble toxic species to the gas-exchange regions of the respiratory system, including species that would be removed by deposition in the upper airways if no particles were present. An extensive review of the equilibrium of water vapor with particle-bound water as it pertains to ambient aerosols was given in Chapter 3 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a).

The interaction of particles with water vapor may be described briefly as follows. As relative humidity increases, particles of crystalline soluble salts, such as $(\text{NH}_4)_2\text{SO}_4$, NH_4HSO_4 , or NH_4NO_3 , undergo a phase transition to become aqueous solution particles. According to the phase rule, for particles consisting of a single component, this phase transition is abrupt, taking place at a relative humidity that corresponds to the vapor pressure of water above the saturated solution (the deliquescence point). With a further increase in relative humidity, the solution particle adds water (and the concentration of the solute decreases) so that the vapor pressure of the solution is maintained equal to that of the surrounding relative humidity; thus, the solution particle tends to follow the equilibrium growth curve. As relative humidity decreases, the solution particle follows the equilibrium curve to the deliquescence point. However, rather than crystallizing at the deliquescence relative humidity, the solute remains dissolved in a supersaturated solution to considerably lower relative humidities. Ultimately, the solution particle abruptly loses its water vapor (efflorescence) and typically returns to the initial crystalline form.

For particles consisting of more than one component, the solid to liquid transition will take place over a range of RHs, with an abrupt onset at the lowest deliquescence point of the several components. Subsequent growth will occur as crystalline material in the particle dissolves

according to the phase diagram for the particular multicomponent system. Under such circumstances, a single particle may undergo several more or less abrupt phase transitions until the soluble material is fully dissolved. At decreasing relative humidity, such particles tend to remain in solution to relative humidities well below the several deliquescence points. In the case of the sulfuric acid-ammonium sulfate-water system, the phase diagram is fairly well understood. For particles of composition intermediate between NH_4HSO_4 and $(\text{NH}_4)_2\text{SO}_4$, this transition occurs in the range from 40% to below 10% relative humidity, indicating that for certain compositions the solution cannot be fully dried in the atmosphere. At low relative humidities, particles of this composition would likely be present in the atmosphere as supersaturated solution droplets (liquid particles) rather than as solid particles. Thus, they would exhibit hygroscopic rather than deliquescent behavior during relative humidity cycles.

Other pure compounds, such as sulfuric acid, are hygroscopic (i.e., they form aqueous solutions at any relative humidity and maintain a solution vapor pressure over the entire range of relative humidity). Soluble organic compounds may also contribute to the hygroscopicity of the atmospheric aerosol (Saxena et al., 1995; Saxena and Hildeman, 1996), but the equilibria involving organic compounds and water vapor, and, especially for mixtures of salts, organic compounds, and water, are not so well understood. These equilibrium processes may cause an ambient particle to significantly increase its diameter at relative humidities above about 40% (Figure 2-16a). A particle can grow to five times its dry diameter as the RH approaches 100% (Figure 2-16b). The Federal Reference Methods, for filter measurements of $\text{PM}_{2.5}$ and PM_{10} mass, require equilibration at a specified, low relative humidity after collection (for $\text{PM}_{2.5}$, between 30 and 40% RH with control of $\pm 5\%$ RH [Code of Federal Regulations, 2001a]). This equilibration removes much of the particle-bound water and provides a relatively stable PM mass for gravimetric measurements (see Section 2.2 for details and references). Otherwise, particle mass would be a function of relative humidity, and the particle mass would be largely particle-bound water at higher relative humidities. However, some particle-bound water may be retained even after equilibration. Recent studies have shown that significant amounts of particle-bound water are retained in particles collected on impaction surfaces even after equilibration and that the amount of retained particle-bound water increases with relative humidity during collection (Hitzenberger et al., 1997).




Figure 2-16a. Particle growth curves showing fully reversible hygroscopic growth of sulfuric acid (H₂SO₄) particles, deliquescent growth of ammonium sulfate [(NH₄)₂SO₄] particles at the deliquescent point (A, about 80% relative humidity [RH]), reversible hygroscopic growth of ammonium sulfate solution droplets at RH greater than 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached.

Source: National Research Council (1993) adapted from Tang (1980).

The retention of particle-bound water is a greater problem for continuous monitors, which measure changes in mass collected on a filter over long sampling times. If particle-bound water is not removed, changes in relative humidity would cause changes in the mass of PM collected over previous hours or days. These changes could be much greater than amount of PM mass added in one hour. Therefore, continuous monitoring techniques generally attempt to remove particle-bound water before measurement either by heating or dehumidification. However, other semivolatile materials (e.g., ammonium nitrate and organic compounds) that may be partially




Figure 2-16b. Theoretical predictions and experimental measurements of growth of NH_4HSO_4 particles at relative humidities between 95 and 100%.

Source: Li et al. (1992).

lost during sampling or equilibration of an unheated filter are certainly lost when the collected sample is heated above ambient temperature. These changes in particle size with relative humidity also mean that particle measurements such as surface area or volume, or composition as a function of size, should be made at the same RH in order for the results to be comparable. These problems are addressed in more detail in Section 2.2, *Measurement of Particulate Matter*.

Particle-Bound Water as a Carrier

Water vapor is not a pollutant, and particle-bound water (PBW) is not included in the mass of PM subject to regulation and control. However, while water is not itself a pollutant, PBW may act as a carrier for pollutants, as stated by Wilson (1995):

“It is possible that water-soluble gases, which would be removed by deposition to wet surfaces in the upper respiratory system during inhalation, could dissolve in particle-bound water and be carried with the particles into the deep lung. Water-soluble gases in polluted air include oxidants such as O₃, H₂O₂, and organic peroxides; acid gases such as SO₂, HCl, HNO₃, HONO, and formic acid; and polar organic species such as formaldehyde. Some of these species may have biological effects, but current techniques do not measure particle-bound water or the species dissolved in it.”

Kao and Friedlander (1995) also suggest that aerosols could carry adsorbed free radicals or dissolved H₂O₂ into the lung. Friedlander and Yeh (1998) point out that the aqueous component of the atmospheric aerosol carries short-lived, reactive chemical species, including hydrogen peroxide and organic peroxides. Equilibrium calculations and limited experimental data suggest that aerosol-phase concentrations of hydrogen peroxide will fall within a range (around 1 mM) in which significant biochemical effects have been observed when respiratory tract epithelial cells were treated with hydrogen peroxide solutions.

Wexler and Sarangapani (1998) used a physical model of “gas-particle-mucus heat and mass transport in the human airways,” developed in order to investigate the hypothesis that endogenous ammonia neutralizes aerosol acidity (Sarangapani and Wexler, 1996), to investigate the transport by particles of soluble vapors to the tracheobronchial and air exchange (also called alveolar or pulmonary) regions of the lung. Wexler and Sarangapani (1998) provides a concise description of this process:

“Air pollutants are deposited in the human airway via two pathways — particle and vapor deposition. In the absence of particles, vapors deposit at different locations in the lung depending on their solubility in mucus, which is over 99% water. High-solubility compounds, such as nitric acid or hydrogen peroxide, are rapidly removed in the upper airways while low-solubility compounds, such as oxygen or ozone, are less well removed and so penetrate to the alveoli. Pollutant deposition in the upper airways is less harmful than in the lower airways because upper airways clearances is more rapid and the epithelium is protected by a mucus layer. As a result, low-solubility pollutants, such as ozone, may harm the alveoli while high-solubility pollutants, such as nitric acid, do not reach these tissues.

“In the presence of aerosol particles, this scenario changes. Under most ambient conditions, aerosol particles contain some liquid water so that soluble compounds are partitioned between the gas phase and the aerosol liquid-water phase. The degree of deposition via the gas compared to that via the particles is a function of a number of factors including the solubility of the compound and the liquid-water content of the aerosol. Since highly soluble compounds deposit in the upper

airways, particles may provide a vector for deposition of these compounds in the lower airways. Lower-solubility compounds may persist in the vapor phase and so may deposit in lower airway segments.

“When both a water-soluble vapor and an aerosol liquid phase are present together, some of the vapor will dissolve in the liquid-water phase until an equilibrium is established. The equilibrium condition is given by the Henry’s law coefficient, H ($M \text{ atm}^{-1}$), which gives the relationship between the partial pressure of the vapor, typically in units of atmospheres, and the concentration of the dissolved compound in the aqueous phase, typically in units of molal.

“Direct deposition of the vapor is primarily a function of its solubility. Since the airways are lined with mucus, deposition is higher for compounds with a higher solubility than for those with lower solubility. Compounds with Henry’s law constant (H) greater than $100 M \text{ atm}^{-1}$ deposit relatively uniformly over the first 10 generations (e.g., formaldehyde and hydrogen peroxide [and sulfur (IV) $\{\text{SO}_2$ (aq) plus H_2SO_3 (aq) plus HSO_3^- (aq)]}, so they affect health via interactions with the upper airway epithelia. All the vapor is deposited if H is greater than about 0.1. Low-solubility compounds do not deposit effectively in the mucus until past the 20th generation so are still present in significant concentrations in the pulmonary region. These compounds (e.g., ozone, nitrogen dioxide) affect human health via interactions in the air-exchange regions.

“As aerosols are inhaled, soluble vapors deposit on the mucus, disrupting the gas-particle equilibrium, and the compound begins to evaporate from the aerosol particles. If the evaporation is rapid, the pattern of deposition of the compound will not be influenced by the presence of the particles, i.e., the deposition pattern is essentially that of the vapor alone. If the evaporation is very slow compared to a breathing cycle, a significant amount of the compound will remain in the particle phase and the pattern of deposition may be shifted toward the pulmonary region by the particle.”

Hygroscopic particles, present in the inhaled aerosol disrupt the gas-particle equilibria. The soluble gas will deposit on the mucus, reducing the gas phase vapor pressure, and reducing the equilibrium concentration of the gas in solution. However, in the higher relative humidity of the lung, hygroscopic particles will add PBW and grow in volume. Therefore, even though the concentration of the soluble gas may be lower, there will be a greater amount of the soluble gas in solution. The amount of the soluble gas in the particle will depend on the Henry’s Law coefficient of the gas, the size of the particle, and the position of the particle in the respiratory tract. Figure 2-17 shows how the concentration of the soluble gas ($H = 10^4 M \text{ atm}^{-1}$) in the PBW of the aerosol relative to that in the gas phase changes as the particle moves deeper into the lung.




Figure 2-17. Concentration of the dissolved gas in the particle normalized by its initial concentration as a function of airway generation for $H = 10^4 \text{ M atm}^{-1}$ for particle diameters of 0.1, 0.3, 0.7, and 1 μm .

Source: Wexler and Sarangapani (1998).

In similar figures for other values of H , Wexler and Sarangapani (1998) show that, regardless of particle size, the greatest relative increase in the amount of soluble gas dissolved in PBW (a factor of 10 to 100) occurs for lower solubility gases ($H < 0 \text{ M atm}^{-1}$). However, the deposition of such gases by particles will be small since the concentration of dissolved gas will be low. For higher solubility gases ($H = 10^2 \text{ M atm}^{-1}$), the enhancement may be 2- to 20-fold, depending on particle size. There may be a sufficient amount of the soluble gas in the PBW to influence the deposition pattern, with more soluble gas being transported to higher generations (deeper into the lung). For the highest solubility gases, the enhancement is less than 2, but a larger amount of gas is dissolved in the PBW. The highest solubility gases ($H > 10^4 \text{ M atm}^{-1}$) will evaporate from particles $< 0.1 \mu\text{m}$ before the particles reach the air exchange region of the lung. However, as shown in Figure 2-17, particles $> 0.3 \mu\text{m}$ can efficiently transport high solubility gases into the air exchange region.

Wexler and Sarangapani (1998) observed that even for a high liquid water content (PBW = 100 $\mu\text{g}/\text{m}^3$) and for a highly soluble gas such as H_2O_2 ($H = 10^5 \text{ M atm}^{-1}$), only about 1% of the soluble gas will dissolve in the PBW. Therefore, any influence of the particles on the gas concentration will cause little change in the quantity of soluble gas deposited on the upper airways. The Wexler and Sarangapani (1998) analysis was conducted under the assumption that equilibrium with the gas phase is obtained quickly and considered only the “physical” solubility of the soluble gas. For example, their discussion of SO_2 assumes an H of 1 M atm^{-1} which is correct for the equilibrium of SO_2 (aq) with SO_2 gas. However, SO_2 reacts chemically with water to yield SO_2 (aq), H_2SO_3 (aq), and HSO_3^- (aq). The H for sulfur (IV) in solution in equilibrium with SO_2 gas is dependent on the pH of the solution but may reach 10^4 M atm^{-1} at a pH of 6 (Schwartz, 1984). Thus, if SO_2 (aq), H_2SO_3 (aq), and HSO_3^- (aq) are in equilibrium in the liquid phase, and any one of the three species is toxic, significantly more toxic sulfur (IV) will be carried to the air exchange region than predicted by Wexler and Sarangapani (1998). Friedlander and Yeh (1998) point out that H_2O_2 forms adducts called peroxohydrates with many substances in which it acts like water of hydration (Elvers et al., 1991). If such adducts form in the PBW of atmospheric aerosols, a much higher percentage of the atmospheric peroxide would dissolve in the PBW and be carried into the air exchange region of the lungs.

Thus, soluble gases, like H_2O_2 , SO_2 , and formaldehyde, which would be completely removed in the upper airways in the absence of particles, can dissolve in the PBW of hygroscopic particles and be transported by the particles into the air exchange (alveolar or pulmonary) region of the lungs.

Friedlander and Yeh (1998) also suggested that the epidemiologic associations found between adverse health effects and sulfate or PM mass may represent a response to atmospheric peroxides, or other toxic substances dissolved in PBW. Friedlander and Yeh (1998) stated,

“This hypothesis supports reduction of the total submicron aerosol mass as a way to reduce adverse health effects, because the total submicron mass is closely linked to the aqueous component that carries the reactive species. To test this hypothesis, studies are needed of the effects of exposures of cellular layers and/or animals to submicron hydrogen peroxide-containing aerosols that also contain salts such as ammonium sulfate.”

Morio et al. (2001) conducted an inhalation study in which rats were exposed to $(\text{NH}_4)_2\text{SO}_4$ (429 or 215 $\mu\text{g}/\text{m}^3$; 0.3 to 0.4 μm mass median diameter) or H_2O_2 (10, 20, or 100 ppb) alone or in combination for 2 h. The authors summarized the study as follows.

“In summary, the present studies find that exposure of rats to $(\text{NH}_4)_2\text{SO}_4 + \text{H}_2\text{O}_2$ results in tissue damage and altered alveolar macrophage activity. Our findings suggest that macrophage-derived TNF- α and peroxyxynitrite are potential factors contributing to the toxicity of $(\text{NH}_4)_2\text{SO}_4 + \text{H}_2\text{O}_2$. It is possible that alterations in macrophage functioning following inhalation of $(\text{NH}_4)_2\text{SO}_4 + \text{H}_2\text{O}_2$ underlie increases in individual susceptibility to infection following the fine PM exposure.”

The potential for PBW to carry toxic gases into the air exchange region of the lung suggests that it would be useful to measure both the amount of PBW in the atmospheric aerosol and the composition of the material dissolved in it. Although measurements have been made of the amount of PBW (discussed in 2.2.3.1), there is little information on the composition or concentration of soluble species that would evaporate as the PBW evaporates. Hung and Wang (2001) measured the concentrations of reactive oxygen species (ROS) in various size fractions of atmospheric aerosols and vehicle emissions in Taipei using a technique which would respond to ROS adsorbed on particle surfaces or dissolved in PBW. The ROS were not identified, but the ROS concentration was correlated with the atmospheric ozone concentration. The ROS concentrations were highest in the 0.18 to 1.0 μm particle size range and decreased by over 50% between analyses at 1 h and 115 h after collection. While this study does not demonstrate that PBW carries H_2O_2 , it does demonstrate the ability of particles to act as carriers of reactive species and offers a technique for the measurement of ROS absorbed on particles or dissolved in PBW.

PBW Effects on Visibility

Light scattering, and the resultant effects on visibility, depend on the size and refractive index of the particle. PBW contributes to light scattering just like any other component of fine PM. As discussed in detail in Chapter 4, the light scattering due to a given mass of PM can be estimated from the composition and the relative humidity using mass scattering coefficients for the chemical components that are higher for hygroscopic components.

PBW Effects on Dosimetry

As particles are inhaled, they will be exposed to the high relative humidities of the respiratory tract. Although supersaturation may occur in the early generations of the airways, by the time the aerosol reaches the air exchange region, the relative humidity will have reached 99.5% and the particles will have attained their equilibrium size (Sarangapani and Wexler, 1996). As discussed in Chapter 6, these size changes can have significant effects on the fractions of inhaled particles deposited in the various portions of the respiratory tract.

PBW as a Reaction Medium

If the air becomes saturated with water vapor, hygroscopic particles will grow into cloud droplets. The liquid water content of cloud droplets is much higher than that of aerosols at relative humidities below 100%. This liquid water provides a reaction medium where reactions can take place that do not take place in the air. For example, dissolved SO₂ can react with dissolved O₃ or dissolved H₂O₂ to form H₂SO₄ in cloud droplets (Lazrus et al., 1983; Schwartz, 1984, 1986; Martin, 1994; Hegg and Hobbs, 1982, 1986).

The chemistry of fog and cloud droplets is important in the formation of secondary pollutants. An extensive discussion was given in Section 3.3.1.4 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). The possibility of chemical reactions in PBW is controversial because of the much lower quantity of liquid water at lower relative humidities (Brock and Durham, 1984; McMurray and Wilson, 1982, 1983; McMurray et al., 1981).

2.1.3.4 Atmospheric Lifetimes and Removal Processes

The lifetimes of particles vary with size. Nuclei-mode particles rapidly grow into the accumulation mode. However, the accumulation mode does not grow into the coarse mode. Accumulation-mode fine particles are kept suspended by normal air motions and have very low deposition rates to surfaces. They can be transported thousands of km and remain in the atmosphere for a number of days. Coarse particles can settle rapidly from the atmosphere (within hours) and normally travel only short distances. However, when mixed high into the atmosphere, as in dust storms, the smaller-sized coarse-mode particles have longer lives and travel greater distances. Dry deposition rates are expressed in terms of a deposition velocity that varies with particle size, reaching a minimum between 0.1 and 1.0 μm aerodynamic diameter

(e.g., Lin et al., 1994). Accumulation-mode particles are removed from the atmosphere primarily by cloud processes. Fine particles, especially particles with a hygroscopic component, grow as the relative humidity increases, serve as cloud condensation nuclei, and grow into cloud droplets. If the cloud droplets grow large enough to form rain, the particles are removed in the rain. Falling rain drops impact coarse particles and remove them. Ultrafine or nuclei-mode particles are small enough to diffuse to the falling drop, be captured, and be removed in rain. Falling rain drops, however, are not nearly as effective in removing accumulation-mode particles as the cloud processes mentioned above. A more detailed discussion of particle deposition, including acid deposition, especially as it applies to deposition to vegetation, soil, and water surfaces, is given in Chapter 4 (*Environmental Effects of Airborne Particulate Matter*). Acid deposition and PM are intimately related, first, because particles contribute to the acidification of rain and, secondly, because the gas-phase species that lead to dry deposition of acidity are also precursors of particles. Therefore, reductions in SO₂ and NO_x emissions will decrease both acidic deposition and PM concentrations.

Sulfate, nitrate, and some partially oxidized organic compounds are hygroscopic and act as nuclei for the formation of cloud droplets. These droplets serve as chemical reactors in which (even slightly) soluble gases can dissolve and react. Thus, SO₂ can dissolve in cloud droplets and be oxidized to sulfuric acid by dissolved ozone or hydrogen peroxide. These reactions take place only in aqueous solution, not in the gas phase. Sulfur dioxide may also be oxidized by dissolved oxygen. This process will be faster if metal catalysts such as iron or manganese are present in solution. If the droplets evaporate, larger particles are left behind. If the droplets grow large enough, they will fall as rain; and the particles will be removed from the atmosphere with potential effects on the materials, plants, or soil on which the rain falls. (Similar considerations apply to dew.) Atmospheric particles that nucleate cloud droplets also may contain other soluble or nonsoluble materials such as metal salts and organic compounds that may add to the toxicity of the rain. Sulfuric acid, ammonium nitrate, ammonium sulfates, and organic particles also are deposited on surfaces by dry deposition. The utilization of ammonium by plants leads to the production of acidity. Therefore, dry deposition of particles can also contribute to the ecological impacts of acid deposition. These effects are discussed in Chapter 4 (Section 4.1).

2.1.4 Comparison of Fine and Coarse Particles

The physical and chemical properties of fine particles (including ultrafine particles and accumulation-mode particles) and coarse particles are summarized for comparison purposes in Table 2-2. These include important differences in sources, formation mechanisms, composition, atmospheric residence time, removal processes, and travel distances. Ensuing chapters in this document will also show that fine and coarse particles differ in aspects of concentrations, exposure, dosimetry, toxicology, and epidemiology. Collectively, these differences continue to warrant consideration of fine particles as a separate air pollutant class from coarse particles and the setting of separate standards for fine and coarse particles.

2.2 MEASUREMENT OF PARTICULATE MATTER

The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) summarized the sampling and analytical techniques for PM and acid deposition that had appeared in the literature since the earlier 1982 PM AQCD (U.S. Environmental Protection Agency, 1982). Excellent reviews have also been published by Chow (1995) and McMurry (2000). This section discusses problems in measuring PM; new techniques that attempt to alleviate these problems or measure problem species; Federal Reference Methods, speciation monitors, analytical methods for inorganic elements, organic and elemental carbon, and ionic species; and continuous and multiday monitors.

2.2.1 Particle Measurements of Interest

There are many PM components and parameters that are of interest across the various types of uses to which PM measurement data are applied. These uses include analyses of compliance with air quality standards and trends; source category apportionment studies related to the development of pollution reduction strategies and the validation of air quality models; studies related to health, ecological, and radiative effects; and characterization of current air quality for presentation to the public in the context of EPA's Air Quality Index. Particulate matter measurement components and parameters of specific interest for these various purposes are noted below and summarized in Table 2-3.

**TABLE 2-2. COMPARISON OF AMBIENT PARTICLES,
FINE PARTICLES (Ultrafine plus Accumulation-Mode) AND COARSE PARTICLES**

	Fine		
	Ultrafine	Accumulation	Coarse
Formation Processes:	Combustion, high-temperature processes, and atmospheric reactions		Break-up of large solids/droplets
Formed by:	Nucleation Condensation Coagulation	Condensation Coagulation Reactions of gases in or on particles Evaporation of fog and cloud droplets in which gases have dissolved and reacted	Mechanical disruption (crushing, grinding, abrasion of surfaces) Evaporation of sprays Suspension of dusts Reactions of gases in or on particles
Composed of:	Sulfate Elemental carbon Metal compounds Organic compounds with very low saturation vapor pressure at ambient temperature	Sulfate, nitrate, ammonium, and hydrogen ions Elemental carbon Large variety of organic compounds Metals: compounds of Pb, Cd, V, Ni, Cu, Zn, Mn, Fe, etc. Particle-bound water	Suspended soil or street dust Fly ash from uncontrolled combustion of coal, oil, and wood Nitrates/chlorides/sulfates from HNO ₃ /HCl/SO ₂ reactions with coarse particles Oxides of crustal elements (Si, Al, Ti, Fe) CaCO ₃ , CaSO ₄ , NaCl, sea salt Pollen, mold, fungal spores Plant and animal fragments Tire, brake pad, and road wear debris
Solubility:	Probably less soluble than accumulation mode	Largely soluble, hygroscopic, and deliquescent	Largely insoluble and nonhygroscopic
Sources:	Combustion Atmospheric transformation of SO ₂ and some organic compounds High temperature processes	Combustion of coal, oil, gasoline, diesel fuel, wood Atmospheric transformation products of NO _x , SO ₂ , and organic compounds, including biogenic organic species (e.g., terpenes) High-temperature processes, smelters, steel mills, etc.	Resuspension of industrial dust and soil tracked onto roads and streets Suspension from disturbed soil (e.g., farming, mining, unpaved roads) Construction and demolition Uncontrolled coal and oil combustion Ocean spray Biological sources
Atmospheric half-life:	Minutes to hours	Days to weeks	Minutes to hours
Removal Processes:	Grows into accumulation mode Diffuses to raindrops	Forms cloud droplets and rains out Dry deposition	Dry deposition by fallout Scavenging by falling rain drops
Travel distance:	< 1 to 10s of km	100s to 1000s of km	< 1 to 10s of km (small size tail, 100s to 1000s in dust storms)

Source: Adapted from Wilson and Suh (1997).

TABLE 2-3. PARTICULATE MATTER COMPONENTS/PARAMETERS OF INTEREST FOR HEALTH, ECOLOGICAL, OR RADIATIVE EFFECTS; FOR SOURCE CATEGORY APPORTIONMENT STUDIES; OR FOR AIR QUALITY MODELING EVALUATION STUDIES

- Particle number
 - Particle surface area
 - Particle size distribution
 - PM mass (fine PM mass [PM_{2.5}] and coarse thoracic PM mass [PM_{10-2.5}]) including both nonvolatile mass as measured by the current Federal Reference method and total mass (including semivolatile components such as ammonium nitrate and semivolatile organic compounds, but not particle-bound water)
 - Ions (sulfate, nitrate, and ammonium)
 - Strong acidity (H⁺)
 - Elemental carbon
 - Organic carbon (total, nonvolatile, and semivolatile; functional groups and individual species)
 - Transition metals (water soluble, bioavailable, oxidant generation)
 - Specific toxic elements and organic compounds
 - Crustal elements
 - Bioaerosols
 - Particle refractive index (real and imaginary)
 - Particle density
 - Particle size change with changes in relative humidity
-
-

Particle measurements are needed to determine if a location is in compliance with air quality standards, to determine long-term trends in air quality patterns, and for use in epidemiologic studies. For these purposes, the precision of the measurements made by a variety of measurement instruments in use is a critical consideration. Therefore, the intercomparisons of various samplers under a variety of atmospheric and air quality conditions are important.

In order to reduce pollution to attain a standard, pollution control agencies and national research organizations need measurements to identify source categories and to develop and validate air quality models. For these purposes, PM parameters other than mass, such as chemical composition and size distribution, must also be measured. Moreover, measurements are needed with shorter time resolutions in order to match changes in pollution with the diurnal changes in the boundary layer.

A number of PM measurements are needed for use in epidemiologic and exposure studies and to determine components of PM to guide the planning and interpretation of toxicologic studies. Thus, size and chemical composition measurements are important, as are measurement across different time intervals. For epidemiologic studies of acute (i.e., short-term) PM exposures, 1-h or continuous measurements can provide important information beyond that provided by 24-h measurements. However, for epidemiologic studies of chronic PM exposures, measurements that permit integration over longer intervals (e.g., a week to a month) are more relevant. For dosimetric studies and modeling, information will be needed on the particle size distribution and on the behavior of particles as the relative humidity and temperature changes found in the atmosphere are increased to those found in the respiratory system.

For studies of ecological effects and materials damage, measurements of particles and of the chemical components of PM in rain, fog, and dew are needed to understand the contributions of PM to the soiling of surfaces and damage to materials and to understand the wet and dry deposition of acidity and toxic substances to surface water, soil, and plants. Some differentiation into particle size is needed to determine dry deposition.

For studies of visibility impairment and radiative effects, information is needed regarding how particles scatter and absorb light, including data on refractive index, ratio of scattering to absorption, size distribution, and change in particle size with change in relative humidity.

2.2.2 Issues in Measurement of Particulate Matter

The EPA decision to revise the PM standards by adding daily and yearly standards for PM_{2.5} has led to a renewed interest in the measurement of atmospheric particles and to a better understanding of the problems in developing precise and accurate measurements of particles. It is very difficult to measure and characterize particles suspended in the atmosphere; however, numerous improvements in PM monitoring are in use and others are in development.

EPA's PM standards are based, in part, on epidemiologic relationships between health effects and PM concentrations as measured with existing monitoring methods. As understanding of suspended PM has advanced and new monitoring information has become available, EPA has changed the indicator for the PM NAAQS from TSP to PM₁₀, and added PM_{2.5}. During the current PM NAAQS review, consideration will be given to a standard for coarse thoracic PM.

The U.S. Federal Reference Methods (FRM) for PM_{2.5} and PM₁₀ provide relatively precise ($\pm 10\%$) methods for determining the mass of material remaining on a Teflon filter after equilibration. However, numerous uncertainties remain as to the relationship between the mass and composition of material remaining on the filter as determined by the FRM measurement procedure and the mass and composition of material that existed in the atmosphere as suspended PM. As a result, the EPA defines accuracy for PM measurements in terms of agreement of a candidate sampler with a reference sampler. Therefore, intercomparison of samplers is very important in determining how well various samplers agree and how various design choices influence what is actually measured.

There are eight general areas where choices are made in the design and use of an aerosol sampler. These include (1) consideration of positive artifacts due to chemical reaction or adsorption, (2) treatment of semivolatile components, (3) selection of particle size cut characteristics for the upper cut point, (4) separation of fine and coarse PM, (5) treatment of pressure, temperature, and relative humidity, (6) time resolution, (7) assessment of the reliability of the measurement technique, and (8) operation and maintenance procedures needed to maintain consistent measurements over time. In many cases, choices have been made without adequate recognition of their consequences. As a result, measurement methods developed by different organizations may give different results when sampling the same atmosphere even though the techniques appear to be similar.

2.2.2.1 Artifacts Due to Chemical Reactions

When TSP was collected on glass-fiber filters, the reaction of SO₂ (and other acid gases) with basic sites on the glass fiber or with basic coarse particles on the filter led to the formation of sulfate (or other nonvolatile salts, e.g., nitrate, chloride). These positive artifacts led to the overestimation of mass, sulfate, and probably also of nitrate. The metal impurities in the glass fiber resulted in high background levels that led to low precision in the measurement of trace

metals. These problems were largely overcome by changing to quartz fiber or Teflon filters and by the separate collection of $PM_{2.5}$. However, the possible reaction of acidic gases with basic coarse particles remains a possibility, especially with PM_{10} and $PM_{10-2.5}$ measurements. The reaction of NH_3 with acidic particles, either during sampling or during transportation, storage, and equilibration remains a problem in areas such as the eastern United States where PM is frequently acidic. One technique that has been developed to overcome this problem makes use of a denuder to remove NH_3 during sampling and to protect the collected PM from NH_3 (Suh et al., 1992, 1994; Brauer et al., 1991; Koutrakis et al., 1988a,b). However, this technique has been applied primarily for measurement of particle strong acidity, not for the measurement of artifact-free ammonium or mass concentrations. In the measurement of particle strong acidity, basic coarse particles must be separated from acidic fine particles (Koutrakis et al., 1992).

2.2.2.2 Treatment of Semivolatile Components of Particulate Matter

Current filtration-based mass measurements can result in significant evaporative losses, during and possibly after collection, of a variety of semivolatile components (i.e., species that exist in the atmosphere in dynamic equilibrium between the condensed phase and gas phase). Important examples include ammonium nitrate, semivolatile organic compounds, and particle-bound water. This problem is illustrated in Figure 2-18.

The following approaches, that have been used to address the problem of potentially lost semivolatile components, will be discussed in more detail in subsequent sections.

- (1) Collect/measure all components present in the atmosphere in the condensed phase except particle-bound water. Examples: Brigham Young absorptive sampler and Harvard pressure drop monitor. Both require preconcentration of the accumulation mode and reduction of ambient humidity by diffusion denuder techniques.
- (2) Stabilize PM at a specified temperature high enough to remove all, or almost all, particle-bound water. This results in loss of much of the semivolatile PM. Examples: tapered element oscillating microbalance (TEOM) operated at 50 °C beta gauge with heated inlet.
- (3) Equilibrate collected material at fixed, near-room temperature and moderate relative humidity to reduce particle-bound water. Accept the loss of an unknown but possibly significant fraction of semivolatile PM. Examples: U.S. Federal Reference Method and most filter-weighing techniques. Equilibration originally was designed to remove adsorbed water vapor from glass fiber filters in order to maintain a stable filter weight.




Figure 2-18. Schematic showing major nonvolatile and semivolatile components of PM_{2.5}. Semivolatile components are subject to partial-to-complete loss during equilibration or heating. The optimal technique would be to remove all particle-bound water but no ammonium nitrate or semivolatile organic PM.

The designated RH ($35 \pm 5\%$) was a compromise. If the RH is too low, electrostatic charging becomes a problem. The equilibration process does help provide a stable and reproducible mass. The equilibrium process reduces the amount of particle-bound water but may not remove all particle-bound water. Moreover, the equilibration process may lead to the loss of other semivolatile PM components.

2.2.2.3 Upper Cut Point

The upper cut point of the high volume sampler varies with wind speed and direction. However, it is usually desirable to have an upper cut point that is independent of these factors. Considerations in selecting an upper cut point are discussed in Section 2.1.2.2.

2.2.2.4 Cut Point for Separation of Fine and Coarse Particulate Matter

As shown in Table 2-2, fine and coarse particles differ not only in size but also in formation mechanisms, sources, and chemical, physical, and biological properties. They also differ in concentration-exposure relationships, dosimetry (deposition and retention in the respiratory system), toxicity, and health effects as observed in epidemiologic studies. Thus, it is desirable to measure fine and coarse PM separately in order to properly allocate health effects to either fine or coarse PM and to correctly determine sources by receptor modeling approaches. Considerations in selecting a cut point to separate fine and coarse particles are discussed in Section 2.1.2.2.

2.2.2.5 Treatment of Pressure, Temperature, and Relative Humidity

There are a variety of techniques for defining (or ignoring) the pressure, temperature, and relative humidity during and after sampling. For example, the sample volume may be based on the mass or volumetric flow corrected to standard temperature and pressure (298 K and 1 atm) (as in the current FRM for PM_{10}), or it may be based on the volumetric flow at ambient conditions of temperature and pressure (current FRM for $PM_{2.5}$).

There are also a variety of options for the control of temperature during collection. The particles may be heated enough to remove much of the particle-bound water (i.e., TEOM at 50 °C); the particles may be heated several degrees, just enough to prevent condensation of water in the sampling system; the particles and the sampler may be maintained near ambient temperature (± 5 °C of ambient temperature is required for EPA FRM samplers for $PM_{2.5}$); or the particles and sampler may be maintained at a constant temperature inside a heated or air conditioned shelter. There are also options for controlling of temperature after collection: (a) no control (room temperature) or (b) ship and store at cool temperature (4 °C is the current EPA FRM requirement).

Consideration must also be given to relative humidity (RH). Changes in RH cause changes in particle size of hygroscopic or deliquescent particles. Changing RH by adding or removing water vapor affects particle number, particle surface area, and particle size distribution measurements and the amount of overlap of accumulation-mode and coarse-mode particles. Changing RH by intentional or inadvertent changes in temperature also affects the relative loss of ammonium nitrate and semivolatile organic compounds. Monitoring personnel should be

aware of the various options for the treatment of pressure, temperature, and RH factors; make appropriate selections; and document which options are used.

Studies of relationships between personal/indoor/outdoor measurements present special problems. Indoor environments are typically dryer than outdoors and may be warmer or, if air-conditioned, cooler. These differences may change particle size and the amount of volatilization of semivolatile components. Such changes between indoors and outdoors will complicate the comparison of indoor to outdoor PM concentrations, the modeling of personal exposure to all particles, and apportionment of personal exposure into particles of ambient origin, particles of indoor origin, and particles originating from personal activity.

2.2.2.6 Time Resolution

The classic 24-h filter collection technique is being supplemented by a variety of continuous monitors for various PM constituents. This process is being accelerated by the lower operational cost of continuous monitors and the availability of new continuous monitors for mass, number, and certain chemical components, as well as refinements of older methods based on beta attenuation or light scattering. Most epidemiologic studies have used 24-hour concentrations as exposure indicators. However, one epidemiologic study of chronic effects uses a filter sampler with a 2-week collection period (Gauderman et al., 2000). Another recent study used 1 to 2 h concentrations (see Peters et al., 2000). Continuous methods are discussed in Section 2.2.5.

2.2.2.7 Accuracy and Precision

Precision is typically determined by comparing measurements obtained with collocated samplers or through replicate analyses; whereas accuracy is determined through the use of traceable calibration standards. Unfortunately, no standard reference calibration material or procedure has been developed for suspended, atmospheric PM. It is possible to determine the accuracy of certain components of the PM measurement system (e.g., flow control, inlet aspiration, PM_{2.5} cut, weighing, etc.). The absolute accuracy for collecting a test aerosol can also be determined by isokinetic sampling in a wind tunnel. However, it is not currently feasible to provide a simulated atmospheric aerosol with naturally occurring semivolatile components. It is particularly challenging to develop an atmospheric aerosol calibration standard suitable for

testing samplers in the field. Therefore, it is not possible at the present time to establish the absolute accuracy of a PM monitoring technique. Intercomparison studies may be used to establish the precision of identical monitors and the extent of agreement between different types of monitors. Such studies are important for establishing the reliability of PM measurements. Intercomparison studies have contributed greatly to our understanding of the problems in PM measurement. Such studies will be discussed as they apply to specific measurement problems, monitoring instruments, or analytical techniques.

Some measurement errors of concern in PM_{10} sampling, including those that arise due to uncertainty tolerances in cut point, particle bounce and reentrainment; impactor surface overloading; and losses to sampler internal surfaces, were discussed in detail in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). Other measurement errors of concern in $PM_{2.5}$ sampling arise because of our inability to assess accuracy in an absolute sense due to a lack of an atmospheric aerosol calibration standard, the inclusion in $PM_{2.5}$ of a small amount of coarse particles as discussed in Section 2.2.1.3, and problems associated with the definition of $PM_{2.5}$ as what remains on a filter after collection and equilibration, rather than as the mass of particles as they exist in the air. Still, it is possible to measure PM indicators with high precision. Detailed information on precision and quality assurance may be found on the EPA's Technology Transfer Network website (U.S. Environmental Protection Agency, 2002a). See discussion in Section 2.2.4.

Because of the difficulties associated with determining the accuracy of PM measurements, EPA has sought to make FRM measurements equivalent by specifying operating conditions and, in the case of $PM_{2.5}$ samplers, by specifying details of the sampler design. Thus, both the PM_{10} and the $PM_{2.5}$ standards are defined with consistency of measurement technique rather than with the accuracy of the true mass concentration measurement in mind (McMurry, 2000). It is acknowledged in the Federal Register (1997) that, "because the size and volatility of the particles making up ambient particulate matter vary over a wide range and the mass concentration of particles varies with particle size, it is difficult to define the accuracy of $PM_{2.5}$ measurements in an absolute sense. . . ." Thus, accuracy is defined as the degree of agreement between a field $PM_{2.5}$ sampler and a collocated $PM_{2.5}$ reference method audit sampler (McMurry, 2000). The Federal Reference Method for $PM_{2.5}$ is discussed in Section 2.2.3.3. As mentioned earlier, volatilization of organic compounds and ammonium nitrate during sampling or post-sampling

handling can lead to significant underestimation of the fine particulate mass concentration in some locations. Sources of error in the measurement of mass of $PM_{2.5}$ suspended in the atmosphere also arise because of adsorption or desorption of semivolatile vapors onto or from collected PM, filter media, or other sampler surfaces; neutralization of acid or basic vapors on either filter media or collected PM; and artifacts associated with particle-bound water.

During the past 25 years, there have been advancements in the generation and classification of monodisperse aerosols as well as in the development of electron microscopy and imaging analyses that have contributed to the advancement in aerosol calibration (Chen, 1993). Still, one of the limitations in PM sampling and analysis remains the lack of primary calibration standards for evaluating analytical methods and for intercomparing laboratories. Klouda et al. (1996) examined the possibility of resuspending the National Institute of Standards and Technology (NIST) Standard Reference Material 1649 (Urban Dust) in air for collection on up to 320 filters simultaneously using SRI's dust generation and collection system. However, the fine component does not resuspend and the semivolatile component evaporates. Consequently, this material is not a suitable standard for suspended PM. NIST is continuing work in this area with EPA support.

Method validation was discussed in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a), and the usefulness of intercomparisons and "internal redundancy" was emphasized. For example, a number of internal consistency checks are applied to the IMPROVE network (Malm et al., 1994). These include mass balances, sulfur measurements by both proton-induced X-ray emission (PIXE) and ion chromatography (IC), and comparison of organic matter by combustion and by proton elastic scattering (PESA) of hydrogen. Mass balances compare the gravimetrically determined mass with the mass calculated from the sum of the major chemical components (i.e., crustal elements plus associated oxygen, organic carbon, elemental carbon, sulfate, nitrate, ammonium, and hydrogen ions). Mass balances are useful validation techniques; however, they do not check for, or account for, artifacts associated with the absorption of gases during sampling, the loss of semivolatile material during sampling, or errors in assumptions regarding unmeasured "associated species." The mass balance check may appear reasonable even if such artifacts are present, because only the material collected on the filter is included in the balance. Mass balance checks may also suffer from errors due to some particle-bound water remaining in the PM even after equilibration and from the use of an

arbitrary factor to account for the amount of oxygen and hydrogen atoms per carbon atom in the organic carbon fractions (Andrews et al., 2000).

2.2.3 Measurement of Semivolatile Particulate Matter

Some atmospheric species, such as ammonium nitrate, water, and certain organic compounds have vapor pressures such that, for typical ambient concentrations and temperatures, they may exist in an equilibrium state with some material in the gas phase and some in the condensed (particulate) phase. Such species are known as semivolatile materials (SVM) and may partition to varying degrees between the gas and particle phases (Pankow, 1994a). Organic compounds with vapor pressure $< 10^{-6}$ Torr are nonvolatile, i.e., primarily in the particle phase. Organic compounds with vapor pressure $> 10^{-5}$ Torr will be primarily in the gas phase. Organic compounds with vapor pressures on the order of 10^{-5} to 10^{-6} are semivolatile. Semivolatile material, originally in the atmosphere in the particulate phase and collected on a filter, may subsequently be lost from the filter. For example, SVM may evaporate during sampling due to a reduction in its concentration in the atmosphere being sampled or due to the pressure drop across the filter. Or, SVM may evaporate after sampling; during intentional equilibration at a low relative humidity; or during transport, handling, and storage if exposed to an atmosphere in which the vapor pressure of one or more semivolatile components is lower than in the atmosphere sampled. Since water is not a pollutant, it is necessary to remove most of the particle-bound water before weighing (Chow, 1995). However, the collection and measurement of ammonium nitrate and semivolatile organic compounds in suspended atmospheric PM represents a major analytical challenge (McMurry, 2000).

2.2.3.1 Particle-Bound Water

It is generally desirable to collect and measure ammonium nitrate and semivolatile organic compounds. However, for many measurements of suspended particle mass, it is desirable to remove the particle-bound water before determining the mass. The mass of particle-bound water is strongly dependent on the relative humidity and the particle composition. However, the dependence on relative humidity is not linear, because there is significant hysteresis in the water adsorption-desorption pathways (Seinfeld and Pandis, 1998). Water vapor cannot be controlled, and particle-bound water is not included in the mass of PM subject to regulation and control.

components, a measurement of PM mass including particle-bound water would depend more on relative humidity than pollution. For all these reasons, it is usually desirable to remove most, if not all, particle-bound water before weighing collected PM. On the other hand, in some situations it may be important to know how much of the suspended particle mass or volume results from particle-bound water. Figures 2-16a and 2-16b show the change in diameter of sulfate particles as a function of relative humidity. Figure 2-16a also shows hysteresis resulting from the difference between deliquescence and crystallization points.

Pilinis et al. (1989) calculated the water content of atmospheric PM above and below the deliquescent point. They predicted that aerosol water content is strongly dependent on composition and concluded from their calculations that liquid water could represent a significant mass fraction of aerosol concentration at relative humidities above 60%. Since then, a few researchers have attempted to measure the water content of atmospheric aerosols. Most techniques have focused on tracking changes in the particle mass as the relative humidity is changes. Only a few demonstrations have been carried out using atmospheric particles. Of particular interest is the development of the Tandem Differential Mobility Analyzer (TDMA) and its applications in investigations of the effects of relative humidity on particle growth.

Lee et al. (1997) examined the influence of relative humidity on the size of atmospheric aerosols using a TDMA coupled with a scanning mobility particle sizer (SMPS). They reported that the use of the TDMA/SMPS system allowed for the abrupt size changes of aerosols at the deliquescence point to be observed precisely. They also reported that at relative humidities between 81 and 89% the water content of ammonium sulfate aerosols (by mass) ranged from 47 to 66%.

Andrews and Larson (1993) investigated the interactions of single aerosol particles coated with an organic film within a humid environment. Using an electrodynamic balance, they conducted laboratory experiments in which sodium chloride and carbon black particles were coated with individual organic surfactants (intended to simulate the surface-active, organic films that many atmospheric aerosol particles may exhibit) and their water sorption curves were examined. Their results showed that when ordinarily hydrophobic carbon black particles were coated with an organic surfactant, they sorbed significant amounts of water (20 to 40% of the dry mass of the particle).

Liang and Chan (1997) developed a fast technique using the electrodynamic balance to measure the water activity of atmospheric aerosols. In their technique, the mass of a levitated particle is determined as the particle either evaporates or grows in response to a step change in the relative humidity. Their technique was demonstrated using laboratory experiments with NaCl, $(\text{NH}_4)_2\text{SO}_4$, NaNO_3 , and $(\text{NH}_4)_2\text{SO}_4/\text{NH}_4\text{NO}_3$ solutions. They concluded that one of the advantages of their fast method is the ability to measure the water activity of aerosols containing volatile solutes such as ammonium chloride and some organics.

McInnes et al. (1996) measured aerosol mass concentration, ionic composition, and the associated water mass of marine aerosol over the remote Pacific Ocean. The mass of PBW was determined by taking the difference between the mass obtained at 48% RH and at 19% RH, assuming the aerosol particles were dry at 19% RH. Based on a comparison of the remote Pacific aerosol to an aerosol collected at a site at the marine/continental interface of the Washington coast, the amount of water associated with the aerosol was observed to be a function of the ammonium to sulfate ratio. They found that the amount of water associated with the submicrometer aerosol comprised 29% of the total aerosol mass collected at 47% RH and 9% of the total mass at 35% RH.

Ohta et al. (1998) characterized the chemical composition of atmospheric fine particles (50% cut point of 2 μm) in Sapporo, Japan and, as part of their measurements, determined the water content using the Karl Fischer method (Meyer and Boyd, 1959). After exposing a Teflon filter, a portion of the filter was equilibrated at 30% RH for 24 h. Then the filter piece was placed in a water evaporator and heated at 150 °C, vaporizing the particle-bound water. The evolved vapor was analyzed for water in an aqua-counter where it was titrated coulometrically in Karl Fischer reagent solution (containing iodine, sulfur, and methanol). The accuracy of the aqua-counter is ± 1 mg. Using this technique, they determined that the water content of the particles ranged from 0.4 to 3.2% of the total particulate mass (at RH < 30%). This represents a smaller portion of water compared to their previous reported values (Ohta and Okita, 1990), which were determined by calculation at 50% RH.

Speer et al. (1997) developed an aerosol liquid water content analyzer (LWCA) in which aerosol samples are collected on Teflon filters and then placed in a closed chamber in which the relative humidity is closely controlled. The aerosol mass is monitored using a beta-gauge, first as the relative humidity is increased from low RH to high RH, and then as the RH is

decreased again. They demonstrated the LWCA on a laboratory-generated aerosol and on an ambient $PM_{2.5}$ sample collected in Research Triangle Park, NC. The ambient aerosol sample was also analyzed for chemical constituents. It is interesting to note that, although their laboratory-generated $(NH_4)_2SO_4$ aerosol demonstrated a sharp deliquescent point, their atmospheric aerosol, which was essentially $(NH_4)_2SO_4$, did not show a sharp deliquescent point.

Hygroscopic properties of aerosols have been studied from the viewpoint of their ability to act as condensation nuclei. The hygroscopic properties of fresh and aged carbon and diesel soot particles were examined by Weingartner et al. (1997) who found that fresh, submicron-size particles tended to shrink with increasing relative humidity because of a structural change. Lammel and Novakov (1995) found, in laboratory studies, that the hygroscopicity of soot particles could be increased by chemical modification and that the cloud condensation nucleation characteristics of diesel soot were similar to those of wood smoke aerosol.

The results of several of the above studies in which aerosol water content was determined as a function of relative humidity are summarized in Figure 2-19, which includes the results of Lee et al. (1997), McInnes et al. (1996), and Ohta et al. (1998). Relative humidity ranged from 9%, at which the aerosol water content was assumed to be zero (McInnes et al., 1996), to 89%, at which the aerosol water content was determined to be 66% by mass (Lee et al., 1997). Koutrakis et al. (1989) and Koutrakis and Kelly (1993) also have reported field measurements of the equilibrium size of atmospheric sulfate particles as a function of relative humidity and acidity.




Figure 2-19. Aerosol water content expressed as a mass percentage, as a function of relative humidity.

The effects of relative humidity on particle growth were also examined in several studies. Fang et al. (1991) investigated the effects of flow-induced relative humidity changes on particle cut sizes for aqueous sulfuric acid particles in a multi-nozzle microorifice uniform deposit impactor (MOUDI). Laboratory experiments were conducted in which polydisperse sulfuric acid aerosols were generated and the RH was adjusted. The aerosols were analyzed by a differential mobility analyzer. Fang et al. (1991) observed that for inlet RH less than 80%, the cut sizes for the sulfuric acid aerosols were within 5% of that for nonhygroscopic particles except at the stage for which the cut size was 0.047 μm , (10.7% larger than the nonhygroscopic particle cut size). They concluded that flow-induced RH changes would have only a modest effect on MOUDI cut sizes at $\text{RH} < 80\%$.

Hitzenberger et al. (1997) collected atmospheric aerosol in the size range of 0.06 to 15 μm in Vienna, Austria using a nine-stage cascade impactor and measured the humidity-dependent water uptake when the individual impaction foils were exposed to high RH. They observed particle growth with varying growth patterns. Calculated extinction coefficients and single scattering albedo increased with humidity.

The hygroscopic properties, along with the mixing characteristics, of the submicrometer particles sampled in Los Angeles, CA during the summer of 1987 SCAQS study and at the Grand Canyon, AZ during the 1990 Navajo Generating Station Visibility Study were reported by Zhang et al. (1993). They used a tandem differential mobility analyzer (TDMA; McMurry and Stolzenburg, 1989) to measure the hygroscopic properties for particles in the 0.05 to 0.5 μm range. In their experimental technique, monodisperse particles of a known size are selected from the atmospheric aerosol with the first analyzer. Then, the relative humidity of the monodisperse aerosol is adjusted, and the new particle size distribution is measured with the second analyzer. At both sites, they observed that monodisperse particles could be classified according to “more” hygroscopic and “less” hygroscopic. Aerosol behavior observed at the two sites differed markedly. Within the experimental uncertainty ($\pm 2\%$) the “less” hygroscopic particles sampled in Los Angeles did not grow when the RH was increased to 90%; whereas at the Grand Canyon, the growth of the “less” hygroscopic particles varied from day to day, but ranged from near 0 to 40% when the RH was increased to 90%. The growth of the “more” hygroscopic particles in Los Angeles was dependent on particle size (15% at 0.05 μm to 60% at 0.5 μm); whereas at the Grand Canyon, the “more” hygroscopic particles grew by about 50% with the growth not

varying significantly with particle size. After comparing the TDMA data to impactor data, Zhang et al. (1993) surmised that the more hygroscopic particles contained more sulfates and nitrates, while the less hygroscopic particles contained more carbon and crustal components.

Although most of the work to date on the hygroscopic properties of atmospheric aerosols has focused on the inorganic fraction, the determination of the contribution of particle-bound water to atmospheric particulate mass is greatly complicated by the presence of organics. The effect of RH on the adsorption of semivolatile organic compounds is discussed elsewhere in this chapter. Saxena et al. (1995) observed that particulate organic compounds can also affect the hygroscopic behavior of atmospheric particles. They idealized the organic component of aerosol as containing a hydrophobic fraction (e.g., high-molecular weight alkanes, alkenoic acids, alkenoic acids, aldehydes, and ketones) and a hydrophilic fraction (e.g., lower molecular weight carboxylic acids, dicarboxylic acids, alcohols, aldehydes, etc.) that would be likely to absorb water. They then analyzed data from a TDMA in conjunction with particle composition observations from an urban site (Claremont, CA) and from a nonurban site (Grand Canyon) to test the hypothesis that, by adding particulate organics to an inorganic aerosol, the amount of water absorbed would be affected, and the effect could be positive or negative, depending on the nature of the organics added. They further presumed that the particulate organic matter in nonurban areas would be predominantly secondary, and thus hydrophilic, compared to the urban aerosol that was presumed to be derived from primary emissions and thus hydrophobic in nature. Their observations were consistent with their hypothesis, in that at the Grand Canyon, the presence of organics tended to increase the water uptake by aerosols; whereas at the Los Angeles site, the presence of organics tended to decrease water uptake.

Peng and Chan (2001) also recently studied the hygroscopic properties of nine water soluble organic salts of atmospheric interest using an electrodynamic balance operated at 25 °C. Salts studied included sodium formate, sodium acetate, sodium succinate, sodium pyruvate, sodium methanesulfonate, sodium oxalate, ammonium oxalate, sodium malonate, and sodium maleate. They observed that hygroscopic organic salts have a growth factor of 1.76 to 2.18 from RH = 10 to 90%, which is similar to that of typical hygroscopic inorganic salts such as NaCl and $(\text{NH}_4)_2\text{SO}_4$.

Nonequilibrium issues may be important for the TDMA, as well as for other methods of measuring water content. Although the approach to equilibrium as the RH is increased is

expected to be rapid for pure salts, it may be much slower for aerosols containing a complex mix of components (Saxena et al., 1995). For example, if an aerosol contains an organic film or coating, that film may impede the transport of water across the particle surface, thus increasing the time required for equilibrium (Saxena et al., 1995). Insufficient time to achieve equilibrium in the TDMA could result in underestimation of the water content.

2.2.3.2 Nitrate and Organic Species

Particulate Nitrates

It is now well known that volatilization losses of particulate nitrates occur during sampling on Teflon filters (e.g., Zhang and McMurry [1992]; see also Hering and Cass [1999] and Babich et al. [2000]). The effect on the accuracy of atmospheric particulate measurements from these volatilization losses is more significant for $PM_{2.5}$ than for PM_{10} . The FRM for $PM_{2.5}$ will likely suffer a loss of nitrates similar to that experienced with other simple filter collection systems. Sampling artifacts resulting from the loss of particulate nitrates represents a significant problem in areas such as southern California that experience high amounts of nitrates. Hering and Cass (1999) reported on errors in $PM_{2.5}$ mass measurements due to the volatilization of particulate nitrate (Figure 2-20). They examined data from two field measurement campaigns that were conducted in southern California: (1) the Southern California Air Quality Study (SCAQS) (Lawson, 1990) and (2) the 1986 CalTech study (Solomon et al., 1992). In both these studies, side-by-side sampling of $PM_{2.5}$ was conducted. One sampler collected particles directly onto a Teflon filter. The second sampler consisted of a denuder to remove gaseous nitric acid followed by a nylon filter that absorbed the HNO_3 as it evaporated from ammonium nitrate. In both studies, the denuder consisted of MgO-coated glass tubes (Appel et al., 1981). Fine particulate nitrate collected on the Teflon filter was compared to fine particulate nitrate collected on the denuded nylon filter. In both studies, the $PM_{2.5}$ mass lost because of ammonium nitrate volatilization represented a significant fraction of the total $PM_{2.5}$ mass. The fraction of mass lost was higher during summer than during fall (17% versus 9% during the SCAQS study, and 21% versus 13% during the CalTech study; Figure 2-20). In regard to percentage loss of nitrate, as opposed to percentage loss of mass discussed above, Hering and Cass (1999) found that the amount of nitrate remaining on the Teflon filter samples was on average 28% lower than that on the denuded nylon filters.




Figure 2-20. Amount of ammonium nitrate volatilized from Teflon filters, expressed as a percentage of the measured $PM_{2.5}$ mass, for the SCAQS and CalTech studies, for spring and fall sampling periods.

Source: Hering and Cass (1999).

Hering and Cass (1999) also analyzed these data by extending the evaporative model developed by Zhang and McMurry (1987). The extended model used by Hering and Cass (1999) takes into account the dissociation of collected particulate ammonium nitrate on Teflon filters into nitric acid and ammonia via three mechanisms: (1) the scrubbing of nitric acid and ammonia in the sampler inlet (John et al. [1988] showed that clean PM_{10} inlet surfaces serve as an effective denuder for nitric acid); (2) the heating of the filter substrate above ambient temperature by sampling; and (3) the pressure drop across the Teflon filter. For the sampling systems modeled, the flow-induced pressure drop was measured to be less than 0.02 atm, and the corresponding change in vapor pressure was 2%, so losses driven by pressure drop were not considered to be significant in this work. Losses from Teflon filters were found to be higher during the summer compared to the winter, higher during the day compared to night, and reasonably consistent with modeled predictions.

Finally, during the SCAQS (Lawson, 1990) study, particulate samples also were collected using a Berner impactor and greased Tedlar substrates in size ranges from 0.05 to 10 μm in aerodynamic diameter. The Berner impactor $PM_{2.5}$ nitrate values were much closer to those

from the denuded nylon filter than those from the Teflon filter, the impactor nitrate values being ~2% lower than the nylon filter nitrate for the fall measurements and ~7% lower for the summer measurements. When the impactor collection was compared to the Teflon filter collection for a nonvolatile species (sulfate), the results were in agreement. Chang et al. (2000) discuss reasons for reduced loss of nitrate from impactors.

It should be noted that filters or collection surfaces were removed immediately after sampling and placed into vials containing a basic extraction solution during these intercomparison studies to preclude losses that might occur during their handling, storage, and equilibration. The loss of nitrate observed from Teflon filters and impaction surfaces in this study, therefore, is a lower limit compared to losses that might occur during the normal processes involved in equilibration and weighing of filters and impaction surfaces. Brook and Dann (1999) observed much higher nitrate losses during a study in which they measured particulate nitrate in Windsor and Hamilton, Ontario, Canada, by three techniques: (1) a single Teflon filter in a dichotomous sampler, (2) the Teflon filter in an annular denuder system (ADS), and (3) total nitrate including both the Teflon filter and the nylon back-up filter from the ADS. The Teflon filter from the dichotomous sampler averaged only 13% of the total nitrate, whereas the Teflon filter from the ADS averaged 46% of the total nitrate. The authors concluded that considerable nitrate was lost from the dichotomous sampler filters during handling, which included weighing and X-ray fluorescence (XRF) measurement in a vacuum.

Kim et al. (1999) also examined nitrate sampling artifacts by comparing denuded and undenuded quartz and nylon filters, during the PM₁₀ Technical Enhancement Program (PTEP) in the South Coast Air Basin of California. They observed negative nitrate artifacts (losses) for most measurements; however, for a significant number of measurements, they observed positive nitrate artifacts. Kim et al. (1999) pointed out that random measurement errors make it difficult to measure true amounts of nitrate loss.

Diffusion denuder samplers, developed primarily to measure particle strong acidity (Koutrakis et al., 1988a,b, 1992), also can be used to study nitrate volatilization. Such techniques were used to measure loss of particulate nitrate from Teflon filters in seven U.S. cities (Babich et al., 2000). Measurements were made with two versions of the Harvard-EPA Annular Denuder System (HEADS). Nitric acid vapor was removed by a Na₂CO₃-coated denuder. Particulate nitrate was the sum of nonvolatile nitrate collected on a Teflon filter and

volatilized nitrate collected on a Na_2CO_3 -coated filter downstream of the Teflon filter (full HEADS) or on a Nylon filter downstream of the Teflon filter (Nylon HEADS). It was found that the full HEADS (using a Na_2CO_3 filter) consistently underestimated the total particulate nitrate by approximately 20% compared to the nylon HEADS. Nonvolatilized nitrate was also measured on a Teflon filter from collocated Harvard Impactors (HIs). The $\text{PM}_{2.5}$ HI sampler, like the $\text{PM}_{2.5}$ FRM, uses impactors with a 50% cut at 2.5 μm . The HI uses a 37-mm filter and a flow rate of 10 L/min (0.465 L/min/cm²), while the FRM uses a 47-mm filter and a flow rate of 16.7 L/min (0.481 L/min/cm²). Therefore, the flow rate and pressure drop across the filters are similar and the loss of nitrate should be similar for both types of samples. Babich et al. (2000) found significant nitrate losses in Riverside, CA; Philadelphia, PA; and Boston, MA but not in Bakersfield, CA; Chicago, IL; Dallas, TX; or Phoenix, AZ where measurements were made only during the winter. Tsai and Huang (1995) used a diffusion denuder to study the positive and negative artifacts on glass and quartz filters. They found positive artifacts attributed to SO_2 and HNO_3 reaction with basic sites on glass fibers and basic particles and negative artifacts attributed to loss of HNO_3 and HCl due to volatilization of NH_4NO_3 and NH_4Cl and reaction of these species with acid sulfates.

Eatough et al. (1999a) developed a high-volume diffusion denuder system that combined diffusion denuder and particle concentrator techniques (see Section 2.2.3.2). In this system, the particle concentrator reduces the flow through the denuder so that the denuder can be operated for weeks without a loss of collection efficiency, making the sampler suitable for routine field sampling. The system was evaluated for the collection of fine particulate sulfate and nitrate in Riverside, CA (Eatough et al., 1999b). Concentrations of $\text{PM}_{2.5}$ nitrate obtained from the PC-BOSS agreed with those obtained using the Harvard-EPA Annular Denuder Sampler, HEADS (Koutrakis et al., 1988b).

Particulate Organic Compounds

Many semivolatile organic compounds (SVOCs) are of interest because of their possible health effects. Semivolatile organic compounds include products of incomplete combustion such as polycyclic aromatic hydrocarbons (PAHs) and polycyclic organic matter, which has been identified as a hazardous air pollutant. Polycyclic aromatic hydrocarbons also have been suggested as alternative particulate tracers for automobile emissions, because lead is no longer a

good tracer for automobile emissions sources due to the phase-out of organo-lead additives in gasoline (Venkataraman et al., 1994). PAHs also are emitted during biomass burning, i.e., burning of cereal crop residues and wood fuels (Jenkins et al., 1996; Roberts and Corkill, 1998).

A number of studies have used absorbing material following quartz filters to determine phase equilibria of specific organic compounds (Liang et al., 1997; Gundel et al., 1995; Kamens et al., 1995). Much work has also gone into the development of a theory to help understand the phase relationships (Yamasaki et al., 1982; Rounds and Pankow, 1990; Pankow, 1987, 1994a,b; Pankow et al., 1993; Rounds et al., 1993; Odum et al., 1994). The development of a theory describing phase equilibria of SVOC and the gas/particle partitioning of SVOC on inorganic, organic, and ambient smog aerosols is ongoing (Liang et al., 1997; Jang et al., 1997; Strommen and Kamens, 1997; Jang and Kamens, 1998, 1999, 2001; Leach et al., 1999; Kamens et al., 1999; Kamens and Jaoui, 2001; Chandramouli and Kamens, 2001; Chandramouli et al., 2003).

The mass of OC and EC is usually determined by the analysis of PM collected on a quartz filter. However, quartz fiber filters have a large specific surface area on which adsorption of gases can also occur. Possible artifacts associated with adsorption of organic gases onto quartz filters have been examined in experiments in which two quartz-fiber filters were deployed in series. The second quartz filter may indicate gaseous volatile organic compounds (VOCs) adsorbed on both filters (positive artifact), SVOCs evaporated from particles on the first filter and subsequently adsorbed on the second filter (negative artifact) or a combination of both effects. Unless the individual compounds are identified, the investigator does not know what to do with the loading value on the second filter (i.e., to add or subtract from the first filter loading value). Moreover, even if the individual compounds were identified on the back-up filter, the decision concerning adding or subtracting the back-up filter loading would not be straightforward.

The positive quartz filter artifact has been discussed by Gundel et al. (1995) and Turpin et al. (2000). It is also possible that some SVOCs may desorb from the filter resulting in a negative artifact (Eatough et al., 1993, 1995; Tang et al., 1994; Gundel et al., 1995; Cui et al., 1998; Pang et al., 2001; Finn et al., 2001). Semivolatile organic compounds can similarly be lost from Teflon filters because of volatilization, causing the PM_{2.5} mass to be significantly underestimated (negative artifact). Like particulate nitrates, the FRM for PM_{2.5} will suffer loss of SVOC similar to the losses experienced with other simple filter collection systems. Most

studies that have focused on the positive and negative sampling artifacts associated with SVOCs have used either diffusion denuder technology (Eatough et al., 1995; Mader et al., 2001) or placed an adsorbent media, such as a back-up quartz filter or a polyurethane foam adsorbent, behind the main filter (Wallace and Hites, 1995). Further information on denuder techniques is given in Appendix 2A.

Using their multichannel diffusion denuder sampling system (BOSS), Eatough et al. (1995) reported that, for samples collected at the South Coast Air Quality Management District sampling site at Azusa, CA, changes in the phase distribution of SVOCs could result in an average loss of 35% of the particulate organic material. Cui et al. (1998) found that losses of SVOCs from particles in the Los Angeles Basin during the summer were greater during the night (average = 62%) than during the day (average = 42%).

The percent SVOC lost from the front filter in a filter-denuder system may be greater than that lost in a filter-only system such as the FRM. In a filter-denuder system, the gas-phase component of the SVOCs is removed. The absence of the gas-phase causes the gas-particle equilibrium to shift, so the SVOCs collected on the filter may evaporate more rapidly in a filter-denuder system than in a filter-only collection system. To determine the fraction of SVOCs lost from a Teflon filter in a filter-only system, it is necessary to compare the amount measured by a nondenuder system with that measured by a denuder system. (Similar considerations apply to the collection of ammonium nitrate. However, in the case of ammonium nitrate, the total particulate nitrate is easily obtainable from the sum of nitrate on the Teflon front filter and the back-up Nylon filter. In the case of SVOCs, the existence of the positive artifact [since the organic denuder is less efficient than the nitric acid denuder], makes it much more difficult to determine the total OC.) At present, little information is available on the volatilization losses of SVOCs. However, in one study (Pang et al., 2001), the total mass on denuded and undenuded filters were compared and found to be identical within error limits ($R^2 = 0.816$, slope = 0.961 ± 0.027 for total mass compared to $R^2 = 0.940$, slope = 0.986 ± 0.020 for sulfate). Pang et al. (2001) interpreted this result as suggesting that the major cause of loss of SVOCs is the pressure drop across the filter.

Positive artifacts may occur during sample collection because of the adsorption of gases onto the filter materials (e.g., Gundel et al., 1995). Using a quartz filter behind a Teflon filter, Kim et al. (2001a) estimated that on an annual average basis 30% of the $PM_{2.5}$ organic carbon

concentration resulted from positive artifacts. Quartz fiber filters have a larger positive artifact than Teflon filters because of a greater adsorption of organic vapor onto quartz fibers (Turpin et al., 1994; Chow et al., 1994a,b, 1996; Eatough et al., 1996; Finn et al., 2001; Kirchstetter et al., 2001).

Kirchstetter et al. (2001) reported that adsorptive properties of quartz fiber filters vary with lot number; therefore, front and back-up filters should be taken from the same lot. The literature suggests that a Teflon filter followed by a quartz back-up filter provides a better estimate of the adsorption of gases on a quartz fiber front filter than does a quartz filter followed by a quartz backup, and that the difference between these two adsorption estimates can be substantial for short durations (Novakov et al., 1997; Kirchstetter et al., 2001; Turpin et al., 2000). The typically lower organic carbon loadings on concurrently collected quartz followed by quartz filters relative to Teflon followed by quartz filters are believed to occur because adsorption on the quartz front filter acts to reduce the gas-phase concentration downstream until adsorption equilibrium has been achieved in the vicinity of the front quartz filter surface. Because Teflon filters have little affinity for organic vapors, this equilibrium occurs almost instantaneously for Teflon filters; and the Teflon-quartz back-up filter is exposed to the ambient concentration of organic vapors from the beginning of the sampling period. It might be expected that the quantity of organic vapor adsorbed on quartz filters would depend on the organic composition and would vary by season and location. However, it is also possible that the quartz possesses a limited number of adsorption sites that are rapidly occupied, so that the quantity of OC on the back up filter would be relatively constant and depend on the pretreatment of the quartz.

Combined Measurement of Semivolatile Nitrate and Organic Carbon and Nonvolatile (Organic Carbon, Elemental Carbon, Nitrate, and Sulfate)

Fine particles in urban atmospheres contain substantial quantities of semivolatile material (e.g., NH_4NO_3 and SVOCs) that are lost from particles during collection on a filter. Several diffusion denuder samplers have been developed for the determination of both NO_3^- and organic semivolatile fine particulate components as well as nonvolatile nitrate, organic compounds, and nonvolatile sulfate (Pang et al., 2001; Eatough et al., 1993). The combination of technology used in the BOSS diffusion denuder sampler and the Harvard particle concentrator has resulted in the Particle Concentrator-Brigham Young University Organic Sampling System (PC-BOSS) for the 24-h integrated collection of $\text{PM}_{2.5}$, including NH_4NO_3 and SVOC. Modifications of the

BOSS sampler allow for the determination of these same species on a time scale from a few hours to weekly (Ding et al., 2002; Eatough et al., 1999a,b, 2001). Episodic studies have been conducted in Riverside, CA, and Bakersfield, CA, (Obeidi et al., 2002) and Provo, UT (Obeidi and Eatough, 2002). The average concentration of semivolatile and nonvolatile components in the three cities, during the study periods, are shown in Figure 2-21.


2.2.3.3 Continuous Measurement of Semivolatile and Nonvolatile Mass

Techniques for the continuous measurement of PM mass are needed both to provide real-time information on pollution levels (Long et al., 2002) and to reduce the costs involved in visiting sites to change filters and in the equilibration and weighing of filters. Two techniques are currently in use. The TEOM is normally operated at 50 °C in order to remove particle-bound water. However, at 50 °C most semivolatile material is also evaporated. Therefore, the TEOM, operated at 50 °C, may be considered to measure the mass of nonvolatile PM. Since the filter in the beta gauge mass monitor is changed more frequently than the filter in the TEOM, the beta gauge is less sensitive to changes in mass caused by changes in relative humidity. However, most beta gauge monitors heat the inlet, but do not otherwise control the temperature at the filter. This heating causes evaporation of a substantial fraction of the particle-bound water and an unknown fraction of the semivolatile PM. Thus, the beta gauge may be considered to measure the nonvolatile PM plus a small fraction of the particle-bound water and an unknown fraction of the semivolatile PM. Three new techniques have been developed to address the issue of lost semivolatile PM mass: the real-time ambient mass sampler (RAMS), the differential TEOM, and the continuous ambient mass monitor (CAMM).


Real-Time Total Ambient Mass Sampler (RAMS)

The RAMS, a monitor based on diffusion denuder and TEOM monitor technology, has been developed, validated, and field tested for the real-time determination of total fine PM mass, including semivolatile PM (Eatough et al., 1999a; Obeidi and Eatough, 2002; Obedi et al., 2002; Pang et al., 2001). The RAMS measures the total mass of collected particles including semivolatile species with a TEOM monitor using a “sandwich filter.” The sandwich contains a Teflon coated particle collection filter followed by a charcoal-impregnated filter (CIF) to collect any semivolatile species lost from the particles during sampling. Because the instrument

Riverside, CA; September 1997



Bakersfield, CA; March 1998



Provo, UT; September 1998




Figure 2-21. Average concentration (µg/m³) of nonvolatile and semivolatile PM components in three cities.

Source: Obeidi and Eatough (2002); Obeidi et al. (2002).

measures total mass collected by the sandwich filter, all gas phase compounds that can be adsorbed by a CIF must be removed from the sampling stream prior to the TEOM monitor. Laboratory and field validation data indicate that the precision of fine PM mass determination is better than 10%. The RAMS uses a Nafion dryer to remove particle-bound water from the suspended particles and a particle concentrator to reduce the amount of gas phase organics that must be removed by the denuder. Examples of $PM_{2.5}$ mass as measured by the RAMS, the TEOM, the PC-BOSS, and an FRM sampler are shown in Figure 2-22. This figure also shows that the TEOM recorded the $PM_{2.5}$ mass as being negative for the hours of 16 to 19. This likely results from the loss of volatile materials from the heated filter.




Figure 2-22. Comparison of mass measurements with collocated RAMS (real-time data), PC-BOSS (1-h data), FRM $PM_{2.5}$ sampler (average of 24-h data), and a conventional TEOM monitor (real-time data). The semivolatile fine particulate matter is sampled with the RAMS and PC-BOSS, but not with the TEOM monitor or the FRM $PM_{2.5}$ single filter sampler. The PC-BOSS provides information on both the nonvolatile component (NV) and the semivolatile organic component (SVOC).

Source: Eatough et al. (1999).

Differential TEOM

Patashnick et al. (2001) developed a differential TEOM system that is based on a pair of TEOM sensors, each of which is preceded by its own electrostatic precipitator (ESP) and downstream from a common size selective inlet. By alternately switching the ESPs on and off and out of phase with each other, the two sensors measure “effective mass” that includes both the nonvolatile component and the volatile component sampled by the TEOM, less the volatile component that vaporized during the sampling interval. On the sensor side with the ESP turned on, there is no particle collection on that filter so that only volatilization of previously collected particles continues. This allows a correction for the effective mass as measured by the first sensor by subtracting out the volatilization artifact and leaving the nonvolatile and volatile components of the PM. This system has yet to be well characterized for other biases or interferences such as reactions on the filters, particle collection efficiency of the ESPs, and particle and semivolatile material losses.

Continuous Ambient Mass Monitor (CAMM)

Koutrakis and colleagues (Koutrakis et al., 1996; Wang, 1997) developed CAMM, a technique for the continuous measurement of ambient particulate matter mass concentration based on the measurement of pressure drop increase with particle loading across a membrane filter. Recently, Sioutas et al. (1999) examined the increase in pressure drop with increasing particle loading on Nucleopore filters. They tested filters with two pore diameters (2 and 5 μm) and filter face velocities ranging from 4 to 52 cm s^{-1} and examined the effects of relative humidity in the range of 10 to 50%. They found that, for hygroscopic ammonium sulfate particles, the change in pressure drop per unit time and concentration was a strong function of relative humidity, decreasing with increasing relative humidity. These results suggest that particulate concentration measurements made with the pressure drop method (Koutrakis et al., 1996) may be subject to additional uncertainties if used in an environment where the ambient RH is quite variable and the RH where the particles are measured cannot be controlled accurately. The current version of the CAMM (Wang, 1997) uses a particle concentrator and a Nafion dryer and frequently moves the filter tape to avoid artifacts due to evaporation of semivolatile components from the active portion of the filter tape which would occur if the atmospheric concentration of the semivolatile components decreased.

The CAMM was recently operated alongside a gravimetric PM method (the Harvard Impactor, or HI) in seven U.S. cities selected for their distinctly different ambient particulate compositions and densities. The correlation between the two methods was high, with an overall r^2 of 0.90 and average CAMM/HI ratio of 1.07 (Babich et al., 2000).

2.2.4 U.S. Environmental Protection Agency Monitoring Methods

2.2.4.1 The Federal Reference Methods for Measurement of Equilibrated Mass for PM_{10} , $PM_{2.5}$, and $PM_{10-2.5}$

In 1997, the EPA promulgated new standards for $PM_{2.5}$ to address fine-fraction thoracic particles and retained with minor revisions, the 1987 PM_{10} standards to continue to address coarse-fraction thoracic particles (Federal Register, 1997). In partial response to numerous challenges to these standards, the U.S. Court of Appeals for the District of Columbia Circuit in *American Trucking Association v. EPA*, 175 F. 3d 1027 (U.S. Court of Appeals, D.C. Cir. 1999) found “ample support” for regulating coarse-fraction particles. However, the court revoked the revised PM_{10} standards (leaving in effect the 1987 PM_{10} standards) on the basis of PM_{10} being a “poorly matched indicator for coarse particulate pollution” because PM_{10} includes fine particles. Consistent with this specific aspect of the Court’s ruling, which the EPA did not appeal, the EPA is now considering use of $PM_{10-2.5}$ as the indicator for coarse-fraction thoracic particles, in conjunction with $PM_{2.5}$ standards that address fine-fraction thoracic particles. Thus, the EPA is now developing a Federal Reference Method for the measurement of $PM_{10-2.5}$.

2.2.4.1.1 PM_{10}

The FRM for measuring PM_{10} is specified in the Code of Federal Regulations (2001a,b). The PM_{10} FRM defines performance specifications for samplers in which particles are inertially separated with a penetration efficiency of 50% at an aerodynamic diameter (D_a) of $10 \pm 0.5 \mu\text{m}$. The collection efficiency increases to $\approx 100\%$ for smaller particles and drops to $\approx 0\%$ for larger particles. Particles are collected on filters and mass concentrations are determined gravimetrically. Instrument manufacturers are required to demonstrate through field tests a measurement precision for 24-h samples of $\pm 5 \mu\text{g}/\text{m}^3$ for PM_{10} concentrations below $80 \mu\text{g}/\text{m}^3$ and 7% above this value. A number of samplers have been designated as PM_{10} reference samplers. The TEOM and several beta gauge samplers with 1-h time resolution have been designated as automated equivalent methods (U.S. Environmental Protection Agency, 2001).

2.2.4.1.2 $PM_{2.5}$

In contrast to the performance-based FRM standard for PM_{10} , the FRM for $PM_{2.5}$ (Code of Federal Regulations, 2001a) specifies certain details of the sampler design, as well as of sample handling and analysis, whereas other aspects have performance specifications (Noble et al., 2001). The $PM_{2.5}$ FRM sampler consists of a PM_{10} inlet/impactor, a $PM_{2.5}$ impactor with an oil-soaked impaction substrate to remove particles larger than $2.5 \mu m D_a$, and a 47-mm PTFE filter with a particle collection efficiency greater than 99.7%. The sample duration is 24 h, during which time the sample temperature is not to exceed ambient temperatures by more than $5 \text{ }^\circ\text{C}$. A schematic diagram of the $PM_{2.5}$ FRM sample collection system is shown in Figure 2-23. After collection, samples are equilibrated for 24 h at temperatures in the range of 20 to $23 \text{ }^\circ\text{C}$ ($\pm 2 \text{ }^\circ\text{C}$) and at relative humidities in the range of 30 to 40% ($\pm 5\%$). The equilibration tends to reduce particle-bound water and stabilizes the filter plus sample weight. Filters are weighed before and after sampling under the same temperature and relative humidity conditions. For sampling conducted at ambient relative humidity $< 30\%$, mass measurements at relative humidities down to 20% are permissible (Code of Federal Regulations, 2001a).

The PM_{10} inlet specified for the $PM_{2.5}$ FRM is modified from a previous low flow rate PM_{10} inlet that was acceptable in both EPA-designated reference and equivalent PM_{10} methods. The modification corrects a flaw that was reported for the previous sampler in that under some meteorological conditions the inlet may allow precipitation to penetrate the inlet. The modification includes a larger drain hole, a one-piece top plate, and louvers. Tolocka et al. (2001a) evaluated the performance of this modified inlet in a series of wind tunnel experiments. The modified inlet was found to provide a size cut comparable to the original inlet, for both $PM_{2.5}$ and PM_{10} sampling. Because the modification did not change the characteristics of the size cut, the modified inlet may be substituted for the original inlet as part of a reference or equivalent method for PM_{10} and $PM_{2.5}$ (Tolocka et al., 2001a).

WINS Impactor. Design and calibration of the EPA $PM_{2.5}$ Well Impactor Ninety-Six (WINS) is given by Peters et al. (2001a). The WINS impactor was designed to be deployed downstream of the Graseby-Andersen 246B PM_{10} inlet as part of a sampler operating at a flow rate of 16.7 L/m. A schematic diagram of the WINS is shown in Figure 2-24. The $PM_{2.5}$ inlet consists of a single jet directed toward a round hole with a jet exit impaction surface made




Figure 2-23. Schematic diagram of the sample collection portion of the $PM_{2.5}$ FRM sampler.

Source: Noble et al. (2001).

up of a 37-mm diameter glass fiber filter immersed in 1 ml of low volatility diffusion pump oil (i.e., the well). Particles not having enough inertia to be removed by the impactor are captured downstream on the sample collection filter. This design was selected to minimize impactor overloading that would otherwise result in particle bounce. The oil wicks through the particulate deposit on the impactor to provide a continuously wetted surface for impaction. The penetration curve indicated a 50% cutpoint of $2.48 \mu m D_a$ with a geometric standard deviation of 1.18 (Figure 2-25).

The WINS separator was evaluated for its loading characteristics (Vanderpool et al., 2001) by monitoring the performance after repeated operation in an artificially generated, high concentration, coarse-mode aerosol composed of Arizona Test Dust, as well as in the field in Rubidoux, Phoenix, Philadelphia, Research Triangle Park, and Atlanta. In the wind tunnel experiments, the WINS performance was found to be a monotonic function of loading.




Figure 2-24. Schematic view of the final design of the WINS.

Source: Peters et al. (2001a).

A -5% bias in the $PM_{2.5}$ measurement resulted from a coarse particulate loading of approximately 16 mg. This negative bias was due to a slight reduction in the separator cut point. It was also found that the predictable results from the controlled laboratory experiments could not be extrapolated to field settings and that the WINS performance was more sensitive to the impactor loading in the field than it was in experiments with the single component aerosol. Significant particle bounce was not observed in either the laboratory or the field experiments. Vanderpool et al. (2001) concluded that their study supports the recommendation that the FRM WINS wells should be replaced after every 5 days of 24-h operation (U.S. Environmental Protection Agency, 1998).

A detailed sensitivity study of the WINS impactor was conducted (Vanderpool et al., 2001) in which the effects on the impactor performance of a number of parameters were examined. The results of this study are summarized in Table 2-4.




Figure 2-25. Evaluation of the final version of the WINS.

Source: Peters et al. (2001a).

The regulations also allow for Class I, II, and III equivalent methods for $\text{PM}_{2.5}$ (Code of Federal Regulations, 2001c). Class I equivalent methods use samplers with relatively small deviations from the sampler described in the FRM. Class II equivalent methods include “all other $\text{PM}_{2.5}$ methods that are based upon 24-h integrated filter samplers that are subjected to subsequent moisture equilibration and gravimetric mass analysis.” Class III equivalent methods include non-filter-based methods such as beta attenuation, harmonic oscillating elements, or nephelometry (McMurry, 2000). As of July 2001, 11 $\text{PM}_{2.5}$ samplers (listed in Table 2-5) had been tested, leading to the conclusion that the PM_{10} sampling systems can be designed to produce concentration measurements that are precise to $\pm 10\%$. Cut point tolerances are not expected to affect the mass concentration for $\text{PM}_{2.5}$ as much as for PM_{10} , because the $2.5 \mu\text{m } D_a$ cut point generally occurs near the minimum in a mass distribution (e.g., Figure 2-6).

**TABLE 2-4. SUMMARY OF SENSITIVITY STUDIES OF
WINS IMPACTOR PERFORMANCE**

Parameter	Amount of variance	Cutpoint variation	PM_{2.5} mass concentration bias
Manufacturing tolerances on WINS components	Specified tolerances	0.05 micrometers	< 1%
Flow control biases	4%	0.05 micrometers	Cutpoint shift partially offset volume bias
T and P measurement	Allowable ambient	± 0.02 micrometers	± 0.4%
Diffusion oil volume	0.75 mL to 3 mL	No effect	
Impactor loading	After 5 24 h events	-0.07 micrometers	< 1.5%
Ambient P variations		Negligible	Negligible
Air Properties	0 °C	2.40 micrometers	NA
Impactor oil crystallization		No effect	No effect
Impactor oil viscosity	-20 °C	No effect	No effect
	-35 °C	Need to change WINS more frequently than every 5 days	

Source: Vanderpool et al. (2001).

On the other hand, the PM_{2.5} mass concentration will be affected by other sampling issues mentioned, but not discussed extensively, in the previous 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). These issues (described earlier in this chapter) include gas/particle, particle/particle, and particle/substrate interactions for sulfates and nitrates (e.g., Appel et al., 1984), volatilization losses of nitrates (Zhang and McMurry, 1992), SVOC artifacts (e.g., Eatough et al., 1993), and relative humidity effects (e.g., Keeler et al., 1988).

Several studies have been reported in which the FRM was collocated with other PM_{2.5} samplers in intercomparison studies. During the Aerosol Research and Inhalation Epidemiology Study (ARIES), several PM_{2.5} samplers were collocated at a mixed industrial-residential site near Atlanta, GA (Van Loy et al., 2000). These samplers included a standard PM_{2.5} FRM, a TEOM with Nafion drier, a particulate composition monitor (PCM; Atmospheric Research and Analysis, Cary, NC), a medium-volume (113 L/min flow rate) fine particle (PM_{2.5}) and SVOC sampler (i.e., a filter followed by a solid adsorbent) operated by the Desert Research Institute,

**TABLE 2-5. PM_{2.5} SAMPLERS CURRENTLY DESIGNATED AS FRMs FOR
PM_{2.5} MASS CONCENTRATIONS**

Sampler	Manufacturer	Description	Federal Register Reference
RAAS2.5-100	Andersen Instruments	FRM single	Vol. 63, p. 31991, 6/11/98
RAAS2.5-300	Andersen Instruments	FRM sequential	Vol. 63, p. 31991, 6/11/98
RAAS2.5-200	Andersen Instruments	FRM audit	Vol. 64, p. 12167, 3/11/99
Partisol 2000	Rupprecht & Patashnick	FRM single	Vol. 63, p. 18911, 4/16/98
Partisol-Plus 2025	Rupprecht & Patashnick	FRM sequential	Vol. 63, p. 18911, 4/16/98
Partisol 2000 audit	Rupprecht & Patashnick	FRM audit	Vol. 64, p. 19153, 4/19/99
PQ 200	BGI, Inc.	FRM single	Vol. 63, p. 18911, 4/16/98
PQ 200A	BGI, Inc.	FRM audit	Vol. 63, p. 18911, 4/16/98
605 CAPS	ThermoEnvironmental Instruments	FRM single	Vol. 63, p. 58036, 10/29/98
MASS 100	URC	FRM single	Vol. 65, p. 26603, 05/08/00
MASS 300	URC	FRM sequential	Vol. 65, p. 26603, 05/08/00

Source: Peters et al. (2001b); U.S. Environmental Protection Agency (2001).

a HEADS sampler, and a dichotomous sampler for coarse PM. The PCM sampler has three channels, all of which have PM₁₀ cyclone inlets. The first two channels have two denuders preceding a 2.5- μ m WINS impactor and filter packs. The first denuder is coated with Na₂CO₃ to remove acid gases, and the second is coated with citric acid to remove ammonia. The third channel has a carbon-coated, parallel-plate denuder preceding the WINS impactor. Measurements of 24-h mass from the FRM, PCM, and TEOM samplers, as well as reconstructed PM_{2.5} mass (RPM), were compared for a 12-month period. The slopes for the TEOM-FRM, PCM-FRM, and RPM-FRM correlations were 1.01, 0.94, and 0.91, respectively; whereas the y-intercepts for each were 0.68, 0.04, and 0.98. Particulate sulfate measurements on the FRM Teflon filter, the PCM Teflon filter, and PCM Nylon filter were nearly identical. Nitrate results from the three filters were much less consistent, with the FRM collecting substantially less nitrate than was collected on either the denuded nylon filter or denuder followed by a Teflon-nylon filter sandwich. Particulate ammonia measurements were also

compared and showed more scatter than the sulfate measurements but less than the nitrate measurements.

An intercomparison of both PM_{10} and $PM_{2.5}$ mass measurements was conducted during the 1998 Baltimore PM Study (Williams et al., 2000). PM monitors were collocated at a residential indoor, residential outdoor, and ambient monitoring site within Baltimore County, MD. PM samplers included TEOMs, $PM_{2.5}$ FRMs, cyclone-based inlets manufactured by University Research Glassware (URG), and Versatile Air Pollution Samplers (VAPSs). The VAPS is a dichotomous sampler operating at 33 L/min (one coarse particle channel at 3 L/min, and two fine particle channels at 15 L/min, each). In the configuration employed during this study, one fine particle channel was operated with a Teflon filter backed by a nylon filter and preceded by a Na_2CO_3 -coated annular denuder; the second fine particle channel had a quartz filter preceded by a citric acid-coated annular denuder; and the coarse particle channel had a polycarbonate filter followed by a Zefluor filter for flow distribution. Differences in $PM_{2.5}$ mass concentrations between the samplers, although not large, were attributed to potential particle nitrate losses, denuder losses, and losses of SVOCs for some samplers. Differences between coarse PM concentrations, on the other hand, varied widely between the instruments.

In another intercomparison study, Tolocka et al. (2001b) examined the magnitude of potential sampling artifacts associated with the use of the FRM by collocating FRMs alongside other chemical speciation samplers at four U.S. cities. The locations included a high-nitrate, high-carbon, low-sulfate site (Rubidoux, CA); a high-crustal, moderate-carbon, moderate-nitrate site (Phoenix); a high-sulfate, moderate-carbon, low-nitrate site (Philadelphia); and a low- $PM_{2.5}$ mass site (Research Triangle Park, NC). The use of Teflon versus heat-treated quartz filters was also examined in this study. The Teflon filters collected less nitrate than the heat-treated quartz filters. Filters in samplers using denuders to remove organic gases collected less organic PM than filters in samplers without denuders.

Peters et al. (2001b) compiled the results of several field studies in which a number of FRM and other $PM_{2.5}$ samplers were intercompared. In addition to the FRM samplers listed in Table 2-5, other $PM_{2.5}$ samplers included in the evaluation were the Sierra Instruments dichotomous sampler, the Harvard Impactor, the IMPROVE sampler, and the Air Metrics saturation monitor. Results were compiled from $PM_{2.5}$ field studies conducted in Birmingham, Denver, Bakersfield, Phoenix, Research Triangle Park, Atlanta, and Rubidoux. Limited studies

on precision for the non-FRM samplers suggest that the Harvard Impactor and dichotomous samplers had the lowest coefficients of variation (CVs), with both under 10%. The CV for this study was calculated by dividing the sample standard deviation by the average concentration. The IMPROVE samplers had CVs between 10 and 12%, and the Air Metrics samplers had the highest observed CVs, over 15%. In intercomparisons with FRM samplers, the non-FRM samplers showed strong linear relationships with the FRM sampler; however, none of the comparisons passed the current EPA Subpart C equivalent method criteria, which EPA is in the process of revising.

Detailed information on precision of PM samplers used in monitoring networks may be found at EPA's Technology Transfer Network website (U.S. Environmental Protection Agency, 2002a).

2.2.4.1.3 PM_{10-2.5}

Measurement techniques for PM_{10-2.5} are somewhat more complex than those for PM_{2.5} or PM₁₀, because it is necessary to isolate a size fraction between an upper 50% cut point of 10 μm D_a and a lower 50% cut point of 2.5 μm D_a for PM_{10-2.5}. EPA is currently developing an FRM for PM_{10-2.5}. Several candidate techniques are discussed below.

The Difference Method. One approach to measuring PM_{10-2.5} is to make separate measurements of PM₁₀ and PM_{2.5} and take the difference of the resulting equilibrated masses. One problem is that, if either the PM_{2.5} or the PM₁₀ sampler fails, no PM_{10-2.5} measurement can be obtained. In addition, errors in cut point, flow rate, and filter weights (both before use and after collection and equilibration of particles) and uncertainties due to the loss of semivolatile components of PM may occur for each size cut. Careful control of flow rate and equivalent treatment of PM₁₀ and PM_{2.5} filters in terms of pressure drop across the filter and temperature of the filter during and after collection can improve precision and accuracy. Allen et al. (1999a) summarized several sampling issues that should be considered in measuring coarse particulate mass by difference, including the use of identical instrumentation (except cut points), filter media, filter face velocity, and ambient-filter temperature differences; common flow measurement devices; use of higher sampler flow rates (a 10 L/min minimum for 24-h sample is recommended); and avoiding excessive filter loading. The concern, expressed by Allen et al.

(1999a), that the “pie-plate” inlet required by the final version of the PM_{2.5} FRM might have a different cut point than the “flat-top” inlet of the PM₁₀ FRM, has been addressed by a wind tunnel study, which found each to have an appropriate PM₁₀ cut point (Tolocka et al., 2001a).

Since the difference method requires weighing two filters, the key to obtaining high precision in the coarse mass measurement is precise measurements of filter weights. Allen et al. (1999a) discussed techniques for increasing the precision of the difference method by reducing errors in filter weights. These include proper temperature and humidity controls, use of a high quality microbalance, 100% replicate weighings, control of static charge, aging of new filters, weighing of a sufficient number of laboratory blank filters, and accounting for buoyancy errors caused by variability in barometric pressure. Allen et al. (1999a) emphasized the necessity of replicate weighing of filters and of using a third weighing if the difference between the first two weights exceeds a specified minimum. Lawless et al. (2001) investigated the magnitude of uncertainties attributed to fluctuations in some of these parameters (humidity, temperature, drafts, vibration, and electrostatic charges) and recommended methods for improving their control. Koistinen et al. (1999) and Hänninen et al. (2002) gave an excellent discussion of the procedures developed to overcome problems associated with gravimetric measurements of PM_{2.5} mass in the EXPOLIS (Air Pollution Exposure Distributions Within Adult Urban Populations in Europe) Study. They discussed factors such as corrections for buoyancy, elimination of static charge, and increases in the mass of blank filters with time. The establishment of a temperature and humidity controlled room required for the equilibration and weighing of filters for the FRM is expensive. Allen et al. (2001) described a more cost-effective technique that uses a chamber with relative humidity controlled at 34% relative humidity by a saturated aqueous solution of MgCl₂.

Allen et al. (1999a) recommended that, in reporting precision from collocated samplers both the CV and the square of the correlation coefficient (r^2) be reported. For a study in Boston with 27 pairs of mass data from collocated PM₁₀ and PM_{2.5} using standard weighing methods, they obtained a CV of 4.7% with an r^2 of 0.991 for PM_{2.5}, a CV of 4.4% with an r^2 of 0.994 for PM₁₀, and a CV of 15% with an r^2 of 0.88 for PM_{10-2.5}. By using duplicate weighings and other techniques suggested for improving precision, they obtained a CV of 1.3% with an r^2 of 0.998 for PM_{2.5} in a study in Chicago with 38 collocated measurements. On the basis of the improvement in the CV for PM_{2.5}, they estimated that use of the recommended techniques

for $PM_{10-2.5}$ by difference would have yielded a CV of 3.8% with an r^2 of 0.98 if they had been applied in the Chicago study.

Multistage Impaction. A second technique involves the use of impaction to isolate the size fraction between 2.5 and 10 $\mu m D_a$. In the impaction process, the air stream is first accelerated through a small hole (nozzle) or slit. The air stream is directed so that it “impacts” on a surface. Depending on the velocity and pressure of the air stream, particles smaller than a certain size will follow the air stream around the impactor surface. Larger particles will impact on the surface. In practice, impactors have 50% cut points similar to the 50% cut point for the rejection of larger particles in $PM_{2.5}$ and PM_{10} samples (see Figure 2-7).

Multistage impactors are used to separate particles into several size fractions for the determination of mass and chemical composition as a function of size (Wang and John, 1988; Marple et al., 1991). The major problem with the use of impactors to separate the $PM_{10-2.5}$ $\mu m D_a$ fraction of coarse particles (thoracic coarse PM) is bounce. Coarse particles tend to be dry, solid particles. When they hit a hard surface, they can bounce and be carried away with the air stream (e.g., Dzubay et al., 1976; Wesolowski et al., 1977; Rao and Whitby, 1978; Cheng and Yeh, 1979; Wang and John, 1987; John and Sethi, 1993). Various techniques have been used to reduce bounce. One technique is to use a porous substance such as a glass- or quartz-fiber filter (Chang et al., 1999) material or a polyurethane foam (Breum, 2000; Kavouras and Koutrakis, 2001). However, this technique may result in less precise separation and yield a sample that must be extracted before chemical analyses can be performed. Another technique is to coat the impactor with a soft wax or grease (Rao and Whitby, 1977; Turner and Hering, 1987; Pak et al., 1992). This can cause problems with weighing and chemical analyses, and as the deposit of particles builds up, incoming particles may not hit the soft surface, but instead hit a previously collected hard particle and bounce off of it. The WINS impactor discussed earlier uses a filter in a well of low volatility oil to ensure a wetted surface at all times. However, such a technique, while appropriate for removing unwanted particles, would not yield a particle sample suitable for weighing or for chemical analyses.

Virtual Impaction. The problems of bounce and blow off of particles from impactors, especially for the collection of large quantities of particles, was addressed by aerosol scientists in the mid-1960s by the development of what is now known as “virtual” impaction (Hounam and Sherwood, 1965; Conner, 1966).

In a virtual impactor, a hole is placed in the impaction plate just below the accelerating jet. Two controlled flows allow a fraction, e.g., 10% (or another predetermined fraction, typically 5 to 20%), of the air containing the coarse particles to go through the hole and through a filter (minor flow). A 10% minor flow gives a coarse channel enrichment factor of 10. The remaining fraction (e.g., 90% of the airflow) containing the fine particles follows a different path and goes through a second filter (major flow). The upper cut point is usually set by the inlet (e.g., $10 \mu\text{m } D_a$). The flow rates, pressures, and distance from the nozzle to the virtual impactor surface can be varied to direct particles with an D_a greater than the lower cut point (i.e., $> 2.5 \mu\text{m}$) to go through the hole and be collected on the first filter and to direct smaller particles (i.e., $< 2.5 \mu\text{m}$) to flow around the impactor and be collected on the second filter. Large particles “impact” into the hole with a small amount of the air flow. The smaller particles follow the major air flow around the impactor plate. This technique overcomes the problem of bounce. An example of the separation into fine and coarse particles is shown in Figure 2-26.

The usefulness of this technique for collecting samples of fine and coarse particles for chemical analysis was recognized by the EPA in the mid-1970s and led to the development of the now well-known “dichotomous sampler” (a virtual impactor that separates particles into two size fractions) and an associated XRF analyzer (Dzubay and Stevens, 1975; Loo et al., 1976; Jaklevic et al., 1977; Dzubay et al., 1977). The dichotomous sampler was originally developed for use in the Regional Air Monitoring Study (RAMS), part of the Regional Air Pollution Study (RAPS), conducted in St. Louis, MO in the mid-1970s. Dichotomous samplers were operated at 10 RAMS sites from March 1975 to March 1977; and 33,695 filters were collected and analyzed by XRF with an overall sampling efficiency of 97.25% (Strothmann and Schiermeier, 1979; Loo et al., 1976; Loo et al., 1978; Dzubay, 1980; Lewis and Macias, 1980). Dichotomous samplers were a novel concept at that time. Concern over particle losses and other problems at cut point sizes $< 2.5 \mu\text{m } D_a$ influenced the decision to choose 2.5 instead of 1.0 as the cut point diameter.

Since the use of the dichotomous sampler in RAPS, considerable progress has been made in the theory and practice of designing virtual impactors, especially in how to reduce losses and




Figure 2-26. Schematic diagram showing the principle of virtual impaction. The initial flow, Q_0 , is split into a minor flow, Q_1 , which carries the larger particles that impact into the hole, to the coarse particle filter and a major flow, Q_2 , which carries the smaller particles that can follow the airflow, to the fine particle filter.

Source: Loo et al. (1976).

provide a sharp cut (Masuda et al., 1979; Marple and Chien, 1980; Chen et al., 1985, 1986; Loo and Cork, 1988). Now virtual impactors, with rectangular slits or round holes, are used to (a) provide cut point sizes as low as $0.15 \mu\text{m } D_a$ and (b) concentrate coarse, accumulation, and ultrafine mode particles for use in health studies (Solomon et al., 1983; Marple et al., 1990; Sioutas et al., 1994a,b,c). Dichotomous samplers were also used in a national network to measure $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ in the Harvard Six City Study (Spengler and Thurston, 1983; Dockery et al., 1993) and the Inhalable Particulate Network (Suggs and Burton, 1983). A trichotomous high volume sampler has also been developed that provides samples of particles $< 1.0 \mu\text{m}$, $1.0 \mu\text{m}$ to $2.5 \mu\text{m}$, and $> 2.5 \mu\text{m}$ (Marple and Olsen, 1995). This sampler was intended to aid in the study of the composition of the intermodal mass in the range of 1.0 to $2.5 \mu\text{m } D_a$.

More recently, two dichotomous sequential PM air samplers were collocated with a manual FRM air sampler and operated for over a year at a waterfront site on Tampa Bay (Poor et al., 2002). The FRM sampler was alternately configured as a PM_{2.5}, then as a PM₁₀ sampler. For the dichotomous sampler measurements, daily 24-h integrated PM_{2.5} and PM_{10-2.5} ambient air samples were collected at a total flow rate of 16.7 L/min. As was the case for earlier versions of the dichotomous sampler, a virtual impactor split the air into flow rates of 1.67 and 15.01 L/min and collected PM_{10-2.5} and PM_{2.5} on 47-mm diameter Teflon filters. Between the two dichotomous air samplers, the average concentration, relative bias, and relative precision for PM_{2.5} were 13.3 µg/m³, 0.02% and 5.2% (n = 282), and for PM_{10-2.5} were 12.3 µg/m³, 3.9% and 7.7% (n = 282). Federal Reference Method measurements were alternate-day 24-h integrated PM_{2.5} or PM₁₀ ambient air samples collected onto 47-mm diameter PTFE filters at a flow rate of 16.71 L/min. Between a dichotomous and a PM_{2.5} FRM air sampler, the average concentration, relative bias, and relative precision were 12.4 µg/m³, -5.6%, and 8.2% (n = 43). Between a dichotomous and a PM₁₀ FRM air sampler, the average concentration, relative bias and relative precision were 25.7 µg/m³, -4.0%, and 5.8% (n = 102). The PM_{2.5} concentration measurement standard errors for the two dichotomous and one FRM samplers were 0.95, 0.79, and 1.02 µg/m³; and for the PM₁₀ sampler, the standard errors were 1.06, 1.59, and 1.70 µg/m³. The authors (Poor et al., 2002) concluded that their results indicated that “the dichotomous samplers have superior technical merit” and demonstrate “the potential for the dichotomous sequential air sampler to replace the combination of the PM_{2.5} and PM₁₀ FRM air samplers, offering the capability of making simultaneous, self-consistent determinations of these particulate matter fractions in a routine ambient monitoring mode.”

The dichotomous sampler provides two separate samples. However, a fraction of the smaller particles, equal to the minor flow, will go through the virtual impaction opening with the air stream and be collected on the coarse particle filter. In the dichotomous sampler used in the RAPS program, 10% of the fine particles were collected with the coarse particles. Thus, in order to determine the mass or composition of the coarse particles, it is necessary to determine the mass and composition of the fine particles and subtract the appropriate fraction from the mass or composition of the particles collected on the coarse particle filter. Allen et al. (1999b) discussed potential errors in the dichotomous sampler caused by uncertainties in the coarse mass channel enrichment factor. Virtual impactors have also been designed with a clean air jet in the center of

the round nozzle. This reduces the contamination of coarse particles by fine particles, while maintaining low losses and sharp cuts (Chen and Yeh, 1987; Chein and Lundgren, 1993). The fine particle intrusion into the coarse particle sample can also be reduced by operating two virtual impactors in series (Dzubay and Stevens, 1975).

Aerosol physicists have also conducted theoretical and experimental investigations of virtual impaction using slits instead of round holes (Forney et al., 1978, 1982; Ravenhall et al., 1978; Masuda and Nakasita, 1988; Sioutas et al., 1994b,c,d; Ding and Koutrakis, 2000). The slit virtual impactor permits a much higher flow rate than round hole virtual impactors and resolves problems that occur with multihole virtual impactors (Marple et al., 1990; Fang et al., 1991). The slit technique has been used to develop virtual impaction systems for concentrating particles in the size range 0.1 to 2.5 μm D_a for exposure studies using animals and humans (Sioutas et al., 1995a, b). The slit impactor can also be used to concentrate coarse particles for measurement (Misra et al., 2001) or exposure studies (Chang et al., 2002). In addition, ultrafine particles ($> 0.1 \mu\text{m}$) can be concentrated by first separating ultrafine particles from larger particles, adding water vapor to saturate the air containing the ultrafine particles, cooling the air to cause supersaturation and growth of the ultrafine particles into the 1.0 to 4.0 μm size range, then concentrating these particles with a slit virtual impactor, and finally, heating the air to return the particles to their original size (Sioutas and Koutrakis, 1996; Sioutas et al., 1999; Sioutas et al., 2000; Kim et al., 2001b,c; Geller et al., 2002).

2.2.5 Speciation Monitoring

Speciation Network and Monitoring

In addition to FRM sampling to determine compliance with PM standards, the EPA requires states to conduct chemical speciation sampling primarily to determine source categories and trends (Code of Federal Regulations, 2001b). Source category apportionment calculations are discussed in Chapter 3. A $\text{PM}_{2.5}$ chemical Speciation Trends Network (STN) has been deployed that consists of 54 core National Ambient Monitoring Stations (NAMS) and approximately 250 State and Local Air Monitoring Stations (SLAMS). In addition, over 100 IMPROVE (Interagency Monitoring of Protected Visual Environments) samplers located at regional background and transport sites can be used to fulfill SLAMS requirements. The overall goal of the speciation program is “to provide ambient data that support the Nation’s air quality

program objectives” (U.S. Environmental Protection Agency, 1999). Information and reports on the EPA’s speciation monitoring program may be found on the EPA’s Technology Transfer Network at <http://www.epa.gov/ttn/amtic/pmspec.html>. The NAMS speciation sites will provide routine chemical speciation data to be used in developing annual and seasonal aerosol characterization, air quality trends analysis, and emission control strategies. The SLAMS speciation sites will further support the NAMS network and provide information for development of state implementation plans (SIPs).

At both NAMS and SLAMS sites, aerosol samples will be collected for the analysis of trace elements, ions (sulfate, nitrate, ammonium, sodium, and potassium), and total carbon (TC). The NAMS speciation sites will operate on a 1-in-3-day schedule, with 10 of these sites augmented with continuous speciation analyses for everyday operation. The SLAMS speciation sites will generally operate on a 1-in-6-day basis; however, many sites may be operated on a 1-in-3-day basis in locations where increased data collection is needed. There are several samplers that are suitable for use in the NAMS/SLAMS network. These samplers include an inlet cut point with size cut characteristics comparable to the WINS FRM, proven denuder technology for nitrate, and sampler face velocity and sample volume similar to that of the FRM. The current samplers include three filters: (1) Teflon for equilibrated mass and elemental analysis by energy dispersive X-ray fluorescence (EDXRF), (2) a nitric acid denuded nylon filter for ion analysis (ion chromatography), (3) a quartz-fiber filter for elemental and organic carbon. Elemental carbon and OC are determined by thermal-optical analysis via a modification of the NIOSH (National Institute for Occupational Safety and Health) method 5040 (thermal-optical transmission [TOT]). However, no corrections are made for positive artifacts caused by adsorption on organic gases or the quartz filters or negative artifacts due to the evaporation of SVOCs from collected particles.

Since 1987, the IMPROVE network has provided measurements of ambient PM and associated light extinction in order to quantify PM chemical components that affect visibility at Federal Class 1 areas that include designated national parks, national monuments, and wilderness areas. Management of this network is a cooperative effort between the EPA, federal land management agencies, and state governments. The IMPROVE program has established analytical protocols for measurements of ambient concentrations of PM_{10} , $PM_{2.5}$,

sulfates, nitrates, organic and elemental carbon, crustal material, and a number of other elements. Information on the IMPROVE program may be found at <http://vista.cira.colostate.edu/improve>.

IMPROVE aerosol monitoring consists of a combination of particle sampling and sample analysis. The IMPROVE sampler, which collects two 24-h samples per week, simultaneously collects one sample of PM₁₀ on a Teflon filter, and three samples of PM_{2.5} on Teflon, nylon, and quartz filters. PM₁₀ mass concentrations are determined gravimetrically from the PM₁₀ filter sample, while PM_{2.5} mass concentrations are determined gravimetrically from the PM_{2.5} Teflon filter sample. The PM_{2.5} Teflon filter sample is also used to determine concentrations of selected elements using PIXE, XRF, and PESA. The PM_{2.5} nylon filter sample, which is preceded by a denuder to remove acidic gases, is analyzed to determine nitrate and sulfate aerosol concentrations using IC. Finally, the PM_{2.5} quartz filter sample is analyzed for OC and EC using the thermal-optical reflectance (TOR) method. Corrections are made for positive artifacts but not for negative artifacts.

The STN and the IMPROVE networks represent a major advance in the measurement of nitrate, because the combination of a denuder to remove nitric acid vapor and a Nylon filter to adsorb nitric acid vapor volatilizing from the collected ammonium nitrate particles overcomes the loss of nitrate from Teflon filters. However, different techniques used for measurement of OC and EC lead to significant differences between their measurements when the two techniques are intercompared (Chow et al., 2001). IMPROVE yields higher EC and lower OC, although there is good agreement for TC. Another difference arises from the treatment of the positive artifacts due to the absorption of organic gases by the quartz filters used in IMPROVE and STN samplers. More information on these differences is given in Section 2.2.7 and details are discussed in Appendix 2B.

Several of the PM_{2.5} size selectors developed for use in the EPA National PM_{2.5} STN were recently evaluated by comparing their penetration curves under clean room experiments with that of the WINS impactor (Peters et al., 2001c). The corresponding speciation monitors were then compared to the FRM in four cities. The PM_{2.5} inlets tested were the SCC 2.141 cyclone (6.7 L/min) that is in the Met One Instruments SASS sampler, the SCC 1.829 cyclone (5.0 L/min) that is proposed for use in the Rupprecht and Patashnik real-time sulfate/nitrate monitor, the AN 3.68 cyclone (24.0 L/min) that is in the Andersen RAAS, and the spiral separator (7.0 L/min) that was previously in the Met One SASS. The cutpoints of the SCC

cyclones compared reasonably well with the WINS (2.52 and 2.44 μm for the SCC 2.141 and SCC 1.829, respectively, at their design flow rates), but both demonstrated a tail extending into the coarse-particle mode. The AN inlet had the sharpest cut point curve, but the 50% cut point diameter was 2.7 μm D_a at its design flow rate. The spiral inlet had the shallowest cut point curve: the 50% cut point was 2.69 and 2.67 μm D_a for an ungreased and greased inlet, respectively. The speciation samplers were also compared to the FRM sampler with WINS inlet under ambient conditions in four cities. The Andersen RAAS equipped with the AN 3.68 cyclone compared well to the FRM in all four cities when compared on the basis of $\text{PM}_{2.5}$ mass, sulfate, and crustal concentrations. Greasing the spiral inlet in the Met One sampler improved the performance of that sampler, which tended to give much higher $\text{PM}_{2.5}$ concentrations than the FRM in cities with high crustal PM.

2.2.6 Inorganic Elemental Analyses

In addition to the lighter elements (hydrogen, carbon, oxygen and nitrogen), the following 40 heavier elements are commonly found in ambient air samples: sodium, magnesium, aluminum, silicon, phosphorus, sulfur, chlorine, potassium, calcium, titanium, vanadium, chromium, manganese, iron, cobalt, nickel, copper, zinc, gallium, arsenic, selenium, bromine, rubidium, strontium, yttrium, zirconium, molybdenum, palladium, silver, cadmium, indium, tin, antimony, barium, lanthanum, gold, mercury, thallium, lead, and uranium. These heavier elements are often indicators of air pollution sources, and several of them are considered to be toxic (such as transition metals, water-soluble metals, and metals in certain valence states [e.g., Fe(II), Fe(III), Cr(III), Cr(VI), As(III), As(V)]). Various measurement methods for inorganic elements are listed in Table 2-6. These methods differ with respect to detection limits, sample preparation, and cost (Chow, 1995). EDXRF and PIXE are the most commonly applied methods because they quantify more than 40 detectable elements, are nondestructive, and are relatively inexpensive. Both were discussed in the previous 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a). TRXRF and S-XRF are newer techniques with lower detection limits. AAS, ICP-AES, and ICP-MS are also appropriate for ionic measurements of elements that can be dissolved. PESA provides a means for measuring elements with lower atomic numbers (i.e., the elements from hydrogen to carbon). More detailed information on each technique is given in Appendix 2B.1.

TABLE 2-6. MEASUREMENT METHODS FOR INORGANIC ELEMENTS

#	Acronym	Full Name	Comments
1	EDXRF	Energy Dispersive X-ray Fluorescence	Heavier elements
2	S-XRF	Synchrotron Induced X-ray Fluorescence	Heavier elements; lower detection limits than EDXRF
3	PIXE	Proton Induced X-ray Emission	Heavier elements; lower detection limits than EDXRF
4	PESA	Proton (or particle) Elastic Scattering Analysis	Lighter elements
5	TRXRF	Total Reflection X-ray Fluorescence	Heavier elements; lower detection limits than EDXRF
6	INAA	Instrumental Neutron Activation Analysis	Many elements; sensitivity different than EDXRF
7	AAS	Atomic Absorption Spectrophotometry	Many elements that can be dissolved
8	ICP-AES	Inductively Coupled Plasma with Atomic Emission Spectroscopy	Many elements that can be dissolved
9	ICP-MS	Inductively Coupled Plasma with Mass Spectroscopy	Many elements that can be dissolved
10	SEM	Scanning Electron Microscopy	Heavier elements

2.2.7 Elemental and Organic Carbon in Particulate Matter

Ambient particles from combustion sources contain carbon in several chemically and optically distinct forms. Health- and visibility-related studies of these particles require information about the relative contributions to total particle mass by these different forms of carbon. With the exception of carbonate-based carbon, however, a clear classification scheme has not yet been established to distinguish OC, light-absorbing carbon, black carbon, soot and EC. The absence of clear, physically-based definitions results in confusion in the interpretation of speciation data. For example, depending on the radiation wavelength specified, “light-absorbing” carbon can include compounds that volatilize without oxidation. “Black” carbon includes various mixtures containing “elemental” (graphitic) carbon; partially degraded, oxidized graphitic fragments; and partially oxidized amorphous aromatic carbon. For studying

visibility reduction, a measurement of light-absorbing carbon may be more useful than one of EC. For source apportionment by receptor models, several consistent, but distinct, fractions of carbon in both source and receptor samples are desired, regardless of their light-absorbing or chemical properties. Differences in the ratios of the carbon concentrations in these fractions form part of the source profiles that distinguish the contribution of one source from another (Watson et al., 1994a,b).

Three method-dependent operational classes of carbon are commonly measured in ambient aerosol samples collected on quartz fiber filters: (1) semivolatile organic or non-visible light-absorbing carbon, termed “organic carbon (OC)”; (2) elemental carbon, soot, black carbon, or light-absorbing carbon, termed “elemental carbon (EC)”; and (3) carbon present as K_2CO_3 , Na_2CO_3 , $MgCO_3$, $CaCO_3$, termed “carbonate carbon (CC).” The sum of OC, EC, and CC in PM gives the total carbon (TC).

The thermal-optical reflectance (TOR), thermal-optical transmission (TOT), and thermal-manganese oxidation (TMO) methods are most commonly used for the analysis of OC and EC in atmospheric PM. In thermal separation methods, OC is vaporized and the EC remaining on the filter is then oxidized to CO_2 and quantified by nondispersive infrared detection, by electrochemical techniques or by reducing the CO_2 to CH_4 and the quantifying CH_4 via flame ionization detection (FID). Organic carbon that does not vaporize below $550\text{ }^\circ\text{C}$ can pyrolyze at higher temperatures to form additional black carbon. Thermal optical methods must correct for this effect in order to correctly distinguish OC from EC. The various methods give similar results for TC, but not for EC or OC, due to differing assumptions and analytical procedures regarding the thermal behavior of ambient aerosol carbon. These methods are discussed in detail in Appendix 2B.2.

Carbonate carbon can be determined thermally, or on a separate filter section by measuring the carbon dioxide (CO_2) evolved after acidification (Johnson et al., 1980). It is usually on the order of 5% or less of TC for ambient particulate samples collected in urban areas (Appel, 1993).

The forms of carbon present in natural materials that may lead to the formation of atmospheric aerosol tend to be poorly defined. Thus, the pyrolysis products of these materials during thermal-optical analysis cannot be predicted. The Geochemical Society convened an international steering committee in 1999 to define a set of representative black carbon and black carbon-containing benchmark materials to be used to support ambient aerosol sample analysis.

Such materials can be used to provide thermal-optical “fingerprints” for deducing primary aerosol sources, and to establish characteristic analytic interferences or artifacts associated with such sources. The materials recommended to date include n-hexane soot, lignocellulosic chars, soils, marine sediments, and the NIST urban dust standard reference material (SRM 1649a). The committee has also recommended a set of standard materials that may potentially interfere with black carbon analyses: shale, natural organic matter, coals, and melanoidin (an amino acid-based material). These recommendations are discussed on the steering committee’s website: <http://www.du.edu/~dwismith/bcsteer.html>.

2.2.8 Ionic Species

Ion chromatography is widely used for analyzing ionic species in the water-soluble portion of suspended PM. Ion chromatography is the method of choice for the measurement of sulfate, nitrate, ammonium, sodium, and potassium ions for the NAMS program. Aerosol strong acidity, H^+ , is determined by titrating an aqueous solution of PM that is collected after passage through a series of annular denuders to remove acid and basic gases with back-up filters to collect the NH_3 and HNO_3 that volatilize from the PM during collection. The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) discussed measurement of ions by ion chromatography (Section 4.3.3.1) and of strong acidity (Sections 3.3.1.1 and 4.3.3.1); so, no further details will be discussed here.

2.2.9 Continuous Monitoring

The EPA expects that many local environmental agencies will operate continuous PM monitors. All currently available continuous measurements of suspended particle mass share the problem of dealing with semivolatile PM components. So as not to include particle-bound water as part of the mass, the particle-bound water must be removed by heating or dehumidification. However, heating also causes the loss of ammonium nitrate and SVOCs. A variety of potential candidates for the continuous measurement of particle mass and related properties are listed in Table 2-7. These techniques are discussed in more detail in Appendix 2B.3.

**TABLE 2-7. METHODS FOR CONTINUOUS MEASUREMENT OF PM MASS,
PM COMPONENTS, ETC.**

#	Acronym	Name	Comments
1	TEOM	Tapered Element Oscillating Microbalance	Measures only nonvolatile components of PM.
2	TEOM with SES	TEOM with Sample Equilibration System	By using dehumidification instead of heat for drying and a lower temperature for stabilization, includes some of the semivolatile components of PM.
3	—	Differential TEOM	Research instrument designed to measure both the nonvolatile and semivolatile components of PM.
4	FDMS	Filter Dynamics Measurement System	Commercial version of the differential TEOM.
5	RAMS	Real-time total Ambient Mass Sampler	Research instrument designed to measure total PM using denuders and an adsorptive filter.
6	CAMM	Continuous Ambient Mass Monitor	Measures total PM by pressure drop across a frequently changed filter.
7	BAM	Beta gauge	Measures PM mass by beta attenuation. Unless dried by heat or dehumidification will also measure particle-bound water.
8	—	Piezoelectric microbalance	Measures mass by change in resonance frequency when particles are deposited on a crystal.
9	CCPM	Continuous Coarse Particle Monitor	Virtual impaction is used to concentrate PM _{10-2.5} which is then measured by a TEOM.
10	—	Semi-continuous EC/OC	Several commercially available instruments automate the thermal technique for EC/OC and provide hourly measurements.
11	—	Semi-continuous nitrate	Collection of PM followed by flash vaporization and determination of NO _x provides 10 minute measurements of particulate nitrate.
12	—	Semi-continuous sulfate	Several techniques are available in which particulate sulfate is converted to SO ₂ which is measured by a pulsed fluorescence analyzer.
13	—	Continuous ion chromatography of water-soluble ions	Particles are grown by mixing with water vapor, collected in water, and injected into an automatic ion chromatography.
14	—	Mass spectroscopy of individual particles	Single particles are evaporated, ionized and the components analyzed by mass spectroscopy. Several different systems are in use in research studies.
15	EAD	Electrical Aerosol Detector	This instrument measures charge collected by particles and gives a continuous signal that is proportional to the integral of the particle diameter.
16	—	Integrating nephelometer	Light scattering by suspended particles, collected over a large solid angle, provides an indicator for particle mass including particle-bound water unless the air sample is dried by heating or dehumidification.

2.2.10 Measurements of Individual Particles

Recently, several researchers have developed instruments for real-time in situ analysis of single particles (e.g., Noble and Prather, 1996; Gard et al., 1997; Johnson and Wexler, 1995; Silva and Prather, 1997; Thomson and Murphy, 1994). Although the technique varies from one laboratory to another, the underlying principle is to fragment each particle into ions, using either a high-power laser or a heated surface and, then, a time-of-flight mass spectrometer (TOFMS) to measure the ion fragments in a vacuum. Each particle is analyzed in a suspended state in the air stream (i.e., without collection), avoiding sampling artifacts associated with impactors and filters. The technique is called aerosol time-of-flight mass spectrometry (ATOFMS). By measuring both positive and negative ions from the same particle, information can be obtained about the composition of individual particles of known aerodynamic diameter. This information is especially useful in determining the sources of particles. Noble and Prather (1996) used ATOFMS to provide compositionally resolved particle-size distributions. Their instrument is capable of analyzing size and chemical composition of 50 to 100 particles/min at typical ambient concentrations and up to 600/min at high particle concentrations. Four systems for measurement of single particles using mass spectroscopy are reviewed by Middlebrook et al. (2003). An example of the type of information that can be determined is shown in Figure 2-27.

Because particles are analyzed individually, biases in particle sampling (including biases in the efficiency of particle transmission into the sensor chamber as a function of size; particle size measurement, and detection of particles prior to fragmentation) represent a major challenge for these instruments. Moreover, laser ablation has a relatively large variability in ion yields, i.e., identical samples would yield relatively large differences in mass spectrometer signals (Thomson and Murphy, 1994). Therefore, quantitation is inherently difficult (Murphy and Thomson, 1997). Quantitation will be even more challenging for complex organic mixtures because of the following two reasons: (1) a large number of fragments are generated from each molecule, and (2) ion peaks for organics can be influenced or obscured by inorganic ions (Middlebrook et al., 1998). Nonetheless, scientists have been successful in using these techniques to identify the presence of organics in atmospheric particles and laboratory-generated particles (i.e., as contaminants in laboratory-generated sulfuric acid droplets) as well as the identification of specific compound classes such as PAHs in combustion emissions (Castaldi and Senkan, 1998; Hinz et al., 1994; Middlebrook et al., 1998; Murphy and Thomson, 1997;




Figure 2-27. Size distribution of particles divided by chemical classification into organic, marine, and soil (or crustal).

Source: Noble and Prather (1998).

Neubauer et al., 1998; Noble and Prather, 1998; Reilly et al., 1998; Silva and Prather, 1997). A new multivariate technique for calibrating of ATOFMS using microorifice impactors shows promise for simplifying the calibration process (Ferguson et al., 2001). This calibration technique has been applied to gasoline and diesel particles to demonstrate the feasibility of using this technique for the source apportionment of gasoline and diesel particles in an atmospheric mixture (Song et al., 2001).

Previously, ATOFMS systems have only been able to characterize particles larger than ~0.2 to 0.3 μm in diameter. Recently, Wexler and colleagues (Carson et al., 1997; Ge et al., 1998) developed an ATOFMS instrument that is able to size, count, and provide chemical composition on individual particles ranging in size from 10 nm to 2 μm .

2.2.11 Low Flow Filter Samples for Multiday Collection of Particulate Matter

For some purposes, such as demonstrating the attainment of an annual standard or as an exposure indicator for epidemiologic studies of chronic health effects, 24-h measurements are not essential. Annual or seasonal averages may be adequate and multiday sampling techniques can result in lower costs for weighing, chemical analysis, and travel time to change filters. The multiday sampler serves a second purpose. Most commercially available samplers are optimized for collecting 24-h samples of PM concentrations as found in the United States, Europe, or Japan, but many other parts of the world have significantly higher PM concentrations. Under such conditions, the 16.7 L/min flow through 37- or 47-mm diameter filters may overload the filter, preventing the sampler from maintaining the prescribed flow rate for 24 h. A low-flow sampler with a 0.4 L/min flow rate and a 47-mm diameter filter has been designed by Aerosol Dynamics, Inc. With this sampler, the sample collection time can be chosen to suit the ambient concentration level. This sampler, with a 1-week collection period, has been used to characterize PM_{2.5} in Beijing, PRC (He et al., 2001). It is also being used with a 2-week collection period in a chronic epidemiologic study in southern California (Gauderman, et al., 2000).

The low-flow sampler, as described by He et al. (2001), has three PM_{2.5} channels. One channel collects PM on a Teflon filter for gravimetric mass measurement and elemental analysis by XRF. A second channel collects PM on a quartz filter for OC and EC analysis. A denuder may be added to the second channel to remove organic gases, as may a backup filter to collect SVOCs. The third channel uses a carbonate denuder to remove acid gases such as HNO₃ and SO₂, a Teflon filter to collect PM for analysis of ions by IC, and a nylon filter to collect volatilized nitrate. The Teflon filter can also be weighed prior to extraction; thus, the multiday sampler can provide the information needed for source apportionment by chemical mass balance techniques (Watson et al., 1990a,b; U.S. Environmental Protection Agency, 2002b).

For monitoring sites with high day-to-day variability in PM concentrations, a sample integrated over a week may provide a more accurate measurement of the annual average than can be obtained by 1-in-3- or 1-in-6-day sampling schedules. Daily PM data from Spokane, WA were resampled to simulate common sampling schedules and the error due to less-than-daily sampling was computed (Rumburg et al., 2001). The error in the annual mean concentration for PM_{2.5}, expressed as a percentage difference from the daily sampling mean, was 1.7, 3.4, and

7.7% for 1-in-2-, 1-in-3, and 1-in-6-day sampling, respectively. These results may or may not be representative of other locations and time periods.

2.3 SUMMARY AND KEY POINTS

2.3.1 Chemistry and Physics of Atmospheric Particles

Atmospheric particles originate from a variety of sources and possess a range of morphological, chemical, physical, and thermodynamic properties. The composition and behavior of aerosols are linked with those of the surrounding gas. An aerosol may be defined as a suspension of solid or liquid particles in air and includes both the particles and all vapor or gas phase components of air. However, the term aerosol is also often used to refer to the suspended particles only.

A complete description of atmospheric PM would include an accounting of the chemical composition, morphology, and size of each particle, and the relative abundance of each particle type as a function of particle size. Recent developments in single particle analysis techniques are bringing this description closer to reality.

The diameter of a spherical particle may be determined geometrically, from optical or electron microscopy, by light scattering and Mie theory, or by a particle's behavior (e.g., electrical mobility or its aerodynamic behavior). However, the various types of diameters may be different, and atmospheric particles often are not spherical. Therefore, particle diameters are described by an "equivalent" diameter. Aerodynamic diameter, D_a (the diameter of a unit density sphere that would have the same terminal settling velocity as the real particle), and the Stokes diameter, D_p (the diameter of a sphere of the same density as the particle that would have the same aerodynamic resistance or drag), are the most widely used equivalent diameters.

Atmospheric size distributions show that most atmospheric particles are quite small, $< 0.1 \mu\text{m}$ in diameter; whereas most of the particle volume (and therefore most of the mass) is found in particles $> 0.1 \mu\text{m}$. An important feature of the mass or volume size distributions of atmospheric particles is their multimodal nature. Volume distributions, measured in ambient air in the United States, almost always have a minimum between 1.0 and 3.0 μm . The portion of the size distribution that contains particles that are mostly larger than the minimum is called "coarse" particles or the "coarse" mode. The portion of the size distribution that contains

particles that are mostly smaller than the minimum is called “fine” particles and includes several modes. “Accumulation mode” refers to that portion of fine particles with diameters above about 0.1 μm . The portion of fine particles with diameters $< 0.1 \mu\text{m}$ are usually called “ultrafine” by toxicologists and epidemiologists and “nanoparticles” by aerosol physicists and material scientists. In the number distribution of ultrafine particles, particles in the size range < 0.01 are called the nucleation mode and particles between 0.01 and 0.1 are called the Aitken mode. The Aitken mode can be observed as a separate mode in mass or volume distributions only in clean or remote areas or near sources of new particle formation by nucleation.

The aerosol community uses three different approaches or conventions in the classification of particles by size: (1) modes, based on the observed size distributions and formation mechanisms; (2) dosimetry or occupational sizes, based on the entrance into various compartments of the respiratory system; and (3) cut point, usually based on the 50% cut point of the specific sampling device, including legally specified, regulatory sizes for air quality standards. Over the years, the terms fine and coarse as applied to particle sizes have lost their original precise meanings. In any given article, therefore, the meaning of fine and coarse, unless defined, must be inferred from the author’s usage. In particular, $\text{PM}_{2.5}$ and fine particles are not equivalent. $\text{PM}_{2.5}$ refers to the aggregate sample of PM that is collected following a size-selective inlet with a specified penetration as a function of size and a 50% cutpoint of 2.5 μm D_a . It may also be used to refer to the number (or other measure) of particles suspended in the atmosphere that would be collected by such a sampler). PM_{10} is defined similarly. $\text{PM}_{10-2.5}$ refers to the sample that would be collected if the $\text{PM}_{2.5}$ component could be removed from a PM_{10} sample.

Several processes influence the formation and growth of particles. New particles may be formed by nucleation from gas phase material. Particles may grow by condensation as gas phase material condenses onto existing particles. Particles may also grow by coagulation as two particles combine to form one. Gas phase material condenses preferentially on smaller particles, and the rate constant for coagulation of two particles decreases as the particle size increases. Therefore, nucleation-mode particles grow into Aitken-mode particles and Aitken-mode particles grow into the accumulation mode, but growth of accumulation-mode particles into the coarse mode size range is unusual.

The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental or black carbon; a great variety of organic compounds; and crustal material. Atmospheric PM contains a large number of elements in various compounds and concentrations and hundreds of specific organic compounds. Particulate material can be primary or secondary. It is designated primary if it is in the same chemical form in which it was emitted into the atmosphere. It is called secondary PM if it is formed by chemical reactions in the atmosphere. Primary coarse particles are usually formed by mechanical processes; whereas primary fine particles are emitted from sources either directly as particles or as vapors that rapidly condense to form particles. Water-soluble and potentially toxic gases such as SO₂, H₂O₂, and formaldehyde, which in the absence of particles would be removed by deposition to the mucus on the upper airways of the respiratory tract, may dissolve in particle-bound water and be carried into the air exchange regions of the lungs.

Most of the sulfate and nitrate and a portion of the organic compounds in atmospheric particles are secondary. Secondary aerosol formation depends on numerous factors including the concentrations of precursors; the concentrations of other gaseous reactive species such as ozone, hydroxyl radical, peroxy radicals, and hydrogen peroxide; atmospheric conditions, including solar radiation and relative humidity; and the interactions of precursors and preexisting particles within cloud or fog droplets or on or in the liquid film on solid particles. As a result, it is considerably more difficult to relate ambient concentrations of secondary species to sources of precursor emissions than it is to identify the sources of primary particles.

The lifetimes of particles vary with particle size. Coarse particles can settle rapidly from the atmosphere within minutes or hours and normally travel only short distances. However, when mixed high into the atmosphere, as in dust storms, the smaller-sized, coarse-mode particles may have longer lives and travel greater distances. Accumulation-mode particles are kept in suspension by normal air motions and have a lower deposition velocity than coarse-mode particles; they can be transported thousands of kilometers and remain in the atmosphere for a number of days. They are removed from the atmosphere primarily by cloud processes. Dry deposition rates are expressed in terms of a deposition velocity that varies with the particle size, reaching a minimum between 0.1 and 1.0 μm D_a.

Particulate matter is a factor in acid deposition. Particles serve as cloud condensation nuclei and contribute directly to the acidification of rain. In addition, the gas-phase species that

lead to the dry deposition of acidity are also precursors of particles. Therefore, reductions in SO₂ and NO_x emissions will decrease both acid deposition and PM concentrations, but not necessarily in a linear fashion. Sulfuric acid, ammonium nitrate, and organic particles also are deposited on surfaces by dry deposition and can contribute to ecological effects.

2.3.2 Measurement of Atmospheric Particles

The decision by the EPA to revise the PM standards by adding daily and yearly standards for PM_{2.5} has led to a renewed interest in the measurement of atmospheric particles and to a better understanding of the problems in their precise and accurate measurements. Unfortunately, it is very difficult to measure and characterize particles suspended in the atmosphere.

Particulate matter monitoring is used to develop information to guide the implementation of standards (i.e., to identify particles sources, to determine whether or not a standard has been attained) and to determine health, ecological, and radiative effects. Federal Reference Methods (FRM) specify techniques for measuring PM₁₀ and PM_{2.5}. Particles are collected on filters and mass concentrations are determined gravimetrically. The PM₁₀ FRMs consist of low flow rate (1 L/h) and high flow rate (67.92 L/h) samplers, each with a PM₁₀ inlet/impactor. The high flow rate sampler uses a 20.3 × 35.4 cm² quartz filter whereas the low flow rate sampler uses a 47-mm Teflon filter, each with a collection efficiency of ≥ 99% as described in Appendix M to 40 CFR, Part 50 (Code of Federal Regulations, 2001e). The PM_{2.5} FRM is similar to the low flow rate PM₁₀ FRM except that it includes a PM_{2.5} impactor with an oil-covered impaction substrate to remove particles > 2.5 μm and a filter with a collection efficiency of greater than 99.7% as described in Appendix L, 40 CFR, Part 50 (Code of Federal Regulations, 2001c). Both techniques provide relatively precise (± 10%) methods for determining the mass of material remaining on a filter after equilibration.

Despite considerable progress in measuring the atmospheric PM mass concentration, numerous uncertainties continue to exist as to the relationship between the mass and composition of material remaining on the filter as measured by the FRMs and the mass and composition of material that exists in the atmosphere as suspended PM. There is no reference standard for particles suspended in the atmosphere, nor is there an accepted way to remove particle-bound water without losing some of the semivolatile components of PM such as ammonium nitrate and semivolatile organic compounds. It also is difficult to cleanly separate fine and coarse PM. As a

result, EPA defines accuracy for PM measurements in terms of agreement of a candidate sampler with a reference sampler. Therefore, intercomparisons of samplers become very important in determining how well various samplers agree and how various design choices influence what is actually measured.

Current filtration-based mass measurements lead to significant evaporative losses of a variety of semivolatile components (i.e., species that exist in the atmosphere in dynamic equilibrium between the condensed phase and gas phase) during and possibly after collection. Important examples include ammonium nitrate, SVOCs, and particle-bound water. Loss of these components may significantly affect the quality of the measurement and can lead to either positive or negative sampling artifacts. Negative artifacts resulting from loss of ammonium nitrate and SVOC may occur during sampling because of changes in temperature, relative humidity, or the composition of the aerosol, or because of the pressure drop across the filter. Negative artifacts also may occur during handling and storage because of evaporation. Positive artifacts occur when gas-phase compounds (H_2O , HNO_3 , SO_2 , and organic compounds) absorb onto or react with filter media or collected PM, or when some particle-bound water is not removed.

Sampling systems for semivolatile PM components make use of denuders to remove the gas-phase fraction and absorptive filters to remove the condensed phase while retaining any material that subsequently evaporates from the collected PM. The loss of particulate nitrate may be determined by comparing nitrate collected on a Teflon filter to that collected on a nylon filter (which absorbs nitric acid as it evaporates from ammonium nitrate particles) preceded by a denuder to remove gas-phase nitric acid. In two studies in southern California, the $\text{PM}_{2.5}$ mass lost because of volatilization of ammonium nitrate was found to represent 10 to 20% of the total $\text{PM}_{2.5}$ mass and almost a third of the nitrate. Denuder/absorptive filter sampling systems also have been developed for measuring particulate phase organic compounds. This technique is an improvement over the filter/adsorbent collection method. However, the denuder systems currently discussed in the literature are not straightforward in their use, and the user must have a thorough understanding of the technology. The FRM for $\text{PM}_{2.5}$ will likely suffer loss of particulate nitrates and SVOCs, similar to the losses experienced with other single-filter collection systems.

It is generally desirable to collect and measure ammonium nitrate and SVOCs as part of PM mass. However, it is usually desirable to remove the particle-bound water before determining the mass. In some situations, it may be important to know how much of the suspended particle's mass or volume results from particle-bound water. Calculations and measurements indicate that aerosol water content is strongly dependent on relative humidity and composition. Particle-bound water can represent a significant mass fraction of the PM concentration at relative humidities above 60%. A substantial fraction of accumulation-mode PM is hygroscopic or deliquescent. The more hygroscopic particles tend to contain more sulfates, nitrates, and secondary organic compounds, while the less hygroscopic particles tend to contain more elemental carbon, primary organic compounds, and crustal components. Fresh, submicrometer-size soot particles may tend to shrink with increasing relative humidity because of a structural change. The effects of relative humidity on the sorption of semivolatile organic compounds on particles are not well understood. The amount of water sorbed to an atmospheric aerosol may be affected by the presence of an organic film on the particle, which can impede the transport of water across the surface.

Fine and coarse particles differ not only in formation mechanisms and size, but also in sources, composition, and chemical, physical, and biological properties. Fine and coarse particles overlap in the intermodal size range (1 to 2.5 $\mu\text{m D}_a$). As relative humidity increases, fine particles grow into this size range; as relative humidity decreases, more coarse particles may be suspended in this size range. It is desirable to measure fine PM and coarse PM separately in order to properly allocate health effects to either fine PM or coarse PM as well as to correctly determine sources by factor analysis or chemical mass balance. The selection of a cut point of 2.5 μm as a basis for EPA's 1997 NAAQS for fine particles (Federal Register, 1997) and its continued use in many health effects studies reflects the importance placed on more complete inclusion of accumulation-mode particles, while recognizing that intrusion of coarse-mode particles can occur under some conditions with this cut point.

In addition to FRM sampling of equilibrated mass to determine compliance with PM standards, EPA requires states to conduct speciation sampling, primarily to determine source categories and trends. The current speciation samplers collect $\text{PM}_{2.5}$ on three filters: (1) a Teflon filter for gravimetric determination of mass and for analysis of heavy elements by X-ray fluorescence; (2) a nylon filter preceded by a nitric acid denuder for artifact-free

determination of nitrate and measurement of other ionic species by ion chromatography; and (3) a quartz filter for measurement of elemental carbon (EC) and organic carbon (OC). In addition, IMPROVE samplers provide information on regional PM background and transport. IMPROVE samplers, in addition to the three types of filters collected by the speciation samplers, also collect a PM₁₀ sample. The IMPROVE and speciation networks use slightly different methods for determination of EC and OC. The two methods agree on total carbon (TC) but differ in the split of TC into EC and OC. The two methods also differ in their correction for positive artifacts due to absorption of volatile organic compounds on the quartz filters. Neither EC/OC method provides for any correction for negative artifacts due to the evaporation of SVOCs from the collected particles.

The EPA expects that monitoring agencies will operate continuous PM monitors and is in the process of providing guidance regarding appropriate continuous monitoring techniques. All currently available techniques for continuous measurements of suspended particle mass, e.g., the integrating nephelometer, the beta-absorption monitor, and the Tapered Element Oscillating Microbalance (TEOM), share the problem of dealing with semivolatile PM components: that is, in order to not include particle-bound water as part of the mass, the particle-bound water must be removed by heating or dehumidification; however, heating causes ammonium nitrate and SVOCs to evaporate. The TEOM monitor operates at a constant, but higher than ambient, temperature to remove particle-bound water, whereas the FRM is required to operate at no more than 5 °C above the ambient temperature. Subsequently, much of the particle-bound water is removed during equilibration at 40% relative humidity. This difference in techniques for the removal of particle-bound water causes differences in the measured mass concentration between the TEOM and FRMs.

Several new techniques for continuous PM mass measurements are currently being field tested. The RAMS measures the total mass of collected particles including semivolatile species with a TEOM monitor using a “sandwich filter.” The sandwich contains a Teflon-coated particle-collection filter followed by a charcoal-impregnated filter to collect any semivolatile species lost from the particles during sampling. The RAMS uses a Nafion dryer to remove particle-bound water from the suspended particles and a particle concentrator to reduce the quantity of gas phase organic compounds that must be removed by the denuder. The CAMM estimates ambient PM mass by measuring of the increase in the pressure drop across a

membrane filter caused by particle loading. It also uses a Nafion dryer to remove particle-bound water. A new differential TEOM offers the possibility of measuring both nonvolatile and semivolatile PM mass. In addition to continuous mass measurement, a number of techniques for continuous measurement of sulfate, nitrate, or elements are being tested.

2.3.3 Key Points

Fine and Coarse Particles

Particle size distributions show that atmospheric particles exist in two classes, fine particles and coarse particles. Fine and coarse particles are defined primarily in terms of their formation mechanisms and size; and they also differ in sources, chemical composition, and removal processes (see Table 2-2). Subsequent chapters will show that fine and coarse particles also differ with regard to aspects of concentration, exposure, dosimetry, toxicology, and epidemiology.

These differences support the setting of separate standards for fine and coarse particles. Fine and coarse particles overlap in the size range between 1 and 3 μm D_a where PM concentrations are at a minimum. Coarse particles are generally larger than this minimum and are generally formed by mechanical processes. Coarse particles and coarse-mode particles are equivalent terms. Fine particles are generally smaller than the minimum and are generally formed by coagulation and condensation of gases. Fine particles are subdivided into accumulation, Aitken, and nucleation modes. In earlier texts, nuclei mode referred to the size range now split into the Aitken and nucleation modes (see Figures 2-5 and 2-6).

Measurement of Mass and Composition

Nonvolatile Particulate Matter. Analytical techniques exist for measurement of the mass and chemical composition of PM retained on a filter (nonvolatile mass) in terms of elements (except carbon) and certain key ions (sulfate, nitrate, hydrogen, and ammonium). Acceptable measurements can be made of the TC retained on a filter. However, the split into OC and EC depends on the operational details of the analytical methods and varies somewhat among methods. Determination of the various organic compounds in the organic carbon fraction remains a challenge.

Semivolatile Particulate Matter. Important components of atmospheric PM (particle-bound water, ammonium nitrate, and many organic compounds) are termed semivolatile because significant amounts of both the gaseous and condensed phases may exist in the atmosphere in equilibrium. While particle-bound water is not considered to be a pollutant, it can significantly influence the size and light scattering and absorbing properties of particles and may also act as a carrier to convey dissolved gases or reactive species into the lungs. Most of the particle-bound water is removed by heating the particles or by equilibration of the collected particles at a low relative humidity (40%) for 24 hours. However, these processes also cause the loss of other semivolatile components. Semivolatile components also evaporate from the filter during sampling due to the pressure drop across the filter or due to a reduction in the atmospheric concentration during the sampling time. Filter collection and equilibration techniques for PM, such as prescribed by the FRM, lose a fraction of the semivolatile PM. Continuous methods must dry the PM to remove particle-bound water. If heating is used to dry the particles, more of the semivolatile components may be removed than are lost in filter sampling. Collection and retention of ammonium nitrate and SVOCs represents a major challenge in the effort to move to the continuous measurement of PM mass. The use of diffusion dryers, which dehumidify the air stream without heating, represents a promising approach. Uncertainty in the efficiency of the retention of ammonium nitrate and organic compounds on filters also impacts source category attribution and epidemiologic studies.

Separation of Fine and Coarse Particulate Matter

Satisfactory techniques are available to separate fine particles from coarse particles and to collect the fine particles on a filter. However, no consensus exists as yet on the best technique for collecting a coarse particle sample for determination of mass and composition. Candidates include multistage impaction, virtual impaction, and difference (subtracting PM_{2.5} mass or composition from PM₁₀ mass or composition). Advances in the theory and practice of virtual impaction suggest that it would be possible to design virtual impactors with much less than the 10% of fine PM collected in the coarse PM sample as is now the case for the dichotomous samplers used in air quality studies and with penetration curves as sharp as those used in the current FRM for PM_{2.5}.

REFERENCES

- Aiginger, H. (1991) Historical development and principles of total reflection x-ray fluorescence analysis (TXRF). *Spectrochim. Acta Part B* 46: 1313-1321.
- Aiginger, H.; Strelt, C. (1997) Total reflection x-ray fluorescence spectrometry (TXRF) for trace element analysis. *Spectrosc. Eur.* 9: 16-21.
- Allen, G.; Sioutas, C.; Koutrakis, P.; Reiss, R.; Lurmann, F. W.; Roberts, P. T. (1997) Evaluation of the TEOM® method for measurement of ambient particulate mass in urban areas. *J. Air Waste Manage. Assoc.* 47: 682-689.
- Allen, G. A.; Oh, J. A.; Koutrakis, P.; Sioutas, C. (1999a) Techniques for high-quality ambient coarse particulate mass measurements. *J. Air Waste Manage. Assoc.* 49(special issue): PM133-PM141.
- Allen, J. O.; Paschkewitz, J. S.; Plummer, E. F.; Lafleur, A. L.; Sarofim, A. F.; Smith, K. A. (1999b) Absorption of semi-volatile compounds in oiled impaction substrates: measurement of pyrene absorption. *Aerosol Sci. Technol.* 30: 16-29.
- Allen, G. A.; Lawrence, J.; Koutrakis, P. (1999c) Field validation of a semi-continuous method for aerosol black carbon (aethalometer) and temporal patterns of summertime hourly black carbon measurements in southwestern PA. *Atmos. Environ.* 33: 817-823.
- Allen, R.; Box, M.; Larson, T. V.; Liu, L. J. (2001) A cost-effective weighing chamber for particulate matter filters. *J. Air Waste Manage. Assoc.* 51: 1650-1653.
- American Conference of Governmental Industrial Hygienists. (1994) 1994-1995 threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- Andrews, E.; Larson, S. M. (1993) Effect of surfactant layers on the size changes of aerosol particles as a function of relative humidity. *Environ. Sci. Technol.* 27: 857-865.
- Andrews, E.; Saxena, P.; Musarra, S.; Hildemann, L. M.; Koutrakis, P.; McMurry, P. H.; Olmez, I.; White, W. H. (2000) Concentration and composition of atmospheric aerosols from the 1995 SEAVS experiment and a review of the closure between chemical and gravimetric measurements. *J. Air Waste Manage. Assoc.* 50: 648-664.
- Appel, B. R. (1993) Atmospheric sample analysis and sampling artifacts. In: Willeke, K.; Baron, P. A., eds. *Aerosol measurement: principles, techniques, and applications*. New York, NY: Van Nostrand Reinhold; pp. 233-259.
- Appel, B. R.; Tokiwa, Y.; Haik, M. (1981) Sampling of nitrates in ambient air. *Atmos. Environ.* 15: 283-289.
- Appel, B. R.; Tokiwa, Y.; Haik, M.; Kothny, E. L. (1984) Artifact particulate sulfate and nitrate formation on filter media. *Atmos. Environ.* 18: 409-416.
- Arnott, W. P.; Moosmüller, H.; Rogers, C. F.; Jin, T.; Bruch, R. (1999) Photoacoustic spectrometer for measuring light absorption by aerosol; instrument description. *Atmos. Environ.* 33: 2845-2852.
- Arnott, W. P.; Moosmüller, H.; Walker, J. W. (2000) Nitrogen dioxide and kerosene-flame soot calibration of photoacoustic instruments for measurement of light absorption by aerosols. *Rev. Sci. Instruments* 71: 4545-4552.
- Ayers, G. P.; Keywood, M. D.; Gras, J. L. (1999) TEOM vs. manual gravimetric methods for determination of PM_{2.5} aerosol mass concentrations. *Atmos. Environ.* 33: 3717-3721.
- Babich, P.; Davey, M.; Allen, G.; Koutrakis, P. (2000) Method comparisons for particulate nitrate, elemental carbon, and PM_{2.5} mass in seven U.S. cities. *J. Air Waste Manage. Assoc.* 50: 1095-1105.
- Bacon, J. R.; Chenery, S. R. N.; Ellis, A. T.; Fisher, A.; McMahon, A. W.; Potts, P. J.; Williams, J. G. (1995) Atomic spectrometry update—atomic mass spectrometry and x-ray fluorescence spectrometry. *J. Anal. At. Spectrom.* 10: 253R-310R.
- Baldwin, D. P.; Zamzow, D. S.; D'Silva, A. P. (1994) Aerosol mass measurement and solution standard additions for quantitation in laser ablation-inductively coupled plasma atomic emission spectrometry. *Anal. Chem.* 66: 1911-1917.
- Ballach, J.; Hitzengerger, R.; Schultz, E.; Jaeschke, W. (2001) Development of an improved optical transmission technique for black carbon (BC) analysis. *Atmos. Environ.* 35: 2089-2100.
- Baron, P. A.; Willeke, K., eds. (2002) *Aerosol measurement: principles, techniques, and applications*. 2nd ed. New York, NY: John Wiley & Sons.
- Benner, R. L.; Stedman, D. H. (1989) Universal sulfur detection by chemiluminescence. *Anal. Chem.* 61: 1268-1271.
- Benner, R. L.; Stedman, D. H. (1990) Field evaluation of the sulfur chemiluminescence detector. *Environ. Sci. Technol.* 24: 1592-1596.

- Bergin, M. H.; Ogren, J. A.; Schwartz, S. E.; McInnes, L. M. (1997) Evaporation of ammonium nitrate aerosol in a heated nephelometer: implications for field measurements. *Environ. Sci. Technol.* 31: 2878-2883.
- Berner, A. (1989) Haze and its relation to atmospheric accumulation aerosol. *Sci. Total Environ.* 86: 251-263.
- Bertoni, G.; Febo, A.; Perrino, C.; Possanzini, M. (1984) Annular active diffuse sampler: a new device for the collection of organic vapours. *Ann. Chim. (Rome)* 74: 97-104.
- Billiet, J.; Dams, R.; Hoste, J. (1980) Multielement thin film standards for XRF analysis. *X-Ray Spectrom.* 9: 206-211.
- Binkowski, F. S.; Ching, J. K. S. (1995) Modeling fine particulate mass and visibility using the EPA regional particulate model. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development, National Exposure Research Laboratory; report no. EPA/600/A-95/143. Available from: NTIS, Springfield, VA; PB96-139100.
- Binkowski, F. S.; Shankar, U. (1995) The regional particulate matter model. I. Model description and preliminary results. *J. Geophys. Res. [Atmos.]* 100: 26,191-26,209.
- Birch, M. E. (1998) Analysis of carbonaceous aerosols: interlaboratory comparison. *Analyst (Cambridge, UK)* 123: 851-857.
- Birch, M. E.; Cary, R. A. (1996) Elemental carbon-based method for occupational monitoring of particulate diesel exhaust: methodology and exposure issues. *Analyst* 121: 1183-1190.
- Birmili, W.; Wiedensohler, A. (1998) The influence of meteorological parameters on ultrafine particle production at a continental site. *J. Aerosol Sci.* 29(suppl. 1): S1015-S1016.
- Bowers, W. D.; Chuan, R. L. (1989) Surface acoustic-wave piezoelectric crystal aerosol mass monitor. *Rev. Sci. Instrum.* 60: 1297-1302.
- Brauer, M.; Koutrakis, P.; Keeler, G. J.; Spengler, J. D. (1991) Indoor and outdoor concentrations of inorganic acidic aerosols and gases. *J. Air Waste Manage. Assoc.* 41: 171-181.
- Breum, N. O. (2000) The dust holding capacity of porous plastic foam used in particle size-selective sampling. *J. Aerosol Sci.* 31: 379-385.
- Brock, J. R.; Durham, J. L. (1984) Aqueous aerosol as a reactant in the atmosphere. In: Calvert, J. G., ed. *SO₂, NO and NO₂ oxidation mechanisms: atmospheric considerations*. Boston, MA: Butterworth Publishers; pp. 209-249. (Teasley, J. I., ed. *Acid precipitation series: v. 3*).
- Brook, J. R.; Dann, T. F. (1999) Contribution of nitrate and carbonaceous species to PM_{2.5} observed in Canadian cities. *J. Air Waste Manage. Assoc.* 49: 193-199.
- Brorström, E.; Grennfelt, P.; Lindskog, A. (1983) The effect of nitrogen dioxide and ozone on the decomposition of particle-associated polycyclic aromatic hydrocarbons during sampling from the atmosphere. *Atmos. Environ.* 17: 601-605.
- Bunz, H.; Koyro, M.; Möhler, O.; Saathoff, H. (1996) Temperature dependence of the redistribution mechanisms for volatile components of atmospheric aerosol particles. *J. Aerosol Sci.* 27(suppl. 1): S653-S654.
- Cachier, H.; Bremond, M.-P.; Buat-Ménard, P. (1989) Determination of atmospheric soot carbon with a simple thermal method. *Tellus* 41B: 379-390.
- Cadle, S. H.; Groblicki, P. J. (1982) An evaluation of methods for the determination of organic and elemental carbon in particulate samples. In: Wolff, G. T.; Klimisch, R. L., eds. *Particulate carbon: atmospheric life cycle*. New York, NY: Plenum Press; pp. 89-109.
- Cadle, S. H.; Mulawa, P.A. (1990) Atmospheric carbonaceous species measurement methods comparison study: General Motors results. *Aerosol Sci. Technol.* 12: 128-131.
- Cahill, T. A. (1981) Innovative aerosol sampling devices based upon PIXE capabilities. In: Johansson, S. A. E., ed. *Particle induced x-ray emission and its analytical applications: proceedings of the 2nd international conference on PIXE and its analytical applications; June, 1980; Lund, Sweden*. Amsterdam, The Netherlands: North-Holland Publishing Co. *Nucl. Instr. Meth. Phys. Res.* 181: 473-480.
- Cahill, T. A. (1985) Proton-induced X-ray emission (PIXE). In: Res, J. P., ed. *Metals Handbook: Materials Characterization*. Cleveland, OH: American Society for Metals.
- Cahill, T. A. (1990) Particle-induced x-ray emission. In: Mills, K., ed. *Metals Handbook; v. 10, p. 102*. 9th ed. Cleveland, OH: American Society for Metals.
- Cahill, T. A.; Eldred, R. A.; Motallebi, N.; Malm, W. C. (1989) Indirect measurement of hydrocarbon aerosols across the United States by nonsulfate hydrogen-remaining gravimetric mass correlations. *Aerosol Sci. Technol.* 10: 421-429.
- Cahill, T. A.; Wilkinson, K.; Schnell, R. (1992) Composition analyses of size-resolved aerosol samples taken from aircraft downwind of Kuwait, spring 1991. *J. Geophys. Res. [Atmos.]* 97: 14,513-14,520.

- Cahill, T. A.; Wakabayashi, P.; James, T. A. (1996) Chemical states of sulfate at Shenandoah National Park during summer, 1991. *Nucl. Instrum. Methods Phys. Res. B* 109/110: 542-547.
- Camp, D. C.; Stevens, R. K.; Cobourn, W. G.; Husar, R. B.; Collins, J. F.; Huntzicker, J. J.; Husar, J. D.; Jaklevic, J. M.; McKenzie, R. L.; Tanner, R. L.; Tesch, J. W. (1982) Intercomparison of concentration results from fine particle sulfur monitors. *Atmos. Environ.* 16: 911-916.
- Campbell, A. J.; Humayun, M. (1999) Trace element microanalysis in iron meteorites by laser ablation ICPMS. *Anal. Chem.* 71: 939-946.
- Carson, P. G.; Johnston, M. V.; Wexler, A. S. (1997) Laser desorption/ionization of ultra fine aerosol particles. *Rapid Commun. Mass Spectrom.* 11: 993-996.
- Castaldi, M. J.; Senkan, S. M. (1998) Real-time, ultrasensitive monitoring of air toxics by laser photoionization time-of-flight mass spectroscopy. *J. Air Waste Manage. Assoc.* 48: 77-81.
- Chandramouli, B.; Kamens, R. (2001) The photochemical formation and gas-particle partitioning of oxidation products of decamethyl cyclopentasiloxane and decamethyl tetrasiloxane in the atmosphere. *Atmos. Environ.* 35: 87-95.
- Chandramouli, B.; Jang, M.; Kamens, R. M. (2003) Gas-particle partitioning of semi-volatile organics on organic aerosols using a predictive activity coefficient model: analysis of the effects of parameter choices on model performance. *Atmos. Environ.* 37: 853-864.
- Chang, M.; Kim, S.; Sioutas, C. (1999) Experimental studies on particle impaction and bounce: effects of substrate design and material. *Atmos. Environ.* 33: 2313-2322.
- Chang, M. C.; Sioutas, C.; Kim, S.; Gong, H., Jr.; Linn, W. S. (2000) Reduction of nitrate losses from filter and impactor samplers by means of concentration enrichment. *Atmos. Environ.* 34: 85-98.
- Chang, M.-C.; Geller, M. D.; Sioutas, C.; Fokkens, P. H. B.; Cassee, F. R. (2002) Development and evaluation of a compact, highly efficient coarse particle concentrator for toxicological studies. *Aerosol Sci. Technol.* 36: 492-501.
- Chein, H.; Lundgren, D. A. (1993) A virtual impactor with clean air core for the generation of aerosols with narrow size distributions. *Aerosol Sci. Technol.* 18: 376-388.
- Chen, B. T. (1993) Instrument calibration. In: Willeke, K.; Baron, P. A., eds. *Aerosol measurement: principles, techniques, and applications*. New York, NY: Van Nostrand Reinhold; pp. 493-520.
- Chen, B. T.; Yeh, H. C. (1987) An improved virtual impactor: design and performance. *J. Aerosol Sci.* 18: 203-214.
- Chen, B. T.; Yeh, H. C.; Cheng, Y. S. (1985) A novel virtual impactor: calibration and use. *J. Aerosol Sci.* 16: 343-354.
- Chen, B. T.; Yeh, H. C.; Cheng, Y. S. (1986) Performance of a modified virtual impactor. *Aerosol Sci. Technol.* 5: 369-376.
- Cheng, Y.-S.; Yeh, H.-C. (1979) Particle bounce in cascade impactors. *Environ. Sci. Technol.* 13: 1392-1396.
- Chow, J. C. (1995) Measurement methods to determine compliance with ambient air quality standards for suspended particles. *J. Air Waste Manage. Assoc.* 45: 320-382.
- Chow, J. C.; Watson, J. G. (1999) Ion chromatography in elemental analysis of airborne particles. In: Landsberger, S.; Creatchman, M., eds. *Elemental Analysis of Airborne Particles*. Newark, NJ: Gordon and Breach Science Publishers; pp. 97-137.
- Chow, J. C.; Watson, J. G.; Pritchett, L. C.; Pierson, W. R.; Frazier, C. A.; Purcell, R. G. (1993) The DRI thermal/optical reflectance carbon analysis system: description, evaluation and applications in U.S. air quality studies. In: Lodge, J. P., Jr. *Fourth international conference on carbonaceous particles in the atmosphere; April 1991; Vienna, Austria*. *Atmos. Environ. Part A* 27: 1185-1201.
- Chow, J. C.; Fujita, E. M.; Watson, J. G.; Lu, Z.; Lawson, D. R.; Ashbaugh, L. L. (1994a) Evaluation of filter-based aerosol measurements during the 1987 Southern California Air Quality Study. *Environ. Monit. Assess.* 30: 49-80.
- Chow, J. C.; Watson, J. G.; Fujita, E. M.; Lu, Z.; Lawson, D. R.; Ashbaugh, L. L. (1994b) Temporal and spatial variations of PM_{2.5} and PM₁₀ aerosol in the Southern California Air Quality Study. *Atmos. Environ.* 28: 2061-2080.
- Chow, J. C.; Watson, J. G.; Lu, Z.; Lowenthal, D. H.; Frazier, C. A.; Solomon, P. A.; Thuillier, R. H.; Magliano, K. (1996) Descriptive analysis of PM_{2.5} and PM₁₀ at regionally representative locations during SJVAQS/AUSPEX. In: Parrish, D.; Trainer, M.; Rao, S. T.; Solomon, P. A., eds. *Air and Waste Management Association international specialty conference on regional photochemical measurements and modeling, part 2; November 1993; San Diego, CA*. *Atmos. Environ.* 30: 2079-2112.

- Chow, J. C.; Zielinska, B.; Watson, J. G.; Fujita, E. M.; Richards, H. W.; Neff, W. D.; Dietrich, D.; Hering, S. V. (1998) Northern Front Range Air Quality Study. Volume A: Ambient measurements. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere. Available: <http://www.nfraqs.colostate.edu/dri.html> [4 February, 2002].
- Chow, J. C.; Watson, J. G.; Crow, D.; Lowenthal, D. H.; Merrifield, T. (2000) Comparison of IMPROVE and NIOSH carbon measurements. *Aerosol Sci. Technol.* 34: 23-34.
- Chow, J. C.; Watson, J. G.; Crow, D.; Lowenthal, D. H.; Merrifield, T. (2001) Comparison of IMPROVE and NIOSH carbon measurements. *Aerosol Sci. Technol.* 34: 23-34.
- Claiborn, C. S.; Finn, D.; Larson, T. V.; Koenig, J. Q. (2000) Windblown dust contributes to high PM_{2.5} concentrations. *J. Air Waste Manage. Assoc.* 50: 1440-1445.
- Clark, W. E.; Whitby, K. T. (1975) Measurements of aerosols produced by the photochemical oxidation of SO₂ in air. *J. Colloid Interface Sci.* 51: 477-490.
- Clarke, A. D. (1992) Atmospheric nuclei in the remote free-troposphere. *J. Atmos. Chem.* 14: 479-488.
- Clarke, A. D.; Varner, J. L.; Eisele, F.; Mauldin, R. L.; Tanner, D.; Litchy, M. (1998) Particle production in the remote marine atmosphere: cloud outflow and subsidence during ACE 1. *J. Geophys. Res. A* 103: 16,397-16,409.
- Cobourn, W. G.; Husar, R. B. (1982) Diurnal and seasonal patterns of particulate sulfur and sulfuric acid in St. Louis, July 1977-June 1978. *Atmos. Environ.* 16: 1441-1450.
- Cobourn, W. G.; Husar, R. B.; Husar, J. D. (1978) Continuous *in situ* monitoring of ambient particulate sulfur using flame photometry and thermal analysis. In: Husar, R. B.; Lodge, J. P., Jr.; Moore, D. J., eds. *Sulfur in the atmosphere: proceedings of the international symposium; September 1977; Dubrovnik, Yugoslavia.* *Atmos. Environ.* 12: 89-98.
- Code of Federal Regulations. (2001a) Appendix J to Part 50—reference method for the determination of particulate matter as PM₁₀ in the atmosphere. *C. F. R.* 40: pt. 50, app. J.
- Code of Federal Regulations. (2001b) Ambient air monitoring reference and equivalent methods. *C. F. R.* 40: §53.
- Code of Federal Regulations. (2001c) Appendix L to Part 50—reference method for the determination of fine particulate matter as PM_{2.5} in the atmosphere. *C. F. R.* 40: pt. 50, app. L.
- Code of Federal Regulations. (2001d) Ambient air quality surveillance. *C. F. R.* 40: §58.
- Code of Federal Regulations. (2001e) Appendix M to Part 50—reference method for the determination of particulate matter as PM₁₀ in the atmosphere. *C. F. R.* 40: pt. 50, app. M.
- Conner, W. D. (1966) An inertial-type particle separator for collecting large samples. *J. Air Pollut. Control Assoc.* 16: 35-38.
- Conner, T. L.; Norris, G. A.; Landis, M. S.; Williams, R. W. (2001) Individual particle analysis of indoor, outdoor, and community samples from the 1998 Baltimore particulate matter study. *Atmos. Environ.* 35: 3935-3946.
- Cotham, W. E.; Bidleman, T. F. (1992) Laboratory investigations of the partitioning of organochlorine compounds between the gas phase and atmospheric aerosols on glass fiber filters. *Environ. Sci. Technol.* 26: 469-478.
- Countess, R. J. (1990) Interlaboratory analyses of carbonaceous aerosol samples. *Aerosol Sci. Technol.* 12: 114-121.
- Courtney, W. J.; Shaw, R. W.; Dzubay, T. G. (1982) Precision and accuracy of a β gauge for aerosol mass determinations. *Environ. Sci. Technol.* 16: 236-239.
- Coutant, R. W.; Callahan, P. J.; Chuang, J. C.; Lewis, R. G. (1992) Efficiency of silicone-grease-coated denuders for collection of polynuclear aromatic hydrocarbons. *Atmos. Environ. Part A* 26: 2831-2834.
- Covert, D. S.; Kapustin, V. N.; Quinn, P. K.; Bates, T. S. (1992) New particle formation in the marine boundary layer. *J. Geophys. Res. [Atmos.]* 97: 20,581-20,589.
- Covert, D. S.; Wiedensohler, A.; Aalto, P.; Heintzenberg, J.; McMurry, P. H.; Leck, C. (1996) Aerosol number size distributions from 3 to 500 nm diameter in the arctic marine boundary layer during summer and autumn. *Tellus* 48B: 197-212.
- Cui, W.; Machir, J.; Lewis, L.; Eatough, D. J.; Eatough, N. L. (1997) Fine particulate organic material at Meadview during the project MOHAVE summer intensive study. *J. Air Waste Manage. Assoc.* 47: 357-369.
- Cui, W.; Eatough, D. J.; Eatough, N. L. (1998) Fine particulate organic material in the Los Angeles Basin - I: assessment of the high-volume Brigham Young University organic sampling system, BIG BOSS. *J. Air Waste Manage. Assoc.* 48: 1024-1037.
- Dagnall, R. M.; Thompson, K. C.; West, T. S. (1967) Molecular-emission in cool flames. Part I. the behaviour of sulphur species in a hydrogen-nitrogen diffusion flame and in a shielded air-hydrogen flame. *Analyst* 92: 506-512.
- Dams, R.; Robbins, J. A.; Rahn, K. A.; Winchester, J. W. (1970) Nondestructive neutron activation analysis of air pollution particulates. *Anal. Chem.* 42: 861-867.

- Dane, A. D.; Timmermans, P. A. M.; Van Sprang, H. A.; Buydens, L. M. C. (1996) A genetic algorithm for model-free x-ray fluorescence analysis of thin film. *Anal. Chem.* 68: 2419-2425.
- Davis, D. W.; Reynolds, R. L.; Tsou, G. C.; Zafonte, L. (1977) Filter attenuation corrections for x-ray fluorescence analysis of atmospheric aerosols. *Anal. Chem.* 49: 1990-1993.
- Davis, C. S.; Fellin, P.; Otson, R. (1987) A review of sampling methods for polyaromatic hydrocarbons in air. *J. Air Pollut. Control Assoc.* 37: 1397-1408.
- DeBoer, D. K. G.; Leenaers, A. J. G.; Van Den Hoogenhof, W. W. (1995) Glancing-incidence x-ray analysis of thin-layered materials: a review. *X-Ray Spectrom.* 24: 91-102.
- Ding, Y.; Koutrakis, P. (2000) Development of a dichotomous slit nozzle virtual impactor. *J. Aerosol Sci.* 31: 1421-1431.
- Ding, Y.; Lee, M. L.; Eatough, D. J. (1998a) The determination of total nitrite and *N*-nitroso compounds in atmospheric samples. *Int. J. Environ. Anal. Chem.* 69: 243-255.
- Ding, Y.; Cui, W.; Lee, M. L.; Eatough, D. J. (1998b) Fine particulate *N*-nitroso and nitrite organic compounds in the atmosphere. *Appl. Occup. Environ. Hyg.* 13: 432-438.
- Ding, Y.; Pang, Y.; Eatough, D. J.; Eatough, N. L.; Tanner, R. L. (2002) High volume diffusion denuder sampler for the routine monitoring of fine particulate matter: II. field evaluation of the PC BOSS. *Aerosol Sci. Technol.* 36: 383-396.
- Dockery, D. W.; Pope, C. A., III; Xu, X.; Spengler, J. D.; Ware, J. H.; Fay, M. E.; Ferris, B. G., Jr.; Speizer, F. E. (1993) An association between air pollution and mortality in six U.S. cities. *N. Engl. J. Med.* 329: 1753-1759.
- Durham, J. L.; Wilson, W. E.; Bailey, E. B. (1978) Application of an SO₂-denuder for continuous measurement of sulfur in submicrometric aerosols. *Atmos. Environ.* 12: 883-886.
- Dzubay, T. G. (1980) Chemical element balance method applied to dichotomous sampler data. *Ann. N. Y. Acad. Sci.* 338: 126-144.
- Dzubay, T. G.; Stevens, R. K. (1975) Ambient air analysis with dichotomous sampler and X-ray fluorescence spectrometer. *Environ. Sci. Technol.* 9: 663-668.
- Dzubay, T. G.; Hines, L. E.; Stevens, R. K. (1976) Particle bounce errors in cascade impactors. *Atmos. Environ.* 10: 229-234.
- Dzubay, T. G.; Stevens, R. K.; Peterson, C. M. (1977) Application of the dichotomous sampler to the characterization of ambient aerosols. In: Dzubay, T. G., ed. *X-ray fluorescence analysis of environmental samples*. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.
- Easter, R. C.; Peters, L. K. (1994) Binary homogeneous nucleation: temperature and relative humidity fluctuations, nonlinearity, and aspects of new particle production in the atmosphere. *J. Appl. Meteorol.* 33: 775-784.
- Eatough, D. J. (1999) BOSS, the Brigham Young University Organic Sampling System: determination of particulate carbonaceous material using diffusion denuder sampling technology. In: Lane, D., ed. *Gas and particle phase measurements of atmospheric organic compounds*. New York, NY: Gordon and Breach Science Publishers. (*Advances in environmental, industrial and process control technologies*: v. 2).
- Eatough, D. J.; Wadsworth, A.; Eatough, D. A.; Crawford, J. W.; Hansen, L. D.; Lewis, E. A. (1993) A multiple-system, multi-channel diffusion denuder sampler for the determination of fine-particulate organic material in the atmosphere. *Atmos. Environ. Part A* 27: 1213-1219.
- Eatough, D. J.; Tang, H.; Cui, W.; Machir, J. (1995) Determination of the size distribution and chemical composition of fine particulate semi-volatile organic material in urban environments using diffusion denuder technology. In: Phalen, R. F.; Bates, D. V., eds. *Proceedings of the colloquium on particulate air pollution and human mortality and morbidity, part II*; January 1994; Irvine, CA. *Inhalation Toxicol.* 7: 691-710.
- Eatough, D. J.; Eatough, D. A.; Lewis, L.; Lewis, E. A. (1996) Fine particulate chemical composition and light extinction at Canyonlands National Park using organic particulate material concentrations obtained with a multisystem, multichannel diffusion denuder sampler. *J. Geophys. Res. [Atmos.]* 101: 19,515-19,531.
- Eatough, D. J.; Obeidi, F.; Pang, Y.; Ding, Y.; Eatough, N. L.; Wilson, W. E. (1999a) Integrated and real-time diffusion denuder sampler for PM_{2.5}. *Atmos. Environ.* 33: 2835-2844.
- Eatough, D. J.; Pang, Y.; Eatough, N. L. (1999b) Determination of PM_{2.5} sulfate and nitrate with a PC-BOSS designed for routine sampling for semi-volatile particulate matter. *J. Air Waste Manage. Assoc.* 49(special issue): PM69-75.
- Eatough, D. J.; Eatough, N. L.; Obeidi, F.; Pang, Y.; Modey, W.; Long, R. (2001) Continuous determination of PM_{2.5} mass, including semi-volatile species. *Aerosol Sci. Technol.* 34: 1-8.
- Elvers, B.; Hawkins, S.; Schulz, G., eds. (1991) Peroxo compounds. *Inorganic*. In: *Ullmann's Encyclopedia of Industrial Chemistry*. V. A19. 5th ed. New York, NY: VCH Publishers; pp. 177-197.

- Fairchild, C. I.; Wheat, L. D. (1984) Calibration and evaluation of a real-time cascade impactor. *Am. Ind. Hyg. Assoc. J.* 45: 205-211.
- Falconer, R. L.; Bidleman, T. F.; Cotham, W. E. (1995) Preferential sorption of non- and mono-ortho-polychlorinated biphenyls to urban aerosols. *Environ. Sci. Technol.* 29: 1666-1673.
- Fang, C. P.; McMurry, P. H.; Marple, V. A.; Rubow, K. L. (1991) Effect of flow-induced relative humidity changes on size cuts for sulfuric acid droplets in the microorifice uniform deposit impactor (MOUDI). *Aerosol Sci. Technol.* 14: 266-277.
- Farwell, S. O.; Rasmussen, R. A. (1976) Limitations of the FPD and ECD in atmospheric analysis: a review. *J. Chromatogr. Sci.* 14: 224-234.
- Fassel, V. A.; Kniseley, R. N. (1974) Inductively coupled plasma-optical emission spectroscopy. *Anal. Chem.* 46: 1110A-1120A.
- Federal Register. (1987) Revisions to the national ambient air quality standards for particulate matter. F. R. (July 1) 52: 24,634-24,669.
- Federal Register. (1997) National ambient air quality standards for particulate matter; final rule. F. R. (July 18) 62: 38,652-38,752.
- Federal Register. (1998) National ambient air quality standards for particulate matter and revised requirements for designation of reference and equivalent methods for PM_{2.5} and ambient air quality surveillance for particulate matter; correction. F. R. (February 17) 63: 7710-7718.
- Fergenson, D. P.; Song, X.-H.; Ramadan, Z.; Allen, J. O.; Hughes, L. S.; Cass, G. R.; Hopke, P. K.; Prather, K. A. (2001) Quantification of ATOFMS data by multivariate methods. *Anal. Chem.* 73: 3535-3541.
- Fernandez, F. J. (1989) Atomic absorption spectroscopy. In: Lodge, J. P., Jr., ed. *Methods of Air Sampling and Analysis*. 3rd ed. Chelsea, MI: Lewis Publishers, Inc.; pp. 83-89.
- Finn, D.; Rumburg, B.; Claiborn, C.; Bamesberger, L.; Siems, W. F.; Koenig, J.; Larson, T.; Norris, G. (2001) Sampling artifacts from the use of denuder tubes with glycerol based coatings in the measurement of atmospheric particulate matter. *Environ. Sci. Technol.* 35: 40-44.
- Fitz, D. R. (1990) Reduction of the positive organic artifact on quartz filters. *Aerosol. Sci. Technol.* 12: 142-148.
- Flocchini, R. G.; Cahill, T. A.; Shadoan, D. J.; Lange, S. J.; Eldred, R. A.; Feeney, P. J.; Wolfe, G. W.; Simmeroth, D. C.; Suder, J. K. (1976) Monitoring California's aerosols by size and elemental composition. *Environ. Sci. Technol.* 10: 76-82.
- Forney, L. J.; Ravenhall, D. G.; Winn, D. S. (1978) Aerosol impactors: a study of a fluid jet impinging upon a void. *J. Appl. Phys.* 49: 2339-2345.
- Forney, L. J.; Ravenhall, D. G.; Lee, S. S. (1982) Experimental and theoretical study of a two-dimensional virtual impactor. *Environ. Sci. Technol.* 16: 492-497.
- Friedlander, S. K. (1970) The characterization of aerosols distributed with respect to size and chemical composition. *J. Aerosol Sci.* 1: 295-307.
- Friedlander, S. K. (2000). *Smoke, dust and haze: fundamentals of aerosol dynamics*. 2nd ed. New York, NY: Oxford University Press.
- Friedlander, S. K.; Yeh, E. K. (1998) The submicron atmospheric aerosol as a carrier of reactive chemical species: case of peroxides. *Appl. Occup. Environ. Hyg.* 13: 416-420.
- Fuchs, N. A. (1964) *The mechanics of aerosols*. New York, NY: Pergamon Press.
- Fuchs, N. A. (1989) *The mechanics of aerosols*. New York, NY: Dover Publications.
- Fung, K. (1990) Particulate carbon speciation by MnO₂ oxidation. *Aerosol Sci. Technol.* 12: 122-127.
- Fung, K.; Chow, J. C.; Watson, J. G. (2002) Evaluation of OC/EC speciation by thermal manganese dioxide oxidation and the IMPROVE method. *J. Air Waste Manage. Assoc.* 52: 1333-1341.
- Gallorini, M. (1995) Trace element monitoring in atmospheric pollution processes by neutron activation analysis. *Microchem. J.* 51: 127-137.
- Gard, E.; Mayer, J. E.; Morrical, B. D.; Dienes, T.; Fergenson, D. P.; Prather, K. A. (1997) Real-time analysis of individual atmospheric aerosol particles: design and performance of a portable ATOFMS. *Anal. Chem.* 69: 4083-4091.
- Gauderman, W. J.; McConnell, R.; Gilliland, F.; London, S.; Thomas, D.; Avol, E.; Vora, H.; Berhane, K.; Rappaport, E. B.; Lurmann, F.; Margolis, H. G.; Peters, J. (2000) Association between air pollution and lung function growth in southern California children. *Am. J. Respir. Crit. Care Med.* 162: 1383-1390.
- Ge, Z.; Wexler, A. S.; Johnston, M. V. (1998) Laser desorption/ionization of single ultrafine multicomponent aerosols. *Environ. Sci. Technol.* 32: 3218-3223.
- Geller, M. D.; Kim, S.; Misra, C.; Sioutas, C.; Olson, B. A.; Marple, V. A. (2002) Methodology for measuring size-dependent chemical composition of ultrafine particles. *Aerosol Sci. Technol.* 36: 748-763.

- Goss, K.-U.; Eisenreich, S. J. (1997) Sorption of volatile organic compounds to particles from a combustion source at different temperatures and relative humidities. *Atmos. Environ.* 31: 2827-2834.
- Gray, A. L.; Williams, J. G. (1987a) Oxide and doubly charged ion response of a commercial inductively coupled plasma mass spectrometry instrument. *J. Anal. At. Spectrom.* 2: 81-82.
- Gray, A. L.; Williams, J. G. (1987b) System optimisation and the effect on polyatomic, oxide and doubly charged ion response of a commercial inductively coupled plasma mass spectrometry instrument. *J. Anal. At. Spectrom.* 2: 599-606.
- Grosjean, D. (1983) Polycyclic aromatic hydrocarbons in Los Angeles air from samples collected on Teflon, glass and quartz filters. *Atmos. Environ.* 17: 2565-2573.
- Grosjean, D.; Parmar, S. S. (1990) Interferences from aldehydes and peroxyacetyl nitrate when sampling urban air organic acids on alkaline traps. *Environ. Sci. Technol.* 24: 1021-1026.
- Gundel, L. A.; Lane, D. A. (1999) Sorbent-coated diffusion denuders for direct measurement of gas/particle partitioning by semi-volatile organic compounds. In: Lane, D. A., ed. *Gas and particle phase measurements of atmospheric organic compounds*. Amsterdam, The Netherlands: Gordon and Breach Science Publishers. (Advances in environmental, industrial and process control technologies: v. 2).
- Gundel, L. A.; Lee, V. C.; Mahanama, K. R. R.; Stevens, R. K.; Daisey, J. M. (1995) Direct determination of the phase distributions of semi-volatile polycyclic aromatic hydrocarbons using annular denuders. *Atmos. Environ.* 29: 1719-1733.
- Haller, L.; Claiborn, C.; Larson, T.; Koenig, J.; Norris, G.; Edgar, R. (1999) Airborne particulate matter size distributions in an arid urban area. *J. Air Waste Manage. Assoc.* 49: 161-168.
- Hänninen, O. O.; Koistinen, K. J.; Kousa, A.; Keski-Karhu, J.; Jantunen, M. J. (2002) Quantitative analysis of environmental factors in differential weighing of blank teflon filters. *J. Air Waste Manage. Assoc.* 52: 134-139.
- Hansen, A. D. A.; McMurry, P. H. (1990) An intercomparison of measurements of aerosol elemental carbon during the 1986 carbonaceous species method comparison study. *J. Air Waste Manage. Assoc.* 40: 894-895.
- Hansen, A. D. A.; Rosen, H.; Novakov, T. (1984) The aethalometer - an instrument for the real-time measurement of optical absorption by aerosol particles. *Sci. Total Environ.* 36: 191-196.
- Harman, J. N. (1989) Inductively coupled plasma emission spectroscopy. In: Lodge, J. P., Jr., ed. *Methods of air sampling and analysis*. 3rd ed. Chelsea, MI: Lewis Publishers, Inc.; pp. 89-92.
- Haupt, O.; Klane, B.; Schaefer, C.; Dannecker, W. (1995) Preparation of quartz fibre filter standards for x-ray fluorescence analysis of aerosol samples. *X-Ray Spectrom.* 24: 267-275.
- He, K.; Yang, F.; Ma, Y.; Zhang, Q.; Yao, X.; Chan, C. K.; Cadle, S.; Chan, T.; Mulawa, P. (2001) The characteristics of PM_{2.5} in Beijing, China. *Atmos. Environ.* 35: 4959-4970.
- Hegg, D. A.; Hobbs, P. V. (1982) Measurements of sulfate production in natural clouds. *Atmos. Environ.* 16: 2663-2668.
- Hegg, D. A.; Hobbs, P. V. (1986) Sulfate and nitrate chemistry in cumuliform clouds. *Atmos. Environ.* 20: 901-909.
- Hegg, D. A.; Radke, L. F.; Hobbs, P. V. (1990) Particle production associated with marine clouds. *J. Geophys. Res.* [Atmos.] 95: 13,917-13,926.
- Hegg, D. A.; Radke, L. F.; Hobbs, P. V. (1991) Measurements of Aitken nuclei and cloud condensation nuclei in the marine atmosphere and their relation to the DMS-cloud-climate hypothesis. *J. Geophys. Res.* [Atmos.] 96: 18,727-18,733.
- Hering, S.; Cass, G. (1999) The magnitude of bias in the measurement of PM_{2.5} arising from volatilization of particulate nitrate from Teflon filters. *J. Air Waste Manage. Assoc.* 49: 725-733.
- Hering, S. V.; Friedlander, S. K. (1982) Origins of aerosol sulfur size distributions in the Los Angeles basin. *Atmos. Environ.* 16: 2647-2656.
- Hering, S. V.; Appel, B. R.; Cheng, W.; Salaymeh, F.; Cadle, S. H.; Mulawa, P. A.; Cahill, T. A.; Eldred, R. A.; Surovik, M.; Fitz, D.; Howes, J. E.; Knapp, K. T.; Stockburger, L.; Turpin, B. J.; Huntzicker, J. J.; Zhang, X.-Q.; McMurry, P. H. (1990) Comparison of sampling methods for carbonaceous aerosols in ambient air. *Aerosol Sci. Technol.* 12: 200-213.
- Hinds, W. C. (1999) *Aerosol technology: properties, behavior, and measurement of airborne particles*. 2nd ed. New York, NY: John Wiley & Sons.
- Hinz, K.-P.; Kaufmann, R.; Spengler, B. (1994) Laser-induced mass analysis of single particles in the airborne state. *Anal. Chem.* 66: 2071-2076.
- Hitzenberger, R.; Dusek, U.; Berner, A. (1996) Black carbon measurements using an integrating sphere. *J. Geophys. Res.* [Atmos.] 101: 19,601-19,606.

- Hitzenberger, R.; Berner, A.; Dusek, U.; Alabashi, R. (1997) Humidity-dependent growth of size-segregated aerosol samples. *Aerosol Sci. Technol.* 27: 116-130.
- Hitzenberger, R.; Jennings, S. G.; Larson, S. M.; Dillner, A.; Cachier, H.; Galambos, Z.; Rouc, A.; Spain, T. G. (1999) Intercomparison of measurement methods for black carbon aerosols. *Atmos. Environ.* 33: 2823-2833.
- Holyńska, B.; Ptasiński, J.; Maenhaut, W.; Annegarn, H. J. (1997) Energy-dispersive x-ray fluorescence spectrometer with capillary optics for the chemical analysis of atmospheric aerosols with high time resolution. *J. Aerosol Sci.* 28: 1455-1463.
- Hoppel, W. A.; Frick, G. M.; Fitzgerald, J. W.; Larson, R. E. (1994) Marine boundary layer measurements of new particle formation and the effects nonprecipitating clouds have on aerosol size distribution. *J. Geophys. Res. [Atmos.]* 99: 14,443-14,459.
- Houck, J. E.; Chow, J. C.; Watson, J. G.; Simons, C. A.; Pritchett, L. C.; Goulet, J. M.; Frazier, C. A. (1989) Determination of particle size distribution and chemical composition of particulate matter from selected sources in California: volume I and executive summary (final report). Sacramento, CA: California Air Resources Board.
- Hounam, R. F.; Sherwood, R. J. (1965) The cascade centripeter: a device for determining the concentration and size distribution of aerosols. *Am. Ind. Hyg. Assoc. J.* 26: 122-131.
- Hung, H.-F.; Wang, C.-S. (2001) Experimental determination of reactive oxygen species in Taipei aerosols. *J. Aerosol Sci.* 32: 1201-1211.
- Huntzicker, J. J.; Hoffman, R. S.; Ling, C.-S. (1978) Continuous measurement and speciation of sulfur-containing aerosols by flame photometry. *Atmos. Environ.* 12: 83-88.
- Huntzicker, J. J.; Johnson, R. L.; Shah, J. J.; Cary, R. A. (1982) Analysis of organic and elemental carbon in ambient aerosols by a thermal-optical method. In: Wolff, G. T.; Klimisch, R. L., eds. *Particulate carbon - atmospheric life cycle: proceedings of an international symposium; October 1980; Warren, MI.* New York, NY: Plenum Press; pp. 79-88.
- Husar, R. B.; Tratt, D. M.; Schichtel, B. A.; Falke, S. R.; Li, F.; Jaffe, D.; Gassó, S.; Gill, T.; Laulainen, N. S.; Lu, F.; Reheis, M. C.; Chun, Y.; Westphal, D.; Holben, B. N.; Gueymard, C.; McKendry, I.; Kuring, N.; Feldman, G. C.; McClain, C.; Frouin, R. J.; Merrill, J.; DuBois, D.; Vignola, F.; Murayama, T.; Nickovic, S.; Wilson, W. E.; Sassen, K.; Sugimoto, N.; Malm, W. C. (2001) The Asian dust events of April 1998. *J. Geophys. Res.* 106: 18,317-18,330.
- Jackson, K. W.; Mahmood, T. M. (1994) Atomic absorption, atomic emission, and flame emission spectrometry. *Anal. Chem.* 66: 252R-279R.
- Jaklevic, J. M.; Loo, B. W.; Goulding, F. S. (1977) Photon-induced X-ray fluorescence analysis using energy-dispersive detector and dichotomous sampler. In: Dzubay, T. G., ed. *X-ray fluorescence analysis of environmental samples.* Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 3-18.
- Jaklevic, J. M.; Gatti, R. C.; Goulding, F. S.; Loo, B. W. (1981a) A β -gauge method applied to aerosol samples. *Environ. Sci. Technol.* 15: 680-686.
- Jaklevic, J. M.; Loo, B. W.; Fujita, T. Y. (1981b) Automatic particulate sulfur measurements with a dichotomous sampler and on-line x-ray fluorescence analysis. *Environ. Sci. Technol.* 15: 687-690.
- Jang, M.; Kamens, R. M. (1998) A thermodynamic approach for modeling partitioning of semivolatile organic compounds on atmospheric particulate matter: humidity effects. *Environ. Sci. Technol.* 32: 1237-1243.
- Jang, M.; Kamens, R. M. (1999) Exploratory model approach for adsorption partitioning of SOCs on inorganic particles. *Environ. Sci. Technol.*
- Jang, M.; Kamens, R. M. (2001) Atmospheric secondary aerosol formation by heterogeneous reactions of aldehydes in the presence of a sulfuric acid aerosol catalyst. *Environ. Sci. Technol.* 35: 4758-4766.
- Jang, M.; Kamens, R. M.; Leach, K. B.; Strommen, M. R. (1997) A thermodynamic approach using group contribution methods to model the partitioning of semivolatile organic compounds on atmospheric particulate matter. *Environ. Sci. Technol.* 31: 2805-2811.
- Jenkins, B. M.; Jones, A. D.; Turn, S. Q.; Williams, R. B. (1996) Particle concentrations, gas-particle partitioning, and species intercorrelations for polycyclic aromatic hydrocarbons (PAH) emitted during biomass burning. *Atmos. Environ.* 30: 3825-3835.
- Johansson, T. B.; Askelsson, R.; Johansson, S. A. E. (1970) X-ray analysis: elemental trace analysis at the 10^{-12} g level. *Nucl. Instrum. Methods* 84: 141-143.
- John, W.; Sethi, V. (1993) Threshold for resuspension by particle impaction. *Aerosol Sci. Technol.* 19: 69-79.
- John, W.; Wall, S. M.; Ondo, J. L. (1988) A new method for nitric acid and nitrate aerosol measurement using the dichotomous sampler. *Atmos. Environ.* 22: 1627-1635.

- John, W.; Wall, S. M.; Ondo, J. L.; Winklmayr, W. (1990) Modes in the size distributions of atmospheric inorganic aerosol. *Atmos. Environ. Part A* 24: 2349-2359.
- Johnson, M V.; Wexler, A. S. (1995) MS of individual aerosol particles. *Anal. Chem.* 67: 721A-726A.
- Johnson, R. L.; Shah, J. J.; Huntzicker, J. J. (1980) Analysis of organic, elemental, and carbonate carbon in ambient aerosols. In: Verner, S. S. *Sampling and analysis of toxic organics in the atmosphere: proceedings of a symposium*; August, 1979; Boulder, CO. Philadelphia, PA: American Society for Testing and Materials, pp. 111-119. [ASTM Special Technical Publication 721].
- Kamens, R. M.; Coe, D. L. (1997) A large gas-phase stripping device to investigate rates of PAH evaporation from airborne diesel soot particles. *Environ. Sci. Technol.* 31: 1830-1833.
- Kamens, R. M.; Jaoui, M. (2001) Modeling aerosol formation from α -pinene + NO_x in the presence of natural sunlight using gas phase kinetics and gas-particle partitioning theory. *Environ. Sci. Technol.* 35: 1394-1405.
- Kamens, R.; Odum, J.; Fan, Z.-H. (1995) Some observations on times to equilibrium for semivolatile polycyclic aromatic hydrocarbons. *Environ. Sci. Technol.* 29: 43-50.
- Kamens, R.; Jang, M.; Chien, C.-J.; Leach, K. (1999) Aerosol formation from the reaction of α -pinene and ozone using a gas-phase kinetics-aerosol partitioning model. *Environ. Sci. Technol.* 33: 1430-1438.
- Kao, A. S.; Friedlander, S. K. (1995) Temporal variations of particulate air pollution: a marker for free radical dosage and adverse health effects? *Inhalation Toxicol.* 7: 149-156.
- Kavouras, I. G.; Koutrakis, P. (2001) Use of polyurethane foam as the impaction substrate/collection medium in conventional inertial impactors. *Aerosol Sci. Technol.* 34: 46-56.
- Keeler, G. J.; Pirrone, N. (1996) Atmospheric transport and deposition of trace elements to Lake Erie from urban areas. *Water Sci. Technol.* 33: 259-265.
- Keeler, G. J.; Brachaczek, W. W.; Gorse, R. A., Jr.; Japar, S. M.; Pierson, W. R. (1988) Effect of ambient humidity on dichotomous sampler coarse/fine ratios. *Atmos. Environ.* 22: 1715-1720.
- Kegler, S. R.; Wilson, W. E.; Marcus, A. H. (2001) PM₁, intermodal (PM_{2.5-1}) mass, and the soil component of PM_{2.5} in Phoenix, AZ, 1995-96. *Aerosol Sci. Technol.* 35: 914-920.
- Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. *Atmos. Environ.* 33: 2907-2913.
- Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (2000) Size distribution and sources of aerosol in Launceston, Australia, during winter 1997. *J. Air Waste Manage. Assoc.* 50: 418-427.
- Khlystov, A.; Wyers, G. P.; Slanina, J. (1995) The steam-jet aerosol collector. *Atmos. Environ.* 29: 2229-2234.
- Kim, Y. J.; Boatman, J. F.; Gunter, R. L.; Wellman, D. L.; Wilkison, S. W. (1993) Vertical distribution of atmospheric aerosol size distribution over south-central New Mexico. *Atmos. Environ. Part A* 27: 1351-1362.
- Kim, B. M.; Lester, J.; Tisopulos, L.; Zeldin, M. D. (1999) Nitrate artifacts during PM_{2.5} sampling in the South Coast Air Basin of California. *J. Air Waste Manage. Assoc.* 49(special issue): PM142-153.
- Kim, B. M.; Cassmassi, J.; Hogo, H.; Zeldin, M. D. (2001a) Positive organic carbon artifacts on filter medium during PM_{2.5} sampling in the South Coast Air Basin. *Aerosol Sci. Technol.* 34: 35-41.
- Kim, S.; Jaques, P. A.; Chang, M. C.; Froines, J. R.; Sioutas, C. (2001b) Versatile aerosol concentration enrichment system (VACES) for simultaneous in vivo and in vitro evaluation of toxic effects of ultrafine, fine and coarse ambient particles Part I: development and laboratory characterization. *J. Aerosol Sci.* 32: 1281-1297.
- Kim, S.; Jaques, P. A.; Chang, M. C.; Barone, T.; Xiong, C.; Friedlander, S. K.; Sioutas, C. (2001c) Versatile aerosol concentration enrichment system (VACES) for simultaneous in vivo and in vitro evaluation of toxic effects of ultrafine, fine and coarse ambient particles Part II: field evaluation. *J. Aerosol Sci.* 32: 1299-1314.
- Kirchstetter, T. W.; Corrigan, C. E.; Novakov, T. (2001) Laboratory and field investigation of the adsorption of gaseous organic compounds onto quartz filters. *Atmos. Environ.* 35: 1663-1671.
- Kittelson, D. B. (1998) Engines and nanoparticles: a review. *J. Aerosol Sci.* 29: 575-588.
- Kittelson, D. B.; McKenzie, R.; Vermeersch, M.; Dorman, F.; Pui, D.; Linne, M.; Liu, B.; Whitby, K. (1978) Total sulfur aerosol concentration with an electrostatically pulsed flame photometric detector system. *Atmos. Environ.* 12: 105-111.
- Klouda, G. A.; Klinedinst, D. B.; Steel, E. B.; Benner, B. A., Jr.; Parish, H. J. (1996) Exploring a method to produce an urban dust particle filter standard. *J. Aerosol Sci.* 27(suppl. 1): S351-S352.
- Koistinen, K. J.; Kousa, A.; Tenhola, V.; Hänninen, O.; Jantunen, M. J.; Oglesby, L.; Kuenzli, N.; Georgoulis, L. (1999) Fine particle (PM_{2.5}) measurement methodology, quality assurance procedures and pilot results of the EXPOLIS study. *J. Air Waste Manage. Assoc.* 49: 1212-1220.
- Korhonen, P.; Kulmala, M.; Laaksonen, A.; Viisanen, Y.; McGraw, R.; Seinfeld, J. H. (1999) Ternary nucleation of H₂SO₄, NH₃, and H₂O in the atmosphere. *J. Geophys. Res.* 104: 26,349-26,353.

- Koutrakis, P.; Kelly, B. P. (1993) Equilibrium size of atmospheric aerosol sulfates as a function of particle acidity and ambient relative humidity. *J. Geophys. Res. [Atmos.]* 98: 7141-7147.
- Koutrakis, P.; Wolfson, J. M.; Slater, J. L.; Brauer, M.; Spengler, J. D.; Stevens, R. K.; Stone, C. L. (1988a) Evaluation of an annular denuder/filter pack system to collect acidic aerosols and gases. *Environ. Sci. Technol.* 22: 1463-1468.
- Koutrakis, P.; Wolfson, J. M.; Spengler, J. D. (1988b) An improved method for measuring aerosol strong acidity: results from a nine-month study in St. Louis, Missouri and Kingston, Tennessee. *Atmos. Environ.* 22: 157-162.
- Koutrakis, P.; Wolfson, J. M.; Spengler, J. D.; Stern, B.; Franklin, C. A. (1989) Equilibrium size of atmospheric aerosol sulfates as a function of the relative humidity. *J. Geophys. Res. [Atmos.]* 94: 6442-6448.
- Koutrakis, P.; Thompson, K. M.; Wolfson, J. M.; Spengler, J. D.; Keeler, G. J.; Slater, J. L. (1992) Determination of aerosol strong acidity losses due to interactions of collected particles: results from laboratory and field studies. *Atmos. Environ. Part A* 26: 987-995.
- Koutrakis, P.; Sioutas, C.; Ferguson, S. T.; Wolfson, J. M.; Mulik, J. D.; Burton, R. M. (1993) Development and evaluation of a glass honeycomb denuder/filter pack system to collect atmospheric gases and particles. *Environ. Sci. Technol.* 27: 2497-2501.
- Koutrakis, P.; Wang, P.-Y.; Wolfson, J. M.; Sioutas, C., inventors. (1996) Method and apparatus to measure particulate matter in gas. U.S. patent 5,571,945. November 5. Available: www.patents.ibm.com [1999, November 24].
- Krieger, M. S.; Hites, R. A. (1992) Diffusion denuder for the collection of semivolatile organic compounds. *Environ. Sci. Technol.* 26: 1551-1555.
- Kulmala, M.; Toivonen, A.; Mäkelä, J. M.; Laaksonen, A. (1998) Analysis of the growth of nucleation mode particles observed in Boreal forest. *Tellus* 50B: 449-462.
- Kulmala, M.; Pirjola, L.; Mäkelä, J. M. (2000) Stable sulphate clusters as a source of new atmospheric particles. *Nature (London, U. K.)* 404: 66-69.
- Kusko, B. H.; Cahill, T. A.; Eldred, R. A.; Matsuda, Y.; Miyake, H. (1989) Nondestructive analysis of total nonvolatile carbon by Forward Alpha Scattering Technique (FAST). *Aerosol Sci. Technol.* 10: 390-396.
- Lammel, G.; Novakov, T. (1995) Water nucleation properties of carbon black and diesel soot particles. *Atmos. Environ.* 29: 813-823.
- Landsberger, S. (1988) Improved methodology for the determination of the seven elemental tracer long-distance pollution signatures using thermal and epithermal neutron activation analysis. *Anal. Chem.* 60: 1842-1845.
- Landsberger, S.; Wu, D. (1993) Improvement of analytical sensitivities for the determination of antimony, arsenic, cadmium, indium, iodine, molybdenum, silicon and uranium in airborne particulate matter by epithermal neutron activation analysis. *J. Radioanal. Nucl. Chem.* 167: 219-225.
- Landsberger, S.; Wu, D.; Vermette, S. J.; Cizek, W. (1997) Intercomparison of IAEA airborne particulate matter reference material. *J. Radioanal. Nucl. Chem.* 215: 117-127.
- Lane, D. A.; Johnson, N. D.; Barton, S. C.; Thomas, G. H. S.; Schroeder, W. H. (1988) Development and evaluation of a novel gas and particle sampler for semivolatile chlorinated organic compounds in ambient air. *Environ. Sci. Technol.* 22: 941-947.
- Lavanchy, V. M. H.; Gäggeler, H. W.; Nyeki, S.; Baltensperger, U. (1999) Elemental carbon (EC) and black carbon (BC) measurements with a thermal method and an aethalometer at the high-alpine research station Jungfraujoch. *Atmos. Environ.* 33: 2759-2769.
- Lawless, P. A.; Rodes, C. E.; Evans, G.; Sheldon, L.; Creason, J. (2001) Aerosol concentrations during the 1999 Fresno Exposure Studies as functions of size, season and meteorology. *Aerosol Sci. Technol.* 34: 66-74.
- Lawrence, J. E.; Koutrakis, P. (1994) Measurement of atmospheric formic and acetic acids: methods evaluation and results from field studies. *Environ. Sci. Technol.* 28: 957-964.
- Lawrence, J.; Koutrakis, P. (1996a) Measurement and speciation of gas and particulate phase organic acidity in an urban environment: 1. analytical. *J. Geophys. Res. [Atmos.]* 101: 9159-9169.
- Lawrence, J.; Koutrakis, P. (1996b) Measurement and speciation of gas and particulate phase organic acidity in an urban environment: 2. speciation. *J. Geophys. Res. [Atmos.]* 101: 9171-9184.
- Lawson, D. R. (1990) The Southern California Air Quality Study. *J. Air Waste Manage. Assoc.* 40: 156-165.
- Lazrus, A. L.; Haagenson, P. L.; Kok, G. L.; Huebert, B. J.; Kreitzberg, C. W.; Likens, G. E.; Mohnen, V. A.; Wilson, W. E.; Winchester, J. W. (1983) Acidity in air and water in a case of warm frontal precipitation. *Atmos. Environ.* 17: 581-591.
- Leach, K. B.; Kamens, R. M.; Strommen, M. R.; Jang, M. (1999) Partitioning of semivolatile organic compounds in the presence of a secondary organic aerosol in a controlled atmosphere. *J. Atmos. Chem.* 33: 241-264.

- Lee, W.-M. G.; Shih, P.-M.; Wang, C.-S. (1997) The influence of relative humidity on the size of atmospheric aerosol. *J. Environ. Sci. Health Part A: Environ. Sci. Eng. Toxic Hazard. Subst. Control* 32: 1085-1097.
- Lewis, C. W.; Macias, E. S. (1980) Composition of size-fractionated aerosol in Charleston, West Virginia. *Atmos. Environ.* 14: 185-194.
- Lewtas, J.; Pang, Y.; Booth, D.; Reimer, S.; Eatough, D. J.; Gundel, L. A. (2001) Comparison of sampling methods for semi-volatile organic carbon associated with PM_{2.5}. *Aerosol Sci. Technol.* 34: 9-22.
- Li, W.; Montassier, N.; Hopke, P. K. (1992) A system to measure the hygroscopicity of aerosol particles. *Aerosol Sci. Technol.* 17: 25-35.
- Liang, Z.; Chan, C. K. (1997) A fast technique for measuring water activity of atmospheric aerosols. *Aerosol Sci. Technol.* 26: 255-268.
- Liang, C.; Pankow, J. F.; Odum, J. R.; Seinfeld, J. H. (1997) Gas/particle partitioning of semivolatile organic compounds to model inorganic, organic, and ambient smog aerosols. *Environ. Sci. Technol.* 31: 3086-3092.
- Lin, J. J.; Noll, K. E.; Holsen, T. M. (1994) Dry deposition velocities as a function of particle size in the ambient atmosphere. *Aerosol Sci. Technol.* 20: 239-252.
- Lindskog, A.; Brorström-Lundén, E.; Alheim, I.; Hagen, I. (1987) Chemical transformation of PAH on airborne particles by exposure to NO₂ during sampling: a comparison between two filter media. *Sci. Total Environ.* 61: 51-57.
- Liousse, C.; Cachier, H.; Jennings, S. G. (1993) Optical and thermal measurements of black carbon aerosol content in different environments: variation of the specific attenuation cross-section, sigma σ . *Atmos. Environ. Part A* 27: 1203-1211.
- Long, R. W.; Smith, R.; Smith, S.; Eatough, N. L.; Mangelson, N. F.; Eatough, D. J.; Pope, C. A., III; Wilson, W. E. (2002) Sources of fine particulate material along the Wasatch Front. *Ener. Fuels* 16: 282-293.
- Loo, B. W.; Cork, C. P. (1988) Development of high efficiency virtual impactors. *Aerosol Sci. Technol.* 9: 167-176.
- Loo, B. W.; Jaklevic, J. M.; Goulding, F. S. (1976) Dichotomous virtual impactors for large scale monitoring of airborne particulate matter. In: Liu, B. Y. H., ed. *Fine particles: aerosol generation, measurement, sampling, and analysis*. New York, NY: Academic Press; pp. 311-350.
- Loo, B. W.; French, W. R.; Gatti, R. C.; Goulding, F. S.; Jaklevic, J. M.; Llacer, J.; Thompson, A. C. (1978) Large-scale measurement of airborne particulate sulfur. *Atmos. Environ.* 12: 759-771.
- Lundgren, D. A.; Burton, R. M. (1995) Effect of particle size distribution on the cut point between fine and coarse ambient mass fractions. In: Phalen, R. F.; Bates, D. V., eds. *Proceedings of the colloquium on particulate air pollution and human mortality and morbidity; January 1994; Irvine, CA. Inhalation Toxicol.* 7: 131-148.
- Lundgren, D. A.; Hausknecht, B. J. (1982) Ambient aerosol size distribution determination using a mobile wide range aerosol classifier. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; EPA contract no. CR-808606-01-1.
- Lundgren, D. A.; Hausknecht, B. J.; Burton, R. M. (1984) Large particle size distribution in five U.S. cities and the effect on a new ambient particulate matter standard (PM₁₀). *Aerosol Sci. Technol.* 7: 467-473.
- Lynch, A. J.; McQuaker, N. R.; Brown, D. F. (1980) ICP/AES analysis and the composition of airborne and soil materials in the vicinity of a lead/zinc smelter complex. *J. Air Pollut. Control Assoc.* 30: 257-260.
- Mader, B. T.; Flagan, R. C.; Seinfeld, J. H. (2001) Sampling atmospheric carbonaceous aerosols using a particle trap impactor/denuder sampler. *Environ. Sci. Technol.* 35: 4857-4867.
- Mahanama, K. R. R.; Gundel, L. A.; Daisey, J. M. (1994) Selective fluorescence detection of polycyclic aromatic hydrocarbons in environmental tobacco smoke and other airborne particles. *Int. J. Environ. Anal. Chem.* 56: 289-309.
- Mäkelä, J. M.; Aalto, P.; Jokinen, V.; Pohja, T.; Nissinen, A.; Palmroth, S.; Markkanen, T.; Seitsonen, K.; Lihavainen, H.; Kulmala, M. (1997) Observations of ultrafine aerosol particle formation and growth in boreal forest. *Geophys. Res. Lett.* 24: 1219-1222.
- Malm, W. C.; Sisler, J. F.; Huffman, D.; Eldred, R. A.; Cahill, T. A. (1994) Spatial and seasonal trends in particle concentration and optical extinction in the United States. *J. Geophys. Res. [Atmos.]* 99: 1347-1370.
- Mamane, Y.; Willis, R.; Conner, T. (2001) Evaluation of computer-controlled scanning electron microscopy applied to an ambient urban aerosol sample. *Aerosol Sci. Technol.* 34: 97-107.
- Marple, V. A.; Chien, C. M. (1980) Virtual impactors: a theoretical study. *Environ. Sci. Technol.* 14: 976-985.
- Marple, V. A.; Olson, B. A. (1995) A high volume PM₁₀/PM_{2.5}/PM_{1.0} trichotomous sampler. In: *Particulate matter: health and regulatory issues: proceedings of an international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; pp. 237-261. (A&WMA publication VIP-49).*
- Marple, V. A.; Liu, B. Y. H.; Burton, R. M. (1990) High-volume impactor for sampling fine and coarse particles. *J. Air Waste Manage. Assoc.* 40: 762-767.

- Marple, V. A.; Rubow, K. L.; Behm, S. M. (1991) A microorifice uniform deposit impactor (MOUDI): description, calibration, and use. *Aerosol Sci. Technol.* 14: 434-446.
- Martin, L. R. (1994) Aqueous sulfur(IV) oxidation revisited. In: Nriagu, J. O.; Simmons, M. S., eds. *Environmental oxidants*. New York, NY: John Wiley & Sons, Inc.; pp. 221-268. (Nriagu, J. O., ed. *Advances in environmental science and technology*: v. 28).
- Masuda, H.; Nakasita, S. (1988) Classification performance of a rectangular jet virtual impactor-effects of nozzle width ratio of collection nozzle to acceleration nozzle. *J. Aerosol Sci.* 19: 243-252.
- Masuda, H.; Hochrainer, D.; Stöber, W. (1979) An improved virtual impactor for particle classification and generation of test aerosols with narrow size distributions. *J. Aerosol Sci.* 10: 275-287.
- McGovern, F. M. (1999) An analysis of condensation nuclei levels at Mace Head, Ireland. *Atmos. Environ.* 33: 1711-1723.
- McGovern, F. M.; Jennings, S. G.; O'Connor, T. C.; Simmonds, P. G. (1996) Aerosol and trace gas measurements during the Mace Head experiment. *Atmos. Environ.* 30: 3891-3902.
- McInnes, L. M.; Quinn, P. K.; Covert, D. S.; Anderson, T. L. (1996) Gravimetric analysis, ionic composition, and associated water mass of the marine aerosol. *Atmos. Environ.* 30: 869-884.
- McMurry, P. H. (2000) A review of atmospheric aerosol measurements. *Atmos. Environ.* 34: 1959-1999.
- McMurry, P. H.; Friedlander, S. K. (1979) New particle formation in the presence of an aerosol. *Atmos. Environ.* 13: 1635-1651.
- McMurry, P. H.; Stolzenburg, M. R. (1989) On the sensitivity of particle size to relative humidity for Los Angeles aerosols. *Atmos. Environ.* 23: 497-507.
- McMurry, P. H.; Wilson, J. C. (1982) Growth laws for the formation of secondary ambient aerosols: implications for chemical conversion mechanisms. *Atmos. Environ.* 16: 121-134.
- McMurry, P. H.; Wilson, J. C. (1983) Droplet phase (heterogeneous) and gas phase (homogeneous) contributions to secondary ambient aerosol formation as functions of relative humidity. *J. Geophys. Res.* 88: 5101-5108.
- McMurry, P. H.; Rader, D. J.; Stith, J. L. (1981) Studies of aerosol formation in power plant plumes--I. growth laws for secondary aerosols in power plant plumes: implications for chemical conversion mechanisms. In: White, W. H.; Moore, D. J.; Lodge, J. P., Jr., eds. *Plumes and visibility: measurements and model components: proceedings of the symposium; November 1980; Grand Canyon National Park, AZ*. *Atmos. Environ.* 15: 2315-2327.
- McMurry, P. H.; Woo, K. S.; Weber, R.; Chen, D.-R.; Pui, D. Y. H. (2000) Size distributions of 3 to 10 nm atmospheric particles: implications for nucleation mechanisms. *Philos. Trans. R. Soc. London Ser. A* 358: 2625-2642.
- McQuaker, N. R.; Kluckner, P. D.; Chang, G. N. (1979) Calibration of an inductively coupled plasma-atomic emission spectrometer for the analysis of environmental materials. *Anal. Chem.* 51: 888-895.
- Meyer, A. S., Jr.; Boyd, C. M. (1959) Determination of water by titration with coulometrically generated Karl Fischer reagent. *Anal. Chem.* 31: 215-219.
- Meyer, M. B.; Rupprecht, E. G. (1996) Particulate matter sampling methods: the importance of standardization. *J. Aerosol. Sci.* 27(suppl. 1): S349-S350.
- Meyer, M. B.; Rupprecht, E.; Patashnick, H. (1995) Considerations for the sampling and measurement of ambient particulate matter. In: *Particulate matter: health and regulatory issues: proceedings of an international specialty conference; April; Pittsburgh, PA*. Pittsburgh, PA: Air & Waste Management Association; pp. 204-210. (A&WMA publication VIP-49).
- Meyer, M. B.; Patashnick, H.; Ambs, J. L.; Rupprecht, E. (2000) Development of a sample equilibration system for the TEOM continuous PM monitor. *J. Air Waste Manage. Assoc.* 50: 1345-1349.
- Middlebrook, A. M.; Murphy, D. M.; Thomson, D. S. (1998) Observations of organic material in individual marine particles at Cape Grim during the first aerosol characterization experiment (ACE 1). *J. Geophys. Res.* 103: 16475-16483.
- Middlebrook, A. M.; Murphy, D. M.; Lee, S.-H.; Thomson, D. S.; Prather, K. A.; Wenzel, R. J.; Liu, D.-Y.; Phares, D. J.; Rhoads, K. P.; Wexler, A. S.; Johnston, M. V.; Jimenez, J. L.; Jayne, J. T.; Worsnop, D. R.; Yourshaw, I.; Seinfeld, J. H.; Flagan, R. C. (2003) A comparison of particle mass spectrometers during the 1999 Atlanta Supersites project. *J. Geophys. Res.* [Atmos.] 108(D7): 10.1029/2001JD000660.
- Mignacca, D.; Stubbs, K. (1999) Effects of equilibration temperature on PM₁₀ concentrations from the TEOM method in the Lower Fraser Valley. *J. Air Waste Manage. Assoc.* 49: 1250-1254.
- Miller, F. J.; Gardner, D. E.; Graham, J. A.; Lee, R. E., Jr.; Wilson, W. E.; Bachmann, J. D. (1979) Size considerations for establishing a standard for inhalable particles. *J. Air Pollut. Control Assoc.* 29: 610-615.

- Misra, C.; Geller, M. D.; Shah, P.; Sioutas, C.; Solomon, P. A. (2001) Development and evaluation of a continuous coarse (PM₁₀-PM_{2.5}) particle monitor. *J. Air Waste Manage. Assoc.* 51: 1309-1317.
- Modey, W. K.; Pang, Y.; Eatough, N. L.; Eatough, D. J. (2001) Fine particulate (PM_{2.5}) composition in Atlanta, USA: assessment of the particle concentrator-Brigham Young University organic sampling system, PC-BOSS, during the EPA supersite study. *Atmos. Environ.* 35: 6493-6502.
- Moosmüller, H.; Arnott, W. P.; Rogers, C. F. (1998) Photoacoustic and filter measurements related to aerosol light absorption during the Northern Front Range Air Quality Study (Colorado 1996/1997). *J. Geophys. Res.* 103: 28,149-28,157.
- Morawska, L.; Bofinger, N. D.; Kocis, L.; Nwankwoala, A. (1998) Submicrometer and supermicrometer particles from diesel vehicle emissions. *Environ. Sci. Technol.* 32: 2033-2042.
- Morio, L. A.; Hooper, K. A.; Brittingham, J.; Li, T.-H.; Gordon, R. E.; Turpin, B. J.; Laskin, D. L. (2001) Tissue injury following inhalation of fine particulate matter and hydrogen peroxide is associated with altered production of inflammatory mediators and antioxidants by alveolar macrophages. *Toxicol. Appl. Pharmacol.* 177: 188-199.
- Mueller, P. K.; Collins, J. F. (1980) Development of a particulate sulfate analyzer. Palo Alto, CA: Electric Power Research Institute; report no. P-1382F.
- Mueller, P. K.; Fung, K. K.; Heisler, S. L.; Grosjean, D.; Hidy, G. M. (1982) Atmospheric particulate carbon observations in urban and rural areas of the United States. In: Wolff, G. T.; Klimisch, R. L., eds. *Particulate carbon: atmospheric life cycle: proceedings of an international symposium; October 1980; Warren, MI.* New York, NY: Plenum Press; pp. 343-370.
- Mueller, P. K.; Hansen, D. A.; Watson, J. G., Jr. (1986) The subregional cooperative electric utility, Department of Defense, National Park Service, and EPA study (SCENES) on visibility: an overview. Palo Alto, CA: Electric Power Research Institute; report no. EA-4664-SR.
- Mukerjee, S.; Ellenson, W. D.; Lewis, R. G.; Stevens, R. K.; Somerville, M. C.; Shadwick, D. S.; Willis, R. D. (1997) An environmental scoping study in the lower Rio Grande Valley of Texas—III. residential microenvironmental monitoring for air, house dust, and soil. *Environ. Int.* 23: 657-673.
- Munksgaard, N. C.; Parry, D. L. (1998) Lead isotope ratios determined by ICP-MS: monitoring of mining-derived metal particulates in atmospheric fallout, Northern Territory, Australia. *Sci. Total Environ.* 217: 113-126.
- Murphy, D. M.; Thomson, D. S. (1997) Chemical composition of single aerosol particles at Idaho Hill: negative ion measurements. *J. Geophys. Res.* 102: 6353-6368.
- Nadkarni, R. A. (1975) Multielement analysis of coal and coal fly ash standards by instrumental neutron activation analysis. *Radiochem. Radioanal. Lett.* 21: 161-176.
- Nam, S.-H.; Masamba, W. R. L.; Montaser, A. (1993) Investigation of helium inductively coupled plasma-mass spectrometry for the detection of metals and nonmetals in aqueous solutions. *Anal. Chem.* 65: 2784-2790.
- NARSTO. (2003) Fine particle assessment (review draft). Oak Ridge, TN: U.S. Department of Energy, Oak Ridge National Laboratory, North American Research Strategy for Tropospheric Ozone, Quality Systems Science Center. Available: <http://www.cgenv.com/Narsto/> (22 April 2003).
- National Research Council. (1993) *Protecting visibility in national parks and wilderness areas.* Washington, DC: National Academy Press. 3v.
- Nelson, M. P.; Zugates, C. T.; Treado, P. J.; Casucio, G. S.; Exline, D. L.; Schlaegle, S. F. (2000) Combining Raman chemical imaging scanning electron microscopy (SEM) to characterize ambient fine particulate. *Aerosol Sci. Technol.* 34: 108-117.
- Neubauer, K. R.; Johnston, M. V.; Wexler, A. S. (1998) Humidity effects on the mass spectra of single aerosol particles. *Atmos. Environ.* 32: 2521-2529.
- Nham, T. T. (1996) Cool plasma technique for the ultra-trace level determination of ⁵⁶Fe, ⁴⁰Ca, ³⁹K, ⁷Li, ²⁴Mg and ²³Na for semi-conductor applications. Mulgrave, Victoria, Australia: Varian Australia Pty Ltd; ICP-MS instruments at work no. ICP-MS-7. Available: <http://www.varianinc.com/cgi-bin/nav?varinc/docs/osi/icpms/atwork/icpms7&cid=680052> [18 January, 2002].
- Noble, C. A.; Prather, K. A. (1996) Real-time measurement of correlated size and composition profiles of individual atmospheric aerosol particles. *Environ. Sci. Technol.* 30: 2667-2680.
- Noble, C. A.; Prather, K. A. (1998) Single particle characterization of albuterol metered dose inhaler aerosol in near real-time. *Aerosol Sci. Technol.* 29: 294-306.
- Noble, C. A.; Vanderpool, R. W.; Peters, T. M.; McElroy, F. F.; Gemmill, D. B.; Wiener, R. W. (2001) Federal reference and equivalent methods for measuring fine particulate matter. *Aerosol Sci. Technol.* 34: 457-464.

- Noël, M. A.; Topart, P. A. (1994) High-frequency impedance analysis of quartz crystal microbalances. 1. General considerations. *Anal. Chem.* 66: 484-491.
- Novakov, T.; Corrigan, C. E.; Penner, J. E.; Chuang, C. C.; Rosario, O.; Mayol Bracero, O. L. (1997) Organic aerosols in the Caribbean trade winds: a natural source? *J. Geophys. Res. [Atmos.]* 102: 21307-21313.
- O'Dowd, C. D.; Aalto, P.; Hämeri, K.; Kulmala, M.; Hoffmann, T. (2002) Atmospheric particles from organic vapours. *Nature (London, U. K.)* 416: 497-498.
- Obeidi, F. D.; Eatough, D. J. (2002) Continuous measurement of semi-volatile fine particulate mass in Provo, UT. *Aerosol Sci. Technol.* 36: 191-203.
- Obeidi, F.; Eatough, N. L.; Eatough, D. J. (2002) Use of the RAMS to measure semivolatile fine particulate matter at Riverside and Bakersfield, California. *Aerosol Sci. Technol.* 36: 204-216.
- Odum, J. R.; Yu, J.; Kamens, R. M. (1994) Modeling the mass transfer of semivolatile organics in combustion aerosols. *Environ. Sci. Technol.* 28: 2278-2285.
- Ohta, S.; Okita, T. (1990) A chemical characterization of atmospheric aerosol in Sapporo. *Atmos. Environ. Part A* 24A: 815-822.
- Ohta, S.; Hori, M.; Yamagata, S.; Murao, N. (1998) Chemical characterization of atmospheric fine particles in Sapporo with determination of water content. *Atmos. Environ.* 32: 1021-1025.
- Olin, J. G.; Sem, G. J. (1971) Piezoelectric microbalance for monitoring the mass concentration of suspended particles. *Atmos. Environ.* 5: 653-668.
- Olmez, I. (1989) Trace element signatures in groundwater pollution. In: Watson, J. G., ed. *Receptor models in air resources management: an international specialty conference; February 1988; San Francisco, CA.* Pittsburgh, PA: Air & Waste Management Association; pp. 3-11. (APCA transactions series: no. 14).
- Ondov, J. M. (1996) Particulate tracers for source attribution: potential for application to California's San Joaquin Valley. *J. Aerosol Sci.* 27(suppl. 1): S687-S688.
- Ondov, J. M.; Divita, F., Jr. (1993) Size spectra for trace elements in urban aerosol particles by instrumental neutron activation analysis. *J. Radioanal. Nucl. Chem.* 167: 247-258.
- Ondov, J. M.; Kelly, W. R.; Holland, J. Z.; Lin, Z. C.; Wight, S. A. (1992) Tracing fly ash emitted from a coal-fired power plant with enriched rare-earth isotopes: an urban scale test. *Atmos. Environ. Part B* 26B: 453-462.
- Ono, D. M.; Hardebeck, E.; Parker, J.; Cox, B. G. (2000) Systematic biases in measured PM₁₀ values with U.S. Environmental Protection Agency-approved samplers at Owens Lake, California. *J. Air Waste Manage. Assoc.* 50: 1144-1156.
- Pak, S. S.; Liu, B. Y. H.; Rubow, K. L. (1992) Effect of coating thickness on particle bounce in inertial impactors. *Aerosol Sci. Technol.* 16: 141-150.
- Pang, Y.; Ren, Y.; Obeidi, F.; Hastings, R.; Eatough, D. J.; Wilson, W. E. (2001) Semi-volatile species in PM_{2.5}: comparison of integrated and continuous samplers for PM_{2.5} research or monitoring. *J. Air Waste Manage. Assoc.* 51: 25-36.
- Pang, Y.; Eatough, N. L.; Wilson, J.; Eatough, D. J. (2002a) Effect of semivolatile material on PM_{2.5} measurement by the PM_{2.5} federal reference method sampler at Bakersfield, California. *Aerosol Sci. Technol.* 36: 289-299.
- Pang, Y.; Eatough, N. L.; Eatough, D. J. (2002b) PM_{2.5} semivolatile organic material at Riverside, CA: implications for the PM_{2.5} federal reference method sampler. *Aerosol Sci. Technol.* 36: 277-288.
- Pankow, J. F. (1987) Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere. *Atmos. Environ.* 21: 2275-2283.
- Pankow, J. F. (1994a) An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol. *Atmos. Environ.* 28: 189-193.
- Pankow, J. F. (1994b) An absorption model of gas/particle partitioning of organic compounds in the atmosphere. *Atmos. Environ.* 28: 185-188.
- Pankow, J. F.; Bidleman, T. F. (1991) Effects of temperature, *TSP* and per cent non-exchangeable material in determining the gas-particle partitioning of organic compounds. *Atmos. Environ. Part A* 25: 2241-2249.
- Pankow, J. F.; Storey, J. M. E.; Yamasaki, H. (1993) Effects of relative humidity on gas/particle partitioning of semivolatile organic compounds to urban particulate matter. *Environ. Sci. Technol.* 27: 2220-2226.
- Parmar, S. S.; Grosjean, D. (1990) Laboratory tests of KI and alkaline annular denuders. *Atmos. Environ. Part A* 24: 2695-2698.
- Patashnick, H.; Rupprecht, G.; Ambs, J. L.; Meyer, M. B. (2001) Development of a reference standard for particulate matter mass in ambient air. *Aerosol Sci. Technol.* 34: 42-45.
- Pellizzari, E. D.; Krost, K. J. (1984) Chemical transformations during ambient air sampling for organic vapors. *Anal. Chem.* 56: 1813-1819.

- Peng, C.; Chan, C. K. (2001) The water cycles of water-soluble organic salts of atmospheric importance. *Atmos. Environ.* 35: 1183-1192.
- Penner, J. E.; Novakov, T. (1996) Carbonaceous particles in the atmosphere: a historical perspective to the fifth international conference on carbonaceous particles in the atmosphere. *J. Geophys. Res. [Atmos.]* 101: 19,373-19,378.
- Perry, K. D.; Hobbs, P. V. (1994) Further evidence for particle nucleation in clear air adjacent to marine cumulus clouds. *J. Geophys. Res. [Atmos.]* 99: 22,803-22,818.
- Perry, K. D.; Cahill, T. A.; Schnell, R. C.; Harris, J. M. (1999) Long-range transport of anthropogenic aerosols to the National Oceanic and Atmospheric Administration baseline station at Mauna Loa Observatory, Hawaii. *J. Geophys. Res. Atmos.* 104: 18,521-18,533.
- Peters, A.; Liu, E.; Verrier, R. L.; Schwartz, J.; Gold, D. R.; Mittleman, M.; Baliff, J.; Oh, J. A.; Allen, G.; Monahan, K.; Dockery, D. W. (2000) Air pollution and incidence of cardiac arrhythmia. *Epidemiology* 11: 11-17.
- Peters, T. M.; Vanderpool, R. W.; Wiener, R. W. (2001a) Design and calibration of the EPA PM_{2.5} well impactor ninety-six (WINS). *Aerosol Sci. Technol.* 34: 389-397.
- Peters, T. M.; Norris, G. A.; Vanderpool, R. W.; Gemmill, D. B.; Wiener, R. W.; Murdoch, R. W.; McElroy, F. F.; Pitchford, M. (2001b) Field performance of PM_{2.5} federal reference method samplers. *Aerosol Sci. Technol.* 34: 433-443.
- Peters, T. M.; Gussman, R. A.; Kenny, L. C.; Vanderpool, R. W. (2001c) Evaluation of PM_{2.5} size selectors used in speciation samplers. *Aerosol Sci. Technol.* 34: 422-429.
- Petzold, A.; Niessner, R. (1996) Photoacoustic soot sensor for in-situ black carbon monitoring. *Appl. Phys. B: Lasers Opt.* 63: 191-197.
- Petzold, A.; Kopp, C.; Niessner, R. (1997) The dependence of the specific attenuation cross-section on black carbon mass fraction and particle size. *Atmos. Environ.* 31: 661-672.
- Pilinis, C.; Seinfeld, J. H.; Grosjean, D. (1989) Water content of atmospheric aerosols. *Atmos. Environ.* 23: 1601-1606.
- Piorek, S. (1994) Modern, PC based, high resolution portable EDXRF analyzer offers laboratory performance for field, in-situ analysis of environmental contaminants. *Nucl. Instr. Meth. Phys. Res. A* 353: 528-533.
- Pirrone, N.; Keeler, G. J. (1996) The Rouge River watershed pollution by trace elements: atmospheric depositions and emission sources. *Water Sci. Technol.* 33: 267-275.
- Plantz, M. (1996) Common molecular ion interferences in ICP-MS. Mulgrave, Victoria, Australia: Varian Australia Pty Ltd; ICP-MS instruments at work no. ICP-MS-6. Available: <http://www.varianinc.com/cgi-bin/nav?varinc/docs/osi/icpms/atwork/index&cid=680052> [18 January, 2002].
- Poor, N.; Clark, T.; Nye, L.; Tamanini, T.; Tate, K.; Stevens, R.; Atkeson, T. (2002) Field performance of dichotomous sequential PM air samplers. *Atmos. Environ.* 36: 3289-3298.
- Pope, C. A., III; Thun, M. J.; Namboodiri, M. M.; Dockery, D. W.; Evans, J. S.; Speizer, F. E.; Heath, C. W., Jr. (1995) Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *Am. J. Respir. Crit. Care Med.* 151: 669-674.
- Pope, C. A., III; Hill, R. W.; Villegas, G. M. (1999) Particulate air pollution and daily mortality on Utah's Wasatch Front. *Environ. Health Perspect.* 107: 567-573.
- Potts, P. J.; Webb, P. C. (1992) X-ray fluorescence spectrometry. *J. Geochem. Explor.* 44: 251-296.
- Radke, L. F.; Hobbs, P. V. (1991) Humidity and particle fields around some small cumulus clouds. *J. Atmos. Sci.* 48: 1190-1193.
- Raes, F.; Van Dingenen, R.; Cuevas, E.; Van Velthoven, P. F. J.; Prospero, J. M. (1997) Observations of aerosols in the free troposphere and marine boundary layer of the subtropical Northeast Atlantic: discussion of processes determining their size distribution. *J. Geophys. Res. [Atmos.]* 102: 21315-21328.
- Ranweiler, L. E.; Moyers, J. L. (1974) Atomic absorption procedure for analysis of metals in atmospheric particulate matter. *Environ. Sci. Technol.* 8: 152-156.
- Rao, A. K.; Whitby, K. T. (1977) Nonideal collection characteristics of single stage and cascade impactors. *Am. Ind. Hyg. Assoc. J.* 38: 174-177.
- Rao, A. K.; Whitby, K. T. (1978) Non-ideal collection characteristics of inertial impactors—I. single-stage impactors and solid particles. *J. Aerosol Sci.* 9: 77-86.
- Ravenhall, D. G.; Forney, L. J.; Jazayeri, M. (1978) Aerosol sizing with a slotted virtual impactor. *J. Colloid Interface Sci.* 65: 108-117.

- Reid, J. S.; Cahill, T. A.; Wakabayashi, P. H.; Dunlap, M. R. (1994) Geometric/aerodynamic equivalent diameter ratios of ash aggregate aerosols collected in burning Kuwaiti well fields. *Atmos. Environ.* 28: 2227-2234.
- Reilly, P. T. A.; Gieray, R. A.; Whitten, W. B.; Ramsey, J. M. (1998) Real-time characterization of the organic composition and size of individual diesel engine smoke particles. *Environ. Sci. Technol.* 32: 2672-2679.
- Reist, P. C. (1984) Introduction to aerosol science. New York, NY: Macmillan Publishing Company.
- Reist, P. C. (1993) Introduction to aerosol science. New York, NY: McGraw-Hill.
- Risse, U.; Flammenkamp, E.; Kettrup, A. (1996) Determination of aromatic hydrocarbons in air using diffusion denuders. *Fresenius' J. Anal. Chem.* 356: 390-395.
- Roberts, R. A.; Corkill, J. (1998) Grass seed field smoke and its impact on respiratory health. *J. Environ. Health* 60: 10-16.
- Rounds, S. A.; Pankow, J. F. (1990) Application of a radial diffusion model to describe gas/particle sorption kinetics. *Environ. Sci. Technol.* 24: 1378-1386.
- Rounds, S. A.; Tiffany, B. A.; Pankow, J. F. (1993) Description of gas/particle sorption kinetics with an intraparticle diffusion model: desorption experiments. *Environ. Sci. Technol.* 27: 366-377.
- Rumburg, B.; Alldredge, R.; Claiborn, C. (2001) Statistical distributions of particulate matter and the error associated with sampling frequency. *Atmos. Environ.* 35: 2907-2920.
- Rupprecht, G.; Patashnick, H.; Beeson, D. E.; Green, R. N.; Meyer, M. B. (1995) A new automated monitor for the measurement of particulate carbon in the atmosphere. In: Particulate matter: health and regulatory issues: proceedings of an international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; pp. 309-319. (A&WMA publication VIP-49).
- Sakata, K.; Kawabata, K. (1994) Reduction of fundamental polyatomic ions in inductively coupled plasma mass spectrometry. *Spectrochim. Acta* 49B: 1027-1038.
- Sarangapani, R.; Wexler, A. S. (1996) Growth and neutralization of sulfate aerosols in human airways. *J. Appl. Physiol.* 81: 480-490.
- Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. *J. Atmos. Chem.* 24: 57-109.
- Saxena, P.; Hildemann, L. M.; McMurry, P. H.; Seinfeld, J. H. (1995) Organics alter hygroscopic behavior of atmospheric particles. *J. Geophys. Res. [Atmos.]* 100: 18,755-18,770.
- Schmid, H.; Laskus, L.; Abraham, J. H.; Baltensperger, U.; Lavanchy, V.; Bizjak, M.; Burba, P.; Cachier, H.; Crow, D.; Chow, J.; Gnauk, T.; Even, A.; Ten Brink, H. M.; Giesen, K.-P.; Hitzengerger, R.; Hueglin, C.; Maenhaut, W.; Pio, C.; Carvalho, A.; Putaud, J.-P.; Toom-Sauntry, D.; Puxbaum, H. (2001) Results of the "carbon conference" international aerosol carbon round robin test stage I. *Atmos. Environ.* 35: 2111-2121.
- Schröder, F.; Ström, J. (1997) Aircraft measurements of submicrometer aerosol particles (> 7 nm) in the midaltitude free troposphere and tropopause region. *Atmos. Res.* 44: 333-356.
- Schwartz, S. E. (1984) Gas-aqueous reactions of sulfur and nitrogen oxides in liquid-water clouds. In: Calvert, J. G., ed. *SO₂, NO and NO₂ oxidation mechanisms: atmospheric considerations*. Boston, MA: Butterworth Publishers; pp. 173-208. (Teasley, J. I., ed. *Acid precipitation series*: v. 3).
- Schwartz, S. E. (1986) Chemical conversions in clouds. In: Lee, S. D.; Schneider, T.; Grant, L. D.; Verkerk, P., eds. *Aerosols: research, risk assessment and control strategies, proceedings of the second U.S.-Dutch international symposium*; May 1985; Williamsburg, VA. Chelsea, MI: Lewis Publishers, Inc.; pp. 349-375.
- Schwartz, J.; Dockery, D. W.; Neas, L. M. (1996) Is daily mortality associated specifically with fine particles? *J. Air Waste Manage. Assoc.* 46: 927-939.
- Seinfeld, J. H.; Pandis, S. N. (1998) *Atmospheric chemistry and physics: from air pollution to climate change*. New York, NY: John Wiley & Sons, Inc.
- Sem, G. J.; Tsurubayashi, K.; Homma, K. (1977) Performance of the piezoelectric microbalance respirable aerosol sensor. *Am. Ind. Hyg. Assoc. J.* 38: 580-588.
- Shah, J. J.; Rau, J. A. (1990) Carbonaceous methods comparison study: interlaboratory round robin interpretation results. Sacramento, CA: California Air Resources Board. Available: <http://www.arb.ca.gov/research/apr/past/pm.htm> (21 March 2002).
- Silva, P. J.; Prather, K. A. (1997) On-line characterization of individual particles from automobile emissions. *Environ. Sci. Technol.* 31: 3074-3080.
- Simon, P. K.; Dasgupta, P. K. (1995) Continuous automated measurement of the soluble fraction of atmospheric particulate matter. *Anal. Chem.* 67: 71-78.
- Sioutas, C.; Koutrakis, P. (1996) Inertial separation of ultrafine particles using a condensational growth/virtual impaction system. *Aerosol Sci. Technol.* 25: 424-436.

- Sioutas, C.; Koutrakis, P.; Burton, R. M. (1994a) Development of a low cutpoint slit virtual impactor for sampling ambient fine particles. *J. Aerosol Sci.* 25: 1321-1330.
- Sioutas, C.; Koutrakis, P.; Burton, R. M. (1994b) A high-volume small cutpoint virtual impactor for separation of atmospheric particulate from gaseous pollutants. Part. *Sci. Technol.* 12: 207-221.
- Sioutas, C.; Koutrakis, P.; Olson, B. A. (1994c) Development and evaluation of a low cutpoint virtual impactor. *Aerosol Sci. Technol.* 21: 223-235.
- Sioutas, C.; Koutrakis, P.; Wolfson, J. M. (1994d) Particle losses in glass honeycomb denuder samplers. *Aerosol Sci. Technol.* 21: 137-148.
- Sioutas, C.; Koutrakis, P.; Burton, R. M. (1995a) A technique to expose animals to concentrated fine ambient aerosols. *Environ. Health Perspect.* 103: 172-177.
- Sioutas, D.; Koutrakis, P.; Ferguson, S. T.; Burton, R. M. (1995b) Development and evaluation of a prototype ambient particle concentrator for inhalation exposure studies. *Inhalation Toxicol.* 7: 633-644.
- Sioutas, C.; Koutrakis, P.; Wolfson, J. M.; Azaroff, L. S.; Mulik, J. D. (1996a) Field evaluation of a glass honeycomb denuder/filter pack system to collect atmospheric gases and particles. In: Gay, B. W.; Fuerst, R. G.; Jayanty, R. K. M., eds. *Measurement of toxic and related air pollutants: proceedings of a conference; May, 1994; Durham, NC. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development; report no. EPA/600/R-94/136; pp. 426-435.*
- Sioutas, C.; Wang, P. Y.; Ferguson, S. T.; Koutrakis, P.; Mulik, J. D. (1996b) Laboratory and field evaluation of an improved glass honeycomb denuder/filter pack sampler. *Atmos. Environ.* 30: 885-895.
- Sioutas, C.; Koutrakis, P.; Wang, P.-Y.; Babich, P.; Wolfson, J. M. (1999) Experimental investigation of pressure drop with particle loading in Nuclepore filters. *Aerosol Sci. Technol.* 30: 71-83.
- Sioutas, C.; Kim, S.; Chang, M.; Terrell, L. L.; Gong, H., Jr. (2000) Field evaluation of a modified DataRAM MIE scattering monitor for real-time PM_{2.5} mass concentration measurements. *Atmos. Environ.* 34: 4829-4838.
- Solomon, P. A.; Moyers, J. L.; Fletcher, R. A. (1983) High-volume dichotomous virtual impactor for the fractionation and collection of particles according to aerodynamic size. *Aerosol Sci. Technol.* 2: 455-464.
- Solomon, P. A.; Salmon, L. G.; Fall, T.; Cass, G. R. (1992) Spatial and temporal distribution of atmospheric nitric acid and particulate nitrate concentrations in the Los Angeles area. *Environ. Sci. Technol.* 26: 1596-1601.
- Song, X.-H.; Faber, N. M.; Hopke, P. K.; Suess, D. T.; Prather, K. A.; Schauer, J. J.; Cass, G. R. (2001) Source apportionment of gasoline and diesel by multivariate calibration based on single particle mass spectral data. *Anal. Chim. Acta* 446: 329-343.
- Speer, R. E.; Barnes, H. M.; Brown, R. (1997) An instrument for measuring the liquid water content of aerosols. *Aerosol Sci. Technol.* 27: 50-61.
- Spengler, J. D.; Thurston, G. D. (1983) Mass and elemental composition of fine and coarse particles in six U.S. cities. *J. Air Pollut. Control Assoc.* 33: 1162-1171.
- Stahlschmidt, T.; Schulz, M.; Dannecker, W. (1997) Application of total-reflection X-ray fluorescence for the determination of lead, calcium and zinc in size-fractionated marine aerosols. *Spectrochim. Acta B* 52: 995-1001.
- Stein, S. W.; Turpin, B. J.; Cai, X.; Huang, P.-F.; McMurry, P. H. (1994) Measurements of relative humidity-dependent bounce and density for atmospheric particles using the DMA-impactor technique. *Atmos. Environ.* 28: 1739-1746.
- Stevens, R. K.; O'Keefe, A. E.; Ortman, G. C. (1969) Absolute calibration of a flame photometric detector to volatile sulfur compounds at sub-part-per-million levels. *Environ. Sci. Technol.* 3: 652-655.
- Stevens, R. K.; Mulik, J. D.; O'Keefe, A. E.; Krost, K. J. (1971) Gas chromatography of reactive sulfur gases in air at the parts-per-billion level. *Anal. Chem.* 43: 827-831.
- Stolzenburg, M. R.; Hering, S. V. (2000) Method for the automated measurement of fine particle nitrate in the atmosphere. *Environ. Sci. Technol.* 34: 907-914.
- Streit, N.; Weingartner, E.; Zellweger, C.; Schwikowski, M.; Gäggeler, H. W.; Baltensperger, U. (2000) Characterization of size-fractionated aerosol from the Jungfraujoch (3580 m asl) using total reflection x-ray fluorescence (TXRF). *Int. J. Environ. Anal. Chem.* 76: 1-16.
- Strobel, H. A.; Heineman, W. R. (1989) *Chemical instrumentation: a systematic approach*. 3rd ed. New York, NY: Wiley and Sons, Inc.
- Strommen, M. R.; Kamens, R. M. (1997) Development and application of a dual-impedance radial diffusion model to simulate the partitioning of semivolatile organic compounds in combustion aerosols. *Environ. Sci. Technol.* 31: 2983-2990.

- Strothmann, J. A.; Schiermeier, F. A. (1979) Documentation of the regional air pollution study (RAPS) and related investigations in the St. Louis air quality control region. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Science Research Laboratory; report no. EPA-600/4-79-076.
- Suarez, A. E.; Caffrey, P. F.; Ondov, J. M.; Thaug, K. C. (1996) Size distribution of Ir-tagged soot from diesel sanitation trucks. *J. Aerosol Sci.* 27(suppl. 1): S697-S698.
- Suggs, J. C.; Burton, R. M. (1983) Spatial characteristics of inhalable particles in the Philadelphia metropolitan area. *J. Air Pollut. Control Assoc.* 33: 688-691.
- Suh, H. H.; Spengler, J. D.; Koutrakis, P. (1992) Personal exposures to acid aerosols and ammonia. *Environ. Sci. Technol.* 26: 2507-2517.
- Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1994) The relationship between airborne acidity and ammonia in indoor environments. *J. Exposure Anal. Environ. Epidemiol.* 4: 1-23.
- Sutherland, J. L.; Bhardwaja, P. S. (1987) Composition of the aerosol in northern Arizona and southern Utah. In: Bhardwaja, P. S., ed. *Transactions, Visibility Protection: Research and Policy Aspects*. Pittsburgh, PA: Air Pollution Control Association; pp. 373-385.
- Tan, S. H.; Horlick, G. (1986) Background spectral features in inductively coupled plasma/mass spectrometry. *Appl. Spectrosc.* 40: 445-460.
- Tang, I. N. (1980) Deliquescence properties and particle size change of hygroscopic aerosols. In: Willeke, K., ed. *Generation of aerosols and facilities for exposure experiments*. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 153-167.
- Tang, H.; Lewis, E. A.; Eatough, D. J.; Burton, R. M.; Farber, R. J. (1994) Determination of the particle size distribution and chemical composition of semi-volatile organic compounds in atmospheric fine particles with a diffusion denuder sampling system. *Atmos. Environ.* 28: 939-947.
- Tanner, R. L.; D'Ottavio, T.; Garber, R.; Newman, L. (1980) Determination of ambient aerosol sulfur using a continuous flame photometric detection system. I. Sampling system for aerosol sulfate and sulfuric acid. *Atmos. Environ.* 14: 121-127.
- Thomson, D. S.; Murphy, D. M. (1994) Analyzing single aerosol particles in real time. *CHEMTECH* 24: 30-35.
- Tobias, H. J.; Beving, D. E.; Ziemann, P. J.; Sakurai, H.; Zuk, M.; McMurry, P. H.; Zarling, D.; Wayrulis, R.; Kittelson, D. B. (2001) Chemical analysis of diesel engine nanoparticles using a nano-DMA/thermal desorption particle beam mass spectrometer. *Environ. Sci. Technol.* 35: 2233-2243.
- Tolocka, M. P.; Peters, T. M.; Vanderpool, R. W.; Chen, F. L.; Wiener, R. W. (2001a) On the modification of the low flow-rate PM₁₀ dichotomous sampler inlet. *Aerosol Sci. Technol.* 34: 407-415.
- Tolocka, M. P.; Solomon, P. A.; Mitchell, W.; Norris, G. A.; Gemmill, D. B.; Wiener, R. W.; Vanderpool, R. W.; Homolya, J. B.; Rice, J. (2001b) East vs. west in the US: chemical characteristics of PM_{2.5} during the winter of 1999. *Aerosol Sci. Technol.* 34: 88-96.
- Török, S. B.; Lábár, J.; Schmeling, M.; Van Grieken, R. E. (1998) X-ray spectrometry. *Anal. Chem.* 70: 495R-517R.
- Tsai, C.-J.; Cheng, Y.-H. (1996) Comparison of two ambient beta gauge PM₁₀ samplers. *J. Air Waste Manage. Assoc.* 46: 142-147.
- Tsai, C.-J.; Huang, H.-Y. (1995) Atmospheric aerosol sampling by an annular denuder system and a high-volume PM₁₀ sampler. *Environ. Int.* 21: 283-291.
- Turner, D. B. (1994) *Workbook of atmospheric dispersion estimates: an introduction to dispersion modeling*. 2nd ed. Boca Raton, FL: Lewis Publishers.
- Turner, J. R.; Hering, S. V. (1987) Greased and oiled substrates as bounce-free impaction surfaces. *J. Aerosol Sci.* 18: 215-224.
- Turpin, B. J.; Huntzicker, J. J. (1991) Secondary formation of organic aerosol in the Los Angeles basin: a descriptive analysis of organic and elemental carbon concentrations. *Atmos. Environ. Part A* 25: 207-215.
- Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. *Atmos. Environ.* 29: 3527-3544.
- Turpin, B. J.; Cary, R. A.; Huntzicker, J. J. (1990) An in situ, time-resolved analyzer for aerosol organic and elemental carbon. *Aerosol Sci. Technol.* 12: 161-171.
- Turpin, B. J.; Liu, S.-P.; Podolske, K. S.; Gomes, M. S. P.; Eisenreich, S. J.; McMurry, P. H. (1993) Design and evaluation of a novel diffusion separator for measuring gas/particle distributions of semivolatile organic compounds. *Environ. Sci. Technol.* 27: 2441-2449.
- Turpin, B. J.; Huntzicker, J. J.; Hering, S. V. (1994) Investigation of organic aerosol sampling artifacts in the Los Angeles basin. *Atmos. Environ.* 28: 3061-3071.
- Turpin, B. J.; Saxena, P.; Andrews, E. (2000) Measuring and simulating particulate organics in the atmosphere: problems and prospects. *Atmos. Environ.* 34: 2983-3013.

- Tyler, G. (1992) ICP vs. AA: how do you choose? *Environ. Lab.* (October/November): 31-34.
- U.S. Court of Appeals for the District of Columbia. (1999) *American Trucking Associations, Inc. vs. U.S. Environmental Protection Agency*. 175 F.3d 1027 (D.C. Cir. 1999).
- U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.
- U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1996b) Review of the national ambient air quality standards for particulate matter: policy assessment of scientific and technical information. OAQPS staff paper. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA/452/R-96-013. Available from: NTIS, Springfield, VA; PB97-115406REB.
- U.S. Environmental Protection Agency. (1998) Monitoring PM_{2.5} in ambient air using designated reference of class I equivalent methods. In: *Quality Assurance Handbook for Air Pollution Measurement Systems, V. II, Part II, Section 2.12*. Research Triangle Park, NC: Research and Development, Environmental Monitoring Systems Laboratory.
- U.S. Environmental Protection Agency. (1999) Particulate matter (PM_{2.5}) speciation guidance. Final draft. Edition 1. Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/amtic/files/ambient/pm25/spec/specfinl.pdf [2000, October 11].
- U.S. Environmental Protection Agency. (2001) List of designated reference and equivalent methods. Available: <http://www.epa.gov/ttn/amtic/files/ambient/criteria/repmlist.pdf> [05 March 02].
- U.S. Environmental Protection Agency. (2002a) Ambient monitoring technology information center (AMTIC). Washington, DC: Office of Air Quality Planning and Standards; Monitoring and Quality Assurance Group. Available: <http://www.epa.gov/ttn/amtic/> (20 March 2002).
- U.S. Environmental Protection Agency. (2002b) Support center for regulatory air models. Washington, DC: Office of Air Quality Planning and Standards. Available: <http://www.epa.gov/scram001> (21 March 2002).
- Van Dingenen, R.; Raes, F.; Jensen, N. R. (1995) Evidence for anthropogenic impact on number concentration and sulfate content of cloud-processed aerosol particles over the North Atlantic. *J. Geophys. Res. [Atmos.]* 100: 21,057-21,067.
- Van Loy, M.; Bahadori, T.; Wyzga, R.; Hartsell, B.; Edgerton, E. (2000) The Aerosol Research and Inhalation Epidemiology Study (ARIES): PM_{2.5} mass and aerosol component concentrations and sampler intercomparisons. *J. Air Waste Manage. Assoc.* 50: 1446-1458.
- Vanderpool, R. W.; Peters, T. M.; Natarajan, S.; Gemmill, D. B.; Wiener, R. W. (2001) Evaluation of the loading characteristics of the EPA WINS PM_{2.5} separator. *Aerosol Sci. Technol.* 34: 444-456.
- Vasiliou, J. G.; Sorenson, D.; McMurry, P. H. (1999) Sampling at controlled relative humidity with a cascade impactor. *Atmos. Environ.* 30: 1049-1056.
- Venkataraman, C.; Lyons, J. M.; Friedlander, S. K. (1994) Size distributions of polycyclic aromatic hydrocarbons and elemental carbon. 1. Sampling, measurement methods, and source characterization. *Environ. Sci. Technol.* 28: 555-562.
- Vincent, J. H. (1989) *Aerosol sampling: science and practice*. New York, NY: John Wiley & Sons.
- Vincent, J. H. (1995) *Aerosol science for industrial hygienists*. Oxford, United Kingdom: Pergamon.
- Wallace, J. C.; Hites, R. A. (1995) Computer-controlled low-volume air sampler for the measurement of semivolatile organic compounds. *Environ. Sci. Technol.* 29: 2099-2106.
- Wallace, L.; Howard-Reed, C. (2002) Continuous monitoring of ultrafine, fine, and coarse particles in a residence for 18 months in 1999-2000. *J. Air Waste Manage. Assoc.* 52: 828-844.
- Wang, P. Y. (1997) *Continuous aerosol mass measurement by flow obstruction*. Boston, MA: Harvard University, School of Public Health.
- Wang, H.-C.; John, W. (1987) Comparative bounce properties of particle materials. *Aerosol Sci. Technol.* 7: 285-299.
- Wang, H.-C.; John, W. (1988) Characteristics of the Berner impactor for sampling inorganic ions. *Aerosol Sci. Technol.* 8: 157-172.
- Ward, M. D.; Buttry, D. A. (1990) In situ interfacial mass detection with piezoelectric transducers. *Science (Washington, DC)* 249: 1000-1007.

- Watson, J. G.; Chow, J. C.; Richards, L. W.; Neff, W. D.; Andersen, S. R.; Dietrich, D. L.; Houck, J. E.; Olmez, I. (1988a) The 1987-88 metro Denver brown cloud study volume II: measurements. Reno, NV: Desert Research Institute; final report no. 8810.1F2 (2 of 3).
- Watson, J. G.; Chow, J. C.; Richards, L. W.; Neff, W. D.; Andersen, S. R.; Dietrich, D. L.; Houck, J. E.; Olmez, I. (1988b) The 1987-88 metro Denver brown cloud study volume III: data interpretation. Reno, NV: Desert Research Institute; final report no. 8810.1F3 (3 of 3).
- Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990a) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. *Environ. Software* 5: 38-49.
- Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B.; Nguyen, Q. T.; Meyer, E. L.; Pace, T. G. (1990b) Receptor model technical series, v. III (1989 revision): CMB7 user's manual. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-450/4-90-004. Available: <http://www.epa.gov/scram001/> [19 February 2002].
- Watson, J. G.; Chow, J. C.; Lu, Z.; Gertler, A. W. (1993) Particulate and gaseous organic receptor modeling for the Southern California Air Quality Study. Sacramento, CA: California Air Resource Board; report no. 8733.1F. Available: <http://www.arb.ca.gov/research/apr/past/atmospheric.htm#Air%20Quality%20Study> (5 June 2003).
- Watson, J. G.; Chow, J. C.; Lu, Z.; Fujita, E. M.; Lowenthal, D. H.; Lawson, D. R.; Ashbaugh, L. L. (1994a) Chemical mass balance source apportionment of PM₁₀ during the Southern California Air Quality Study. *Aerosol Sci. Technol.* 21: 1-36.
- Watson, J. G.; Chow, J. C.; Lowenthal, D. H.; Pritchett, L. C.; Frazier, C. A.; Neuroth, G. R.; Robbins, R. (1994b) Differences in the carbon composition of source profiles for diesel- and gasoline-powered vehicles. *Atmos. Environ.* 28: 2493-2505.
- Watson, J. G.; Fujita, E. M.; Chow, J. C.; Zielinska, B.; Richards, L. W.; Neff, W.; Dietrich, D. (1998) Northern front range air quality study. Final report. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere. Available: <http://www.nfraqs.colostate.edu/index2.html> (16 Jan 2002).
- Watson, J. G.; Chow, J. C.; Frazier, C. A. (1999) X-ray fluorescence analysis of ambient air samples. In: Landsberger, S.; Creatchman, M., eds. *Elemental Analysis of Airborne Particles*. Newark, NJ: Gordon and Breach Science Publishers; pp. 67-96.
- Weber, R. J.; McMurry, P. H.; Eisele, F. L.; Tanner, D. J. (1995) Measurement of expected nucleation precursor species and 3-500-nm diameter particles at Mauna Loa Observatory, Hawaii. *J. Atmos. Sci.* 52: 2242-2257.
- Weber, R. J.; Marti, J. J.; McMurry, P. H.; Mauldin, L.; Tanner, D.; Eisele, F.; Brechtel, F.; Kreidenweis, S.; Kok, G.; Schillawski, R.; Baumgardner, D. (1997) Measurements of new particle formation and ultrafine particle growth rates at a clean continental site. *J. Geophys. Res.* 102: 4375-4385.
- Weber, R. J.; McMurry, P. H.; Mauldin, L.; Tanner, D. J.; Eisele, F. L.; Brechtel, F. J.; Kreidenweis, S. M.; Kok, G. L.; Schillawski, R. D.; Baumgardner, D. (1998) A study of new particle formation and growth involving biogenic and trace gas species measured during ACE 1. *J. Geophys. Res. A* 103: 16,385-16,396.
- Weber, R. J.; McMurry, P. H.; Mauldin, R. L., III; Tanner, D. J.; Eisele, F. L.; Clarke, A. D.; Kapustin, V. N. (1999) New particle formation in the remote troposphere: a comparison of observations at various sites. *Geophys. Res. Lett.* 26: 307-310.
- Weber, R. J.; Orsini, D.; Daun, Y.; Lee, Y.-N.; Kotz, P. J.; Brechtel, F. (2001) A particle-into-liquid collector for rapid measurement of aerosol bulk chemical composition. *Aerosol Sci. Technol.* 35: 718-727.
- Wedding, J. B.; Weigand, M. A. (1993) An automatic particle sampler with beta gauging. *J. Air Waste Manage. Assoc.* 43: 475-479.
- Weingartner, E.; Burtscher, H.; Baltensperger, U. (1997) Hygroscopic properties of carbon and diesel soot particles. *Atmos. Environ.* 31: 2311-2327.
- Weschler, C. J.; Shields, H. C. (1999) Indoor ozone/terpene reactions as a source of indoor particles. *Atmos. Environ.* 33: 2301-2312.
- Wesolowski, J. J.; John, W.; Devor, W.; Cahill, T. A.; Feeney, P. J.; Wolfe, G.; Flocchini, R. (1977) Collection surfaces of cascade impactors. In: Dzubay, T. G., ed. *X-ray fluorescence analysis of environmental samples*. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 121-131.
- Wexler, A. S.; Sarangapani, R. (1998) Particles do not increase vapor deposition in human airways. *J. Aerosol Sci.* 29: 197-204.
- Whitby, K. T. (1978) The physical characteristics of sulfur aerosols. *Atmos. Environ.* 12: 135-159.

- Whitby, K. T.; Sverdrup, G. M. (1980) California aerosols: their physical and chemical characteristics. In: Hidy, G. M.; Mueller, P. K.; Grosjean, D.; Appel, B. R.; Wesolowski, J. J., eds. *The character and origins of smog aerosols: a digest of results from the California Aerosol Characterization Experiment (ACHEX)*. New York, NY: John Wiley & Sons, Inc.; pp. 477-517. (*Advances in environmental science and technology*: v. 9).
- Whitby, K. T.; Charlson, R. E.; Wilson, W. E.; Stevens, R. K. (1974) The size of suspended particle matter in air. *Science* (Washington, DC) 183: 1098-1099.
- White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. *J. Air Waste Manage. Assoc.* 48: 454-458.
- Wiedensohler, A.; Hansson, H.-C.; Orsini, D.; Wendisch, M.; Wagner, F.; Bower, K. N.; Choullarton, T. W.; Wells, M.; Parkin, M.; Acker, K.; Wieprecht, W.; Facchini, M. C.; Lind, J. A.; Fuzzi, S.; Arends, B. G.; Kulmala, M. (1997) Night-time formation and occurrence of new particles associated with orographic clouds. *Atmos. Environ.* 31: 2545-2559.
- Willeke, K.; Baron, P. A., eds. (1993) *Aerosol measurement: principles, techniques and applications*. New York, NY: Van Nostrand Reinhold Publishers.
- Willeke, K.; Whitby, K. T. (1975) Atmospheric aerosols: size distribution interpretation. *J. Air Pollut. Control Assoc.* 25: 529-534.
- Williams, E. L., II; Grosjean, D. (1990) Removal of atmospheric oxidants with annular denuders. *Environ. Sci. Technol.* 24: 811-814.
- Williams, R.; Suggs, J.; Zweidinger, R.; Evans, G.; Creason, J.; Kwok, R.; Rodes, C.; Lawless, P.; Sheldon, L. (2000) The 1998 Baltimore Particulate Matter Epidemiology-Exposure Study: Part I. Comparison of ambient, residential outdoor, indoor and apartment particulate matter monitoring. *J. Exposure Anal. Environ. Epidemiol.* 10: 518-532.
- Wilson, W. E. (1995) Aerosol exposure, physics, and chemistry. *Inhalation Toxicol.* 7: 769-772.
- Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. *J. Air Waste Manage. Assoc.* 47: 1238-1249.
- Wilson, W. E.; Spiller, L. L.; Ellestad, T. G.; Lamothe, P. J.; Dzubay, T. G.; Stevens, R. K.; Macias, E. S.; Fletcher, R. A.; Husar, J. D.; Husar, R. B.; Whitby, K. T.; Kittelson, D. B.; Cantrell, B. K. (1977) General Motors sulfate dispersion experiment: summary of EPA measurements. *J. Air Pollut. Control Assoc.* 27: 46-51.
- Woo, K.-S.; Chen, D.-R.; Pui, D. Y. H.; McMurry, P. H. (2001a) Measurement of Atlanta aerosol size distributions: observations of ultrafine particle events. *Aerosol Sci. Technol.* 34: 75-87.
- Woo, K.-S.; Chen, D.-R.; Pui, D. Y. H.; Wilson, W. E. (2001b) Use of continuous measurements of integral aerosol parameters to estimate particle surface area. *Aerosol Sci. Technol.* 34: 57-65.
- Wu, C. C.; Suarez, A. E.; Lin, Z.; Kidwell, C. B.; Borgoul, P. V.; Caffrey, P. F.; Ondov, J. M.; Sattler, B. (1998) Application of an Ir tracer to determine soot exposure to students commuting to school on Baltimore public buses. *Atmos. Environ.* 32: 1911-1919.
- Yamasaki, H.; Kuwata, K.; Miyamoto, H. (1982) Effects of ambient temperature on aspects of airborne polycyclic aromatic hydrocarbons. *Environ. Sci. Technol.* 16: 189-194.
- Yatin, M.; Tuncel, S. G.; Tuncel, G.; Aras, N. K. (1994) Trace element composition of atmospheric aerosols in Ankara, Turkey, determined by instrumental neutron activation analysis. *J. Radioanal. Nucl. Chem.* 181: 401-411.
- Zeng, X.; Wu, X.; Yao, H.; Yang, F.; Cahill, T. A. (1993) PIXE-induced XRF with transmission geometry. *Nucl. Instr. Meth. Phys. Res. B* 75: 99-104.
- Zhang, X. Q.; McMurry, P. H. (1987) Theoretical analysis of evaporative losses from impactor and filter deposits. *Atmos. Environ.* 21: 1779-1789.
- Zhang, X.; McMurry, P. H. (1992) Evaporative losses of fine particulate nitrates during sampling. *Atmos. Environ. Part A* 26: 3305-3312.
- Zhang, X. Q.; McMurry, P. H.; Hering, S. V.; Casuccio, G. S. (1993) Mixing characteristics and water content of submicron aerosols measured in Los Angeles and at the Grand Canyon. *Atmos. Environ. Part A* 27: 1593-1607.
- Zhu, Y.; Hinds, W. C.; Kim, S.; Shen, S.; Sioutas, C. (2002a) Study of ultrafine particles near a major highway with heavy-duty diesel traffic. *Atmos. Environ.* 36: 4323-4335.
- Zhu, Y.; Hinds, W. C.; Kim, S.; Sioutas, C. (2002b) Concentration and size distribution of ultrafine particles near a major highway. *J. Air Waste Manage. Assoc.* 52: 1032-1042.
- Zoller, W. H.; Gordon, G. E. (1970) Instrumental neutron activation analysis of atmospheric pollutants utilizing Ge(Li) γ -ray detectors. *Anal. Chem.* 42: 257-265.

APPENDIX 2A. TECHNIQUES FOR MEASURING OF SEMIVOLATILE ORGANIC COMPOUNDS

Use of Denuder Systems To Measure Semivolatile Organic Compounds

Phase distribution of semivolatile organic species has been the subject of several studies that have employed denuder technology (see Gundel et al., 1995; Gundel and Lane, 1999) to directly determine the phase distributions while avoiding some of the positive and negative sampling artifacts associated with back-up quartz filters. In an ideal system with a denuder that is 100% efficient, the gas phase would be collected in the denuder and the particle phase would be the sum of the material collected on the filter and the adsorbent downstream. Denuder collection efficiency depends on the denuder surface area (+), the diffusivity (+) and vapor pressure (-) of the compound, the temperature (-) and flow rate (-) of the air stream, and the presence of competing species (-), including water vapor (Cui et al., 1998; Kamens and Coe, 1997; Lane et al., 1988). (The + and - symbols in parentheses indicate qualitatively the effect increasing each parameter would have on efficiency). In a system with a denuder collection efficiency < 100%, the collection efficiency must be known to accurately attribute adsorbed organics from denuder breakthrough to the gas phase and adsorbed organics volatilized from collected particles to the particle phase. In calculating the overall phase distributions of SVOC PAH from a denuder system, the collection efficiency for each compound is needed.

The efficiency of silicone-grease-coated denuders for the collection of polynuclear aromatic hydrocarbons was examined by Coutant et al. (1992), who examined the effects of uncertainties in the diffusion coefficients and in the collisional reaction efficiencies on the overall phase distributions of SVOC PAH calculated using denuder technology. In their study, they used a single stage, silicone-grease-coated aluminum annular denuder with a filter holder mounted ahead of the denuder and an XAD trap deployed downstream of the denuder. In a series of laboratory experiments, they spiked the filter with a mixture of perdeuterated PAH, swept the system with ultra-high purity air for several hours, and then analyzed the filter and the XAD. They found that the effects of these uncertainties, introduced by using a single compound as a surrogate PAH (in their case, naphthalene) for validation of the denuder collection

efficiency, are less significant than normal variations because of sampling and analytical effects. Field study results using their sampling system have not been published.

For measuring particulate phase organic compounds, the denuder-based sampling system represents an improvement over the filter/adsorbent collection method (Turpin et al., 1993). Some researchers, however, have reported that denuder coatings themselves can introduce contamination (Mukerjee et al., 1997) and that the adsorbed species may be difficult to remove from the coating (Eatough et al., 1993).

In a study conducted in southern California (Eatough et al., 1995), the Brigham Young University Organic Sampling System (BOSS; Eatough et al., 1993) was used to determine POM composition, and a high-volume version (BIG BOSS; flow rate 200 L/min) was used to determine the particulate size distribution and the chemical composition of SVOC in fine particles. The BOSS, a multi-channel diffusion denuder sampling system, consists of two separate samplers (each operating at 35 L/min). The first sampler consists of a multi-parallel plate diffusion denuder with charcoal-impregnated filter papers as the collection surfaces followed by a two-stage quartz filter pack and a two-stage charcoal-impregnated filter pack. The second sampler operating in parallel with the first consists of a two-stage quartz filter pack, followed by the parallel plate denuder, followed by the two-stage charcoal-impregnated filter pack. The filter samples collected by the BOSS sampler were analyzed by temperature-programmed volatilization analysis. The second channel allows calculations of the efficiency of the denuder in removing gas-phase specifics that would be absorbed by the charcoal impregnated filter. Eatough et al. (1995) also operated a two-stage quartz filter pack alongside the BOSS sampler. The BIG BOSS system (Tang et al., 1994) consists of four systems (each with a flow rate of 200 L/min). Particle size cuts of 2.5, 0.8, and 0.4 μm are achieved by virtual impaction, and the sample subsequently flows through a denuder, then is split, with the major flow (150 L/min) flowing through a quartz filter followed by an XAD-II bed. The minor flow is sampled through a quartz filter backed by a charcoal-impregnated filter paper. The samples derived from the major flow (quartz filters and XAD-II traps) were extracted with organic solvents and analyzed by gas chromatography (GC) and GC-mass spectroscopy. The organic material lost from the particles was found to represent all classes of organic compounds.

Eatough et al. (1996) operated the BOSS sampler for a year at the IMPROVE site at Canyonlands National Park, UT alongside the IMPROVE monitor and a separate sampler

consisting of a two-stage quartz filter pack. They found that concentrations of particulate carbon determined from the quartz filter pack sampling system were lower by an average 39%, which was attributed to volatilization losses of SVOC from the quartz filters. In another study conducted with the BOSS in southern California, losses of 35% of the POM, on average, were found and attributed to losses of the SVOC during sampling (Eatough et al., 1995).

The denuder used in the various BOSS samplers consists of charcoal-impregnated cellulose fiber filter material. Denuder collection efficiencies of greater than 95% have been reported for organic gases that adsorb on quartz and charcoal-impregnated filters (Eatough et al., 1999a; Ding et al., 2002; Lewtas et al., 2001). However, because the mass concentration of gas phase species that adsorb on quartz and charcoal-impregnated filters is so much greater than the mass of semivolatile organic material in the particulate phase, it is necessary to measure and account for the inefficiency of the denuder in the BOSS samplers. To address this problem, Brigham Young University (BYU) developed a particle-concentrator (PC)-BOSS system (Ding et al., 2002; Eatough et al., 1999b; Lewtas et al., 2001; Modey et al., 2001; Pang et al., 2001, 2002a,b). The PC-BOSS includes a virtual impactor upstream of the denuder to improve the denuder collection efficiency by removing a majority of the gases from the aerosol flow. With this system, denuder collection efficiencies of > 99% have been reported for organic gases and other species, such as SO₂(g), HNO₃(g), etc., that adsorb on quartz and charcoal-impregnated filters (Pang et al., 2001). Since the concentrations of semivolatile organic and other gases are not altered by the virtual-impaction concentration of accumulation mode particles (except possibly by slight changes in pressure), it is anticipated that the gas-particle distribution will not be significantly altered by the concentration process. The virtual impactor has a 50% cut point at 0.1 μm D_a. As a result, some particles in the 0.05 to 0.2 μm diameter size range will be removed in the major flow along with the majority of the gases. Therefore, the mass collection efficiency of the virtual impactor concentrator will be a function of the particle size distribution in the 0.05 to 0.1 μm size range. This collection efficiency is measured by comparing the concentration of nonvolatile components measured in the concentrated sample with that measured in an unconcentrated sample. The concentration efficiency varies from 50 to 75%. It is relatively constant over periods of weeks, but varies by season and by site, presumably as the particle size distribution changes. Previous studies at Harvard (Sioutas et al., 1995a,b) have shown that the composition of the sampled aerosol is little changed by the concentration process. The BYU studies listed

above have shown that the concentration efficiencies for sulfate, OC, and EC are comparable for given sampling locations. Furthermore, the concentrations of these species and of fine particulate nitrate determined using the PC-BOSS have been shown to be comparable to those determined using more conventional samplers for sulfate or EC or using simpler denuder systems for OC and nitrate.

Ding et al. (1998a) developed a method for the determination of total n-nitroso compounds in air samples and used the method to examine organic compounds formed from NO_x chemistry in Provo, UT (Ding et al., 1998b). In their method, n-nitroso compounds are selectively decomposed to yield nitric oxide, which is then detected using chemiluminescence. Using the Provo samples, they found that the majority of the n-nitroso and nitrite organic compounds that were present in fine particulate matter were SVOCs that could be evaporated from the particles during sampling. They found particulate n-nitroso compound concentrations ranging between < 1 and 3 nmoles/m³ and gas-phase n-nitroso compound concentrations in the same range. Particulate organic nitrite concentrations were found in the range of <1 to ≈5 nmoles/m³, and gas-phase concentrations as high as 10 nmoles/m³ were found.

Turpin et al. (1993) developed a sampling system that corrects for the loss of SVOCs during sampling by removing most of the gas phase material from the particles in a diffusion separator sampling system. Unlike the previously mentioned systems, wherein the particulate phase is measured directly, in the system of Turpin et al. (1993), the gas-phase is measured directly. In the laminar flow system, ambient, particle-laden air enters the sampler as an annular flow. Clean, particle-free air is pushed through the core inlet of the separator. The clean air and ambient aerosol join downstream of the core inlet section, and flow parallel to each other through the diffusion zone. Because of the much higher diffusivities for gases compared to particles, the SVOC in the ambient air diffuses to the clean, core flow. The aerosol exits the separator in the annular flow, and the core flow exiting the separator now contains a known fraction of the ambient SVOC. Downstream of the diffusion separator, the core exit flow goes into a polyurethane foam (PUF) plug where the SVOC is collected. The adsorbed gas phase on the PUF plug is extracted with supercritical fluid CO₂ and then analyzed by gas chromatography/mass-selective detection (GC/MSD). The gas-phase SVOC is thus determined. Ultimately, to determine particulate phase SVOC concentrations, the total compound concentration also has to be measured and the particulate phase obtained by difference.

The system was tested for the collection of PAHs. The diffusional transport of gas-phase PAHs and particle concentrations agreed well with theory. Breakthrough was problematic for low molecular weight PAHs (mol. wt. < 160). Detection limits ranged from 20 to 50 pg of injected mass for all PAHs.

Gundel et al. (1995) recently developed a technique for the direct determination of phase distributions of semivolatile polycyclic aromatic hydrocarbons using annular denuder technology. The method, called the integrated organic vapor/particle sampler (IOVPS), uses a cyclone inlet with a 50% cut point of 2.5 μm at a sampling rate of 10 L/min. The airstream then goes through two or three sandblasted glass annular denuders that are coated with ground adsorbent resin material (XAD-4 was initially examined) that traps vapor-phase organics. The airstream subsequently passes through a filter, followed by a backup denuder. The denuder collection efficiency is high and compares well with predictions based on the diffusivity of the compounds. The denuder can also be extracted to obtain gas-phase concentrations directly (Gundel and Lane, 1999). Particle-phase PAHs are taken to be the sum of material on the filter and XAD adsorbent downstream after correction for denuder collection efficiency. The IOVPS was tested for sampling semivolatile PAH in laboratory indoor air and in environmental tobacco smoke (ETS). After exposure, the denuders, filters, and sorbent traps were extracted with cyclohexane (Gundel et al., 1995) and analyzed for PAHs from naphthalene to chrysene using dual-fluorescence detection (Mahanama et al., 1994). Recoveries from both denuders and filters were ~70% for 30 samples. Detection limits (defined as three times the standard deviation of the blanks) for gas-phase SVOC PAHs ranged from 0.06 ng for anthracene to 19 ng for 2-methylnaphthalene. The 95% confidence interval (CI) for reproduction of an internal standard concentration was 6.5% of the mean value. Relative precision, from a propagation of errors analysis or from the 95% CI from replicate analyses of standard reference material SRM 1649 (urban dust/organics), was 12% on average (8% for naphthalene to 22% for fluorene). Sources of error included sampling flow rate, internal standard concentration, and coeluting peaks. Gundel and Lane (1999) reported that roughly two-thirds of particulate PAH fluoranthene, pyrene, benz[*a*]anthracene, and chrysene were found on the postfilter denuders, so that it is likely that considerable desorption from the collected particles took place.

Solid adsorbent-based denuder systems have been investigated by other researchers as well. Bertoni et al. (1984) described the development of a charcoal-based denuder system for the collection of organic vapors. Risse et al. (1996) developed a diffusion denuder system to sample aromatic hydrocarbons. In their system, denuder tubes with charcoal coating and charcoal paper precede a filter pack for particulate collection and an adsorption tube to capture particle blow-off from the filter sample. Breakthrough curves for benzene, toluene, ortho-xylene, and meta-xylene were developed for 60-, 90-, and 120-cm denuder tubes. The effects of relative humidity on the adsorption capacities of the denuder system were examined, and it was found that the capacity of the charcoal was not affected significantly by increases in relative humidity. The feasibility of outdoor air sampling with the system was demonstrated.

Krieger and Hites (1992) designed a diffusion denuder system that uses capillary gas chromatographic columns as the tubes for SVOC collection. The denuder was followed by a filter to collect particles, which in turn was followed by a PUF plug to collect organic material volatilizing off the filter. Denuder samples were analyzed by liquid solvent extraction (CH_2Cl_2) followed by GC-mass spectrometric analysis. The PUF plugs and filters were fluid extracted with supercritical N_2O . Using this system, an indoor air sample was found to contain primarily chlorinated biphenyls, ranging from trichlorobiphenyls (vapor pressures 10^{-3} to 10^{-4} Torr at 25 °C) to octachlorobiphenyls (10^{-6} – 10^{-7} Torr). This demonstrated that the sampler collects compounds with a wide range of volatility. They also found that on-line desorption is successful in maintaining good chromatographic peak shape and resolution. The entire method, from sample collection to the end of the chromatographic separation, took 2 hours.

Organic acids in both the vapor and particulate phases may be important contributors to ambient acidity, as well as representing an important fraction of organic PM. Lawrence and Koutrakis (1996a,b) used a modified Harvard/EPA annular denuder system (HEADS) to sample both gas and particulate phase organic acids in Philadelphia, PA in the summer of 1992. The HEADS sampler inlet had a 2.1- μm cut point impactor (at 10 L/min), followed by two denuder tubes, and finally a Teflon filter. The first denuder tube was coated with potassium hydroxide (KOH) to trap gas phase organic acids. The second denuder tube was coated with citric acid to remove ammonia to avoid neutralizing particle phase acids collected on the filter. The KOH-coated denuder tube was reported to collect gas phase formic and acetic acids at better than 98.5% efficiency and with precisions of 5% or better (Lawrence and Koutrakis, 1994). It was

noted that for future field measurements of particulate organic acids, a Na₂CO₃-coated filter should be deployed downstream of the Teflon filter to trap organic acids that may evaporate from the Teflon filter during sampling.

Role of the Collection Media

The role of the collection media was recently examined in a study conducted in Seattle (Lewtas et al., 2001). In that study, the influence of denuder sampling methods and filter collection media on the measurement of SVOC associated with PM_{2.5} was evaluated. Activated carbon and XAD collection media were used in diffusion denuders and impregnated back-up filters in two different samplers, the Versatile Air Pollution Sampler (VAPS) and the PC-BOSS. XAD-coated glass annular denuders and charcoal-impregnated cellulose fiber (CIF) filter denuders also were used. CIF filters were also compared to XAD-coated quartz filters as backup filter collection media. Lewtas et al. (2001) found that the two denuder types resulted in an equivalent measurement of particulate OC and particle mass. The carbon-coated denuders in the BOSS sampler were more efficient than the XAD-coated denuders for the collection of the more highly volatile organic compounds (MHVOCs). Lewtas et al. (2001) concluded that this MHVOC fraction that is collected in the carbon-coated BOSS denuder does not contribute substantially to the particle mass or to the SVOC measured as OC on quartz filters. However, this MHVOC fraction would be captured in the carbon-impregnated filters placed behind the quartz filters, so that, in the XAD denuder configuration, the captured MHVOC would cause a higher OC concentration and an overestimation of the SVOC.

Some of the recent research in denuder technology has also focused on reducing the size of the denuder, optimizing the residence time in the denuder, understanding the effect of diffusion denuders on the positive quartz filter artifact, identifying changes in chemical composition that occur during sampling, determining the effects of changes in temperature and relative humidity, and identifying possible losses by absorption in coatings.

Reducing the Size of Denuders

The typical denuder configuration is an annular diffusion denuder tube of significant length (e.g., 26.5 cm for 10 L/min [Koutrakis et al., 1988a,b]). A more compact design based on a honeycomb configuration was shown to significantly increase the capacity (Koutrakis et al.,

1993). However, in intercomparisons with an annular denuder/filter pack system (Koutrakis et al., 1993), significant losses of ammonia and nitric acid were observed for the honeycomb configuration and were attributed to the large inlet surface area and long sample residence time of the honeycomb design relative to the annular denuder system. Sioutas et al. (1996a) subsequently designed a modified glass honeycomb denuder/filter pack sampler (HDS) with an inlet that minimizes vapor losses on the inlet surfaces. The modified HDS has reduced inlet surfaces and decreased residence time for sampled gases (NH_3 and HNO_3) compared to its predecessor (Sioutas et al., 1994d). Sioutas et al. (1996b) also tested various inlet materials (glass, PFA, and polytetrafluoroethylene [PTFE]) in laboratory tests and found that a PTFE Teflon coated inlet minimized loss of sampled gases (1 to 8% loss of HNO_3 observed, and -4 to 2% loss of NH_3 observed). The highest inlet losses were observed for HNO_3 lost to PFA surfaces (14 to 25%). The modified HDS was tested in laboratory and field tests and found to agree within 10% with the annular denuder system.

Residence Time in the Denuder

The efficiency of a diffusion denuder sampler for the removal of gas phase material can be improved by increasing the residence time of the sampled aerosol in the denuder. However, the residence time can only be increased within certain limits. Because the diffusion denuder reduces the concentration of gas-phase semivolatile organic material, semivolatile organic matter present in the particles passing through the denuder will be in a thermodynamically unstable environment and will tend to outgas SVOC during passage through the denuder. The residence time of the aerosol in the denuder, therefore, should be short enough to prevent significant loss of particulate-phase SVOC to the denuder. Various studies have suggested that the residence time in the denuder should be less than about 2 s (Gundel and Lane, 1999; Kamens and Coe, 1997; Kamens et al., 1995). The residence times in the various denuder designs described by Gundel and Lane (1999) are from 1.5 to 0.2 s. The equilibria and evaporation rates are not as well understood for organic components as they are for NH_4NO_3 (Zhang and McMurry, 1987, 1992; Hering and Cass, 1999).

Effect of Diffusion Denuders on the Positive Quartz Filter Artifact

The adsorption of organic compounds by a second quartz filter has been shown to be reduced, but not eliminated, in samples collected in the Los Angeles Basin using a multi-channel diffusion denuder with quartz filter material as the denuder collection surface preceding the quartz filters (Fitz, 1990). This artifact can be further reduced by using activated charcoal as the denuder surface and using a particle concentrator to reduce the amount of gas phase organic compounds relative to condensed phase organic compounds (Cui et al., 1997, 1998; Eatough, 1999). Other experiments (Cotham and Bidleman, 1992; Cui et al., 1998; Eatough et al., 1995, 1996) have shown that the quartz filter artifact can result both from the collection of gas phase organic compounds and from the collection of SVOCs lost from particles during sampling. Thus, available results suggest that both a “positive” and a “negative” artifact can occur with the determination of particulate phase organic compounds using two tandem quartz filters.

The importance of the adsorption of organic vapors on PM or filters relative to the volatilization of organic compounds from PM collected on a filter continues to be a topic of active debate. The relative importance of positive and negative artifacts differ for denuded and undenuded filters; depend on face velocity, sample loading, and the vapor pressures of the compounds of interest; and may vary with season and location because of variations in the composition of volatile and semivolatile organic material. Evidence exists for substantial positive and negative artifacts in the collection of organic PM.

Changes in Chemical Composition During Sampling

The use of sampling systems designed to correctly identify the atmospheric gas and particulate phase distributions of collected organic material has been outlined above. An additional sampling artifact that has received little consideration in the collection of atmospheric samples is the potential alteration of organic compounds by the sampling process itself. These alterations appear to result from the movement of ambient air containing oxidants and other reactive compounds past the collected particles. The addition of NO₂ (< 1 ppm) or O₃ (< 200 ppb) to the sampled air stream (at 0 to 5 °C) for a high-volume sampler reduced the concentrations of benzo[*a*]pyrene and benzo[*a*]anthracene from a few percent to 38%, with the observed reduction increasing with increased concentration of the added gases (Brorström et al., 1983). Spiking a filter with an amine increased the measured concentrations of nitrosamines in

both the filter and following XAD sorbent bed for a mid-volume sampler (Ding et al., 1998a,b). Similar results have been obtained for the exposure of a deuterated amine on a filter to NO_x (Pellizzari and Krost, 1984). When Tenax columns spiked with deuterated styrene and cyclohexene were exposed to ppm concentrations of ozone or halogens, oxygenated and halogenated compounds were formed (Pellizzari and Krost, 1984). Similar oxidation of aldehydes and peroxyacetylnitrate (PAN) during sampling has been observed (Grosjean and Parmar, 1990). Collected PAH compounds can be oxygenated or nitrated on a filter (Davis et al., 1987; Lindskog et al., 1987), but 1-nitropyrene has been shown to be resistant to additional nitration (Grosjean, 1983). These various chemical transformations of collected organic compounds can be eliminated by removing the gas phase oxidants, NO_x , HNO_3 , etc., by reaction or adsorption prior to particle collection (Ding, 1998a,b; Grosjean and Parmar, 1990; Parmar and Grosjean, 1990; Pellizzari and Krost, 1984; Williams and Grosjean, 1990). The BOSS denuder should be effective in eliminating most of the chemical transformation artifacts because reactive gases are removed by the charcoal denuder that precedes the particle collection filter.

Temperature and Relative Humidity Effects

The problems of sampling artifacts associated with SVOC adsorption and evaporation are compounded by temperature and relative humidity (RH) effects (Pankow and Bidleman, 1991; Pankow et al., 1993; Falconer et al., 1995; Goss and Eisenreich, 1997). The effects of temperature on the partitioning of PAHs were examined by Yamasaki et al. (1982), who found that the partition coefficient ($\text{PAH}_{\text{vapor}}/\text{PAH}_{\text{part}}$) was inversely related to temperature and could be described using the Langmuir adsorption concept. The dissociation of ammonium nitrate aerosol is also a function of temperature. Bunz et al. (1996) examined the dissociation and subsequent redistribution of NH_4NO_3 within a bimodal distribution using a nine-stage low-pressure Berner impactor followed by ion chromatography analysis and found a strong temperature dependency on the redistribution. Bunz et al. (1996) found that at lower temperatures (below 10 °C) there was little change in the aerosol size distribution. At temperatures between 25 and 45 °C, however, the lifetime of NH_4NO_3 particles decreased by more than a factor of 10, and size redistribution, as measured by average ending particle diameter, increased more for higher temperatures than for lower temperatures.

The effects of relative humidity on the sorption of SVOC on particles are not well understood. In a series of laboratory experiments, Goss and Eisenreich (1997) examined the sorption of both nonpolar (hydrocarbons and chlorinated hydrocarbons) and polar (ethyl ether and acetone) VOCs onto combustion soot particles as a function of temperature and relative humidity. The soot particles used in their experiments were collected from oil furnaces and contained 60% (w/w) iron sulfate (water-soluble fraction) and 9% (w/w) EC and OC. They found that, for all compounds, the sorption of VOCs onto soot particles decreased with increasing relative humidity over the range of 10 to 95%. They also observed hysteresis in the relative humidity dependency, with sorption coefficients at a given relative humidity higher when the RH is being increased than when the RH is being decreased. The sorption coefficients were fit with an exponential function to the RH so that the slope of the regression line would provide a measure of the influence of relative humidity. Based on the magnitude of the slope, they concluded that the RH-dependency of sorption was stronger for water-soluble organic compounds.

In another study by Jang and Kamens (1998), humidity effects on SVOC gas-particle partitioning were examined using outdoor environmental chambers and the experimentally determined partitioning coefficients were compared to theoretical values. They examined the partitioning of SVOC onto wood soot, diesel soot, and secondary aerosols and concluded that “the humidity effect on partitioning was most significant for hydrophobic compounds adsorbing onto polar aerosols.” Although these two studies seem to be contradictory, on closer examination, it is difficult to compare the two studies for several reasons. The experiments conducted by Jang and Kamens (1998) were conducted in outdoor chambers at ambient temperatures and humidities. Their model was for absorptive partitioning of SVOC on *liquid-like* atmospheric PM. In contrast, the results of Goss and Eisenreich (1997) were obtained using a GC system operated at 70 °C higher than ambient conditions. Goss and Eisenreich (1997) modeled adsorptive partitioning of VOC on *solid-like* atmospheric PM. In the study of Jang and Kamens (1998), calculated theoretical values for water activity coefficients for diesel soot were based on an inorganic salt content of 1 to 2%; whereas, the combustion particles studied by Goss and Eisenreich (1997) contained 60% water-soluble, inorganic salt content. Jang and Kamens (1998) obtained their diesel soot from their outdoor chamber, extracted it with organic solvent (mixtures of hexane and methylene chloride), and measured the organic fraction.

The resulting salt content of 2% of the PM studied in Jang and Kamens (1998) is enough to affect water uptake but presumably not enough to affect the sorption partitioning of organics.

Impactor Coatings

Impactors are used to achieve a size cut point and to collect particles on surfaces. Particles collected on impactors are exposed to smaller pressure drops than filter-collected particles, making them less susceptible to volatile losses (Zhang and McMurry, 1987). However, size resolution can be affected by bounce when samples are collected at low humidities (Stein et al., 1994). Other sources of error inherent in some currently acceptable practices could potentially affect PM concentration measurements and will likely become even more important as more emphasis is placed on chemical speciation. Allen et al. (1999a) reported that the practice of greasing impaction substrates may introduce an artifact from the absorption of semivolatile species from the gas phase by the grease, because the grease could artificially increase the amount of PAHs and other organic compounds attributed to the aerosol. Allen et al. (1999b) offered several criteria to ensure that this absorption artifact is negligible, including selecting impaction oils in which analytes of interest are negligibly soluble and ensuring that species do not have time to equilibrate between the vapor and oil phases (this criterion is met for nonvolatile species). They recommend using oiled impaction substrates only if the absorption artifact is negligible as determined from these criteria. The application of greases and impaction oils for preventing or reducing bounce when sampling with impactors is not suitable for carbon analysis, because the greases contain carbon (Vasilou et al., 1999).

Kavouras and Koutrakis (2001) investigated the use of polyurethane foam (PUF) as a substrate for conventional inertial impactors. The PUF impactor substrate is not rigid like the traditional impactor substrate, so particle bounce and reentrainment artifacts are reduced significantly. Kavouras and Koutrakis (2001) found that the PUF impaction substrate resulted in a much smaller 50% cut point at the same flow rate and Reynolds number. Moreover, the lower 50% cut point was obtained at a lower pressure drop than with the conventional substrate, which could reduce artifact vaporization of semivolatile components.

APPENDIX 2B. ANALYTICAL TECHNIQUES

2B.1 INORGANIC ELEMENTS

2B.1.1 Energy Dispersive X-Ray Fluorescence (EDXRF)

EDXRF has usually been the method of choice for analysis of trace elements on filters. EDXRF is preferred for aerosol analysis over wavelength dispersive X-ray fluorescence (XRF) because its fast analysis over the total spectrum allows simultaneously analysis of numerous elements. EDXRF can accommodate small sample sizes and requires little sample preparation or operator time after the samples are placed in the analyzer. It also leaves the sample intact after analysis, making further analysis possible. XRF irradiates a uniform particle deposit on the surface of a membrane filter with 1 to 50 keV X-rays that eject inner shell electrons from the atoms of each element in the sample (Dzubay and Stevens, 1975; Jaklevic et al., 1977; Billiet et al., 1980; Potts and Webb, 1992; Piorek, 1994; Bacon et al., 1995; deBoer et al., 1995; Holyńska et al., 1997; Török et al., 1998; Watson et al., 1999). When a higher energy electron drops into the vacant lower energy orbital, a fluorescent x-ray photon is released. The energy of this photon is unique to each element, and the number of photons is proportional to the concentration of the element. Concentrations are quantified by comparing photon counts for a sample with those obtained from thin-film standards of known concentration (Dane et al., 1996). The previous 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) included a detailed discussion of EDXRF.

Emitted x-rays with energies less than ~4 keV (affecting the elements sodium, magnesium, aluminum, silicon, phosphorus, sulfur, chlorine, and potassium) are absorbed by the filter, by a thick particle deposit, or even by large particles in which these elements are contained. Very thick filters also scatter much of the excitation radiation or protons, thereby lowering the signal-to-noise ratio for XRF and proton (or particle) induced X-ray emission (PIXE). For this reason, thin membrane filters with deposits in the range of 10 to 50 $\mu\text{g}/\text{cm}^2$ provide the best accuracy and precision for XRF and PIXE analysis (Davis et al., 1977; Haupt et al., 1995).

2B.1.2 Synchrotron Induced X-ray Fluorescence (S-XRF)

S-XRF is a form of EDXRF in which the exciting X-rays are generated in a synchrotron. Bremsstrahlung X-rays are generated when energetic electrons (generally several GeV in energy) are forced by a magnetic field to make a bend in their path. The advantages of the technique are that an extremely high flux of X-rays may be obtained and that the X-rays are 100% polarized in the plane of the electron beam. The former allows for X-ray beams generally of 50 to 500 μm in diameter. However, the beams can be focused into X-ray microprobes, with spot sizes on the order of 1 μm diameter. The X-ray polarization allows for removal of most of the background normally found under characteristic X-ray peaks, greatly improving sensitivity compared to other XRF techniques. The primary disadvantages are the limited number of synchrotrons, and that few synchrotrons have S-XRF capabilities. Thus, the technique has been rarely used for PM, and then generally only for special problems, such as for analysis of the smoke from the Kuwaiti oil fires (Cahill et al., 1992; Reid et al., 1994). However, with the increasing availability of S-XRF facilities dedicated to PM analysis (the first of which was the Advanced Light Source opened at Lawrence Berkeley National Laboratory last year), utilization of S-XRF for PM analysis is increasing.

2B.1.3 Proton (or Particle) Induced X-ray Emission (PIXE)

PIXE differs from XRF analysis in the excitation source for producing fluorescence. The filter deposit is bombarded with high-energy protons to remove inner shell electrons, and the resulting characteristic X-rays are analyzed as in XRF (Johansson, 1970; Cahill, 1981, 1985, 1990; Zeng et al., 1993). Small accelerators, generally Van de Graaffs, generate intense beams of low energy protons, usually of a few MeV in energy. These have the ability to remove electrons from the inner shells of atoms of any element. Thus, PIXE can measure a very wide range of elements in a single analysis. The cross section for producing X-rays using protons of a few MeV in energy tends to favor lighter elements, Na through Ca, but sensitivities for equivalent PIXE and multi-wavelength XRF analysis are roughly comparable. The technique has been widely used in the United States (Flocchini et al., 1976; Malm et al., 1994) and around the world, as many universities have the small accelerators needed for the method. Like S-XRF, the proton beams can be focused into μm size beams, but these have not been widely used for PM. However, the mm-size beams used in both S-XRF and PIXE are well suited to analyzing

the limited mass and small deposits that result from detailed particle size measurements by impactors (Perry et al., 1999).

2B.1.4 Proton Elastic Scattering Analysis (PESA)

With the routine availability of elemental analyses for all elements sodium and heavier, organic components remain the major unmeasured species for mass balance. For programs like IMPROVE (Malm et al., 1994), parallel filters are collected for separate OC and EC determinations. Aerosol programs that use PIXE can directly measure hydrogen simultaneously by scattering protons from Teflon filters, which lack hydrogen (Cahill et al., 1989, 1992). After correction for hydrogen in ammonium sulfate and nitrate, the remaining hydrogen can be used to infer the OC mass. Generally, analyses of organic matter by carbon combustion from quartz filters and analysis of organic matter by hydrogen in particles on Teflon filters are in agreement, assuming certain assumptions about the chemical states of sulfates and nitrates are met (Malm et al., 1994; Cahill et al., 1996).

2B.1.5 Total Reflection X-Ray Fluorescence (TRXRF)

One of the limitations of the EDXRF method is its minimum detection limit, which may be high due to high background values (Streit et al., 2000). However, by using implementation of X-ray optical geometries that use the total reflection of the primary radiation on flat surfaces, scattering on the substrate is lessened, thus reducing detection limits. This is the basis for the total reflection X-ray fluorescence (TRXRF) method (Aiginger and Streli, 1997). This modification to the EDXRF technique improves detection limits and eliminates the need to correct for matrix effects. Despite its apparent advantages, TRXRF has not yet become widely used for atmospheric aerosol analysis, although it has been used in the analysis of marine aerosol (Stahlschmidt et al., 1997) and at a high elevation site (Streit et al., 2000). Streit et al. (2000) sampled ambient air at the High Alpine Research Station (3,580 m above sealevel) in the Bernese Alps, Switzerland, using a nine-stage, single-jet, low-pressure, cascade impactor equipped with quartz impactor plates that were coated with silicon oil diluted in 2-propanol. The typical sample volume for a weekly sample was 10 m³. The quartz plates were analyzed directly by TRXRF. Streit et al. reported that the minimum detection limits, defined by the 3 σ values of the blanks, ranged from 25 ng for S, decreased monotonically with increasing atomic number

down to 5 pg for Rb, and decreased after that. The use of TRXRF is expected to increase as EDXRF users become more aware of the method. A relatively low-cost, add-on unit has been developed that would allow EDXRF users to test the TRXRF technique (Aiginger, 1991).

2B.1.6 Instrumental Neutron Activation Analysis (INAA)

INAA irradiates a sample in the core of a nuclear reactor for few minutes to several hours, depending on the elements being quantified (Dams et al., 1970; Zoller and Gordon, 1970; Nadkarni, 1975; Landsberger, 1988; Olmez, 1989; Ondov and Divita, 1993). The neutron bombardment transforms many elements into radioactive isotopes. The energies of the gamma rays emitted by these isotopes identify them and, therefore, their parent elements. Furthermore, the intensity of these gamma rays is proportional to the amount of the parent element present in the sample. Different irradiation times and cooling periods are used before counting with a germanium detector. In source apportionment studies, it is possible to use a combination of XRF and INAA to develop a relatively complete set of elemental measurements. Between these two analytical techniques, good sensitivity is possible for many elements, including most of the toxic metals of interest. In general, XRF provides better sensitivity for some metals (Ni, Pb, Cu, and Fe); whereas INAA provides better sensitivity for others (Sb, As, Cr, Co, Se, and Cd). Both methods provide similar detection limits for still other elements (V, Zn, and Mn). INAA does not quantify some of the abundant species in ambient PM such as silicon, nickel, tin, and lead. While INAA is technically nondestructive, sample preparation involves folding the sample tightly and sealing it in plastic, and the irradiation process makes the filter membrane brittle and radioactive. These factors limit the use of the sample for subsequent analyses.

INAA has been used to examine the chemical composition of atmospheric aerosols in several studies either as the only method of analysis or in addition to XRF (e.g., Yatin et al., 1994; Gallorini, 1995). INAA has higher sensitivity for many trace species, and it is particularly useful in analyzing for many trace metals. Landsberger and Wu (1993) used INAA to analyze air samples collected near Lake Ontario for Sb, As, Cd, In, I, Mo, Si, and V using INAA. They found that using INAA in conjunction with epithermal neutrons and Compton suppression produces very precise values with relatively low detection limits.

Enriched rare-earth isotopes have been analyzed via INAA and used to trace sources of PM from a coal-fired power plant (Ondov et al., 1992), from various sources in the San Joaquin

Valley (Ondov, 1996), from intentionally tagged (iridium) diesel emissions from sanitation trucks (Suarez et al., 1996; Wu et al., 1998), and from iridium-tagged emissions from school buses (Wu et al., 1998).

An intercomparison was conducted in which 18 pairs of filters were sent to participants in the Coordinated Research Program (CRP) on Applied Research on Waste Using Nuclear Related Analytical Techniques (Landsberger et al., 1997). As part of that study, participants used PIXE, INAA, XRF, or AAS to analyze the samples. Many of the results for XRF and PIXE in the coarse fraction were observed to be biased low compared to INAA. The authors speculated that self-attenuation of the X-rays resulting from the particle size effect could create a systematic error.

2B.1.7 Atomic Absorption Spectrophotometry (AAS)

AAS is applied to the residue of a filter extracted in a strong solvent to dissolve the solid material; the filter or a portion of it is also dissolved during this process (Ranweiler and Moyers, 1974; Fernandez, 1989; Jackson and Mahmood, 1994; Chow et al., 2000). A few milliliters of the extract are injected into a flame where the ions are reduced to elements and vaporized. Elements absorb light at certain wavelengths in the visible spectrum, and a light beam with wavelengths specific to the elements being measured is directed through the flame to be detected by a monochromator. The light absorbed by the flame containing the extract is compared with the absorption from known standards to quantify the elemental concentrations. AAS requires an individual analysis for each element, and a large filter or several filters are needed to obtain concentrations for a large variety of elements. AAS is a useful complement to other methods (such as XRF and PIXE) for species such as beryllium, sodium, and magnesium, which are not well quantified by these methods. Airborne particles are chemically complex and do not dissolve easily into complete solution, regardless of the strength of the solvent. Thus, insoluble residues may be left behind, and soluble species may co-precipitate on them or on container walls.

AAS was used to characterize the atmospheric deposition of trace elements Zn, Ni, Cr, Cd, Pb, and Hg to the Rouge River watershed by particulate deposition (Pirrone and Keeler, 1996). The modeled deposition rates were compared to annual emissions of trace elements that were estimated from the emissions inventory for coal and oil combustion utilities, iron and steel

manufacturing, metal production, cement manufacturing, and solid waste and sewage sludge incinerators. They found generally good agreement between the trend observed in atmospheric inputs to the river (dry + wet deposition) and annual emissions of trace elements, with R^2 s varying from ≈ 0.84 to 0.98. Both atmospheric inputs and emissions for Pb showed downward trends. For the period of 1987 to 1992, steady increases were observed for Cd (major sources are municipal solid waste incineration, coal combustion, sludge incineration, and iron and steel manufacturing), Cr and Ni (major sources are iron and steel production and coal combustion), and Hg (major sources are coal, the contribution from which had decreased from 53 to 45%, as well as municipal, solid, and medical waste incineration, the contribution from which has increased).

2B.1.8 Inductively Coupled Plasma with Atomic Emission Spectroscopy (ICP-AES)

ICP-AES introduces an extracted sample into an atmosphere of argon gas seeded with free electrons induced by high voltage from a surrounding Tesla coil (Fassel and Kniseley, 1974; McQuaker et al., 1979; Lynch et al., 1980; Harman, 1989; Tyler, 1992; Baldwin et al., 1994). The high temperatures in the induced plasma raise valence electrons above their normally stable states. When these electrons return to their stable states, a photon of light is emitted that is unique to the element that was excited. This light is detected at specified wavelengths to identify the elements in the sample. ICP-AES can determine a large number of elemental concentrations using small sample volumes with acceptable detection limits for atmospheric samples. As with AAS, this method requires complete extraction and destruction of the sample.

2B.1.9 Inductively Coupled Plasma with Mass Spectroscopy (ICP-MS)

ICP-MS has been applied in the analysis of personal exposure samples (Tan and Horlick, 1986; Gray and Williams, 1987a,b; Nam et al., 1993; Munksgaard and Parry, 1998; Campbell and Humayun, 1999). Ion species generated from ICP and from the sample matrix can produce a significant background at certain masses resulting in the formation of polyatomic ions that can limit the ability of ICP-MS to determine some elements of interest. Cool plasma techniques have demonstrated the potential to detect elements at the ultra-trace level (Nham et al., 1996) and to minimize common molecular ion interferences (Sakata and Kawabata, 1994; Turner,

1994; Plantz, 1996). Detection limits of ICP-MS using a one-second scan are typically in the range of 10^{-3} ng/m³, which is an order of magnitude lower than other elemental analysis methods. The instrument can also be set up to analyze a wide dynamic range of aerosol concentrations. Isotope analysis can also be performed with ICP-MS. Intercomparison studies are needed to establish the comparability of ICP-MS with other nondestructive filter analysis methods.

Keeler and Pirrone (1996) used ICP-MS to determine trace elements Cd, Mn, V, As, Se, and Pb in atmospheric fine particulate (PM_{2.5}) and total suspended particulate samples collected in two Detroit sites. The results were used in a deposition model with additional measurements using AAS to estimate the dry deposition flux of trace elements to Lake Erie.

2B.1.10 Scanning Electron Microscopy (SEM)

Mamane et al. (2001) investigated the use of computer-controlled scanning electron microscopy (CCSEM) as a way of supplementing XRF analysis and providing automated analysis of particle size, chemistry, and particle classification. An ambient coarse particulate sample from Baltimore was collected on a polycarbonate filter for this analysis. CCSEM analyses were conducted for 2,819 particles in 78 randomly selected fields of view during an unattended 8-h run. Mamane et al. (2001) confirmed the stability of the CCSEM instrument over several hours of operation. The physical properties of the sample such as particle diameter, mass loading per field, and particle number per field were well represented by analyzing approximately 360 particles, and little additional information was gained by analyzing more particles. Teflon filters are not well suited for SEM analyses. Unfortunately, analysis of fine PM is expected to pose analytical challenges not addressed in the present study (Mamane et al., 2001).

Nelson et al. (2000) applied Raman chemical imaging and SEM (Raman/SEM) to study the size, morphology, elemental and molecular composition, and molecular structure of fine PM. In their study, filter compatibility was examined, and Raman/SEM chemical imaging was conducted for several standard materials as well as for ambient PM_{2.5} samples. Polycarbonate was determined to be a suitable substrate for both SEM and Raman chemical imaging analysis.

Conner et al. (2001) used CCSEM with individual X-ray analysis to study the chemical and physical attributes of indoor and outdoor aerosols collected around a retirement home in Baltimore. The CCSEM technique was demonstrated to be capable of identifying spherical

particles typical of combustion or other high temperature (presumably industrial) processes as well as pollens and spores. Indoor particles originating from cosmetics were also identified.

2B.2 ORGANIC AND INORGANIC CARBON

Large scale efforts to characterize carbonaceous aerosol require cost effective methods that can analyze samples rapidly. Commercial thermal-optical (TO) instruments were developed to serve this need. The IMPROVE and STN networks have employed these instruments to accumulate large data sets, including measurements taken over the past 18 years by the IMPROVE network and over the past 3 years by STN. In addition to the protocols developed for the IMPROVE and STN networks, a number of alternative TO-based protocols and techniques have been employed by the academic research community. Protocols vary in temperature range and step size, in the duration of heating at each step, in the timing for introduction of oxygen for the conversion of black carbon into CO₂, in the use of a catalyst to reduce pyrolysis, and in the assignment of OC and EC fractions. These operational differences have complicated efforts to compare and combine data sets from studies using different TO protocols.

Thermal-Optical Reflectance

The thermal-optical reflectance (TOR) method of carbon analysis developed by Huntzicker et al. (1982) has been adapted by several laboratories for the quantification of OC and EC in PM collected on quartz-fiber filters. Although the principle applied by these laboratories is identical to that of Huntzicker et al. (1982), the details differed with respect to calibration standards, analysis time, temperature ramping, and volatilization/combustion temperature. The IMPROVE network employs a version of the TOR method for its OC/EC analyses.

In the most commonly used version of the TOR method (Chow et al., 1993), a punch from the filter sample is heated to temperatures ranging from ambient levels to 550 °C in a pure helium atmosphere. In principle, the OC fraction of the PM contained in the filter punch will vaporize, leaving behind only refractory EC. The organic carbon that evolves at each temperature step is first oxidized to CO₂, then converted to methane and finally quantified with a flame ionization detector (FID). The filter punch is incubated at 550 °C for a period sufficient to allow the flame ionization signal to return to its baseline value. The punch is then exposed to a

2% oxygen and 98% helium atmosphere and heated from 550 °C to 800 °C with several temperature ramping steps. The reflectance from the deposit side of the filter punch is monitored throughout the analysis. This reflectance decreases during OC volatilization in the helium atmosphere owing to the pyrolysis of nonvolatile organic material present in the filter punch. When oxygen is added, the reflectance increases as light-absorbing carbon is combusted and removed. It is assumed that the first light-absorbing carbon compounds to combust are pyrolysis artifacts from the first phase of the analysis. Therefore, the OC mass is defined as that fraction that evolves up to the introduction of oxygen plus the oxidized carbon that evolves up to the point when the filter reflectance signal is restored to its preanalysis value. Once the original reflectance level is re-attained, all further carbon evolving from the sample punch is assigned to the EC fraction. Accordingly, “organic carbon” (OC) is carbon that does not absorb light at the laser wavelength (632.8 nm) typically used by TOR instruments, and all other carbon is defined as “elemental carbon” (EC).

Thermal-Optical Transmission

The primary difference between TOR and thermal-optical transmission (TOT) methods is in the choice of absorption detection: light transmission through the filter punch, rather than its reflectance, is monitored throughout the analysis. The TOT method of Birch and Cary (1996) also uses a pure helium atmosphere for volatilizing OC, but the second stage involves a higher oxygen/helium (10%) gas mixture to oxidize the black carbon remaining on the filter punch. The temperature is raised to approximately 820 °C in the helium phase, during which both organic and carbonate carbon are volatilized from the filter. In the second stage, the oven temperature is reduced, then raised to about 860 °C. During this stage, pyrolysis correction and the EC measurement is made. Figure 2B-1 is an example of a TOT thermogram, showing temperature, transmittance, and FID response traces. The peaks that correspond to the concentrations of CO₂ that evolve from the filter punch during the course of the analysis are assigned to OC, carbonate carbon (CC), pyrolytic carbon (PC), and EC. The high temperature in the first stage of the TOT thermal profile is included in order to decompose CC and to volatilize very high-boiling organic compounds. However, many organic carbon compounds will pyrolyze at this temperature to generate PC. The ability to quantify PC is particularly important in high OC/EC regions such as wood smoke-impacted air sheds. Wood smoke aerosol contains many




Figure 2B-1. Thermogram for a sample containing rock dust (a carbonate source) and diesel exhaust, showing three traces that correspond to temperature, filter transmittance, and Flame Ionization Detector (FID) detector response. Peaks correspond to organic (OC), carbonate (CC), pyrolytic (PC), and elemental (EC) carbon. The final peak is a methane calibration peak.

Source: Birch and Cary (1996).

complex compounds that generate substantial PC. Significant error in the EC fraction can result in the absence of a careful PC correction.

Informal intercomparisons among different filter transmission methods have shown high correlations of absorption, but with differences of up to a factor of two in absolute values (Watson et al., 1988a,b). These differences are functions of the type of filter, filter loading, the chemical and physical nature of the deposit, the wavelengths of light used, calibration standards, and light diffusing methods. Currently, there is no agreement on which combination most accurately represents light absorption in the atmosphere.

The National Institute for Occupational Safety and Health (NIOSH) Method 5040 is based on the TOT method (Birch and Cary, 1996). The NIOSH protocol consists of a two-stage process with the first stage being conducted in a pure helium atmosphere at temperatures of 250,

500, 650, and 850 °C for a total of 4.5 min and the second stage conducted in a 2% oxygen/98% helium mix at temperatures of 650, 750, 850, and 940 °C for 4 min. A pyrolysis base correction is made based on the transmittance measurement. The EPA adopted a modified version of the NIOSH Method 5040 protocol for use in its STN.

IMPROVE versus NIOSH protocols

Although the IMPROVE protocol and the NIOSH protocol in use by the STN network yield closely comparable TC values, the differences in thermal profiles and operational definitions of organic versus black carbon result in very different mass assignments to these fractions. Other methodological differences may make data comparisons difficult, including the different choice of absorption detection, e.g. reflectance versus transmission, the different temperature ranges and incubation periods, and the different approaches used to account for background OC. Examples of thermograms obtained with the IMPROVE and NIOSH protocols are shown in Figure 2B-2.

Using both the IMPROVE and NIOSH protocols, Chow et al. (2000) analyzed 60 quartz filter samples on a prototype reflectance/transmittance analyzer that represented a wide variety of aerosol compositions and concentrations. The two TC data sets possessed no statistically significant differences. However, marked differences were found in the fraction of TC that was attributed to EC as determined by the IMPROVE versus NIOSH thermal evolution protocols. The IMPROVE EC measurements were typically higher than the NIOSH EC measurements. When the NIOSH protocol was modified to exclude the helium-only 850 °C temperature step, however, the OC/EC ratios came into agreement between the two methods. Because OC and EC are operationally defined parameters, Chow et al. (2000) pointed out that it is important to retain ancillary information when reporting EC and OC by these analytical methods, so that comparisons can be made among measurements taken at different sites using these two methods.

The NIOSH and IMPROVE protocols both require correction for positive organic artifacts resulting from the absorption of background organic vapor by the heat-treated quartz filters used for OC/EC measurements. Both the IMPROVE and STN science teams have evaluated the presence of carbon artifacts in their measurements. The IMPROVE team has established that heat-treated quartz filters adsorb OC vapors up to a saturation threshold over the course of a few days in the field. The STN science team observed that positive carbon artifacts can vary with




Figure 2B-2. Examples of thermograms obtained by (a) the IMPROVE protocol, and by (b) the NIOSH protocol. Flame ionization detection is used in both examples. FID denotes the observed signal intensity. The intervals denoted ECR/ECT, i.e., elemental carbon reflectance/transmittance, refer to the time and temperature intervals during which evolved CO₂ is assumed to come from oxidized elemental carbon (EC). Laser Reflect and Laser Trans refer to laser reflectance or transmission measurement of filter “blackness” at the He-Ne wavelength, 633 nm.

Source: Birch and Cary (1996).

sampler type. Total carbon artifacts for the samplers used by the STN range from 9.5 to 33.9% of the carbon collected during a typical ambient measurement. Documents describing the issues regarding blank subtraction and the current protocols used by the IMPROVE and STN networks are available on the network websites:

IMPROVE – <http://vista.cira.colostate.edu/improve>

STN – <http://www.epa.gov/ttn/amtic/files/ambient/pm25/spec>

Pyrolytic Carbon (PC) and Other Confounders

In a methods comparison study, Countess (1990) determined that it is necessary to minimize or correct for the PC and OC found in wood smoke and automobile exhaust samples that pyrolyze to create interferences during the course of some thermal-optical protocols. Pyrolysis of organic compounds occurs during analysis by both TO methods, although the greater temperature to which samples are exposed in the NIOSH protocol is likely to produce larger quantities of pyrolysis-derived EC. During the initial heating phase, pyrolysis is indicated by an increase in optical density (blackness) in the filter sample. Both methods distinguish artifact pyrolysis-derived EC from ambient EC at the point at which the transmittance or reflectance signal is restored to the pre-heating level. The assumption made is that heating does not alter the absorption properties of the material collected on the filter. This assumption is reasonable if the only light absorbing species of carbon is strictly a graphite-like EC that is unlikely to undergo a change in its absorption properties over the temperature ranges used by TO methods. The effects of heat-induced changes to the light absorption and chemical properties of atmospheric organic compounds undergoing TO analysis are being evaluated by NIST. NIST has identified three assumptions that must be met in order for TO methods to reliably measure EC: (1) absorptivity of carbonaceous PM remains constant up to the point of pyrolysis; (2) once formed, pyrolyzed carbon (char) absorbs at the analytic wavelength and its absorptivity remains constant within the high temperature step; (3) pyrolyzed OC has the same absorptivity as the EC that is native to the sample. Using urban dust, forest fire emissions, and ambient laboratory aerosol, they observed changes in the absorptivities of these materials during heating before and during the formation of pyrolysis artifacts, up to the OC/EC split point. NIST, therefore, recommends that standard TOR/T protocols be developed that account for these changes.

Light-absorbing carbon or black carbon (BC) includes a large number of compounds that may be altered, not only in their light-absorbing properties but also in their oxidation chemistry. The materials responsible for defining the original blackness of the filter sample may be altered during the initial heating phase of TO analysis, so that the mix present during the oxidation cycle may not be representative of the original, atmospherically derived BC. An error in determining the mass of EC would arise, for example, if a heavy, but weakly light-absorbing, compound were transformed into a material that possessed a higher absorption coefficient and higher resistance to oxidation than absorbing compounds collected from the atmosphere. The pyrolyzed form of this compound would resist oxidation past the OC-EC split point in the analysis, leading to a falsely high EC fraction.

Chow et al. (2000) noted that neither the IMPROVE nor NIOSH methods were able to accurately detect further blackening on optically dense, i.e., very black, filters that are typical of source profile samples. Predictions of artifact pyrolysis carbon by TOR and TOT differed widely. Note that both transmittance and reflectance methods rely on derivations of optics laws (i.e., the Beer-Lambert and Lambert Laws) that predict a linear transmittance/reflectance signal response with species concentration, but only for optically thin samples (Strobel and Heineman, 1989). Very black filters exceed this limitation; thus, the signal response of these methods may not be a linear or otherwise predictable function of BC concentration.

Another important source of error in any TO measurement of aerosol OC/EC arises when samples contain transition metal oxides, such as iron oxide. Many transition metal oxides are found in crustal material. Fung et al. (2002) reported that such oxides can serve as oxidizing agents for BC at high temperatures. The consequence of such an effect is an elevation of the signal usually assigned to OC and corresponding reduction in apparent BC.

Thermal Manganese Oxidation

The thermal manganese oxidation (TMO) method (Mueller et al., 1982; Fung, 1990) uses manganese dioxide in contact with the sample throughout the analysis as the oxidizing agent. Temperature is relied upon to distinguish between OC and EC. Carbon evolving at 525 °C is classified as OC, and carbon evolving at 850 °C is classified as EC. TMO does not correct for pyrolyzed OC, which may lead to the overestimation of EC. This method has been used in the SCENES (Subregional Cooperative Electric Utility, Department of Defense, National Park

Services, and Environmental Protection Agency Study); (Sutherland and Bhardwaja, 1987; Mueller et al., 1986) visibility network, as well as in the SCAQS (Southern California Air Quality Study) (Chow et al., 1994a,b; Watson et al., 1993, 1994a,b).

OC/EC Methods Intercomparisons

Chow et al. (1993) documented several variations of the thermal (T), thermal-optical reflectance (TOR), thermal-optical transmission (TOT), and thermal manganese oxidation (TMO) methods for OC and EC. Comparisons among the results of the majority of these methods show that they yield comparable quantities of TC in aerosol samples, but the distinctions between OC and EC are quite different (Cadle and Groblicki, 1982; Cadle and Mulawa, 1990; Countess, 1990; Hering et al., 1990; Birch, 1998; Schmid et al., 2001). For the sum of OC and EC, these methods reported agreement within 5 to 15% for ambient and source samples (Houck et al., 1989; Kusko et al., 1989; Countess, 1990; Shah and Rau, 1990) and within 3% on carefully prepared standards. Evaluation of these methods is thus a matter of assessing how they differentiate between OC and EC. The EC:TC ratio for samples is method dependent.

An international methods intercomparison study on the analysis of carbonaceous aerosols on quartz fiber filters was organized by the Vienna University of Technology and involved 17 laboratories and 9 different thermal and optical methods (Schmid et al., 2001). All of the participating laboratories were sent punches from three 150-mm quartz-fiber filters that had been exposed for 24 h near a high-traffic street in Berlin. Five laboratories employed the German official standard VDI 2465 methods. Two of these laboratories used the VDI 2465/1 method that determines extractable OC, non-extractable OC, and EC using a solvent-based extraction protocol. Other laboratories participating in the intercomparison used variations of the VDI 2465 standard that rely upon differences in thermal stability to accomplish the separation of carbonaceous aerosol fractions. A range of thermal protocols, TC determination techniques and CO₂ detection schemes were employed by the participating laboratories.

Good agreement of the TC results was obtained by all laboratories, with only two outliers in the complete data set. The relative standard deviations between laboratories for the TC results, were 6.7, 10.6, and 8.8% for the three samples. In contrast, the EC results were much more variable. The relative standard deviations between laboratories for the EC results, were

36.6, 24.4, and 45.5% for the three samples. The VDI methods, especially the VDI 2465/2, were found to give generally higher amounts of EC than the thermal-optical methods. This trend was detected for all samples. The authors recognized that uncorrected thermal methods are prone to positive artifacts by charring during pyrolysis. They also noted that when using solvent extraction methods, the dissolution of polymeric aerosol constituents may not be successful. Both of these effects would lead to an overestimation of the EC fraction. When the laboratories were grouped according to their methods, the relative standard deviation between laboratories was much smaller. This study demonstrated that the TC measurement can yield similar results from a variety of methods, but the EC measurement is highly dependent upon the method used. The problems associated with the determination of EC are exacerbated by the lack of a standard reference material and consistent definitions of EC.

Measuring Black Carbon (BC) Instead of EC

Light-absorbing or black carbon (BC) can be measured by optical techniques (Penner and Novakov, 1996). Both EC and BC define a similar fraction of aerosol; but EC is defined in terms of both the thermal and light-absorption properties of the sample, whereas BC is based on solely on its light-absorption properties. The aethalometer, the integrating sphere sunphotometer, and photoacoustic spectroscopy (described in Section 2B.3) are example techniques for determining BC.

Hitzenberger et al. (1996) investigated the feasibility of using an integrating sphere photometer as an adequate measurement system for the BC content and the absorption coefficient. In another study (Hitzenberger et al., 1999), the integrating sphere method was compared to an aethalometer (Hansen et al., 1984), the thermal method of Cachier et al. (1989), and the thermal-optical method of Birch and Cary (1996). The absorption coefficients that were obtained from both the integrating sphere and the aethalometer were comparable. The BC mass concentrations obtained from the aethalometer were 23% of those obtained from the integrating sphere. Compared to the thermal method, the integrating sphere method overestimated the BC mass concentrations by 21%. Compared to the thermal-optical method, the integrating sphere was within 5% of the 1:1 line. However, the data were not well correlated. Thermal EC and aethalometer BC measurements have also been compared by Lavanchy et al. (1999).

The Carbonaceous Species Methods Comparison Study (CSMCS), as mentioned in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) was conducted in Los Angeles in 1986. Hansen and McMurry (1990) compared the thermal manganese method with the aethalometer for aerosol EC. The first method involved collection of impactor samples backed by a quartz-fiber after-filter followed by EC analysis by oxidation in helium over a MnO₂ catalyst; the other method involved conducting real-time measurements using an aethalometer (an optical absorption technique). They found good agreement between these two very different methods. The CSMCS interlaboratory precision for TC was 4.2% (Turpin et al., 2000). However, because the split between OC and EC is operationally defined, there was substantial interlaboratory variability in OC and EC (e.g., 34% for EC [Turpin et al., 1990]).

EC/OC Summary. With the limitations and precautions described above, laboratory analyses for the carbonaceous properties of collected particles have matured to the point at which they can be performed with commercially available instruments following established standard operating procedures. However, carbon analysis continues to be a subject of active research within the atmospheric sciences community and EPA, and carbon speciation methods comparisons are being undertaken in such studies as the Atlanta Supersite. The state of the art for carbonaceous PM measurements continues to advance; and, although progress is being made, the definitions of OC, EC, and BC continue to be operationally defined in reference to the method employed. Reports of EC/OC measurements should therefore include mention of the method with which the species were determined. Finally, if possible, all ancillary data should be retained, to allow later comparison to other methods.

2B.3 CONTINUOUS METHODS

2B.3.1 Continuous Measurement of Mass

Tapered Element Oscillating Microbalance (TEOM)

The advantages of continuous PM monitoring and the designation of the TEOM as an equivalent method for PM₁₀, have led to the deployment of the TEOM at a number of air monitoring sites. The TEOM also is being used to measure PM_{2.5}. The TEOM differs from the federal reference methods for PM in that it does not require equilibration of the samples at a

specified temperature and RH. The TEOM samples at a constant temperature and is typically heated to some temperature higher than the ambient temperature (Meyer et al., 1995; Meyer and Rupprecht, 1996), whereas the FRM samples at the ambient temperature. Thus, the TEOM does not provide data equivalent to the FRM because of losses of volatile species. Volatilization losses in the TEOM sampler can be reduced by operating the instrument at 30 °C, rather than the typical 50 °C specified, or by using a Nafion diffusion dryer instead of heating to dehumidify the particles (with a 30 °C temperature).

This difference in operation and implications for fine particle measurements was examined by researchers at Commonwealth Scientific and Industrial Research Organization (CSIRO) Atmospheric Research in Australia (Ayers et al., 1999). That group compared 24-h mean $PM_{2.5}$ mass concentrations as determined by a TEOM and by two manual, gravimetric samplers (a low-volume filter sampler and a MOUDI sampler) in four Australian cities, on 15 days in the wintertime. The TEOM was operated at 50 °C at one location and at 35 °C at the other three locations. A systematically lower TEOM response in comparison to the integrated gravimetric methods was observed. In a comprehensive study, Allen et al. (1997) reported results in which TEOM data collected at 10 urban sites in the United States and Mexico were compared with 24-h integrated mass concentrations for both PM_{10} and $PM_{2.5}$. They collected a large data set that included both winter and summer seasons. Allen et al. (1997) concluded that, especially for urban areas, a significant fraction of PM_{10} may be semivolatile compounds that could be lost from the heated filter in the TEOM leading to a systematic difference between the TEOM and the EPA FRM for PM_{10} . They suggested that this difference is likely to be larger for $PM_{2.5}$ than it is for PM_{10} (Allen et al., 1997).

In a similar study conducted in Vancouver, British Columbia the effect of equilibration temperature on PM_{10} concentrations from the TEOM was examined. Two collocated TEOM monitors, operated at 30 and 50 °C, respectively, collected data for a period of ~17 months in the Lower Fraser Valley in British Columbia (Mignacca and Stubbs, 1999). A third TEOM operating at 40 °C was operated for 2 months during this period. They found that, on average, the 1-h average PM_{10} from the TEOM operating at 30 °C was consistently greater than that from the TEOM operated at 50 °C. For the period during which the third TEOM was operated (at 40 °C), the PM_{10} from that instrument was between the values for the other two instruments. They also found that the differences in masses were proportional to the PM_{10} loading, and more

strongly correlated to the PM₁₀ from the TEOM operated at the lower temperature. They recommended that the TEOM monitors be operated at 40 °C in summer and 30 °C in winter, in order to avoid introducing a methodological seasonal bias.

A new sample equilibration system (SES) was developed to reduce losses of semivolatile species from the PM_{2.5} TEOM by conditioning the sample stream to a lower RH and temperature (Meyer et al., 2000). The SES utilizes humidity sensors and a Nafion dryer designed for low particle loss. The dryer fits between the flow splitter that follows the size-selective inlet and the sensor unit. A dry purge gas flows over the exterior of the Nafion tubing and allows for self-regeneration. A TEOM with PM_{2.5} inlet and equipped with an SES was operated at 30 °C alongside another TEOM operating at 50 °C without the SES in Albany, NY, over a 6-day period during a summertime high-temperature, high-relative-humidity episode. The SES maintained the sample air relative humidity under 30%, and the TEOM with the SES generally measured more mass than the other TEOM. The TEOM with SES also was operated alongside an FRM-type sampler for the period of June 6 through September 25, 1999. The correlation between the FRM and TEOM/SES showed a slope of 1.0293 and R² of 0.9352; whereas the correlation between the FRM and the TEOM without SES and operating at 50 °C showed a slope of 0.8612 and R² of 0.8209. The SES can be installed on existing TEOM monitors.

Beta-Gauge Techniques

Beta radiation absorption, as an indicator of particle mass, has been used effectively to measure the mass of equilibrated PM collected on Teflon filters (Jaklevic et al., 1981a; Courtney et al., 1982). The technique also has been used to provide near real-time measurements with time intervals on the order of an hour (Wedding and Weigand, 1993). However, real-time beta-gauge monitors experience the same problems as other continuous or near real-time PM mass monitoring techniques. Particle-bound water (PBW) must be removed to reduce the sensitivity of the indicated mass to relative humidity. However, the simplest technique, mild heating, will remove a portion of the ammonium nitrate and the semivolatile organic compounds as well as the PBW.

An intercomparison study of two beta gauges at three sites indicated that the Wedding beta gauge and the Sierra Anderson (SA) 1200 PM₁₀ samplers were highly correlated, $r > 0.97$ (Tsai and Cheng, 1996). The Wedding beta gauge was not sensitive to relative humidity but yielded

results approximately 7% lower. This suggests that the mild heating in the beta gauge causes losses comparable to those caused by equilibration, although the differences could result from slight differences in the upper cut points. The Kimoto beta gauge that was operated at ambient temperature was sensitive to relative humidity yielding significantly higher mass concentrations relative to the SA 1200 for $RH > 80\%$ than for $RH < 80\%$ even though the correlation with the SA 1200 was reasonable ($r = 0.94$ for $RH > 80\%$ and 0.83 for $RH < 80\%$).

Piezoelectric Microbalance

Piezoelectric crystals have mechanical resonances that can be excited by applying an alternating electrical voltage to the crystal. As the resonance frequencies are well defined, such crystals (quartz in particular) have found applications as secondary time and frequency standards in clocks and watches. As for all mechanical resonators, the resonance frequency is a function of mass. Therefore, by monitoring the resonance frequency in comparison with a second crystal, one can continuously measure the mass deposited on the crystal (Sem et al., 1977; Bowers and Chuan, 1989; Ward and Buttry, 1990; Noël and Topart, 1994). Comparison with a second crystal largely compensates for the effect of temperature changes on the resonance frequency.

The piezoelectric principle has been used to measure particle mass by depositing the particles on the crystal surface either by electrostatic precipitation or by impaction (Olin and Sem, 1971). The collection efficiency of either mechanism has to be determined as function of particle size to achieve quantitative measurements. In addition, the mechanical coupling of large particles to the crystal is uncertain. Both single and multi-stage impactors have been used (Olin and Sem, 1971; Fairchild and Wheat, 1984). Quartz crystals have sensitivities of several hundred hertz per microgram. This sensitivity results in the ability to measure the mass concentration of a typical $100 \mu\text{g}/\text{m}^3$ aerosol to within a few percent in less than one minute (Olin and Sem, 1971).

Coarse Particle Mass

The RAMS and CAMM are only appropriate for fine particle measurements (e.g., $\text{PM}_{2.5}$ or PM_1). However, the TEOM, beta gauge, and piezoelectric microbalance may be used to measure either $\text{PM}_{2.5}$ or PM_{10} (or a sample with any specified upper 50% size cut). A pair of such samplers may be used to measure thoracic coarse PM mass concentration ($\text{PM}_{10-2.5}$) by

difference between the PM_{10} and $PM_{2.5}$ concentrations. However, concerns have been raised concerning the quality of the data from such difference calculations and the resulting potential biases in exposure assessment and risk determinations (Wilson and Suh, 1997; White, 1998). Misra et al. (2001) describe the development and evaluation of a continuous coarse particle monitor (CCPM) that may provide direct measurements of coarse mode PM mass concentrations at short time intervals (on the order of 5-10 min). The basis of the CCPM is enrichment of the coarse particle concentrations through use of virtual impaction while maintaining fine particle concentrations at ambient levels. The resulting aerosol mixture is analyzed using a standard TEOM for which the response is now dominated by the enriched coarse PM mass. The coarse PM concentrations determined from the CCPM were compared to those obtained with a MOUDI, operating with only the 10- and 2.5- μm cut point stages, and a Partisol dichotomous sampler. The CCPM coarse particulate concentrations were highly correlated with both the MOUDI ($R^2 = 0.88$) and the Partisol ($R^2 = 0.88$) coarse PM concentrations. By operating the CCPM at a coarse particle enrichment factor of 25, the coarse PM concentration can be determined a priori without determination of the fine particle concentration, so long as the fine-to-coarse particle concentration ratios are not unusually high (i.e., 4 to 6). Misra et al. (2001) also concluded from field experiments that the coarse particulate concentrations determined from the CCPM were independent of the ambient fine-to-coarse particulate concentration ratio due to the decrease in particle mass median diameter that should accompany fine-to-coarse particulate concentration ratios during stagnation conditions.

2B.3.2 Continuous Measurement of Organic and/or Elemental Carbon

Testing and refinement of models that simulate aerosol concentrations from gas and particle emissions require air quality measurements of approximately 1-h time resolution to reflect the dynamics of atmospheric transport, dispersion, transformation, and removal. Below instruments are described that have been used to collect and analyze atmospheric organic PM with better than 2-h resolution. These instruments were all present at the Atlanta Supersite experiment during the summer of 1999, and an intercomparison of results is underway.

Automated Carbon Analyses

Turpin et al. (1990) describe an in situ, time-resolved analyzer for particulate organic and EC that can operate on a time cycle as short as 90 min. This analyzer collects PM on a quartz-fiber filter mounted in a thermal-optical transmittance carbon analyzer (Turpin et al., 1990). A second quartz-fiber filter behind a Teflon filter in a second sampling port may also be analyzed to provide an estimate of the positive sampling artifact (i.e., gas adsorption on the quartz sampling filter). The organic material in the collected PM is thermally desorbed from the filter at 650 °C and oxidized at 1000 °C over a MnO₂ catalyst bed. The evolved CO₂ is converted to methane over a nickel catalyst, and the methane is measured in a flame ionization detector. Then the EC is oxidized at 350 °C in a 98% He-2% O₂ atmosphere. Correction is made for pyrolytic conversion of some of the organic PM. The instrument was operated with a 2-h resolution during SCAQS in 1987 (Turpin and Huntzicker, 1991, 1995), as well as during CSMCS in 1986 (Turpin et al., 1990). By using EC as a tracer for primary, combustion-generated OC, these authors estimated the contributions of primary sources (i.e., material emitted in particulate form) and secondary sources (i.e., PM formed in the atmosphere) to the total atmospheric particulate OC concentrations in these locations.

An automated carbon analyzer with 15-min to 1-h resolution is now commercially available (Rupprecht et al., 1995) and has been operated in several locations including the Atlanta Supersite. It collects samples on a 0.1 µm impactor downstream of an inlet with a 2.5 µm cut point. Use of an impactor eliminates the gas adsorption that must be addressed when filter collection is used. However, this collection system may experience substantial particle bounce and loss of a sizable fraction of EC since some EC is in particles < 0.2 µm. It is possible that ongoing research, in which particle size is increased by humidification prior to impaction, may result in an improvement in collection efficiency. In the analysis step, carbonaceous compounds are removed by heating in filtered ambient air. Carbonaceous material removed below 340 °C is reported as OC, and material removed between 340 and 750 °C is reported as EC. Turpin et al. (2000) commented that it would be more appropriate to report carbon values obtained by this method as “low-” and “high-temperature” carbon, because some organics are known to evolve at temperatures greater than 340 °C (e.g., organics from woodsmoke).

Aethalometer for Elemental Carbon

As discussed earlier, black carbon (BC), a carbon fraction very similar to EC, is most commonly measured using an aethalometer, a commercially available, automated, time-resolved instrument (i.e., 5- to 15-min sample duration) that measures the light attenuation of aerosol particles collected on a filter tape (Hansen et al., 1984). The concentration of EC is derived from the light absorption measured on a filter using an estimate of the specific absorption (m^2/g) of EC on the filter; the specific absorption value is derived from laboratory and atmospheric tests and is specified by the manufacturer. The specific absorption value could be expected to vary with location, season, and source mix. Comparisons in atmospheric experiments at some locations with EC values measured by thermal methods confirm that the aethalometer provides a statistically meaningful estimate of EC concentration (Allen et al., 1999c; Liousse et al., 1993). For instance, Allen et al. (1999c) found the following statistical relationship for Uniontown, PA during summer 1990: $\text{black carbon (aethalometer)} = 0.95 * \text{EC (thermal)} - 0.2$ ($r^2 = 0.925$, n not specified but appears to be > 50 , EC range from 0 to $9 \mu\text{g}/\text{m}^3$). Another source of error in aethalometer measurements arises from the sampling procedure. Particles are trapped within a three-dimensional filter matrix. Therefore, scattering of transmitted and reflected light may erroneously be attributed to absorption, thus causing errors in the BC calculation. Ballach et al. (2001) investigated immersing the filter in oil of a similar refractive index as a means to minimize the interferences due to light scattering effects from the filter, similar to a procedure common in microscopy. Black carbon measurements determined using the oil immersion technique were compared to those from an integrating sphere, a polar photometer, and Mie calculations. Aerosol tests included several pure carbon blacks from different generating procedures that were used to calibrate the immersion technique, pure ammonium sulfate aerosol, and external and internal mixtures of ammonium sulfate with varying amounts of carbon blacks. The oil immersion technique was also tested on ambient air samples collected at two different sites in the cities of Frankfurt am Main and Freiburg, Germany. Optical measurements, both of blank and loaded filters, showed that the oil immersion technique minimizes scattering losses. Ballach et al. (2001) found that site-related effects were reduced and that there was reasonably good agreement with the other optical techniques as well as with the Mie calculations.

Photoacoustic Measurement of Elemental Carbon

Measurement of aerosol light absorption utilizing photoacoustic spectroscopy has been examined as a continuous method for measuring EC mass concentrations (Petzold and Niessner, 1996; Petzold et al., 1997; Arnott et al., 1999, 2000). Like the aethalometer, this method measures light absorption; however, unlike most other light absorption methods, the photoacoustic technique does not require a filter. The photoacoustic spectrometer of Arnott and coworkers was demonstrated during the Northern Front Range Air Quality Study and compared to an aethalometer (Moosmüller et al., 1998). Neither the aethalometer nor the photoacoustic spectrometer measure EC mass directly. Because the photoacoustic spectrometer measures the absorption coefficient directly, the specific absorption efficiency must be known or assumed in order to determine EC mass. Assuming a light absorption efficiency of $10 \text{ m}^2/\text{g}$, Arnott et al. (1999) reported a lower detection limit for light absorption of $0.4 \text{ M}/\text{m}$ corresponding to a mass concentration of EC of approximately $40 \text{ ng}/\text{m}^3$.

2B.3.3 Continuous Measurements of Nitrate and Sulfate

Nitrate

An integrated collection and vaporization cell was developed by Stolzenburg and Hering (2000) that provides automated, 10-min resolution monitoring of fine-particulate nitrate. In this system, particles are collected by a humidified impaction process and analyzed in place by flash vaporization and chemiluminescent detection of the evolved nitrogen oxides. In field tests in which the system was collocated with two FRM samplers, the automated nitrate sampler results followed the results from the FRM, but were offset lower. The system also was collocated with a HEADS and a SASS speciation sampler (MetOne Instruments). In all these tests, the automated sampler was well correlated to other samplers with slopes near 1 (ranging from 0.95 for the FRM to 1.06 for the HEADS) and correlation coefficients ranging from 0.94 to 0.996.

During the Northern Front Range Air Quality Study in Colorado (Watson et al., 1998), the automated nitrate monitor captured the 12-min variability in fine-particle nitrate concentrations with a precision of approximately $\pm 0.5 \text{ }\mu\text{g}/\text{m}^3$ (Chow et al., 1998). A comparison with denuded filter measurements followed by ion chromatographic (IC) analysis (Chow and Watson, 1999) showed agreement within $\pm 0.6 \text{ }\mu\text{g}/\text{m}^3$ for most of the measurements, but exhibited a discrepancy of a factor of two for the elevated nitrate periods. More recent intercomparisons took place

during the 1997 Southern California Ozone Study (SCOS97) in Riverside, CA. Comparisons with 14 days of 24-h denuder-filter sampling gave a correlation coefficient of $R^2 = 0.87$ and showed no significant bias (i.e., the regression slope is not significantly different from 1). As currently configured, the system has a detection limit of $0.7 \mu\text{g}/\text{m}^3$ and a precision of $0.2 \mu\text{g}/\text{m}^3$.

Sulfate

Continuous methods for the quantification of aerosol sulfur compounds first remove gaseous sulfur (e.g., SO_2 , H_2S) from the sample stream by a diffusion tube denuder followed by the analysis of particulate sulfur (Cobourn et al., 1978; Durham et al., 1978; Huntzicker et al., 1978; Mueller and Collins, 1980; Tanner et al., 1980). Another approach is to measure total sulfur and gaseous sulfur separately by alternately removing particles from the sample stream. Particulate sulfur is obtained as the difference between the total and gaseous sulfur (Kittelson et al., 1978). The total sulfur content is measured by a flame photometric detector (FPD) by introducing the sampling stream into a fuel-rich, hydrogen-air flame (e.g., Stevens et al., 1969; Farwell and Rasmussen, 1976) that reduces sulfur compounds and measures the intensity of the chemiluminescence from electronically excited sulfur molecules (S_2^*).

Because the formation of S_2^* requires two sulfur atoms, the intensity of the chemiluminescence is theoretically proportional to the square of the concentration of molecules that contain a single sulfur atom. In practice, the exponent is between 1 and 2 and depends on the sulfur compound being analyzed (Dagnall et al., 1967; Stevens et al., 1971). Calibrations are performed using both particles and gases as standards. The FPD can also be replaced by a chemiluminescent reaction with ozone that minimizes the potential for interference and provides a faster response time (Benner and Stedman, 1989, 1990).

Capabilities added to the basic system include in situ thermal analysis and sulfuric acid speciation (Cobourn et al., 1978; Huntzicker et al., 1978; Tanner et al., 1980; Cobourn and Husar, 1982). Sensitivities for particulate sulfur as low as $0.1 \mu\text{g}/\text{m}^3$, with time resolution ranging from 1 to 30 min, have been reported. Continuous measurements of particulate sulfur content have also been obtained by on-line XRF analysis with resolution of 30 min or less (Jaklevic et al., 1981b). During a field-intercomparison study of five different sulfur instruments, Camp et al. (1982) reported four out of five FPD systems agreed to within $\pm 5\%$ during a 1-week sampling period.

2B.4 OTHER CONTINUOUS MEASUREMENTS

Continuous Ion Chromatography of Water-Soluble Ions

Two particle collection systems that grow particles by increasing the relative humidity and collect the particles in an aqueous solution suitable for injection into an ion chromatography have been developed independently (Simon and Dasgupta, 1995; Khlystov et al., 1995). Automation of these systems yield semi-continuous monitors for those ions that can be determined using ion chromatography. A similar system using a particle size magnifier has been reported by Weber et al. (2001).

Determination of Aerosol Surface Area in Real Time

Aerosol surface area is an important aerosol property for health effects research. However, methods for on-line measurement of surface area are not widely available. Woo et al. (2001b) used three continuous aerosol sensors to determine aerosol surface area. They used a condensation particle counter (CPC, TSI, Inc., Model 3020), an aerosol mass concentration monitor (MCM, TSI, Inc., Model 8520), and an electrical aerosol detector (EAD, TSI, Inc., Model 3070) to measure integral moments of the particle size distribution. The three sensor signals were inverted to obtain the aerosol size distribution, using a lognormal size distribution model (by minimizing the difference between the measured signals and the theoretical values based upon a size distribution model, the instrument calibration, and its theoretical responses). The lognormal function was then integrated to calculate the total surface area concentration. Woo et al. (2001b) demonstrated that this method can give near real-time measurements of aerosol surface area.

Light Scattering

A variety of types of nephelometers that integrate aerosol light scattering over various solid angles are available (McMurry, 2000). When used to measure visibility, e.g., to provide pilots with real-time data on visual range, it is desirable to include the light scattering due to PBW. However, when used as an indicator of fine particle mass, it is desirable to exclude PBW. This is frequently done by heating the ambient aerosol to a low reference relative humidity of 40%. However, this heating has the potential of also causing the loss of semivolatile components of the aerosol. The evaporation of ammonium nitrate aerosol in a heated nephelometer has been

examined. Bergin et al. (1997) conducted laboratory experiments at low relative humidity (~10%) and as a function of temperature (27 to 47 °C), mean residence time in the nephelometer, and initial particle-size distribution. The evaporation of ammonium nitrate aerosol was also modeled for comparison and was found to describe accurately the decrease in aerosol scattering coefficient as a function of aerosol physical properties and nephelometer operating conditions. Bergin et al. (1997) determined an upper limit estimate of the decrease in the aerosol light scattering coefficient at 450 nm due to evaporation for typical field conditions. The model estimates for their worst-case scenario suggest that the decrease in the aerosol scattering coefficient could be roughly 40%. Under most conditions, however, they estimate that the decrease in aerosol scattering coefficient is generally expected to be < 20%.

3. CONCENTRATIONS, SOURCES, AND EMISSIONS OF ATMOSPHERIC PARTICULATE MATTER

3.1 INTRODUCTION

This chapter discusses topics that were covered in Chapter 5 (Sources and Emissions of Atmospheric Particles) and Chapter 6 (Environmental Concentrations) of the previous document, *Air Quality Criteria for Particulate Matter* or “1996 PM AQCD” (U.S. Environmental Protection, 1996) and updates the subject information based on new data, as available.

Information about concentrations, the composition, and the spatial and temporal variability of ambient particulate matter (PM) across the United States is presented in Section 3.2. Ambient concentration data obtained during the first 3 years of operation of the recently deployed nationwide network of Federal Reference Method (FRM) PM_{2.5} monitors in 27 metropolitan statistical areas (MSAs) are presented and analyzed in Appendix 3A. Initial data from the pilot method evaluation study for the national speciation network are presented in Appendix 3B. Results of field studies that help to characterize the composition of organic compounds in ambient air particles are summarized in Appendix 3C, to complement data on the inorganic composition of ambient particles presented in Appendix 6A of the 1996 PM AQCD and Appendix 3B of this document. Data useful for characterizing the daily and seasonal variability of PM_{2.5} concentrations are discussed in Section 3.2.1, and the intraday variability of PM_{2.5} concentrations is discussed in Section 3.2.2. Relationships among different size fractions are discussed in Section 3.2.3; interrelationships and correlations among PM components are discussed in Section 3.2.4; and the spatial variability of various PM components is discussed in Section 3.2.5.

Atmospheric PM, unlike gaseous criteria pollutants (SO₂, NO₂, CO, O₃) which are well-defined chemical entities, is composed of a variety of particles differing in size and chemical composition. Therefore, sources of each component of the atmospheric aerosol must be considered in turn. Differences in the composition of particles emitted by different sources also lead to spatial and temporal heterogeneity in the composition of the atmospheric aerosol. The nature of the sources and the composition of the emissions from these sources are discussed in Section 3.3. The chemistry of formation of secondary PM from gaseous precursors is

discussed in Section 3.3.1. Estimates of contributions of various sources to ambient PM levels given by source apportionment studies also are presented in Section 3.3.2. More detailed information about the composition of emissions from various sources is given in Appendix 3D. The determination of “policy-relevant background” (PRB) concentrations, including contributions from natural local sources and the long-range transport of PM from sources outside the United States, is discussed in Section 3.3.3. Reviews of transport of PM and its precursors within the United States can be found in the North American Research Strategy for Tropospheric Ozone (NARSTO) *Fine Particle Assessment* (NARSTO, 2002). More detailed information regarding sulfur and nitrogen species can be found in Hidy (1994). Further information about PM concentrations observed at relatively remote monitoring sites (RRMS), i.e., monitoring sites located in relatively remote areas that are not obviously influenced by local anthropogenic sources, is given in Appendix 3E. Because PM is composed of both primary and secondary constituents, emissions of both the primary components and the gaseous precursors of secondary PM are considered here. Nationwide emissions estimates of primary PM and precursors to secondary PM are discussed in Section 3.3.4, and uncertainties in emissions estimates are discussed in Section 3.3.5.

The organization of topics in this chapter (ambient measurements, source characterization and apportionment, and emissions inventories) reflects, in a broad sense, the order in which these topics are addressed in scientific studies and, arguably, the increasing levels of uncertainty that are associated with these different topics.

3.2 PATTERNS AND TRENDS IN AMBIENT PM CONCENTRATIONS

Considerable data exists for characterizing PM_{10} mass concentrations and trends, and those available at the time were presented in the 1996 PM AQCD. In contrast, less extensive data sets were available for characterizing $PM_{2.5}$ and $PM_{10-2.5}$ mass or trends. Sources of data for $PM_{2.5}$ (fine) and $PM_{10-2.5}$ (coarse) discussed in the 1996 PM AQCD include: EPA’s Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2000a); Interagency Monitoring of Protected Visual Environments (IMPROVE; Eldred and Cahill, 1994; Cahill, 1996); the California Air Resources Board (CARB) Data Base (California Air Resources Board, 1995); the Harvard Six-Cities Data Base (Spengler et al., 1986; Neas, 1996); and the

Harvard-Philadelphia Data Base (Koutrakis, 1995). The Inhalable Particulate Network (IPN) (Inhalable Particulate Network, 1985; Rodes and Evans, 1985) provided TSP, PM₁₅, and PM_{2.5} data but only a small amount of PM₁₀ data.

New sources of PM data include the recently deployed nationwide PM_{2.5} compliance-monitoring network, which provides mass measurements using a FRM. This section summarizes data obtained by this network from 1999 to 2001 and provides an approximate characterization of nationwide PM_{10-2.5} concentrations by comparing PM₁₀ to PM_{2.5} data from sites where both types of compliance monitors are located. Various aspects of these data are presented in greater detail in Appendix 3A. In addition, a small number of recent studies in which daily mass and composition measurements are available for extended periods are discussed in this section. The results of quality-assured aerosol composition data obtained by X-ray fluorescence (XRF) and by analyses of organic carbon and elemental carbon for 13 urban areas from the methods evaluation study for the national PM_{2.5} speciation network are presented in Appendix 3B. The terms organic carbon (OC) and elemental carbon (EC) are subject to some ambiguity, and the meanings of these terms as discussed in Section 2.2.7 and Appendix 2.B.2 are applied here.

Organic compounds contribute from 10 to 70% of the dry fine particle mass in the atmosphere (see Appendix 3C). However, concentrations and the composition of organic PM are poorly characterized, as are its formation mechanisms. Particulate organic matter is an aggregate of hundreds of individual compounds spanning a wide range of chemical and thermodynamic properties (Saxena and Hildemann, 1996). Some of the organic compounds are “semivolatile” (i.e., they have atmospheric concentrations and saturation vapor pressures such that both gaseous and condensed phases exist in equilibrium in the atmosphere). The presence of semivolatile or multiphase organic compounds complicates the sampling process. Organic compounds originally in the gas phase may be absorbed on glass or quartz filter fibers and create a positive artifact. Conversely, semivolatile compounds originally present in the condensed phase may evaporate from particles collected on glass, quartz, or Teflon filters creating a negative artifact. In addition, no single analytical technique is currently capable of analyzing the entire range of organic compounds present in atmospheric PM. Rigorous analytical methods are able to identify only 10 to 20% of the organic PM mass on the molecular level (Rogge et al., 1993), and only about 50% of the condensed phase compounds can be identified in smog

chamber studies of specific compounds (Forstner et al., 1997a,b). Measurement techniques and associated artifacts are discussed in Section 2.2.3.2. Information on the identification and concentration of the many different organic compounds identified in atmospheric samples obtained during the 1990s is given in Appendix 3C.

Summary tables giving the results of 66 field studies that obtained data for the composition of particles in the $PM_{2.5}$, $PM_{10-2.5}$, or PM_{10} size ranges were presented in Appendix 6A of the 1996 PM AQCD. The summary tables included data for mass, OC, EC, nitrate, sulfate, and trace elements. Data from the studies were presented for the eastern, western, and central United States. It should be noted that these studies took place at various times and lasted for various durations over a 20-year period, and there may have been significant changes in the concentrations of many species between the times when these studies were conducted and now. These changes resulted from a number of factors (e.g., pollution controls, technological advances, land use changes, etc).

There were a number of discernible differences in the composition of particles across the United States evident in the data sets listed in Appendix 6A in the 1996 PM AQCD (cf., Figures 6-85a to 6-85c). However, these differences can only be discussed in the context of the uncertainties in the measurements of the main components (sulfate, OC, EC, crustal material, ammonium, and nitrate). Sulfate, followed by crustal materials, has the smallest uncertainty associated with its measurement among all the components listed. Sulfate constituted ~38% of $PM_{2.5}$ in the aerosol composition studies in the eastern United States and was the major identifiable component of $PM_{2.5}$. However, it constituted only ~11% of $PM_{2.5}$ in the studies listed for the western United States. The contribution of crustal materials to $PM_{2.5}$ ranged from ~4% in the East to ~15% in the West. The contribution of unidentified material (possibly consisting mostly of water of hydration) ranged ~23% in the East to 0% in the West. The contribution of EC to $PM_{2.5}$ ranged from ~4% in the East to ~15% in the West. Organic compounds constituted ~21% of $PM_{2.5}$ in the eastern United States, reaching to ~39% for the studies listed in the western United States. However, uncertainties for OC, EC, ammonium, and nitrate are larger than for sulfate and crustal material. Concentrations of OC were multiplied by a factor of 1.4 when calculating mass to account for the presence of oxygen, nitrogen, and hydrogen in the organic compounds. This factor may vary among different areas and may represent the lowest reasonable estimate for an urban aerosol (Turpin and Lim, 2001).

In addition, the samples collected in the studies were subject to a number of sampling artifacts involving the adsorption of gases and the evaporation of volatile components that either formed on the filters or were present in the ambient particles. The values reported for OC and EC in filter samples depend strongly on the specific analysis method used (Chow et al., 2002), as discussed extensively in Chapter 2.

Crustal materials constitute from ~52% of $PM_{10-2.5}$ in the eastern United States to ~70% of $PM_{10-2.5}$ in the western United States, as indicated by studies in Appendix 6A of the 1996 PM AQCD. The fraction of unidentified material in $PM_{10-2.5}$ varied from 41% in the eastern United States to 27% in the western United States. However, in the vast majority of these studies, no attempt was made to characterize organic components or nitrate in the $PM_{10-2.5}$ size fraction. It should also be remembered that a small fraction (typically about 10%) of $PM_{2.5}$ is entrained into the flow of the channel of the dichotomous sampler that collects the $PM_{10-2.5}$ sample and that errors may occur during the procedure used to account for this entrainment. Even if analyses of total OC were available, they would not be able to distinguish between primary biological particles (PBPs), which include microorganisms and fragments of living things, and organic compounds of miscellaneous origin in surface deposits on filters. A clear distinction should be made between PBP and primary OC emitted by organisms (e.g., waxes coating the surfaces of leaves, seeds, fruits, stems, pollen, fungi, and insects). Indeed, the fields of view of many photomicrographs of PM samples obtained by scanning electron microscopy are often dominated by large numbers of pollens, plant and insect fragments, and microorganisms. Bioaerosols such as pollens, fungal spores, and most bacteria are expected to be found mainly in the coarse size fraction. However, allergens from pollens can also be found in respirable particles (Monn, 2001; Taylor et al., 2002). Matthias-Maser (1998) summarized information about the size distribution of PBP in and around Mainz, Germany in what is perhaps the most complete study of this sort. Matthias-Maser found that PBP constituted up to 30% of total particle number and volume in the size range from about 0.35 μm to about 50 μm on an annual basis. Additionally, whereas the contribution of PBP to the total aerosol volume did not change appreciably with season, the contribution of PBP to total particle number ranged from about 10% in December and March to about 25% in June and October.

Data for the chemical composition of particles in a number of national parks and remote areas have been collected for a number of years by the IMPROVE network. Concentrations

have been reported for sulfate, nitrate, light-absorbing carbon, OC, and soil components. With the collection of compositional data by the speciation network, more synoptic (i.e., concurrent) coverage will be obtained for these constituents from relatively remote to urban environments across the United States.

PM₁₀ Concentrations and Trends

Nationwide PM₁₀ annual mean concentrations on a county-wide basis from the AIRS database for calendar years 1999, 2000, and 2001 are shown in Figure 3-1a. Concentrations in most areas of the country from 1999 through 2001 were below the PM₁₀ annual standard of 50 µg/m³. Further information about the attainment status of different areas can be found in EPA's Air Quality Trends Reports. The nationwide median value of county-wide annual average PM₁₀ concentrations for this three year period (1999 to 2001) was ~23 µg/m³. Those concentrations flagged as natural events (e.g., resulting from high winds, wildfires, volcanic eruptions) or exceptional events (e.g., construction, prescribed burning) were not included in the calculations. Procedures used for calculating the annual means at the site level followed 40 Code of Federal Regulations (CFR) Part 50 Appendix K (requiring 75% completeness per quarter for all three years). The 98th percentile concentrations from the PM₁₀ monitor showing the highest value in a given county over the three-year period are shown in Figure 3-1b. As shown by the blank areas on the maps, the picture is not complete, because some monitoring locations did not record valid data for all four quarters, or recorded fewer than 11 samples in one or more quarters, or some counties simply did not have monitors. Similar considerations apply to the maps shown later for PM_{2.5} and PM_{10-2.5}. It should also be noted that the area of counties can be much greater in the West than in the East. As a result, the density of monitors may appear to be greater in the West and air quality may appear to be worse over much larger areas in the West than in the East. Concentrations are shown at the county level because this is the typical scale used in many health outcome studies. Metropolitan statistical area (MSA) or multi-county scales have also been used in some number of epidemiologic studies, e.g., Schwartz et al. (1996) or the NMMAPS studies discussed in Chapter 8.

Nationwide trends in annual mean PM₁₀ concentrations from 1992 through 2001 (based on data obtained at 153 rural sites, 297 suburban sites, and 316 urban sites reporting to AIRS) are shown in Figure 3-2 (U.S. Environmental Protection Agency, 2002a). Although average




Figure 3-1a. 1999-2001 county-wide average annual mean PM_{10} concentrations ($\mu g/m^3$) for counties with PM_{10} monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).




Figure 3-1b. 1999-2001 highest county-wide 98th percentile 24-h average PM_{10} concentrations ($\mu g/m^3$) for counties with PM_{10} monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).




Figure 3-2. Nationwide trend in ambient PM₁₀ concentration from 1992 through 2001.

Source: U.S. Environmental Protection Agency (2002a).

concentration levels differ among types of sites with higher levels at urban and suburban sites the nationwide data set shows a decrease of 14% that occurred mainly during the first half of the record. Concentrations at suburban and urban sites show a consistent decline from 1992 to 2001. However, concentrations at rural sites appear to have leveled off since about 1996. Figure 3-3 shows the annual mean PM₁₀ trend summarized by EPA region. Decreases in annual average PM₁₀ concentrations from 1990 to 1999 were largest in the Northwest (9.6 $\mu\text{g}/\text{m}^3$) and smallest in the south-central United States (1.3 $\mu\text{g}/\text{m}^3$). Analyses of available TSP measurements obtained since 1950 indicate that mean TSP concentrations appear to have declined by about 2- to 3-fold in urban areas between 1950 and 1980 (Lipfert, 1998).

PM_{2.5} Concentrations and Trends

Nationwide annual mean PM_{2.5} concentrations obtained from data collected during 1999, 2000, and 2001 are shown in Figure 3-4a; and 98th percentile concentrations are shown in Figure 3-4b. Quantities shown in Figure 3-4a and 3-4b were calculated for individual counties.




Figure 3-3. Trend in PM₁₀ annual mean concentrations by EPA region, 1992 through 2001 (µg/m³).

Source: U.S. Environmental Protection Agency (2002a).

Annual mean concentrations shown in Figure 3-4a were calculated on the basis of the latest 8 consecutive quarters containing at least 11 samples per quarter, and results from the highest monitor were used to produce Figure 3-4b. Data meeting these completeness criteria were obtained at 897 sites in 592 counties. The median PM_{2.5} concentration nationwide was about 13 µg/m³. Annual mean PM_{2.5} concentrations were above 17 µg/m³ at 5% of the sites, mainly in California and in the southeastern United States. The 98th percentile 24-h average concentrations (as defined earlier for PM₁₀) were below 47 µg/m³ at 95% of the sites sampled. Most of the sites with levels above this value are located in California.

Annual average PM_{2.5} concentrations obtained as part of several health studies conducted in various locations in the United States and Canada from the late 1980s to the early 1990s are shown in Figure 3-5 (Bahadori, 2000a). These studies include the Harvard Six-Cities Study (Steubenville, OH; Watertown, MA; Portage, WI; Topeka, KS; St. Louis, MO; and Kingston-




Figure 3-4a. 1999-2001 county-wide average annual mean $\text{PM}_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for counties with $\text{PM}_{2.5}$ monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).




Figure 3-4b. 1999-2001 highest county-wide 98th percentile 24-h average $\text{PM}_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for counties with $\text{PM}_{2.5}$ monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).




Figure 3-5. Collection of annual distribution of 24-h average PM_{2.5} concentrations observed in U.S. and Canadian health studies conducted during the 1980's and early 1990's.

Source:: Bahadori et al. (2000a).

Harriman, TN); Particle Total Exposure Assessment Methodology (PTEAM) study (Riverside, CA); Metropolitan Acid Aerosol Characterization Study (MAACS; Philadelphia, PA; Washington, DC; and Nashville, TN); South Boston Air Quality and Source Apportionment Study (Boston, MA); and NPMRMN (Phoenix, AZ). The remaining sites were part of the 24-Cities Study (Spengler et al., 1996).

Sufficient data are not yet available to permit the calculation of nationwide trends of $PM_{2.5}$ and $PM_{10-2.5}$; however, some general emerging patterns can be discerned. Darlington et al. (1997) proposed that the consistent reductions in PM_{10} concentrations found in a wide variety of environments ranging from urban to rural may have resulted from common factors or controls that affected fine particles more strongly than coarse particles. This is because fine particles have longer atmospheric lifetimes than coarse particles and can be transported over longer distances and, hence, can affect larger areas. Apart from the IMPROVE network of monitoring sites located mainly in national parks, the longest time series of $PM_{2.5}$ concentration and composition data have been obtained by the California Air Resources Board (CARB). Their data show that annual average $PM_{2.5}$ concentrations decreased from 1990 to 1995 by ~50% in the South Coast Air Basin, 35% in the San Joaquin Valley, 30% in the San Francisco Bay Area, and 35% in the Sacramento Valley (Dolislager and Motallebi, 1999). $PM_{2.5}$ data were collected continuously from 1994 to 1998 as part of the Children's Health Study in 12 southern California communities (Taylor et al., 1998). Data obtained at all sites show decreases in $PM_{2.5}$ ranging from 2% at Santa Maria to 37% at San Dimas/Glendora from 1994 through 1998. These decreases were accompanied by decreases in major components such as nitrate, sulfate, ammonium, and acids. Based on the analysis of $PM_{2.5}$ data sets collected prior to 1990, Lipfert (1998) found that $PM_{2.5}$ concentrations appear to have decreased by about 5% per year from 1970 to 1990 in a number of urban areas. These declines were also found to be consistent with decreases in emissions from combustion sources over that time period.

PM_{10-2.5} Concentrations

By using AIRS data for 1999, 2000, and 2001 obtained by the PM_{10} and $PM_{2.5}$ compliance networks, it is possible to construct a picture of the distribution of coarse PM across the country. This is accomplished by pairing data from 208 compliance monitoring sites in 196 counties where PM_{10} and $PM_{2.5}$ monitors are collocated and subtracting the mass concentrations of $PM_{2.5}$

from PM_{10} . Annual mean concentrations were calculated on the basis of the latest 8 consecutive quarters containing at least 11 samples per quarter. Nationwide annual mean $PM_{10-2.5}$ values calculated by this approach are shown in Figure 3-6a. Annual mean $PM_{10-2.5}$ concentrations ranged from ~ 1 to $\sim 50 \mu\text{g}/\text{m}^3$; the nationwide median concentration was $\sim 10 \mu\text{g}/\text{m}^3$; and 5% of the sites had mean concentrations $> 28 \mu\text{g}/\text{m}^3$. The higher values occur mainly in the western United States, particularly in California. The highest county-wide 98th percentile $PM_{10-2.5}$ concentrations based on this same data set are shown in Figure 3-6b. Highest values in the western United States are caused by dust raised locally either by natural means or by anthropogenic activity. It is not clear what the contribution of PBP to these values may be. Elevated dust levels are also found in southern Florida as the result of dust storms in North Africa (Section 3.3.3) and trans-Atlantic transport. In many areas, combined errors in the $PM_{2.5}$ and PM_{10} measurements may be similar to or even greater than the calculated $PM_{10-2.5}$ concentrations. Because of this and other potential problems with this approach (Section 3.2.1), these results should be viewed with caution.

3.2.1 Seasonal Variability in PM Concentrations

PM_{2.5}

Aspects of the spatial and temporal variability of $PM_{2.5}$ concentrations for 1999, 2000, and 2001 in a number of metropolitan areas across the United States are presented in this and following subsections. Data for multiple sites in 27 urban areas across the United States were obtained from the AIRS data base and analyzed for their seasonal variations and for their spatial correlations and spatial uniformity in concentrations. Selection of these 27 MSAs was based on the criteria that data be available for at least 15 days in each calendar quarter of either a 3-year period (1999, 2000, and 2001) or a 2-year period (2000 and 2001) at three or more sites within that MSA. In addition, a maximum of 11 sites per MSA were included for analysis. (In the Chicago and St. Louis MSAs, the 11 sites having the most observations were selected from a greater number of qualifying sites.) A number of aspects of the spatial and temporal variability of the 1999 $PM_{2.5}$ data set were presented in Rizzo and Pinto (2001), based in part on analyses given in Fitz-Simons et al. (2000).

Information regarding the seasonal variability in $PM_{2.5}$ concentrations in four U.S. MSAs (Philadelphia, PA; Cleveland, OH; Dallas, TX; and Los Angeles-Long Beach, CA)




Figure 3-6a. 1999-2000 estimated county-wide average annual mean $PM_{10-2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for counties with collocated $PM_{2.5}$ and PM_{10} monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

is summarized in Figures 3-7a through 3-7d. These four urban areas were chosen to illustrate some general features of the spatial and temporal variability found in the United States. The figures show lowest, lower quartile, median, upper quartile, and highest concentrations for each calendar quarter of 1999, 2000, and 2001 for the Cleveland, OH MSA and for 2000 and 2001 for Philadelphia, Dallas, and Los Angeles MSAs. For each monitoring site, the AIRS ID numbers, annual mean concentrations, the number of observations, and standard deviations are also shown. Data for multiple sites within these MSAs are shown to provide an indication of the degree of inter-site variability. Data for these MSAs and an additional 23 MSAs, criteria used for site selection, and additional descriptions of the data are given in Appendix 3A.





Figure 3-6b. 1999-2000 estimated county-wide highest 98th percentile 24-h $PM_{10-2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for counties with collocated $PM_{2.5}$ and PM_{10} monitors.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

Annual mean $PM_{2.5}$ concentrations at individual monitoring sites in the MSAs examined ranged from ~ 6 to $\sim 30 \mu\text{g}/\text{m}^3$. The lowest values were found in rural portions of the MSAs examined, typically near the perimeter of the MSA. The two highest mean concentrations were found in the Riverside and Los Angeles-Long Beach MSAs in southern California, whereas the three lowest means were found in the Northwest (Portland, OR; Boise, ID; Seattle, WA). These MSAs along the Eastern seaboard (Washington, DC; Philadelphia, PA; Norfolk, VA) tend to have lower mean $PM_{2.5}$ concentrations than those in the north-central United States (Steubenville, OH; Cleveland, OH; Pittsburgh, PA; Chicago, IL; Detroit, MI; Gary, IN; Appendix 3A). Also, average $PM_{2.5}$ concentrations tended to be lower in 1999, 2000, and 2001 in the urban areas given in Appendix 3A compared to concentrations observed during

a. Philadelphia, PA (2000 - 2001)



b. Cleveland, OH (1999 - 2001)





Figure 3-7a,b. Quarterly distribution of 24-h average PM_{2.5} concentrations for selected monitors in the (a) Philadelphia, PA and (b) Cleveland, OH MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual mean concentration (µg/m³), number of observations, and standard deviation of the data are shown above the figures for each site.

c. Dallas, TX (2000 - 2001)



d. Los Angeles, CA (2000 - 2001)




Figure 3-7c,d. Quarterly distribution of 24-h average PM_{2.5} concentrations for selected monitors in the (c) Dallas, TX and (d) Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual mean concentration (µg/m³), number of observations, and standard deviation of the data are shown above the figures for each site.

pollution-health outcome studies conducted in those five urban areas where these overlap (Figure 3-5). It should be noted that there are no data demonstrating the comparability of the monitors used in the studies shown in Figure 3-5 and the FRM.

The patterns of seasonal variability observed in the MSAs examined are complex. In the Philadelphia, PA MSA, highest median concentrations occur at all but one site in the first quarter (cf., Figure 3-7a). First quarter maxima are also observed at all sites in the Detroit, MI and Chicago, IL MSAs (cf., Figures 3A-11 and 3A-14). The Cleveland, OH MSA (Figure 3-7b), in contrast, does not have a clear seasonal pattern. In several MSAs examined in the southern and eastern United States (Atlanta, GA; Baton Rouge, LA; Birmingham, AL; Columbia, SC; and Washington, DC), highest median concentrations occur at all sites during the third calendar quarter (i.e., summer months; cf., Appendix 3A). Sites in Dallas, TX, (Figure 3-7c), as in the other southern cities mentioned above, generally exhibit third quarter median concentration maxima. Highest median concentrations occur during the fourth calendar quarter in MSAs in the western United States as in the Los Angeles, CA MSA (Figure 3-7d), although there are exceptions at individual sites in the Riverside, CA MSA (Figure 3A-26).

Lowest median concentrations occur mainly during the first or fourth quarters at most sites in the eastern United States, as well as in Cleveland, OH and Dallas, TX (Figures 3-7b and 3-7c), whereas some occur during the second quarter (Philadelphia, PA; Figure 3-7a). Moving westward, the seasonal pattern is not as distinct: lowest median concentrations occur in any quarter, but usually in the second or third quarter, as in the Chicago, Detroit, and Los Angeles-Long Beach (Figure 3-7d) MSAs. With the exception of Los Angeles, CA and Riverside, CA, sites in the West show lowest median concentrations in the second quarter. In most of the MSAs examined, seasonal variations follow a similar pattern at all of the sites within the MSA; but in a few MSAs, there are noticeable differences in the seasonal pattern between sites. The large-scale differences in seasonal variability between MSAs tend to follow differences in the major categories of PM sources affecting the monitoring sites. Local heating by wood burning during the colder months is practiced more widely in the western United States than in the eastern United States. Prolonged winter stagnation events are also more common in western mountain valleys during winter than in sections of the eastern United States located in relatively flat topography. These conditions are also conducive to the formation of ammonium nitrate, which is a major and sometimes the dominant contributor to $PM_{2.5}$ in western valleys during the winter

and in southwestern coastal areas during the fall (Kim et al., 2000). Hence, winter maxima and greater variability in $PM_{2.5}$ concentrations across sites are expected in the West due to the influence of these local sources. On the other hand, photochemical production of secondary PM, especially sulfate, occurs over wide areas in relatively homogeneous air masses during the summer months in the eastern United States. Because sulfates (along with associated cations and water) constitute the major fraction of summertime $PM_{2.5}$ in the East, there is greater uniformity in third quarter PM concentrations within eastern MSAs (cf., Appendix 3A).

Maximum 24-h average concentrations shown in the box plots in Figures 3-7a to 3-7d and in Figures 3A-1 to 3A-27 do not necessarily follow the same seasonal pattern as the median concentrations. There is no clear relationship between the maximum and the median concentrations evident in the Philadelphia, PA data set (Figure 3-7a). In Cleveland, OH (Figure 3-7b), maximum concentrations occur during the second or fourth quarters, and highest median concentrations generally occur during the third or first quarters. In Dallas, TX (Figure 3-7c), maximum concentrations generally occur during the fourth quarter, but highest median concentrations tend to occur during the second or third quarter. In the Los Angeles-Long Beach MSA (Figure 3-7d), the maximum and highest median concentrations occur together in the fourth quarter with the exception of one site. Peak individual concentrations likely reflect the occurrence of transient events such as forest fires (mainly in the West) or episodes of secondary PM production (mainly in the East). However, chemical analyses of filter samples or other evidence should be used to determine specific causes in particular locations.

There have been a few studies that have characterized $PM_{2.5}$ and PM_{10} concentrations in major urban areas. The Metropolitan Aerosol Acidity Characterization Study (MAACS) (Bahadori et al., 2000b) characterized the levels and the spatial and temporal variability of $PM_{2.5}$, PM_{10} , and acidic sulfate concentrations in four cities in the eastern United States (Philadelphia, PA; Washington, D.C.; Nashville, TN; and Boston, MA). Seasonal variations in $PM_{2.5}$ and PM_{10} concentrations obtained during the course of this study are shown in Figure 3-8. The data for the four cities included in MAACS are presented in a box plot showing the lowest, lowest tenth percentile, lowest quartile, median, highest quartile, highest tenth percentile, and highest $PM_{2.5}$ and PM_{10} values. Mean and highest $PM_{2.5}$ and PM_{10} concentrations peak during the summer in all four cities (in contrast with Figure 3-7a), although the seasonal pattern in Boston appears to be more nearly bimodal with an additional winter peak.




Figure 3-8. Seasonal concentrations of PM_{2.5} and PM₁₀ measured in the four MAACS cities. The data show the lowest, lowest tenth percentile, lowest quartile, median highest quartile, highest tenth percentile, and highest PM_{2.5} 24-h average values. The dashed line shows the level of the annual PM_{2.5} standard. SP = spring; SU = summer; F = fall; W = winter.

Source: Bahadori et al. (2000b).

PM_{10-2.5}

Of the 27 MSAs selected for analysis of PM_{2.5} data (Appendix 3A), 17 have multiple sites satisfying the PM_{10-2.5} data completeness criteria. A subset of 53 qualifying sites were selected for analysis of PM_{10-2.5} data. Each qualifying site has 11 or more observations per calendar quarter for 12 consecutive quarters (1999 to 2001, 2 MSAs), eight quarters (1999 + 2000 or 2000 + 2001, 7 MSAs) or four quarters (2000 or 2001, 8 MSAs). In addition, data from sites within the same MSA represent the same year (or years) of observations, so that intersite comparisons within an MSA are based on the same reporting period. However, comparisons between different MSAs can involve different annual reporting periods. As can be seen by

comparing Figures 3-7a,b,c,d and Figures 3-9a,b,c, the number of observations used for summarizing $PM_{10-2.5}$ data is much lower than that measured for $PM_{2.5}$. Unfortunately, fewer qualifying MSAs for $PM_{10-2.5}$ result in geographical coverage that is more spotty than with $PM_{2.5}$. Five of the seven MSAs representing the eastern United States for $PM_{2.5}$, for example, did not meet completeness criteria for $PM_{10-2.5}$, leaving only Tampa, FL and Columbia, SC to represent the East. Secondly, it can readily be seen, e.g., for Cleveland, OH (Figure 3-9a) and, also, in Figures 3A-28 through 3A-44, that a number of $PM_{10-2.5}$ concentrations are negative. For example, in 2001, about one-fourth of all $PM_{10-2.5}$ concentrations at the three qualifying sites in the Gary, IN MSA were negative. (The negative estimates have been included in the calculation of mean concentrations.) There are a number of reasons for the negative concentration estimates, many of which arise because the ratios of $PM_{2.5}$ to PM_{10} are based on two independent measurements. Measurement imprecision plays a role when the ratios are large and concentrations are small. Differences in the behavior of semivolatile components in the two samplers could occur; and the results may also reflect errors in sampler placement, field, laboratory, or data processing procedures. Thus, caution should be exercised when attempting to interpret results for $PM_{10-2.5}$ based on current network collocated $PM_{2.5}$ and PM_{10} monitors.

Annual mean $PM_{10-2.5}$ concentrations at individual monitoring sites range from $\sim 6 \mu\text{g}/\text{m}^3$ (Portland, OR) to $\sim 33 \mu\text{g}/\text{m}^3$ (Riverside, CA). (Gary, IN sites were excluded because of numerous negative $PM_{10-2.5}$ concentrations, and one site in Riverside, CA affected by a local dust event was excluded.) The three highest annual mean concentrations are observed in Riverside, CA; Los Angeles-Long Beach, CA; and Salt Lake City, UT; and the lowest (excluding Gary) were observed in Portland, OR. For the remaining MSAs, there does not appear to be a significant geographical trend associated with the annual mean $PM_{10-2.5}$ concentration. Within MSAs, the lowest concentrations are frequently observed at sites near the perimeter of the MSA, although it must be noted that the number of sites is limited.

Within each MSA, collocated $PM_{2.5}$ and $PM_{10-2.5}$ concentrations, averaged over the same years, were compared. The mean $PM_{2.5}:PM_{10-2.5}$ ratio was calculated for each of the 17 MSAs using as many sites as possible. The median mean $PM_{2.5}:PM_{10-2.5}$ ratio for the 17 MSAs was 1.2. For eight MSAs (Tampa, FL; Columbia, SC; Louisville, KY; Chicago, IL; Gary, IN; Milwaukee, WI; Steubenville, OH; Portland, OR), the mean $PM_{2.5}$ concentration exceeded the mean $PM_{10-2.5}$ concentration. For an additional eight MSAs, the $PM_{2.5}$ and $PM_{10-2.5}$ concentration means were





Figure 3-9a,b. Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the (a) Cleveland, OH; (b) Dallas, TX MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.




Figure 3-9c. Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations for selected sites in the Los Angeles, CA MSAs. Values for the lowest, lower quartile, median, upper quartile and highest concentrations are shown in the figures. The AIRS site ID number, annual concentration, number and standard deviation are shown above the figures for each site.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

the same (within one SD). Salt Lake City was the only MSA for which the mean $PM_{10-2.5}$ concentration exceeded the mean $PM_{2.5}$ concentration.

Information regarding the seasonal variability in $PM_{10-2.5}$ concentrations in three MSAs (Cleveland, OH; Dallas, TX; Los Angeles-Long Beach, CA) is summarized in Figures 3-9a through 3-9c. The figures show lowest, lower quartile, median, upper quartile, and highest concentrations for each calendar quarter of 2000 and 2001 for the Cleveland, OH MSA (six sites); and of 2001 for the Dallas, TX and Los Angeles, CA MSAs (four sites each). Data for

these and an additional 13 MSAs, as well as additional descriptions of the data, are given in Appendix 3A. The seasonal pattern for $PM_{10-2.5}$ median concentrations is different from that of $PM_{2.5}$ (Figures 3-7a,b,c,d, and Appendix 3A). Most seasonal median maxima in $PM_{10-2.5}$ occur during the second or third calendar quarters, i.e., spring and summer months (45% and 36% of all sites, respectively) as seen, for example, at most sites in Figures 3-9. Contributions from bioaerosols during the spring blooming season, which mostly affect $PM_{10-2.5}$ concentrations, might be a significant contributing factor in the second quarter $PM_{10-2.5}$ median maxima in some regions of the United States. Lowest median concentrations occur mainly in the first or fourth quarters (62% and 23% of all sites, respectively). Cleveland, OH (Figure 3-9a) and Tampa, FL, where lowest medians generally are observed in the third quarter, are exceptions (Figure 3A-29). At no site was the lowest median $PM_{10-2.5}$ concentration observed in the second quarter. In comparison to $PM_{10-2.5}$, seasonal $PM_{2.5}$ median maxima mostly occurred in the first or third quarters and $PM_{2.5}$ median minima are mostly observed in the second and fourth quarters. With few exceptions, collocated $PM_{2.5}$ and $PM_{10-2.5}$ seasonal medians at individual sites peak in different quarters. Likewise, at a given site, the lowest median concentrations for $PM_{2.5}$ and $PM_{10-2.5}$ rarely occur in the same quarter. In MSAs with several $PM_{10-2.5}$ sites, the seasonal patterns are typically seen at all sites within the MSA. In the Dallas, TX MSA, the maximum and minimum estimated $PM_{10-2.5}$ concentrations both occur in the first quarter at all four sites.

The ratio of $PM_{2.5}:PM_{10-2.5}$ seasonal median concentrations peak in the first quarter for MSAs in the central and north-central United States and tend to peak in the fourth quarter for western states. The largest ratios of $PM_{2.5}:PM_{10-2.5}$ seasonal median concentrations are observed in the central and north-central MSAs (Chicago, Cleveland, Detroit, and St. Louis), whereas smaller ratios are found in the western and southern United States.

As can be seen from Figure 3-9a and Figures 3A-40 and 3A-43, the maxima for $PM_{10-2.5}$ concentrations exceeded $100 \mu\text{g}/\text{m}^3$ in Cleveland, OH and Salt Lake City, UT, and $500 \mu\text{g}/\text{m}^3$ in Riverside, CA. This latter value was related to a dust storm. In several urban areas (Cleveland, OH; Detroit, MI; Chicago, IL; Dallas, TX; and Riverside, CA) maxima in $PM_{10-2.5}$ concentrations were larger than those for $PM_{2.5}$. However, there is no clear geographic pattern.

The results described above should be viewed with caution because of inherently large errors in the technique used to derive them.

Frequency Distributions for PM_{2.5} Data

Frequency distributions for PM_{2.5} concentrations obtained in Philadelphia from 1992 through 1995 are shown in Figure 3-10 (data obtained by Bahadori et al., 2000b). Also shown are concentrations predicted from the log-normal distribution, using geometric mean values and standard deviations derived from the data. In Philadelphia, the highest PM_{2.5} values were observed when winds were from the southwest during sunny but hazy high pressure conditions. In contrast, the lowest values were found after significant rainstorms during all seasons of the year. Mean \pm SD day-to-day concentration differences in the data set are $6.8 \pm 6.5 \mu\text{g}/\text{m}^3$ for PM_{2.5} and $8.6 \pm 7.5 \mu\text{g}/\text{m}^3$ for PM₁₀. Maximum day-to-day concentration differences are $54.7 \mu\text{g}/\text{m}^3$ for PM_{2.5} and $50.4 \mu\text{g}/\text{m}^3$ for PM₁₀.




Figure 3-10. Frequency distribution of 24-h average PM_{2.5} concentrations measured at the Presbyterian home (PBY) monitoring site in southwestern Philadelphia from 1992 to 1995. Log-normal distribution fit to the data shown as solid line.

Source: Bahadori et al. (2000b).

Different patterns are observed in data collected elsewhere in the United States. $PM_{2.5}$ concentrations obtained in Phoenix, AZ from 1995 through 1997 (Zweidinger et al., 1998) are summarized in Figure 3-11; and frequency distributions of $PM_{2.5}$ concentrations obtained in Phoenix are shown in Figure 3-12. Mean \pm SD day-to-day concentration differences in this data set are $2.9 \pm 3.0 \mu\text{g}/\text{m}^3$; the maximum day-to-day concentration difference was $23 \mu\text{g}/\text{m}^3$. $PM_{2.5}$ and $PM_{10-2.5}$ data were obtained with dichotomous samplers at a number of sites in California on a sampling schedule of every 6 days from 1989 through 1998. Histograms showing the frequency distribution of the entire set of $PM_{2.5}$ and $PM_{10-2.5}$ concentrations obtained by the CARB network of dichotomous samplers from 1989 to 1998 are shown in Figures 3-13 and 3-14. Also shown are log-normal distributions generated by using geometric means and standard deviations derived from the data as input. Although the data for both size fractions appear to be reasonably well simulated by the function, data obtained at individual locations may not be. Data showing the seasonal variability of $PM_{2.5}$ obtained at Riverside-Rubidoux are summarized in box plot form in Figure 3-15. The frequency distribution of $PM_{2.5}$ concentrations obtained at Riverside-Rubidoux from 1989 to 1994 is shown in Figure 3-16. It can be seen that the data are not as well fit by a log-normal distribution as are data shown in Figure 3-10, partly as the result of a significant number of days when $PM_{2.5}$ values exceed $100 \mu\text{g}/\text{m}^3$.




Figure 3-11. Concentrations of 24-h average $PM_{2.5}$ measured at the EPA site in Phoenix, AZ from 1995 to 1997. The data show the lowest, lowest tenth percentile, lowest quartile, median (black circles), highest quartile, highest tenth percentile, and highest $PM_{2.5}$ values.

Source: Zweidinger et al. (1998).




Figure 3-12. Frequency distribution of 24-h average PM_{2.5} concentrations measured at the EPA site in Phoenix, AZ from 1995 to 1997.

Source: Zweidinger et al. (1998).




Figure 3-13. Frequency distribution of 24-h average PM_{2.5} measurements obtained from all California Air Resources Board dichotomous sampler sites from 1989 to 1998.




Figure 3-14. Frequency distribution of 24-h average PM_{10-2.5} concentrations obtained from all California Air Resource Board Dichotomous sampler sites from 1989 to 1998.




Figure 3-15. Concentrations of 24-h average PM_{2.5} measured at the Riverside-Rubidoux site from 1989 to 1998. The data show the lowest, lowest tenth percentile, lowest quartile, median (black squares), highest quartile, highest tenth percentile, and highest PM_{2.5} values.




Figure 3-16. Frequency distribution of 24-h average PM_{2.5} concentrations measured at the Riverside-Rubidoux site from 1989 to 1994.

An examination of the data from the four MAACS cities, Phoenix, AZ, and Riverside, CA, indicates that substantial differences exist in aerosol properties between the eastern cities (MAACS) and the western cities (Phoenix, AZ; Riverside, CA). Fine-mode particles account for most of the PM₁₀ mass observed in the MAACS cities and appear to drive the daily and seasonal variability in PM₁₀ concentrations in the East, whereas coarse-mode particles represent a larger fraction of PM₁₀ mass in Phoenix and Riverside and drive the seasonal variability in PM₁₀ seen in the West. The average ratio of PM_{2.5} to PM₁₀ concentrations is much larger in the MAACS cities of Philadelphia, PA (0.72), Washington, DC (0.74), and Nashville, TN (0.63) than in either Phoenix, AZ (0.34) or Riverside, CA (0.49). Differences between median and maximum concentrations in any size fraction are much larger at the Riverside site than at either the MAACS or Phoenix sites. Many of these differences could reflect the more sporadic nature of dust suspension at Riverside. In addition, the seasonal variability of PM_{2.5} concentrations seen in Phoenix and Riverside appears to be different from that observed in the MAACS cities. These considerations demonstrate the hazards in attempting to extrapolate conclusions about the nature of variability in aerosol characteristics inferred at one location to another.

3.2.2 Diurnal (Circadian) Variability in PM Concentrations

The variability of PM concentrations on time scales shorter than a day can, in principle, be characterized by measurements made by continuous samplers (e.g, TEOMs and β -gauge monitors that are currently used to provide Air Quality Index [AQI] information to the public). A description of these methods was provided in Section 2.2.9. However, as shown in Chapter 2, continuous methods are subject to artifacts in large part because of the heating of their inlets to remove water, which results in the loss of components such as ammonium nitrate and, also semivolatile organic compounds (see Sections 2.2.2.1 and 2.2.3 for further details concerning the chemistry of volatilizable components). Consequently, caution should be used in interpreting results obtained by these techniques. It should be remembered that the FRMs are also subject to artifacts; therefore, caution should also be exercised in interpreting results obtained by them.

The composite diurnal variation of $PM_{2.5}$ concentrations obtained across the continental United States by 31 TEOM and β -gauge monitors reporting to AIRS in 1999 is shown in Figure 3-17. As can be seen in that figure, there is a distinct pattern with maxima occurring during the morning and evening. Notable exceptions to this pattern occur in California, where broad nighttime maxima and daytime minima occur, which may be related to the use of β -gauges with unheated inlets there. It should be noted that, in examining the diurnal variations shown in Figure 3-17, there is substantial day-to-day variability in the diurnal profile of $PM_{2.5}$ measured at the same location, which is smoothed out after a suitably long averaging period is chosen. The large ratio of the interquartile range to the median values supports the view that there is substantial variability in the diurnal profiles.

The diurnal variability of PM components is determined by interactions between variations in emissions, the rates of photochemical transformations, and the vertical extent and intensity of turbulent mixing near the surface. Wilson and Stockburger (1990) characterized the diurnal variability of sulfate and lead (Pb) in Philadelphia. At that time, Pb was emitted mainly by motor vehicles. Pollutants emitted mainly by motor vehicles, such as carbon monoxide (CO), show two distinct peaks occurring during the morning and evening rush hours (see Chapter 3, U.S. Environmental Protection Agency, 2000b). Pollutants, such as sulfate, which are transported long distances in the free troposphere (i.e., above the planetary boundary layer), tend to be mixed downward and have their highest concentrations during the afternoon when the intensity and vertical extent of turbulent mixing (and chemical oxidation) are greatest.




Figure 3-17. Intraday variability of hourly average PM_{2.5} concentrations across the United States. Interquartile ranges, median and mean (+) values are shown. Values above the box plots refer to the number of observations during 1999.

Source: Fitz-Simons et al. (2000).

Secondary aerosol components (such as secondary organic compounds) that are produced by photochemical reactions may have a daily maximum in the afternoon, similar to ozone. PM concentrations produced by residential heating (e.g., from wood burning), on the other hand, reach maximum levels during the night when mixing heights are much lower than during the day.

Although the interquartile ranges for hour-to-hour changes in PM_{2.5} concentrations shown in Figure 3-17 encompass several µg/m³, extreme values for the hour-to-hour variations can be much larger (Fitz-Simons et al., 2000). The 98th percentile values for positive and negative excursions in concentration are all less than 20 µg/m³. Maximum positive excursions were much larger, ranging from 27 µg/m³ in the Northeast up to 220 µg/m³ in the Southwest and with maximum excursions in other regions all less than 125 µg/m³. It should be borne in mind that the hour-to-hour changes that are reported reflect the effects of a number of processes occurring during passage through the sampler inlets and on the TEOM measurement elements. These

factors add uncertainty to the interpretation of the hour-to-hour changes that are observed, as discussed in Chapter 2. However, because of the tendency of these monitoring instruments to lose material by evaporation, the concentrations reported during excursions probably represent lower limits for the true values that were present.

3.2.3 Relationships Among Particulate Matter in Different Size Fractions

Relations Among $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10}

Data obtained in 1999 by collocated $PM_{2.5}$ and PM_{10} FRM monitors have been used to calculate the ratio of $PM_{2.5}$ to PM_{10} concentrations and correlations among $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10} concentrations. Results are shown in Table 3-1 for each of the seven aerosol characteristic regions identified in Chapter 6 of the 1996 PM AQCD. As can be seen from the table, the ratio of $PM_{2.5}$ to PM_{10} concentrations tends to be higher in the eastern United States than in the western United States. This general pattern and the values are consistent with that found for the studies included in Appendix 6A of 1996 PM AQCD. In that compilation based on the results of studies using dichotomous samplers, the mean ratio of $PM_{2.5}$ to PM_{10} was 0.75 in the East, 0.52 in the central United States, and 0.53 in the western United States. Although a large number of paired entries have been included in Table 3-1, seasonal variations and annual averages in a number of regions could not be determined from the data set because of data sparseness, mainly during the early part of 1999. It also can be seen in Table 3-1 that the ratio of $PM_{2.5}$ to PM_{10} was > 1 for a few hundred measurements. There are a number of reasons for these results, as mentioned in Section 3.2.1 in the discussion on $PM_{10-2.5}$ concentrations.

Ultrafine Particle Concentrations

Data for characterizing the concentrations of ultrafine particles ($< 0.10 \mu\text{m } D_a$) and the relations between ultrafine particles and larger particles are sparse. Although ultrafine particles dominate particle number concentrations, they make very minor contributions to $PM_{2.5}$ mass. For example, Cass et al. (2000) found that particles between 0.056 and 0.1 $\mu\text{m } D_a$ contributed only 0.55 to 1.16 $\mu\text{g}/\text{m}^3$ at several sites in southern California. Perhaps the most extensive data set for ultrafine particle properties is that described by Woo et al. (2001) for a site located 10 km to the northwest of downtown Atlanta, GA. Size distributions from 3 to 2000 nm were measured every 12 min for 24 months beginning in August 1998. Approximately 89% of the total number

TABLE 3-1. DISTRIBUTION OF RATIOS OF PM_{2.5} TO PM₁₀ AND CORRELATIONS BETWEEN PM_{2.5} AND PM₁₀, PM_{2.5} AND PM_{10-2.5}, AND PM_{10-2.5} AND PM₁₀ FOUND AT COLLOCATED MONITORING SITES IN SEVEN AEROSOL CHARACTERISTIC (EPA/HEI) REGIONS IN 1999

Region	Mean	Sites	Values	Percentiles							Correlations		
				95	90	75	50	25	10	5	PM _{2.5} :PM ₁₀	PM _{2.5} :PM _{10-2.5}	PM _{10-2.5} :PM ₁₀
Northeast	0.70	45	1433	0.97	0.95	0.77	0.67	0.60	0.51	0.48	0.72 ^a	0.02	0.71 ^a
Southeast	0.70	76	2823	1.27	1.06	0.74	0.63	0.54	0.46	0.43	0.69 ^a	-0.04 ^a	0.69 ^a
Industrial Midwest	0.70	92	4827	1.09	0.88	0.78	0.68	0.59	0.51	0.47	0.71 ^a	0.17 ^a	0.81 ^a
Upper Midwest	0.53	39	1446	0.92	0.84	0.62	0.49	0.44	0.34	0.24	0.35 ^a	-0.02	0.93 ^a
Southwest	0.38	23	701	0.51	0.51	0.47	0.40	0.31	0.23	0.23	0.63 ^a	0.49 ^a	0.99 ^a
Northwest	0.50	73	3300	0.67	0.65	0.56	0.49	0.44	0.39	0.36	0.69 ^a	0.07 ^a	0.77 ^a
Southern California	0.47	36	1813	0.70	0.57	0.55	0.48	0.44	0.31	0.24	0.70 ^a	0.19 ^a	0.83 ^a
Total:		384	16343										

^a Results considered to be significantly different from zero at the $\alpha = 0.01$ level.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

of particles were found to be smaller than 100 nm; whereas 26% were < 10 nm. Concentrations tend to be lower during the summer than during the winter. No correlation was found between number concentration and either volume or surface area for particle sizes up to 2 μm . Because the total number of particles is concentrated in the smallest size ranges, these results also indicate that fine particle mass does not correlate with the number of ultrafine particles. The high time resolution of the measurements allows some inferences to be made about the possible sources of the ultrafine particles. The number of particles > 10 nm tends to peak during the morning rush hour around 8:00 am and then to decrease through the day and to increase again after 6:00 pm, consistent with a traffic-related source. Particles < 10 nm tend to peak during the mid-afternoon, consistent with nucleation involving products of active photochemistry (McMurry et al., 2000). More direct relationships between particle mass observed in different size ranges can be obtained using multi-stage impactors. Keywood et al. (1999) found a correlation between $\text{PM}_{2.5}$ and $\text{PM}_{0.15}$ of ~ 0.7 , whereas they found correlations of ~ 0.96 between PM_1 and $\text{PM}_{2.5}$ and between $\text{PM}_{2.5}$ and PM_{10} based on samples collected by MOUDIs (Multiple Orifice Uniform Deposit Impactors) in six Australian cities.

3.2.4 Relationships Between Mass and Chemical Component Concentrations

Time series of elemental composition data for $\text{PM}_{2.5}$ based on X-ray fluorescence (XRF) analyses have been obtained at a number of locations across the United States. Time series of components of the OC fraction of the aerosol have not yet been obtained. The results of XRF analyses for the composition of the inorganic fraction of $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ are presented in Table 3-2 for Philadelphia, PA and in Table 3-3 for Phoenix, AZ. Frequency distributions for $\text{PM}_{2.5}$ concentration data collected at these sites were shown in Figures 3-10 and 3-11. All XRF analyses were performed at the same U.S. EPA X-ray spectrometry facility in Research Triangle Park, NC. Data shown in the first column of Table 3-2 are based on analyses of filters collected over 3 years (April 1992 to April 1995, labeled *a*) at the Castor Avenue Laboratory at the PBY site in southwestern Philadelphia. These data and data for PM_{10} were collected using Harvard impactors. Data for $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ shown in the second and third columns were obtained by the City of Philadelphia from July 25 to August 14, 1994, using a modified dichotomous sampler (VAPS). The samples at the Phoenix site were collected in 1996 and 1997 using the same type of dichotomous sampler used in the shorter term study in Philadelphia.

TABLE 3-2. CONCENTRATIONS (ng/m³) OF PM_{2.5}, PM_{10-2.5}, AND SELECTED ELEMENTS (ng/m³) IN THE PM_{2.5} AND PM_{10-2.5} SIZE RANGES WITH STANDARD DEVIATIONS (SD) AND CORRELATIONS BETWEEN ELEMENTS AND PM_{2.5} MASS IN PHILADELPHIA, PA*

n = 1105	Conc (ng/m³) ± SD (unc)	r	n = 20	Conc (ng/m³) ± SD (unc)	r	n = 20	Conc (ng/m³) ± SD (unc)	r
PM _{2.5} ¹	17 ± 0.9 (0.8) × 10 ³	1	PM _{2.5} ²	29.8 ± 14.7 (1.1) × 10 ³	1	PM _{10-2.5} ²	8.4 ± 2.9 (0.4) × 10 ³	1
Al	4.0 ± 56 (31)	0.1	Al	109 ± 61 (21)	0.15	Al	325 ± 241 (99)	0.89
Si	116 ± 107 (21)	0.51	Si	191 ± 134 (26)	0.22	Si	933 ± 652 (231)	0.9
P	8.6 ± 14 (10)	0.31	P	15 ± 4.3 (2.7)	0.72	P	28 ± 9.4 (7.1)	0.78
S	2100 ± 1610 (143)	0.92	S	3190 ± 1920 (207)	0.91	S	38 ± 45 (71)	-0.15
Cl	5.1 ± 35 (3.4)	-0.01	Cl	23 ± 28 (5.5)	0.19	Cl	47 ± 48 (5.8)	-0.11
K	60.4 ± 45 (4.7)	0.5	K	68 ± 21 (6.4)	0.31	K	100 ± 66 (10)	0.81
Ca	47 ± 33 (4.2)	0.39	Ca	63 ± 33 (9.0)	-0.02	Ca	421 ± 192 (31)	0.81
Ti	4.9 ± 5.2 (4.1)	0.44	Ti	8.7 ± 4.7 (9.0)	0.47	Ti	30 ± 17 (5.6)	0.9
V	8.8 ± 8.7 (1.8)	0.37	V	9.7 ± 7.1 (2.9)	0.38	V	3.2 ± 2.2 (1.5)	0.66
Cr	0.7 ± 1.1 (0.7)	0.15	Cr	1.4 ± 1.2 (2.9)	0.09	Cr	1.0 ± 5.0 (0.9)	0.43
Mn	3.1 ± 2.2 (0.8)	0.39	Mn	3.2 ± 1.5 (1.6)	0.43	Mn	6.3 ± 4.1 (0.6)	0.9
Fe	109 ± 71 (10.5)	0.5	Fe	134 ± 49 (0.5)	0.48	Fe	352 ± 156 (24)	0.9
Co	0.1 ± 1.8 (1.4)	0.04	Co	0.8 ± 0.7 (8.5)	0.58	Co	-0.2 ± 0.5 (0.3)	-0.10
Ni	7.3 ± 8.4 (1.4)	0.22	Ni	8.5 ± 5.6 (0.3)	0.61	Ni	2.0 ± 1.4 (0.3)	0.08
Cu	4.8 ± 4.9 (1.1)	0.25	Cu	7.7 ± 3.8 (0.7)	0.22	Cu	14 ± 12 (1.1)	-0.05
Zn	36.9 ± 44 (3.7)	0.21	Zn	56 ± 37 (4.8)	0.22	Zn	52 ± 43 (4.7)	-0.03
As	0.6 ± 1.4 (1.2)	0.18	As	0.4 ± 1.0 (1.0)	-0.02	As	0 ± 0.5 (0.5)	0.07
Se	1.5 ± 1.3 (0.6)	0.63	Se	1.3 ± 0.8 (0.4)	0.65	Se	-0.1 ± 0.2 (0.2)	-0.24
Br	5.0 ± 11.7 (0.9)	0.11	Br	14 ± 12 (1.3)	0.21	Br	3.0 ± 2.5 (0.5)	-0.10
Pb	17.6 ± 22 (2.5)	0.19	Pb	28 ± 24 (2.4)	0.26	Pb	13 ± 11 (1.3)	0.1

¹Data obtained at the Presbyterian home (PBY) site in Philadelphia from April 1992 to April 1995 with Harvard impactors.

²Data obtained at the Castor Avenue Laboratory, North Central Philadelphia, from July 25 to August 14, 1994 with a modified dichotomous sampler.

*Note: Values in parentheses refer to analytical uncertainty (unc) in X-ray fluorescence determinations.

TABLE 3-3. CONCENTRATIONS (in ng/m³) OF PM_{2.5}, PM_{10-2.5}, AND SELECTED ELEMENTS IN THE PM_{2.5} AND PM_{10-2.5} SIZE RANGE WITH STANDARD DEVIATIONS (SD) AND CORRELATIONS (r) BETWEEN ELEMENTS AND PM_{2.5} AND PM_{10-2.5} MASS IN PHOENIX, AZ *

n = 164	Conc (ng/m³) ± SD (unc)	r	n = 164	Conc (ng/m³) ± SD (unc)	r
PM _{2.5}	11.2 ± 0.6 (0.6) × 10 ³	1	PM _{10-2.5}	27.6 ± 14.8 × 10 ³	1
Al	125 ± 77 (30)	0.23	Al	1879 ± 979 (547)	0.92
Si	330 ± 191 (48)	0.35	Si	5350 ± 2825 (1347)	0.92
P	11 ± 7.8 (5.7)	0.52	P	37 ± 20 (17)	0.58
S	487 ± 254 (40)	0.16	S	131 ± 47 (26)	0.77
Cl	19 ± 44 (3.0)	0.13	Cl	208 ± 204 (24)	0.28
K	110 ± 63 (9.2)	0.67	K	561 ± 298 (62)	0.92
Ca	129 ± 72 (11)	0.51	Ca	1407 ± 755 (124)	0.9
Ti	11 ± 7.1 (2.7)	0.44	Ti	130 ± 71 (20)	0.9
V	0.7 ± 2.0 (2.2)	-0.28	V	2.0 ± 2.0 (1.5)	0.51
Cr	0.6 ± 0.9 (0.7)	0.41	Cr	2.6 ± 1.7 (0.7)	0.76
Mn	5.7 ± 4.3 (0.7)	0.64	Mn	29 ± 16 (3.0)	0.91
Fe	177 ± 113 (16)	0.8	Fe	1211 ± 674 (133)	0.9
Co	-0.4 ± 1.0 (1.0)	-0.01	Co	1.2 ± 2.2 (1.9)	0.38
Ni	0.6 ± 0.9 (0.5)	0.38	Ni	1.8 ± 1.4 (0.7)	0.7
Cu	5.2 ± 6.1 (1.5)	0.69	Cu	10.3 ± 9.0 (1.5)	0.58
Zn	17 ± 14.7 (1.8)	0.64	Zn	25 ± 16 (3.2)	0.64
As	1.9 ± 3.2 (0.6)	0.5	As	0.6 ± 0.8 (0.6)	0.41
Se	0.4 ± 0.8 (0.4)	0.4	Se	-0.02 ± 0.3 (0.3)	0.21
Br	3.8 ± 2.0 (0.6)	0.57	Br	0.8 ± 0.6 (0.4)	0.48
Pb	6.6 ± 6.6 (1.0)	0.69	Pb	4.6 ± 3.8 (1.1)	0.59

* Values in parenthesis refer to analytical uncertainty (unc) in X-ray fluorescence determinations.

Source: Calculations based on data obtained by Zweidinger et al. (1998).

These data are shown to give an idea of the range of concentrations found in studies conducted more recently than those shown in Appendix 6A of the 1996 PM AQCD. The speciation network should provide more thorough coverage of the composition of particles in the PM_{2.5} size

range across the United States. Results from the pilot study for the speciation network are given in Appendix 3B.

As can be seen from inspection of Tables 3-2 and 3-3, the analytical uncertainty (given in parentheses next to concentrations) as a fraction of the absolute concentration is highly variable. It exceeds the concentration for a number of trace metals whose absolute concentrations are low, whereas it is very small for abundant elements such as sulfur.

Sulfur (S) is the major element analyzed in the $PM_{2.5}$ size fraction in the two Philadelphia studies and is highly correlated with $PM_{2.5}$; however, its abundance is roughly two orders of magnitude lower in the $PM_{10-2.5}$ size range and is negatively correlated with $PM_{10-2.5}$. The concentrations of the crustal elements Al, Si, K, Ca, and Fe are much higher in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range and are well correlated with $PM_{10-2.5}$. A number of trace elements (e.g., Cr, Co, Ni, Cu, Zn, As, Se, and Pb) are detectable in the two $PM_{2.5}$ data sets, and the concentrations of many of these elements are much greater than the uncertainty in their determination. Except for Co, As, and Se (which are not detected in the $PM_{10-2.5}$ samples), the concentrations of many elements (Cr, Zn, and Pb) are comparable in the $PM_{2.5}$ and $PM_{10-2.5}$ size ranges. The concentration of Cu is significantly higher in the $PM_{10-2.5}$ size range, whereas the concentration of Ni is smaller in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range. Problems with detecting and quantifying concentrations of Co, As, and Se can result in artificial reduction of their correlation coefficients with $PM_{2.5}$.

There are a number of distinct differences between the $PM_{2.5}$ sets for Philadelphia and Phoenix. For instance, sulfate and associated cations and water that would be expected to correspond to the measurement of S appear to constitute a major fraction of the composition of the PM in the Philadelphia data set; whereas they appear to constitute a much smaller fraction of the PM in the Phoenix data set. The highest $PM_{2.5}$ values were observed in Philadelphia during episodes driven by high sulfate abundances, whereas those in Phoenix were driven by raised soil dust. The concentration of S in Phoenix is much lower in the Phoenix $PM_{2.5}$ data set than in either Philadelphia $PM_{2.5}$ data set, even though it represents the most abundant element and it is only weakly correlated with $PM_{2.5}$. This is in marked contrast to the data shown for Philadelphia and also to data found at other eastern sites. It is not clear what the reasons are for this finding. As in Philadelphia, the concentration of S in Phoenix is higher in the $PM_{2.5}$ size range than in the $PM_{10-2.5}$ size range. Trace metals (e.g., Cr, Co, Ni, Cu, Zn, As, and Pb) are not well

correlated ($0.04 < r < 0.25$) with $PM_{2.5}$ in the Philadelphia data set, whereas they are more variably correlated ($0.01 < r < 0.69$) with $PM_{2.5}$ in the Phoenix data set. The uncertainty in the concentration measurement most probably plays a role in determining a species' correlation with $PM_{2.5}$, especially when the analytical uncertainty is high relative to concentration as it is for a number of elements in the data shown in Tables 3-2 and 3-3. Concentrations of Al, Si, K, Ca, and Fe are again much higher in the $PM_{10-2.5}$ size range than in the $PM_{2.5}$ size range and are strongly correlated with $PM_{10-2.5}$ in both data sets.

There are also similarities in the $PM_{2.5}$ data sets for Philadelphia and Phoenix. Crustal elements are not as well correlated with $PM_{2.5}$ as they are with $PM_{10-2.5}$ in both data sets. The concentrations of trace metals (Cr, Ni, Cu, and Zn) in $PM_{2.5}$ are similar in Philadelphia and Phoenix. It can also be seen that their concentrations are of the same order of magnitude in both $PM_{2.5}$ and $PM_{10-2.5}$. Concentrations of Cu are noticeably higher in $PM_{10-2.5}$ than in $PM_{2.5}$ in both Philadelphia and Phoenix. These results are consistent with those of many monitoring studies shown in Appendix 6A of the 1996 PM AQCD, which also show that concentrations of these metals are of the same order of magnitude in both size fractions and that concentrations of Cu tend to be higher in $PM_{10-2.5}$ than in $PM_{2.5}$.

One study suggests that the partitioning of trace metals between the fine and coarse fractions varies with PM concentration. Salma et al. (2002) determined the size distribution of a number of trace elements at four sites characterizing environments ranging from the urban background to an urban traffic tunnel in Budapest, Hungary. Although S, K, V, Ni, Cu, Zn, As, and Pb were found mainly in the fine fraction at the urban background site, their mass median aerodynamic diameters increased with increasing PM concentrations until they were all found mainly in the coarse fraction in the traffic tunnel. They also found that Na, Mg, Al, Si, P, Ca, Ti, Fe, Ga, Sr, Zr, Mo, and Ba were concentrated mainly in the coarse fraction at all four sites and that their mass median aerodynamic diameters increased with increasing PM concentrations.

The mean concentration of Pb observed in the methods evaluation study for the speciation network was only about 5 ng/m^3 in Philadelphia during the first half of 2000 (Appendix 3B), whereas its concentration was about three times higher during studies conducted during the early 1990s (Table 3-3). In a study conducted in the greater Philadelphia area during the summer of 1982, Dzubay et al. (1988) found concentrations of Pb of about 250 ng/m^3 , or about fifty times higher than observed in 2000. The mean Pb concentration was about 3 ng/m^3 at the Phoenix site

included as part of the same methods evaluation study for the speciation network; however, the mean Pb concentration was 39 ng/m³ during an earlier study conducted during 1989 and 1990 in Phoenix (Chow et al., 1991). These changes in Pb concentrations are consistent with those in many other urban areas for which monitoring studies have been conducted during the late 1970s and 1980s (cf., Appendix 6A of the 1996 PM AQCD) and for which there are data given in Appendix 3B. It should be remembered that the older studies were conducted while Pb was still used as a gasoline additive. The ratio of Pb in PM_{2.5} to Pb in PM_{10-2.5} was also much higher in the older studies than in the more recent ones, reflecting the importance of combustion as its source. Smaller decreases are apparent in the concentrations of other trace metals such as Cu, Ni, and Zn between studies conducted in the early 1980s and in the methods evaluation study for the speciation network conducted in 2000.

Some indication of the sources of metals such as Pb, Cu, Cd, and Zn in current ambient PM_{2.5} and PM_{10-2.5} samples can be obtained by examining their sources in urban runoff. The sources of these elements in urban runoff were found to be the weathering of building surfaces, motor vehicle brake and tire wear, engine oil and lubricant leakage and combustion, and wet and dry atmospheric deposition (Davis et al., 2001). Once deposited on the ground, these elements can be resuspended with other material as PM_{2.5} and PM_{10-2.5}, although research is needed into the mechanisms of how this is accomplished. Wind-abrasion on building siding and roofs (coatings such as Pb paint and building material such as brick, metal, and wood siding); brake wear (brake pads contain significant quantities of Cu and Zn); tire wear (Zn is used as a filler in tire production); and burning engine oil could all produce particles containing these metals, especially Zn.

Data for the chemical composition of ambient ultrafine particles are sparse. In a study conducted at several urban sites in Southern California, Cass et al. (2000) found that the composition of ultrafine particles ranged from 32 to 67% organic compounds, 3.5 to 17.5% elemental carbon, 1 to 18% sulfate, 0 to 19% nitrate, 0 to 9% ammonium, 1 to 26% metal oxides, 0 to 2% sodium, and 0 to 2% chloride. Thus carbon, in various forms, was found to be the major contributor to the mass of ultrafine particles. However, ammonium was found to contribute 33% of the mass of ultrafine particles at one site in Riverside. Iron was the most abundant metal found in the ultrafine particles. Chung et al. (2001) found that carbon was the major component of the mass of ultrafine particles in a study conducted during January of 1999 in Bakersfield,

CA. However, in the study of Chung et al., the contribution of carbonaceous species (OC and EC; typically 20 to 30%) was much lower than that found in the cities in Southern California. They found that calcium was the dominant cation, accounting for about 20% of the mass of ultrafine particles in their samples. Sizable contributions from silicon (0 to 4%) and aluminum (6 to 14%) were also found. Further studies, including scanning electron microscopy, may be needed to quantify the role of coarse particle bounce from the upper stages of their MOUDI impactor.

Gone et al. (2000) measured the size distribution of trace elements from 0.056 μm to 1.8 μm D_a in Pasadena, CA and in the Great Smoky Mountains National Park, TN. They found that elements identified as being of anthropogenic origin had mass median diameters $< 1 \mu\text{m}$ PM; whereas elements of crustal origin generally had a mass median diameter $> 1 \mu\text{m}$. Concentrations of trace metals were much higher in the accumulation mode than in the ultrafine mode in both study areas. In PM_{10} , 76% of Cr, 95% of Fe, 94% of Zn, 89% of As, and 79% of Cd at the Tennessee site were found in the accumulation mode; and 70% of Fe, 85% of Zn, 92% of As, and 84% of Cd were found in the accumulation mode in Pasadena. Fe was the most abundant metal found in the ultrafine particles. The abundance of crustal elements, such as Al, declined rapidly with decreasing particle size at both locations; and Al in PM_{10} probably represented the lower tail of the coarse PM mode. However, on two days at Pasadena there were increases in the concentration of Al in ultrafine particles that were associated with increases in Sc and Sm. The latter two elements originate exclusively from crustal material (Gone et al., 2000).

3.2.5 Spatial Variability in Particulate Matter and its Components

PM_{2.5}

Aspects of the spatial variability of $\text{PM}_{2.5}$ concentrations on the urban scale are examined in this section. Intersite correlation coefficients for $\text{PM}_{2.5}$ can be calculated based on the results of FRM monitors placed at multiple sites within MSAs across the United States. Pearson correlation coefficients (r) calculated for pairs of monitoring sites in the Philadelphia, PA; Cleveland, OH; Dallas, TX; and Los Angeles, CA MSAs are shown in Table 3-4. The 90th percentile value, P_{90} , of the absolute differences (in $\mu\text{g}/\text{m}^3$) between the two sites is shown in parentheses below r along with the coefficient of divergence (COD), and the number of

**TABLE 3-4a-d. MEASURES OF THE SPATIAL VARIABILITY OF
PM_{2.5} CONCENTRATIONS WITHIN SELECTED METROPOLITAN
STATISTICAL AREAS**

(a) Philadelphia, PA								
Site I.D. #	340071007	340155001	420170012	420450002	420910013	421010004	421010136	
340071007	1	0.91 (6.3, 0.14) 170	0.93 (5.2, 0.15) 167	0.87 (6.9, 0.19) 183	0.88 (5.0, 0.16) 176	0.94 (4.6, 0.15) 163	0.93 (5.1, 0.14) 166	
340155001		1	0.84 (7.5, 0.19) 176	0.88 (7.4, 0.18) 194	0.83 (7.1, 0.18) 184	0.89 (7.4, 0.17) 169	0.85 (6.9, 0.18) 173	
420170012			1	0.85 (7.5, 0.16) 199	0.88 (4.7, 0.13) 200	0.94 (4.9, 0.11) 177	0.89 (5.3, 0.13) 180	
420450002				1	0.87 (6.1, 0.15) 208	0.94 (5.1, 0.11) 187	0.88 (4.0, 0.12) 193	
420910013					1	0.90 (4.6, 0.11) 181	0.87 (4.5, 0.11) 185	
421010004						1	0.96 (3.3, 0.08) 550	
421010136							1	
Mean	14.78	14.59	14.11	15.92	14.2	15.72	15.81	
Obs	197	208	217	230	221	610	616	
SD	9.2	8.65	8.47	8.65	8.93	9.18	9.26	
(b) Cleveland, OH								
Site I.D. #	390350013	390350038	390350060	390350065	390350066	390351002	390851001	390932003
390350013	1	0.91 (7.1, 0.13) 320	0.96 (3.3, 0.12) 322	0.94 (5.4, 0.10) 314	0.92 (7.2, 0.16) 300	0.88 (9.0, 0.18) 308	0.89 (10.7, 0.21) 308	0.92 (8.1, 0.17) 265
390350038		1	0.92 (6.9, 0.14) 306	0.89 (9.4, 0.15) 296	0.85 (13.1, 0.21) 290	0.84 (12.9, 0.21) 304	0.84 (14.3, 0.23) 294	0.892 (11.2, 0.18) 256
390350060			1	0.93 (5.1, 0.14) 309	0.90 (8.15, 0.19) 300	0.87 (8.6, 0.20) 310	0.88 (10.8, 0.22) 307	0.90 (8.9, 0.18) 256
390350065				1	0.96 (4.7, 0.14) 295	0.91 (5.4, 0.16) 310	0.90 (7.8, 0.20) 306	0.91 (7.7, 0.18) 264
390350066					1	0.88 (4.9, 0.16) 304	0.91 (5.8, 0.15) 295	0.91 (5.8, 0.13) 247
390351002						1	0.89 (6.0, 0.18) 303	0.87 (5.3, 0.18) 261
390851001							1	0.90 (6.6, 0.15) 275
390932003								1
Mean	18.34	20.16	18.39	17.47	14.74	15	14	15.22
Obs	368	931	353	340	332	351	342	298
SD	9.59	11.5	9.9	8.95	8.45	8.16	8.44	8.8

Key
Airs Site I.D. #
Pearson r
(90th %-tile difference in concentration,
coefficient of divergence)
number of observations

Key
Airs Site I.D. #
Pearson r
(90th %-tile difference in concentration,
coefficient of divergence)
number of observations

**TABLE 3-4a-d (cont'd). MEASURES OF THE SPATIAL VARIABILITY OF
PM_{2.5} CONCENTRATIONS WITHIN SELECTED METROPOLITAN
STATISTICAL AREAS.**

(c) Dallas, TX							
Site I.D. #	480850005	481130020	481130035	481130050	481130057	480850005	481130020
480850005	1	0.92 (3.5, 0.11) 220	0.94 (3.6, 0.11) 204	0.94 (4.3, 0.13) 213	0.89 (6.3, 0.15) 195	0.94 (3.7, 0.10) 218	0.94 (3.1, 0.10) 189
481130020		1	0.95 (3.2, 0.08) 212	0.94 (3.3, 0.09) 603	0.92 (4.1, 0.11) 205	0.95 (2.5, 0.07) 635	0.97 (2.3, 0.07) 207
481130035			1	0.97 (2.0, 0.06) 203	0.93 (3.9, 0.09) 191	0.97 (1.9, 0.06) 207	0.94 (3.6, 0.10) 185
481130050				1	0.94 (2.7, 0.08) 199	0.98 (2.2, 0.06) 608	0.94 (4.2, 0.12) 196
481130057					1	0.95 (3.1, 0.08) 198	0.91 (5.5, 0.14) 182
480850005						1	0.96 (3.0, 0.09) 198
481130020							1
Mean	11.54	12.42	12.83	13.34	13.67	12.68	11.67
Obs	234	677	222	644	215	687	216
SD	5.62	5.69	5.79	5.79	6.14	5.67	5.43
(d) Los Angeles, CA							
Site I.D. #	60370002	60371103	60371201	60371301	60372005	60374002	
60370002	1	0.87 (10.7, 0.18) 581	0.76 (14.6, 0.23) 208	0.68 (17.9, 0.25) 229	0.95 (6.2, 0.14) 212	0.60 (18.1, 0.26) 553	
60371103		1	0.86 (12.8, 0.20) 205	0.89 (10.1, 0.12) 222	0.93 (7.1, 0.11) 207	0.80 (13.6, 0.17) 563	
60371201			1	0.76 (18.1, 0.24) 212	0.85 (12.1, 0.18) 197	0.66 (18.2, 0.24) 197	
60371301				1	0.78 (13.2, 0.18) 214	0.95 (8.1, 0.11) 216	
60372005					1	0.62 (15, 0.20) 204	
60374002						1	
Mean	20.91	22.48	18.32	23.73	20.09	20.31	
Obs	641	656	217	237	220	621	
SD	13.35	13.47	11.65	13.43	11.92	12.42	

observations used in the calculation of r , P_{90} and COD is given on the third line. The COD was used by Wongphatarakul et al. (1998) as a measure of the degree of similarity between aerosol data sets¹. The annual mean concentrations, the number of observations used to calculate the annual average, and the standard deviation are shown directly beneath the correlation tables for each site. These analyses and those for another 23 MSAs are given along with maps in Appendix 3A. As the concentrations of $PM_{2.5}$ at two sampling sites become more alike, the COD approaches zero; as the concentrations diverge, the COD approaches one.

The four MSAs shown in Table 3-4(a-d) were chosen to illustrate different patterns of spatial variability across the United States. In addition, air-pollution health-outcome studies have been performed in a few of these MSAs. It can be seen from inspection of Table 3-4 that correlation coefficients vary over a wide range in the MSAs shown. Correlations between sites in the Philadelphia, PA, Cleveland, OH, and Dallas, TX MSAs are all high and span a relatively narrow range (0.82 to 0.97). However, correlations between sites in the Los Angeles-Long Beach MSA are lower than in the three other MSAs and span a wider range of values (0.60 to 0.95). If the monitoring site in Lancaster, CA were included, correlations would be even lower. This site was omitted because it did not meet completeness criteria for 2001. The extension of these analyses to include the relevant consolidated MSAs (CMSAs) would also produce a number of sites that are even less well correlated with each other, in part, because some included sites are located outside of urban airsheds.

Correlation coefficients between pairs of sites in the other 23 MSAs given in Appendix 3A for the most part fall within the range of values given in Table 3-4. In four MSAs (Columbia, SC; Norfolk, VA; Grand Rapids, MI; and Baton Rouge, LA), intersite correlations are all greater than 0.9. In nine others (Cleveland, OH; Louisville, KY; Chicago, IL; Milwaukee, WI; Philadelphia, PA; Detroit, MI; Kansas City, MO; Dallas, TX; and Salt Lake City, UT), they are

¹ The COD for this purpose is defined as follows:

$$COD_{jk} = \sqrt{\frac{1}{p} \sum_{i=1}^p \left(\frac{x_{ij} - x_{ik}}{x_{ij} + x_{ik}} \right)^2} \quad (3-1)$$

where x_{ij} and x_{ik} represent the 24-h average $PM_{2.5}$ concentration for day i at site j and site k and p is the number of observations.

all greater than 0.8. Correlations between sites in the other MSAs examined tend to be lower and span a broader range than for the MSAs mentioned above.

Seven pairs of collocated monitors in seven MSAs (Columbia, SC; Dallas, TX; Detroit, MI; Grand Rapids, MI; Louisville, KY; Steubenville, OH; Washington, DC) provide an indication of the performance of collocated monitors (see Table 3A-1). Mean values of r , P_{90} , and COD for these seven pairs of monitors are 0.986, $1.63 \mu\text{g}/\text{m}^3$, and 0.060, thus suggesting that most of the intersite variability seen is not due to sampler imprecision.

There are no strong regional patterns evident in the data given in Appendix 3A except that correlations tend to be higher between monitoring sites in MSAs in the southeastern United States than between monitoring sites in other regions.

A number of factors affect intersite correlations within MSAs. These include field measurement and laboratory analysis errors, placement of monitors close to active sources, placement of monitors in outlying areas, placement of monitors in locations that are isolated topographically from other monitors, placement of monitors in areas outside of local atmospheric circulation regimes (e.g., land-sea breezes), and transient local events (e.g., thunderstorms or sporadic emissions). In several MSAs (i.e., Atlanta, GA, Seattle, WA, and Los Angeles-Long Beach, CA) at least one site is remote from the others (by at least 100 km), is physically separated from them by mountains, and is really neither part of the urban area nor the urban airshed. Correlations between concentrations at these sites and others tend to be lower than among the other sites, and concentration differences tend to be larger. It should be noted that outlying sites such as these are included in many epidemiologic time-series studies without any differential weighting (e.g., with respect to the exposed population or spatial differences in susceptibility or with regard to compositional differences). Although it is frequently the case that distance between sites in urban areas is largely responsible for the spatial variability that is observed, there are a few instances for which correlations are higher and differences in concentrations are lower for sites that are located farthest apart. This situation may arise because these sites are influenced more by the regional background of secondary PM rather than by local sources, but there are not any consistent distances evident below which correlations and differences in concentrations tend towards some limiting values. Still, it is generally the case that outlying sites are characterized by lower annual mean concentrations.

Indications of land use (commercial, industrial, residential, agricultural, forest) and location of sites (urban/city center, suburban, rural) are given in the AIRS data base. Categories such as urban/city center can refer to very different conditions in Columbia, SC, and Chicago, IL. Also, it should not be automatically assumed that concentrations measured at sites categorized as industrial are dominated by local emissions. The PM_{2.5} monitoring sites are generally deployed to capture potential population exposures in a variety of environments as opposed to monitoring for compliance as it exists around local sources. It should be remembered that much of PM_{2.5} is secondary in origin. The widespread formation of secondary PM, coupled with the long atmospheric lifetime of PM_{2.5}, ensures some measure of uniformity in the correlations of PM_{2.5} across urban areas. Correlations between many site pairs classified as industrial can be high, even though they are separated by large distances (as in the Seattle MSA).

Some indication of the variability of primary PM_{2.5} produced by local sources can be obtained by examining the variability of CO, which is produced mainly by mobile sources (U.S. Environmental Protection Agency, 2000b), and by the variability in EC concentrations (Kinney et al., 2000). Carbon monoxide is relatively inert on the urban scale, and its distribution is governed by the spatial pattern of its emissions and the subsequent dispersion of these emissions not by photochemistry. Carbon monoxide concentrations are at least a factor of three higher near urban centers than in surrounding rural areas within the four consolidated metropolitan statistical areas examined in the EPA document, *Air Quality Criteria for Carbon Monoxide* (CO AQCD; U.S. Environmental Protection Agency, 2000b). The correlations of CO within the urban areas examined in that document were all low to moderate. Therefore, it might be expected that primary PM_{2.5} produced by local traffic should be at least as heterogeneous as CO in a given urban area. Elemental carbon is a significant component of diesel exhaust (cf., Appendix 3D). Kinney et al. (2000) measured EC and PM_{2.5} concentrations at four sites located on sidewalks of streets characterized by varying exposures to diesel emissions in upper Manhattan (Harlem, NY). Whereas the mean PM_{2.5} concentrations varied by about one-third from 37 to 47 µg/m³ at the four sites, mean EC concentrations varied by a factor of four from 1.5 to 6.2 µg/m³. The corresponding ratios of EC to PM_{2.5} ranged from 0.039 to 0.14. Although EC constituted a relatively small fraction of PM_{2.5} in this study, spatial variability in its sources (diesel and gasoline fueled vehicles, resuspended road dust, and cooking) contributed, on average, about one-third of the spatial variability observed in PM_{2.5} concentrations. Further

analyses are needed to determine whether the remaining variability could be attributed to other local and city-wide sources. Because the effects of emissions from local point sources on receptor sites depend strongly on wind direction, correlations involving contributions from local sources can be much lower than from area sources (much as motor vehicle traffic) or from regionally dispersed sources (such as the photochemical production of secondary organic PM and sulfate).

The difference in mean PM_{2.5} concentrations between the site with the lowest and the site with the highest mean concentration range in all MSAs included in Appendix 3A ranges from 0.4 µg/m³ (Baton Rouge) to about 8 µg/m³ (Pittsburgh). Six MSAs (Chicago, Seattle, Cleveland, St. Louis, Detroit, and Pittsburgh) show maximum intersite differences in the annual mean larger than 6 µg/m³. In the Seattle MSA, there is one monitoring site (Figure 3A-23a) that is separated from the remaining sites by topography and has much lower mean PM_{2.5} concentrations, much smaller seasonal variability in concentrations, and much lower maximum concentrations than these other sites. However, the annual mean concentrations at all the other sites within the Seattle MSA are within 3 µg/m³ of each other. Differences in annual mean concentrations are also larger between sites located in different MSAs but within the same CMSA. For example, in the consolidated MSA of Los Angeles-Riverside, the range of annual mean PM_{2.5} values is extended from ~20 µg/m³ in the urban area of Los Angeles county to ~29 µg/m³ in Riverside County. Large differences in annual mean concentrations within a given area reflect differences in source or meteorological or unique topographic characteristics affecting sites, whereas very small differences found in some areas may mainly be the result of measurement imprecision.

While high correlations of PM_{2.5} provide an indication of the spatial uniformity in temporal variability (directions of changes) in PM_{2.5} concentrations across urban areas, they do not imply uniformity in the PM_{2.5} concentrations themselves. The 90th percentile difference in concentrations (P₉₀) and the coefficient of divergence (COD) are used here to give a more quantitative indication of the degree of spatial uniformity in PM_{2.5} concentrations across urban areas. A COD of zero implies that both data sets are identical, and a COD of one indicates that two data sets are completely different. The calculation of the Pearson correlation coefficient (r), P₉₀, and COD allows for distinctions between pairs of sites to be made based on various combinations of these parameters. Figure 3-18 shows examples of the varying degree of heterogeneity in concentrations between pairs of sites that are highly correlated (r > 0.9 for all




Figure 3-18. Occurrence of differences between pairs of sites in three MSAs. The absolute differences in daily average $\text{PM}_{2.5}$ concentrations between sites are shown on the x-axis and the number of occurrences on the y-axis. The MSA, years of observations, AIRS site I.D. numbers for the site pairs, Pearson correlation coefficients (r), coefficients of divergence (COD), 90th percentile (P_{90}) difference in concentration between concurrent measurements are also shown.

three site pairs). The increase in the spread of concentrations between the chosen site-pairs is reflected in increases in both P_{90} and COD. Pairs of sites showing high correlations and CODs < 0.1 and $P_{90}s < 4 \mu\text{g}/\text{m}^3$ (as in Columbia, SC, Figure 3-7a) indicate homogeneity in both $\text{PM}_{2.5}$ concentrations and in their temporal variations. Presumably, sites such as these are more strongly affected by regional than local sources. Pairs of sites showing low correlations, values of $P_{90} > 10 \mu\text{g}/\text{m}^3$ and CODs > 0.2 , as in Los Angeles, CA (Table 3-5), indicate heterogeneity in both $\text{PM}_{2.5}$ concentrations and in their temporal variations. Note that the extended urban area or the CMSA includes Riverside County, as well as Los Angeles County. Even lower correlations and a greater degree of heterogeneity in $\text{PM}_{2.5}$ concentrations were found in the extended CMSA. Pairs of sites showing high correlations ($r > 0.9$) and CODs > 0.2 and $P_{90}s > 10 \mu\text{g}/\text{m}^3$ indicate heterogeneity in concentrations but homogeneity in their day-to-day changes. Selected pairs of sites in the Cleveland MSA show moderate to high correlations coupled with CODs > 0.2 and $P_{90}s \geq 10 \mu\text{g}/\text{m}^3$ (Table 3-4), suggesting moderate homogeneity in day-to-day changes but significant spatial heterogeneity in concentrations.

The effect of local point sources on intersite variability can be seen at several sites among those listed in Table 3-4(a-d) and Appendix 3A. Sites 39-035-0038 (Cleveland, OH; Table 3-4 and Figure 3A-8), 18-089-0022 (Gary, IN; Figure 3A-15), 55-079-0043 (Milwaukee, WI; Figure 3A-13), and 17-119-0023 (St. Louis, MO; Figure 3A-17) are designated as “source oriented” in the AIRS data base in contrast to the “population exposure” objective associated with most of the MSA sites. $\text{PM}_{2.5}$ concentrations at these sites are weakly correlated with other sites within the MSA as evidenced by low correlation coefficients and large $P_{90}s$ and CODs even though some of the neighboring sites may be located short distances away. Other sites designated as “source oriented” in Chicago, Milwaukee, and St. Louis do not show clear evidence that local sources are contributing to intersite variability. Conversely, in the Tampa, FL MSA, pairs of sites are only moderately correlated ($0.7 < r < 0.87$), but the distribution of concentrations is rather homogeneous (COD < 0.14 and $P_{90} < 5 \mu\text{g}/\text{m}^3$; Figure 3A-7). Thus, a number of different combinations of spatial uniformity in $\text{PM}_{2.5}$ concentrations, and correlations of these concentrations are found.

Values of P_{90} for absolute differences in concentrations between sites span a wide range in the data set given in Appendix 3A. In many instances they can be quite low, only about a few $\mu\text{g}/\text{m}^3$. Such cases are found mainly in the eastern United States. The largest P_{90} values

**TABLE 3-5. MEASURES OF THE SPATIAL VARIABILITY OF
PM_{10-2.5} CONCENTRATIONS WITHIN SELECTED METROPOLITAN
STATISTICAL AREAS**

(a) Cleveland, OH						
Site	390350013	390350038	390350045	390350060	390350065	390851001
390350013	1	0.67 (23.2, 0.26) 182	0.67 (28.5, 0.28) 95	0.73 (17.9, 0.22) 97	0.62 (27.0, 0.31) 98	0.41 (40.0, 0.60) 94
390350038		1	0.65 (16.1, 0.22) 90	0.73 (11.7, 0.18) 93	0.69 (13.9, 0.62) 90	0.44 (24.9, 0.53) 89
390350045			1	0.66 (18.1, 0.23) 94	0.71 (10.6, 0.31) 102	0.49 (19.9, 0.50) 99
390350060				1	0.74 (15.4, 0.38) 93	0.31 (28.0, 0.59) 94
390350065					1	0.22 (20.4, 0.55) 99
390851001						1
<i>MEAN</i>	26.36	18.63	16.76	21.35	16.79	7.15
<i>Obs</i>	216	614	112	113	111	109
<i>SD</i>	17.38	11.6	8.96	16.39	9.49	4.94
(b) Dallas, TX						
Site	481130020	481130035	481130050	481130057		
481130020	1	0.79 (4.5, 0.17) 54	0.71 (9.3, 0.22) 55	0.66 (16.5, 0.32) 54		
481130035		1	0.69 (7.8, 0.18) 50	0.60 (13.2, 0.30) 50		
481130050			1	0.69 (13.5, 0.24) 50		
481130057				1		
<i>MEAN</i>	11.22	12.86	14.46	19.12		
<i>Obs</i>	60	55	56	55		
<i>SD</i>	5.35	6.66	6.44	10.55		
(c) Los Angeles, CA						
Site	60370002	60371002	60371103	60374002		
60370002	1	0.82 (19.0, 0.24) 49	0.63 (15.5, 0.18) 49	0.58 (17.3, 0.27) 45		
60371002		1	0.74 (11.5, 0.21) 49	0.54 (11.5, 0.25) 47		
60371103			1	0.57 (12.5, 0.22) 45		
60374002				1		
<i>MEAN</i>	24.1	15.33	21.44	16.08		
<i>Obs</i>	56	56	57	53		
<i>SD</i>	11.67011257	6.68	8.65	6.61		

Key
Airs Site I.D. #
Pearson r
(90th %-tile difference in concentration,
coefficient of divergence)
number of observations

Key
Airs Site I.D. #
Pearson r
(90th %-tile difference in
concentration,
coefficient of divergence)
number of observations

Key
Airs Site I.D. #
Pearson r
(90th %-tile difference in
concentration,
coefficient of divergence)
number of observations

were associated with a single site in Pittsburgh and reached as high as $21 \mu\text{g}/\text{m}^3$ (Figure 3A-9). Excluding this site, large P_{90} values are found mainly in the western United States. Values of $P_{90} > 18 \mu\text{g}/\text{m}^3$ are found in the Riverside and Los-Angeles-Long Beach MSAs. Maximum differences in concentrations between sites can be much larger than shown in Figure 3-18 and have been larger than $100 \mu\text{g}/\text{m}^3$ on several occasions in the Atlanta, GA and Los Angeles-Long Beach, CA MSAs. Rizzo and Pinto (2001) and Fitz-Simons et al. (2000) examined correlations between sites located even farther apart than those examined here based on the 1999 AIRS data set for $\text{PM}_{2.5}$. They found that in a number of MSAs, $\text{PM}_{2.5}$ concentrations are still well correlated ($r > 0.7$) up to distances of 100 km or more. Leaderer et al. (1999) found $r = 0.49$ between sites outside of homes and a regional background monitor located from 1 to 175 km away in southwestern Virginia. $\text{PM}_{2.5}$ tends to be correlated over much larger areas in the East than in the West, mainly because the terrain tends to be flatter over wider areas in the East (Rizzo and Pinto, 2001). As a result, there is a greater opportunity for mixing of emissions among dispersed source regions. Many large urban areas in the West are surrounded by mountains. The presence of more rugged terrain in the West leads to greater confinement of emissions from large urban areas. Other factors such as differences in the composition and amount of emissions of precursors and in the rates of photochemical oxidation of these emissions in the atmosphere also play a role.

There is also evidence for inter-annual variability in the spatial variability in $\text{PM}_{2.5}$ concentrations. The median year-to-year changes in intersite r (0.03), P_{90} ($-0.75 \mu\text{g}/\text{m}^3$), and COD (-0.015) from 1999 to 2000 do not differ significantly from zero for all the site pairs considered in Appendix 3A. The year-to-year changes in the spatial variability of $\text{PM}_{2.5}$ concentrations in a number of MSAs (e.g., the Columbia, SC; Grand Rapids, MI; Milwaukee, WI; Baton Rouge, LA; Kansas City, MO; Boise, ID; and Portland, OR MSAs) are similar and are smaller than those found in the Cleveland, OH; Salt Lake City, UT; and San Diego, CA MSAs. The ranges in these parameters are largest for a number of individual site-pairs, especially those involving sites that are remote from the others in their MSAs. In these MSAs (e.g., Atlanta, GA; Los Angeles, CA; and Seattle, WA MSAs), some sites may be located in different airsheds from the remaining sites. Year-to-year changes in parameters describing spatial variability in $\text{PM}_{2.5}$ concentrations tend to be larger when sites in different counties within a given MSA are considered rather than when sites in the same county are considered. There are

a number of factors that can account for inter-annual variability in these parameters, such as changes in patterns in the emissions of primary $PM_{2.5}$; in the transport and rates of transformation of secondary $PM_{2.5}$ precursors in field measurement; and analysis procedures.

Some additional data for indicating the stability with respect to year-to-year changes in spatial variability are available from earlier studies. For example, a comparison between data obtained during the summers of 1992 and 1993 (Wilson and Suh, 1997), shown in Figure 3-19, and data obtained during the summer of 1994 (Pinto et al., 1995) (cf., Table 3-8) in Philadelphia, PA, suggests that inter-site correlations of $PM_{2.5}$ have remained high and have changed very little between the two study periods.




Figure 3-19. Intersite correlation coefficients for $PM_{2.5}$, PM_{10} , and $PM_{10-2.5}$.

Source: Wilson and Suh (1997).

PM_{10-2.5}

Intersite correlations of PM_{10-2.5} concentrations obtained during the summers of 1992 and 1993 in Philadelphia, PA (Wilson and Suh, 1997) are shown in Figure 3-19. As can be seen, correlations of PM_{10-2.5} are substantially lower than those for PM_{2.5}.

Intersite correlation coefficients can also be calculated for PM_{10-2.5} based on the AIRS data set as shown in Table 3-5 for the Cleveland, OH, Dallas, TX, and Los Angeles, CA MSAs. However, data for analyzing the spatial variability of PM_{10-2.5} are more limited than for PM_{2.5}; therefore, fewer urban areas could be characterized in Appendix 3A (Figures 3A-28 to 3A-44). Whereas PM_{2.5} concentrations were found to be highly correlated between sites in the Detroit, MI MSA (Table 3-4), estimated PM_{10-2.5} concentrations are noticeably less well correlated. Likewise, correlations of PM_{10-2.5} in the Chicago, IL MSA are also lower than those for PM_{2.5}. In contrast, correlations of PM_{10-2.5} concentrations between several pairs of sites in the Los Angeles-Long Beach partial MSA are higher than those for PM_{2.5}.

The interpretation of these results is not straightforward, as concentrations of PM_{10-2.5} are generated by taking the difference between collocated PM_{2.5} and PM₁₀ monitors. Consequently, caution must be exercised when viewing them. Errors in the measurement of PM_{2.5} and PM₁₀ may play a large role in reducing apparent correlations of PM_{10-2.5} such that collocated PM_{10-2.5} “measurements” may be expected to be poorly correlated (White, 1998). Indeed, several estimates of concentrations are negative. Negative PM_{10-2.5} concentrations also lead to artifacts in the calculation of CODs. In cases where these artifacts cause a division by zero or a very small number in the calculation of CODs, dashes are used in Figures 3A-28 through 3A-44. These results imply that negative concentrations can be almost equal in absolute magnitude to positive concentrations in the same MSA. The possible causes of these errors are essentially the same as those discussed in Section 3.2.1 with regard to the occurrence of PM_{2.5} to PM₁₀ ratios greater than one. There are also physical bases for expecting that PM_{10-2.5} concentrations may be more variable than those for PM_{2.5}. PM_{10-2.5} is mainly primary in origin, and its emissions are spatially and temporally heterogenous. Similar considerations apply to primary PM_{2.5}, but much of PM_{2.5} is secondary, and sources of secondary PM are much less spatially and temporally variable. Dry deposition rates of particles depend strongly on particle size. Whereas all particles may be brought to the surface by turbulent motions in the atmosphere, gravitational settling becomes more important with increasing particle size. Gravitational settling can effectively

limit the horizontal distance a particle can travel. For example, 10 μm D_a particles suspended in a hypothetical 1 km deep planetary boundary layer can be removed within a few hours, but 1 μm D_a particles can remain suspended in the atmosphere for up to 100 to 1,000 times longer before being dry deposited. (Estimated atmospheric lifetimes were based on deposition velocities given in Lin et al. [1994] for typical wind speeds.) The findings of larger correlations of $\text{PM}_{10-2.5}$ between several site pairs in the Los Angeles basin (cf., Figures 3A-25/26 and Figures 3A-42/43) and one other site pair in the St. Louis, MO-IL MSA (cf., Figures 3A-17 and 3A-37) are anomalous in light of the discussion above. However, these findings could have resulted from differences between the spatial and temporal behavior of sources of $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ in these locations. Because of negative values, CODs were not calculated.

PM Components

Three methods for comparing the chemical composition of aerosol databases obtained at different locations and times were discussed by Wongphatarakul et al. (1998). Log-log plots of chemical concentrations obtained at pairs of sampling sites accompanied by the coefficient of divergence (COD) were examined as a way to provide an easily visualized means of comparing two data sets². Examples comparing downtown Los Angeles with Burbank and with Riverside-Rubidoux are shown in Figures 3-20 and 3-21. As the composition of two sampling sites become more similar, the COD approaches zero; as their compositions diverge, the COD approaches one. Correlation coefficients calculated between components can be used to show the degree of similarity between pairs of sampling sites.

In addition to calculating correlation coefficients for total mass or for individual components, correlation coefficients for characterizing the spatial variation of the contributions from given source types can also be calculated by averaging the correlation coefficients of the set of chemical components that represent the source type. Correlation coefficients showing the

²The COD for two sampling sites is defined as follows:

$$COD_{jk} = \sqrt{\frac{1}{p} \sum_{i=1}^p \left(\frac{x_{ij} - x_{ik}}{x_{ij} + x_{ik}} \right)^2} \quad (3-2)$$

where x_{ij} represents the average concentration for a chemical component i at site j , j and k represent two sampling sites, and p is the number of chemical components.




Figure 3-20. PM_{2.5} chemical components in downtown Los Angeles and Burbank (1986) have similar characteristics. The spread in the data is shown by the bars.

Source: Wongphatarakul et al. (1998).




Figure 3-21. Concentrations of PM_{2.5} chemical components in Rubidoux and downtown Los Angeles (1986). The diagram shows a significant spread in the concentrations for the two sites compared with downtown Los Angeles and Burbank (Figure 3-20).

Source: Wongphatarakul et al. (1998).

spatial relationships among $PM_{2.5}$ (total) and contributions from different source categories obtained at various sites in the South Coast Air Basin (SoCAB) Study are shown in Table 3-6. In Wongphatarakul et al. (1998), crustal material (crustal), motor vehicle exhaust (mv), residual oil emissions (residual oil), and secondary PM (sec) were considered as source categories. Also, in that study, Al, Si, Fe, and Ca were used as markers for crustal material (crustal); V and Ni were used as markers for fuel oil combustion (residual oil); and Pb, Br, and Mn were used as markers for motor vehicle exhaust (mv), based on the lack of other, perhaps more suitable, tracers. NO_3^- , NH_4^+ , and SO_4^{2-} represent secondary PM components. The averages of the correlation coefficients of marker elements within each source category are shown in Table 3-6. Values of r_{sec} and r_{mv} are much higher than those for $r_{crustal}$ and $r_{residual\ oil}$ throughout the SoCAB, suggesting a more uniform distribution of the contributions from secondary PM formation and automobiles than from crustal material and localized stationary sources.

Correlation coefficients in Philadelphia air based on data obtained at four sites for $PM_{2.5}$ (total), crustal components (Al, Si, Ca, and Fe), the major secondary component (sulfate), organic carbon (OC), and elemental carbon (EC) are shown in Table 3-7. Because these data were obtained after Pb had been phased out of gasoline, a motor vehicle contribution could not be estimated from the data. Pb also is emitted by discrete point sources, such as the Franklin smelter. Concentrations of V and Ni were often beneath detection limits; so, the spatial variability in PM due to residual oil combustion was not estimated. Sulfate in aerosol samples collected in Philadelphia arises mainly from long-range transport from regionally dispersed sources (Dzubay et al., 1988). This conclusion is strengthened by the high correlations in sulfate between different monitoring sites and the uniformity in sulfate concentrations observed among the sites. Widespread area sources (e.g., motor vehicle traffic) also may emit pollutants that are correlated between sites provided that traffic patterns and emissions are similar throughout the area under consideration.

Landis et al. (2001) found relatively high correlations between $PM_{2.5}$ ($r = 0.97$), sulfate ($r = 0.99$), OC ($r = 0.97$), EC ($r = 0.83$), NaCl ($r = 0.83$), and nitrate ($r = 0.83$) measured at two sites located several km apart in the Baltimore, MD area. Concentrations of crustal material ($r = 0.63$) and the sum of total metal oxides ($r = 0.76$) were not as well correlated. These results are consistent with those for another eastern city, Philadelphia, PA, given in Table 3-7. The results presented above for Philadelphia, PA; Baltimore, MD; and Los Angeles, CA, indicate

**TABLE 3-6. CORRELATION COEFFICIENTS FOR SPATIAL VARIATION OF
PM_{2.5} MASS AND DIFFERENT SOURCES FOR PAIRS OF SAMPLING
SITES IN THE SOUTH COAST AIR BASIN (1986)**

Sampling Site	r_{total}	r_{crustal}	r_{sec}	r_{mv}	r_{residual oil}
Hawthorne and Rubidoux	-0.027				
Long Beach and Rubidoux	0.051				
Anaheim and Rubidoux	0.066				
Downtown Los Angeles and Rubidoux	0.095				
Burbank and Rubidoux	0.12				
Hawthorne and Anaheim	0.76	0.034	0.768	0.492	0.17
Long Beach and Anaheim	0.852	0.075	0.888	0.504	0.15
Burbank and Anaheim	0.77	0.105	0.749	0.579	0.161
Downtown Los Angeles and Anaheim	0.827	0.143	0.804	0.556	0.233
Downtown Los Angeles and Hawthorne	0.808	0.568	0.854	0.669	0.533
Burbank and Hawthorne	0.704	0.599	0.79	0.688	0.491
Long Beach and Burbank	0.731	0.633	0.737	0.714	0.295
Long Beach and Hawthorne	0.88	0.649	0.909	0.861	0.482
Downtown Long Angeles and Long Beach	0.842	0.653	0.817	0.719	0.378
Downtown Los Angeles and Burbank	0.928	0.825	0.96	0.871	0.606

Source: Wongphatarakul et al. (1998).

**TABLE 3-7. CORRELATION COEFFICIENTS FOR SPATIAL VARIATION OF
PM_{2.5} MASS AND DIFFERENT COMPONENTS FOR PAIRS OF SAMPLING
SITES IN PHILADELPHIA (1994)**

Sampling Site	r_{tot}	r_{crustal}	r_{sec}	r_{OC}	r_{EC}	r_{Pb}
Castor Ave. and Roxboro	0.92	0.52	0.98	0.88	0.84	0.43
Castor Ave. and NE Airport	0.93	0.47	0.99	0.88	0.77	-0.07
Castor Ave. and Broad St.	0.93	0.57	0.99	0.85	0.89	0.11
Roxboro and NE Airport	0.98	0.67	0.98	0.83	0.82	0.2
Roxboro and Broad St.	0.95	0.9	0.98	0.86	0.79	0.47
NE Airport and Broad St.	0.95	0.69	0.99	0.84	0.63	0.11

Source: Pinto et al. (1995).

that secondary PM components are more highly correlated than primary components and may be more highly correlated than total PM_{2.5}. These results suggest that the correlation of PM concentrations across an urban area may depend on the relative proportions of primary and secondary components of PM at individual sites. Sampling artifacts affecting the measurement of nitrate and organic carbon can obscure these relations and may depress correlations between sites.

Kao and Friedlander (1995) examined the statistical properties of a number of PM components in the South Coast Air Basin (Los Angeles area). They found that, regardless of source type and location within their study area, the concentrations of nonreactive, primary components of PM₁₀ had approximately log-normal frequency distributions with constant values of the geometric standard deviations (GSDs). However, aerosol constituents of secondary origin (e.g., SO₄²⁻, NH₄⁺, and NO₃⁻) were found to have much higher GSDs. Surprisingly, the GSDs of organic (1.87) and elemental (1.74) carbon were both found to be within 1 SD (0.14) of the mean GSD (1.85) for nonreactive primary species, compared to GSDs of 2.1 for sulfate, 3.5 for nitrate, and 2.6 for ammonium. These results suggest that most of the OC seen in ambient samples in the South Coast Air Basin was of primary origin. Pinto et al. (1995) found similar results for data obtained during the summer of 1994 in Philadelphia. Further studies are needed to determine if these relations are valid at other locations and to what extent the results might be influenced by sampling artifacts such as the evaporation of volatile constituents during or after sampling.

The use of correlations between OC and EC and between OC to EC ratios based on a comparison between values measured in source emissions and ambient observations has also been suggested as a means to distinguish between secondary and primary sources of OC (Turpin and Huntzicker, 1995; Strader et al., 1999). Ratios of OC to EC from combustion sources are typically < 3 and may even be < 1 in diesel emissions (cf., Appendix 3D). Cabada et al. (2002) concluded that secondary organic PM can contribute from 10 to 35% of total organic PM on an annual basis, with values > 50% during the summer and 0% during the winter months in Pittsburgh, PA based on chemistry-transport model (CTM) results and comparison with emissions inventory values of OC to EC ratios. All of these inferences are subject to considerable uncertainty in the methods for measuring OC, as discussed in Chapter 2. Ambiguity also arises in the ratio method, as the ratios may change due to chemical reactions

occurring during aging of the particles. However, little work has been done on this issue. The ratio can be greater than nine in emissions from wildfires, and transport from distant fires can skew results unless this transport is taken into account (cf., Appendix 3D). Modeling studies that rely on OC to EC ratios in emissions inventories to predict the amount of secondary OC may be biased towards higher ratios of secondary OC because emissions of primary biogenic particles are not included in the inventories. Additional concerns arise from uncertainties in the mechanism of formation of secondary OC from gaseous biogenic and anthropogenic precursor emissions and the uncertainty in those emissions (Section 3.3.1). It is clear, however, that secondary organic PM is being formed in the atmosphere (Blando et al., 1998 and Appendix 3C).

Few studies have compared aerosol composition in urban areas to that in nearby rural areas. One exception is Tanner and Parkhurst (2000), which found that sulfate constituted a larger fraction of fine particle mass at rural sites in the Tennessee Valley PM_{2.5} monitoring network than did OC. For urban sites, the situation was largely reversed: OC constituted a larger fraction of aerosol mass than sulfate. Future systematic comparisons of urban-rural differences in aerosol properties should be facilitated with implementation of the national speciation network and continued operation of the IMPROVE network.

3.3 SOURCES OF PRIMARY AND SECONDARY PM

Information about the nature and relative importance of sources of ambient PM is presented in this section. Table 3-8 summarizes anthropogenic and natural sources for the major primary and secondary aerosol constituents of fine and coarse particles. Anthropogenic sources can be further divided into stationary and mobile sources. Stationary sources include those such as: fuel combustion for electrical utilities, residential space heating, and industrial processes; construction and demolition; metals, minerals, and petrochemicals; wood products processing; mills and elevators used in agriculture; erosion from tilled lands; waste disposal and recycling. Mobile or transportation-related sources include direct emissions of primary PM and secondary PM precursors from highway vehicles and non-road sources as well as fugitive dust from paved and unpaved roads. In addition to fossil fuel combustion, biomass in the form of wood is burned for fuel. Vegetation is burned to clear new land for agriculture and for building construction, to dispose of agricultural and domestic waste, to control the growth of animal or plant pests, and

TABLE 3-8. CONSTITUENTS OF ATMOSPHERIC PARTICLES AND THEIR MAJOR SOURCES¹

Aerosol species	Sources					
	Primary (PM < 2.5 µm)		Primary (PM > 2.5 µm)		Secondary PM Precursors (PM < 2.5 µm)	
	Natural	Anthropogenic	Natural	Anthropogenic	Natural	Anthropogenic
Sulfate (SO ₄ ²⁻)	Sea spray	Fossil fuel combustion	Sea spray	—	Oxidation of reduced sulfur gases emitted by the oceans and wetlands and SO ₂ and H ₂ S emitted by volcanism and forest fires	Oxidation of SO ₂ emitted from fossil fuel combustion
Nitrate (NO ₃ ⁻)	—	—	—	—	Oxidation of NO _x produced by soils, forest fires, and lighting	Oxidation of NO _x emitted from fossil fuel combustion and in motor vehicle exhaust
Minerals	Erosion and reentrainment	Fugitive dust from paved and unpaved roads, agriculture, forestry, construction, and demolition	Erosion and reentrainment	Fugitive dust, paved and unpaved road dust, agriculture, forestry, construction, and demolition	—	—
Ammonium (NH ₄ ⁺)	—	—	—	—	Emissions of NH ₃ from wild animals, and undisturbed soil	Emissions of NH ₃ from animal husbandry, sewage, and fertilized land
Organic carbon (OC)	Wildfires	Prescribed burning, wood burning, motor vehicle exhaust, and cooking	Soil humic matter	Tire and asphalt wear and paved road dust	Oxidation of hydrocarbons emitted by vegetation (terpenes, waxes) and wild fires	Oxidation of hydrocarbons emitted by motor vehicles, prescribed burning, and wood burning
Elemental carbon (EC)	Wildfires	Motor vehicle exhaust, wood burning, and cooking	—	Tire and asphalt wear and paved road dust	—	—
Metals	Volcanic activity	Fossil fuel combustion, smelting, and brake wear	Erosion, reentrainment, and organic debris	—	—	—
Bioaerosols	Viruses and bacteria	—	Plant and insect fragments, pollen, fungal spores, and bacterial agglomerates	—	—	—

¹Dash (—) indicates either very minor source or no known source of component.

to manage forest resources (prescribed burning). Also shown are sources for precursor gases, the oxidation of which forms secondary PM. The atmospheric chemical processes producing secondary PM are described in Section 3.3.1.

In general, the sources of fine PM are very different from those for coarse PM. Some of the mass in the fine size fraction forms during combustion from material that has volatilized in combustion chambers and then recondensed before emission into the atmosphere. Some ambient $PM_{2.5}$ forms in the atmosphere from photochemical reactions involving precursor gases. That PM formed by the first mechanism is referred to as primary, and the PM formed by the second mechanism is referred to as secondary. $PM_{10-2.5}$ is mainly primary in origin, as it is produced by the abrasion of surfaces or by the suspension of biological material. Because precursor gases undergo mixing during transport from their sources, it is difficult to identify individual sources of secondary PM constituents. Transport and transformations of precursors can occur over distances of hundreds of kilometers. The coarse PM constituents have shorter lifetimes in the atmosphere; so their effects tend to be more localized. Only major sources for each constituent within each broad category shown at the top of Table 3-8 are listed. Not all sources are equal in magnitude. Chemical characterizations of primary particulate emissions for a wide variety of natural and anthropogenic sources (as shown in Table 3-8) were given in Chapter 5 of the 1996 PM AQCD. Summary tables of the composition of source emissions presented in the 1996 PM AQCD and updates to that information are provided in Appendix 3D. The profiles of source composition were based in large measure on the results of various studies that collected signatures for use in source apportionment studies.

Natural sources of primary PM include windblown dust from undisturbed land, sea spray, and plant and insect debris. The oxidation of a fraction of terpenes emitted by vegetation and reduced sulfur species from anaerobic environments leads to secondary PM formation. Ammonium (NH_4^+) ions, which play a major role in regulating the pH of particles, are derived from emissions of ammonia (NH_3) gas. Source categories for NH_3 have been divided into emissions from undisturbed soils (natural) and emissions that are related to human activities (e.g., fertilized lands, domestic and farm animal waste). There is ongoing debate about characterizing emissions from wildfires (i.e., unwanted fire) as either natural or anthropogenic. Wildfires have been listed in Table 3-8 as natural in origin, but land management practices and other human actions affect the occurrence and scope of wildfires. For example, fire suppression

practices allow the buildup of fire fuels and increase the susceptibility of forests to more severe and infrequent fires from whatever cause, including lightning strikes. Similarly, prescribed burning is listed as anthropogenic, but can be viewed as a substitute for wildfires that would otherwise occur eventually on the same land.

The transformations that gaseous precursors to secondary PM undergo after being emitted from the sources shown in Table 3-8 are described in Section 3.3.1. Aspects of the transport of primary PM and secondary PM, including the transport of material from outside the United States, are described in Section 3.3.2. A brief introduction to the deposition of particles is also given in Section 3.3.2, and a more detailed discussion of deposition processes is presented in Chapter 4. Methods to infer contributions from different source categories to ambient PM using receptor models and the results of these modeling efforts are given in Section 3.3.3. Estimates of emissions of primary PM and precursors to secondary PM from major sources are presented in Section 3.3.4. A discussion of the uncertainties associated with these emissions is given in Section 3.3.5.

3.3.1 Chemistry of Secondary PM Formation

Precursors to secondary PM have natural and anthropogenic sources, just as primary PM has natural and anthropogenic sources. The major atmospheric chemical transformations leading to the formation of particulate nitrate and sulfate are relatively well understood; whereas those involving the formation of secondary aerosol OC are less so and are still subject to much current investigation. A large number of organic precursors are involved and many of the kinetic details still need to be determined. Also, many of the actual products of the oxidation of hydrocarbons have yet to be identified.

Formation of Sulfates and Nitrates

A substantial fraction of the fine particle mass, especially during the warmer months of the year, is secondary sulfate and nitrate formed as the result of atmospheric reactions. Such reactions involve the gas phase conversion of SO_2 to H_2SO_4 (which forms liquid particles) initiated by reaction with OH radicals and aqueous-phase reactions of SO_2 with H_2O_2 , O_3 , or O_2 (catalyzed by Fe and Mn). These heterogeneous reactions may occur in cloud and fog droplets or in films on atmospheric particles. NO_2 can be converted to gaseous HNO_3 by reaction with

OH radicals during the day. At night, NO_2 also is oxidized to nitric acid by a sequence of reactions initiated by O_3 that produce nitrate radicals (NO_3) and dinitrogenpentoxide (N_2O_5) as intermediates. Both H_2SO_4 and HNO_3 react with atmospheric ammonia (NH_3). Gaseous NH_3 reacts with gaseous HNO_3 to form particulate NH_4NO_3 . Gaseous NH_3 reacts with H_2SO_4 to form acidic HSO_4^- (in NH_4HSO_4) as well as SO_4^{2-} in $(\text{NH}_4)_2\text{SO}_4$. In addition, acid gases such as SO_2 and HNO_3 may react with coarse alkaline particles to form coarse secondary PM containing sulfate and nitrate. Examples include reactions with basic compounds resulting in neutralization (e.g., $\text{CaCO}_3 + 2\text{HNO}_3 \rightarrow \text{Ca}(\text{NO}_3)_2 + \text{H}_2\text{CO}_3 \uparrow$) or with salts of volatile acids resulting in release of the volatile acid (e.g., $\text{SO}_2 + 2\text{NaCl} + \text{H}_2\text{O} \rightarrow \text{Na}_2\text{SO}_3 + 2\text{HCl} \uparrow$).

If particulate NH_4NO_3 coagulates with an acidic sulfate particle (H_2SO_4 or HSO_4^-), gaseous HNO_3 will be released, and the NH_3 will increase the neutralization of the acidic sulfate. Thus, in the eastern United States, where PM tends to be acidic, sulfate usually constitutes a larger fraction of PM mass than nitrate. However, in the western United States, where higher NH_3 and lower SO_2 emissions permit complete neutralization of H_2SO_4 , the concentration of nitrate could be higher than that of sulfate as it is in areas such as the Los Angeles Basin and the San Joaquin Valley. As SO_2 concentrations in the atmosphere in the eastern United States are reduced, the NH_3 left in the atmosphere after neutralization of H_2SO_4 will be able to react with HNO_3 to form NH_4NO_3 . Therefore, a reduction in SO_2 emissions, especially without a reduction in NO_x emissions, could lead to an increase in NH_4NO_3 concentrations (West et al., 1999; Ansari and Pandis, 1998). Thus, the sources, trends, and possible environmental effects of NH_4NO_3 are of interest for both the western and eastern United States.

Chemical reactions of SO_2 and NO_x within plumes are an important source of H^+ , SO_4^{2-} , and NO_3^- . These conversions can occur by gas-phase and aqueous-phase mechanisms. In power-plant or smelter plumes containing SO_2 and NO_x , the gas-phase chemistry depends on plume dilution, sunlight, and VOCs either in the plume or in the ambient air mixing into and diluting the plume. For the conversion of SO_2 to H_2SO_4 in the gas-phase in such plumes during summer midday conditions in the eastern United States, the rate typically varies between 1 and 3%/h, but in the cleaner western United States the rate rarely exceeds 1%/h. For the conversion of NO_x to HNO_3 , the gas-phase rates appear to be approximately three times faster than the SO_2 conversion rates. Winter rates for SO_2 conversion are approximately an order of magnitude lower than summer rates.

The contribution of aqueous-phase chemistry to particle formation in point-source plumes is highly variable, depending on the availability of the aqueous phase (wetted aerosols, clouds, fog, and light rain) and the photochemically generated gas-phase oxidizing agents, especially H_2O_2 for SO_2 chemistry. The in-cloud conversion rates of SO_2 to SO_4^{2-} can be several times larger than the gas-phase rates given above. Overall, it appears that SO_2 oxidation rates to SO_4^{2-} by gas-phase and aqueous-phase mechanisms may be comparable in summer, but aqueous-phase chemistry may dominate in winter. Further details concerning the chemistry of SO_2 and NO_x in power plant plumes can be found in Hewitt (2001).

In the western United States, markedly higher SO_2 conversion rates have been reported in smelter plumes than in power plant plumes. The conversion occurs predominantly by a gas-phase mechanism. This result is attributed to the lower NO_x in smelter plumes. In power plant plumes, NO_2 depletes OH radicals and competes with SO_2 for OH radicals.

In urban plumes, the upper limit for the gas-phase SO_2 conversion rate appears to be about 5%/h under the more polluted conditions. For NO_2 , the rates appear to be approximately three times faster than the SO_2 conversion rates. Conversion rates of SO_2 and NO_x in background air are comparable to the peak rates in diluted plumes. Neutralization of H_2SO_4 formed by SO_2 conversion increases with plume age and background NH_3 concentration. If the NH_3 levels are more than sufficient to neutralize H_2SO_4 to $(\text{NH}_4)_2\text{SO}_4$, the HNO_3 formed from NO_x conversions may be converted to NH_4NO_3 .

Formation of Secondary Organic Particulate Matter (SOPM)

Atmospheric reactions involving VOCs such as alkanes, alkenes, aromatics, cyclic olefins, and terpenes (or any reactive organic gas that contains at least seven carbon atoms) yield organic compounds with low-saturation vapor pressures at ambient temperature. Such reactions may occur in the gas phase, in fog or cloud droplets (Graedel and Goldberg, 1983; Faust, 1994), or possibly in aqueous aerosols (Aumont et al., 2000). Reaction products from the oxidation of reactive organic gases also may nucleate to form new particles or condense on existing particles to form SOPM. Organic compounds with two double bonds or cyclic olefins may react to form dicarboxylic acids, which, with four or more carbon atoms, also may condense. Both biogenic and anthropogenic sources contribute to primary and secondary organic PM (Grosjean, 1992;

Hildemann et al., 1996; Mazurek et al., 1997; Schauer et al., 1996). Oxalic acid was the most abundant organic acid found in PM_{2.5} in California (Poore, 2000).

Although the mechanisms and pathways for forming inorganic secondary PM are fairly well known, those for forming SOPM are not as well understood. Ozone and the OH radicals are thought to be the major initiating reactants. However, HO₂ and NO₃ radicals also may initiate reactions; and organic radicals may be nitrated by HNO₂, HNO₃, or NO₂. Pun et al. (2000) discuss formation mechanisms for highly oxidized, multifunctional organic compounds. The production of such species has been included in a photochemical model by Aumont et al. (2000), for example. Understanding the mechanisms of formation of SOPM is important because SOPM can contribute in a significant way to ambient PM levels, especially during photochemical smog episodes. Experimental studies of the production of secondary organic PM in ambient air have focused on the Los Angeles Basin. Turpin and Huntzicker (1991, 1995) and Turpin et al. (1991) provided strong evidence that secondary PM formation occurs during periods of photochemical ozone formation in Los Angeles and that as much as 70% of the OC in ambient PM was secondary in origin during a smog episode in 1987. Schauer et al. (1996) estimated that 20 to 30%, on an annually averaged basis, of the total organic carbon PM in the < 2.1 μm size range in the Los Angeles airshed is secondary in origin.

Pandis et al. (1992) identified three mechanisms for formation of SOPM: (1) condensation of oxidized end-products of photochemical reactions (e.g., ketones, aldehydes, organic acids, and hydroperoxides), (2) adsorption of semivolatile organic compounds (SVOCs) onto existing solid particles (e.g., polycyclic aromatic hydrocarbons), and (3) dissolution of soluble gases that can undergo reactions in particles (e.g., aldehydes). The first and third mechanisms are expected to be of major importance during the summer when photochemistry is at its peak. The second pathway can be driven by diurnal and seasonal temperature and humidity variations at any time of the year. With regard to the first mechanism, Odum et al. (1996) suggested that the products of the photochemical oxidation of reactive organic gases are semivolatile and can partition themselves onto existing organic carbon at concentrations below their saturation concentrations. Thus, the yield of SOPM depends not only on the identity of the precursor organic gas but also on the ambient levels of OC capable of adsorbing the oxidation products.

Haagen-Smit (1952) first demonstrated that hydrocarbons irradiated in the presence of NO_x produce light scattering aerosols. The aerosol-forming potentials of a wide variety of individual

anthropogenic and biogenic hydrocarbons were compiled by Pandis et al. (1992), based mainly on estimates made by Grosjean and Seinfeld (1989) and on data from Pandis et al. (1991) for β -pinene and from Izumi and Fukuyama (1990) for aromatic hydrocarbons. Zhang et al. (1992) examined the oxidation of α -pinene. Pandis et al. (1991) found no aerosol products formed in the photochemical oxidation of isoprene, although they and Zhang et al. (1992) found that the addition of isoprene to reaction mixtures increased the reactivity of the systems studied. Further details about the oxidation mechanisms and SOPM yields from various reactive organic gases are given in the above studies. Estimates of the production rate of SOPM in the Los Angeles airshed are provided in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996).

More recently, Odum et al. (1997a,b) have found that the aerosol-formation potential of whole gasoline vapor can be accounted for solely by summing the contributions of the individual aromatic compounds in the fuel. In general, data for yields for SOPM formation can be broken into two distinct categories. The oxidation of toluene and aromatic compounds containing ethyl or propyl groups (i.e., ethylbenzene, ethyltoluene, n-propylbenzene) produced higher yields of SOPM than did the oxidation of aromatic compounds containing two or more methyl groups (i.e., xylenes, di-, tri-, tetra-methylbenzenes). Yields in the first group ranged from about 7 to 10%; in the second group, yields ranged from 3 to 4% for OC concentrations between 13 and 100 $\mu\text{g}/\text{m}^3$. Reasons for the differences in SOPM yields found between the two classes of compounds are not clear.

There have been a few recent studies that have examined the composition of secondary organic PM. Edney et al. (2001) carried out a smog chamber study to investigate the formation of multifunctional oxygenates from photooxidation of toluene. The experiments were carried out by irradiating toluene/propylene/ NO_x /air mixtures in a smog chamber operated in the dynamic mode and analyzing the collected aerosol by positive chemical ionization GC-MS after derivatization of the carbonyl oxidation products. The results of the GC-MS analyses were consistent with the formation of semivolatile multifunctional oxygenates, including hydroxy diones as well as triones, tetraones, and pentaones. The authors also suggested that some of these compounds could be present in SOPM in the form of polymers.

Jang and Kamens (2001a) employed a number of analytical approaches, including GC-MS detection of volatile derivatives of carbonyl, hydroxy, and acid compounds in SOPM formed in the irradiation of toluene/ NO_x mixtures. A wide range of substituted aromatics, nonaromatic

ring-retaining and ring-opening products were detected. Newly identified ring-opening oxycarboxylic acids detected included glyoxylic acid; methylglyoxylic acid; 4-oxo-2-butenic acid; oxo-C₅-alkenoic acids; dioxopentenoic acids; oxo-C₇-alkadienoic acids; dioxo-C₆-alkenoic acids; hydroxydioxo-C₇-alkenoic acids; and hydroxytrioxo-C₆-alkanoic acids. Other newly identified compounds included methylcyclohexenetriones; hydroxymethylcyclohexenetriones; 2-hydroxy-3-penten-1,5-dial, hydroxyoxo-C₆-alkenals; hydroxy-C₅-triones, hydroxydioxo-C₇-alkenals; and hydroxy-C₆-tetranones. Included among these compounds were a number of the hydroxy polyketones detected by Edney et al. (2001).

Recent laboratory and field studies support the concept that nonvolatile and semivolatile oxidation products from the photooxidation of biogenic hydrocarbons contribute significantly to ambient PM concentrations in both urban and rural environments. The oxidation of a variety of biogenic hydrocarbons emitted by trees and plants, such as terpenes (α -pinene, β -pinene, Δ^3 -carene, sabinene, α -terpinene, γ -terpinene, terpinolene, myrcene, and ocimene) and sesquiterpenes (β -caryophyllene and α -humulene) could form SOPM. Vegetation also emits oxygenated organic compounds such as alcohols, acetates, aldehyde, ketones, ethers, and esters (Winer et al., 1992). However, their contribution to SOPM remains uncertain. Hoffmann et al. (1997) found SOPM yields of $\approx 5\%$ for open-chain biogenic hydrocarbons such as ocimene and linalool; 5 to 25% for monounsaturated cyclic monoterpenes such as α -pinene, Δ^3 -carene and terpinene-4-ol; and $\approx 40\%$ for a cyclic monoterpene with an endocyclic and an exocyclic double bond such as *d*-limonene. Secondary organic PM yields of close to 100% were observed during the photochemical oxidation of one sesquiterpene, trans-caryophyllene. These results were all obtained for initial hydrocarbon mixing ratios of 100 ppb, which are much higher than found in the atmosphere.

Kamens et al. (1999) observed SOPM yields of 20 to 40% for α -pinene. Using information on the composition of secondary PM formed from α -pinene (Jang and Kamens, 1999), they were able to calculate formation rates with a kinetic model including formation mechanisms for O₃ + α -pinene reaction products.

Griffin et al. (1999) introduced the concept of incremental aerosol reactivity, the change in the secondary organic aerosol mass produced (in $\mu\text{g}/\text{m}^3$) per unit change of parent organic reacted (in ppb), as a measure of the aerosol-forming capability of a given parent organic compound in a prescribed mixture of other organic compounds. They measured the incremental

aerosol reactivity for a number of aromatic and biogenic compounds for four initial mixtures. Incremental aerosol reactivity ranged from 0.133 to 10.352 $\mu\text{g}/\text{m}^3/\text{ppb}$ and varied by almost a factor of two depending on the initial mixture.

A number of multifunctional oxidation products produced by the oxidation of biogenic hydrocarbons have been identified in laboratory studies (Yu et al., 1998; Glasius et al., 2000; Christoffersen et al., 1998; Koch et al., 2000; and Leach et al., 1999). Many of these compounds have subsequently been identified in field investigations (Yu et al., 1999; Kavouras et al., 1998, 1999a,b; Pio et al., 2001; and Castro et al., 1999). Most studies of the formation of secondary organic aerosol formation from terpenes have focused on their reactions with ozone. There have been many fewer studies dealing with the oxidation of terpenes initiated by OH radicals. Larsen et al. (2001) found that the major aerosol products produced ultimately from the reaction of OH radicals with monoterpenes with endocyclic double bonds (α -pinene, 3-carene) were C_{10} keto-carboxylic acids (such as pinonic and caronic acids); whereas the major products from the oxidation of monoterpenes with exocyclic double bonds (β -pinene) were C_9 -dicarboxylic acids (such as pinic acid), and the major product from the oxidation of limonene (which has both endo- and exocyclic double bonds) was 3-acetyl-6-oxo-heptanal (keto-limonaldehyde). A large number of related aldehydes, ketones and acids were also found in their experiments. However, the total yields of condensable products are much lower than for the corresponding reactions with ozone. For example, yields of C_9 -dicarboxylic acids, C_{10} -hydroxy-keto-carboxylic acids, and C_{10} -hydroxy-keto-aldehydes from the reaction of ozone with mono-terpenes with endocyclic double bonds ranged from 3 to 9%, whereas they ranged only from 0.4 to 0.6% in the reaction with OH radicals. Likewise, the reaction of monoterpenes with exocyclic double bonds with O_3 produced much higher yields (1 to 4%) of C_8 - and C_9 -dicarboxylic acids than did their reaction with OH radicals (0.2 to 0.3%). Apart from the complex products noted above, it should be remembered that much simpler products, such as formaldehyde and formic acid, are also formed in much larger yields from the same reactants (e.g., Winterhalter et al., 2000). Compounds such as these also contribute to the formation of secondary organic aerosol according to the mechanisms given in Pandis et al. (1992) and mentioned earlier in this section.

It is worth noting that the dicarboxylic acids and hydroxy-keto-carboxylic acids have very low vapor pressures and may act as nucleating species in OH- and O_3 -terpene reactions (Larsen et al., 2001). The rate coefficient for reaction of α -pinene with OH radicals is approximately a

factor of 10^6 greater than for its reaction with O_3 , based on data given in Atkinson (1994). The daytime average concentration of O_3 is typically a factor of 10^6 greater than that for OH radicals in polluted boundary layers, whereas the above-mentioned yields of aerosol products are roughly a factor of ten greater in the O_3 -initiated reaction than in the corresponding OH radical reaction. The foregoing analysis suggests that the O_3 -initiated reaction may be more important than the OH-initiated reaction for the formation of aerosol products. During the day, new particles may be generated indoors through the infiltration of ambient O_3 ; and, because ambient O_3 is also present at night in lower concentrations, new particles may be generated under these conditions at lower rates. For example, Wainman et al. (2000) found that O_3 can react with limonene released by air fresheners in indoor environments to produce substantial quantities of submicron particles. The corresponding reaction involving OH radicals outdoors at night is expected to be negligible by comparison because of the very low OH concentrations present. Sarwar et al. (2002) estimated indoor OH radical concentrations and suggested that OH in indoor environments is produced mainly by reactions of O_3 transported from outdoors and terpenes emitted from indoor sources. They reported that indoor OH levels ($1-5 \times 10^5$ OH/cm³) are usually lower than typical urban outdoor daytime OH levels ($1-5 \times 10^6$ OH/cm³). However, they can be greater than typical urban outdoor night time OH levels ($1-5 \times 10^4$ OH/cm³). Although much progress has been made in determining the importance of anthropogenic and biogenic hydrocarbons for the formation of SOPM, further investigations are needed to accurately assess their overall contributions to PM_{2.5} concentrations.

Reactions of organic compounds either in particles or on the surface of particles have only come under study during the past 20 years. Tobias and Ziemann (2000) reported evidence for the formation of relatively stable low-volatility peroxy hemiacetals from reactions of hydroperoxides with aldehydes and ketones on the surface of secondary organic particles. Not long after the publication of these results, Jang and Kamens (2001a) suggested, based on results of their outdoor Teflon chamber studies of SOPM formation from irradiation of toluene/propylene/ NO_x /air mixtures, that carbonyls and hydroxy compounds (either within or on the surface of aromatic SOPM) could react together to form larger and less volatile hemiacetals and acetals. They also proposed that dissolved carbonyls could undergo further reactions leading to the formation of a polymer, a mechanism that has also been suggested by Edney et al. (2001). Jang and Kamens (2001b) carried out a series of screening experiments to assess

whether volatile carbonyl compounds adsorbed onto particles could undergo heterogeneous reactions forming low vapor pressure compounds. Experiments were carried out in which aldehydes were introduced in Teflon bags in the dark in the presence of a seed aerosol containing either ammonium sulfate or a mixture of ammonium sulfate and sulfuric acid. The increase in the aerosol volume was then measured using a scanning mobility particle sizer. The aldehydes employed for the study included glyoxal, butanal, hexanal, octanal, and decanal. Increased organic aerosol yields were found in the presence of the ammonium sulfate seed aerosol for each of the carbonyls; the highest yield was found for hexanal followed in decreasing order by glyoxal and then octanal. The presence of the acidified sulfate salt significantly increased the yields even further. In a number of other experiments, 1-decanol was added to the carbonyl-aerosol system to investigate the possible formation of hemiacetals and/or acetals. Again, the volume of aerosol increased in both the presence of ammonium sulfate aerosol and the acidified salt with a significantly larger yield found in the presence of acidity.

To explain their findings for acid-catalyzed carbonyl reactions, Jang and Kamens (2001a,b) proposed a chemical mechanism in which the dissolved carbonyl first undergoes a protonization reaction forming an adduct that can react with water to form its hydrate (1,1-dihydroxy gem-diol). The adducts can then react with OH groups of the gem-diol, forming higher molecular weight and less volatile dimers that are subject to further reactions. In principal, this process, which the authors refer to as a “zipping reaction” can lead to the formation of polymers. However, because the individual reactions are reversible, the process can be reversed by an unzipping reaction. The zipping process could serve as an important mechanism for SOPM formation by converting volatile oxidation products, including glyoxal and methyl glyoxal, into low-volatility compounds. On the other hand, the unzipping process that could take place during the workup of the aerosol samples could be responsible for the detection of highly volatile oxidation products in SOPM (including glyoxal and methyl glyoxal) reported by Edney et al. (2001), Cocker et al. (2001), and Jang and Kamens (2001a). While these processes may take place in the absence of significant acidity, the experimental results suggest that the processes are likely enhanced by acid-catalyzed reactions. Further research is needed to determine the importance of the mechanisms proposed above for the ambient atmosphere.

Sampling and characterizing PM in the ambient atmosphere and in important microenvironments is required to address important issues in exposure, toxicology, and

compliance. Currently, it is not possible to fully quantify the concentration, composition, or sources of the organic components. Many of the secondary organic aerosol components are highly oxidized, difficult-to-measure multifunctional compounds. Additional laboratory studies are needed to identify such compounds, strategies need to be developed to sample and measure such compounds in the atmosphere, and models of secondary organic aerosol formation need to be improved and added to air quality models in order to address issues related to human exposure.

A high degree of uncertainty is associated with all aspects of the calculation of secondary organic PM concentrations. This is compounded by the volatilization of OC from filter substrates during and after sampling as well as potential positive artifact formation from the absorption of gaseous hydrocarbon on quartz filters. Significant uncertainties always arise in the interpretation of smog chamber data because of wall reactions, sampling artifacts, and the use of unrealistically high concentrations of reactants. Limitations also exist in extrapolating the results of smog chamber studies to ambient conditions found in urban airsheds and forest canopies. Concentrations of terpenes and NO_x are much lower in forest canopies (Altshuller, 1983) than the levels commonly used in smog chamber studies. The identification of aerosol products of terpene oxidation has seldom been a specific aim of field studies, making it difficult to judge the results of model calculations of secondary organic PM formation.

Uncertainties also arise because of the methods used to measure biogenic hydrocarbon emissions. Khalil and Rasmussen (1992) found much lower ratios of terpenes to other hydrocarbons (e.g., isoprene) in forest air than were expected based on their relative emissions strengths and rate coefficients for reaction with OH radicals and O_3 . In many cases, reactions with these species are capable of reducing the concentrations of monoterpenes to beneath minimum detection levels as has been found by others in a wide range of North American forest ecosystems (Guenther et al., 1996; Helmig et al., 1998; Geron et al., 2000). Thus, making judgments about the importance of additional loss processes can be a highly problematic exercise, given uncertainties in obtaining relevant OH radical and O_3 concentrations and in the reaction rate coefficients. The ocimenes and sesquiterpenes are estimated to have half lives of seconds to minutes in the presence of ambient O_3 levels, while the pinenes and isoprene have lifetimes ranging from several hours to days. Reaction with hydroxyl radicals is the major sink for isoprene. Khalil and Rasmussen offered two explanations for their findings: (1) either the

terpenes were being removed rapidly by some heterogeneous process, or (2) emissions were enhanced artificially by feedbacks caused by the bag enclosures they used. The isoprene emission rates observed by Khalil and Rasmussen (1992) are reasonably consistent with those found by Geron et al. (2000a and 2001), given inevitable self-shading of much of the foliage within the bag enclosures and also the lower position of the foliage within the accessible portion of the sampled tree crowns. Temperatures were sufficiently warm (≥ 30 °C) and sunny to induce high isoprene emission, although cloudiness from a passing thunderstorm could have markedly reduced these emissions from some of the species. Disturbance or elevated temperatures may also have induced elevated monoterpene emissions from several of the species sampled by Khalil and Rasmussen, because emissions rates reported by them are in many cases 2 to 20 times higher than those reported by others (including those who performed enclosure studies that did not cause elevated temperatures, as well as micrometeorological flux studies that did not disturb the forest canopy). Thus, the somewhat reduced isoprene emissions, combined with elevated monoterpene emissions, can indeed affect the comparison of ambient isoprene versus monoterpene emissions. However, monoterpene compounds recently have been found to undergo heterogeneous reactions on the surface of acid aerosol particles. Further work is needed to assess the importance of these reactions for ambient monoterpene concentrations and for the rate of production of secondary organic PM in forest ecosystems.

3.3.2 Source Contributions to Ambient PM Determined by Receptor Models

Receptor models are perhaps the primary means used to estimate the contributions of different source categories to PM concentrations at individual monitoring sites. Dispersion models (i.e., three-dimensional chemistry and transport models) are formulated in a prognostic manner (i.e., they attempt to predict species concentrations using a tendency equation that includes terms based on emissions inventories, atmospheric transport, chemical transformations, and deposition). Receptor models are diagnostic in their approach (i.e., they attempt to derive source contributions based either on ambient data alone or in combination with data from the chemical composition of sources). These methods have the advantage that they do not invoke all of the uncertainties inherent in emissions inventories or in parameterizing atmospheric transport processes in grid point models.

There are two main approaches to receptor modeling. Receptor models such as the chemical mass balance (CMB) model (Watson et al., 1990a) relate source category contributions to ambient concentrations based on analyses of the composition of ambient PM and source emissions samples. This technique has been developed for apportioning source categories of primary PM and was not formulated to include the processes of secondary PM formation. In the second approach, various forms of factor analysis are used that rely on the varying mix of species present in observations of compositional data to derive both the composition of sources and the source contributions. Few of these techniques actually utilize the information contained in the temporal sequences of the data. Standard approaches such as factor analysis or Principal Component Analysis (PCA) can apportion only the variance, but not the mass, in an aerosol composition data set. The other techniques described below, PMF and UNMIX, do apportion mass, however. Positive matrix factorization (PMF) is a recently developed multivariate technique (Paatero and Tapper, 1993, 1994) that overcomes many of the limitations of standard techniques, such as PCA, by allowing for the treatment of missing data and data near or below detection limits. This is accomplished by weighting elements inversely according to their uncertainties. Standard methods such as PCA weight elements equally regardless of their uncertainty. Solutions also are constrained to yield non-negative factors. Both the CMB and the PMF approaches find a solution based on least squares fitting and minimize an object function. Both methods provide error estimates for the solutions based on estimates of the errors in the input parameters. It should be remembered that the error estimates often contain subjective judgments. For a complete apportionment of mass, all of the major sources affecting a monitoring site must be sampled for analysis by CMB, whereas there is no such restriction in the use of PMF.

Among other approaches, the UNMIX model takes a geometric approach that exploits the covariance of the ambient data to determine the number of sources, the composition and contributions of the sources, and the uncertainties (Henry, 1997). A simple example may help illustrate the approach taken by UNMIX. In a two-element scatter plot of ambient Al and Si, a straight line and a high correlation for Al versus Si can indicate a single source for both species (soil), while the slope of the line gives information on the composition of the soil source. In the same data set, Fe may not plot on a straight line against Si, indicating other sources of Fe in addition to soil. More importantly, the Fe-Si scatter plot may reveal a lower edge. The points

defining this edge represent ambient samples collected on days when the only significant source of Fe was soil. Success of the UNMIX model hinges on the ability to find these “edges” in the ambient data from which the number of source types and the source compositions are extracted. UNMIX uses principal component analysis to find edges in m-dimensional space, where m is the number of ambient species. The problem of finding edges is more properly described as finding hyperplanes that define a simplex. The vertices at which the hyperplanes intersect represent pure sources from which source compositions can be determined. However, there are measurement errors in the ambient data that “fuzz” the edges, making them difficult to find. UNMIX employs an “edge-finding” algorithm to find the best edges in the presence of error. UNMIX does not make explicit use of errors or uncertainties in the ambient concentrations, unlike the methods outlined above. This is not to imply that the UNMIX approach regards data uncertainty as unimportant, but rather that the UNMIX model results implicitly incorporate error in the ambient data. The underlying philosophy is that the uncertainties are often unquantifiable and, hence, it is best to make no *a priori* assumptions about what they are.

In addition to chemical speciation data, Norris et al. (1999) showed that meteorological indices could prove useful in identifying sources of PM responsible for observed health effects (specifically asthma) associated with exposure to ambient PM. They examined meteorology associated with elevated pollution events in Spokane and Seattle, WA and identified a “stagnation index” that was associated with low wind speeds and increases in concentrations of combustion-related pollutants. Their factor analysis also identified a meteorological index (low relative humidity and high temperatures) that was associated with increases in soil-derived PM as well as a third factor (high temperatures and low relative humidity) that was associated with increases in concentrations of particulate sulfate and nitrate species (Norris, 1998). In addition to these variables, past investigations have also used cooling and heating degree days.

Ondov (1996) examined the feasibility of using sensitive isotopic and elemental tracer materials to determine the contributions of petroleum-fueled sources of PM₁₀ in the San Joaquin Valley, CA. Costs of these experiments are affected not only by the expense for tracer materials, but also by the sensitivities of the analytical methods for each tracer, as well as their background levels. Suarez et al. (1996) used iridium as a tracer to tag emissions from diesel-burning sanitation trucks in Baltimore and determined the size distribution of soot from the trucks.

A number of specialty conference proceedings, review articles, and books have been published that provide greater detail about source category apportionment receptor models than described in the 1996 PM AQCD. A review of the various methods used to apportion PM in ambient samples among its source categories was given in Section 5.5.2 of the 1996 PM AQCD. The collection of the source category characterization profiles shown in Appendix 3D has been motivated in many cases by the need to use them in receptor modeling applications.

The results of several source apportionment studies are discussed in this section to provide an indication of the relative importance of different sources of ambient PM across the United States. First, results obtained mainly by using the CMB approach for estimating contributions to $PM_{2.5}$ from different source categories at monitoring sites in the United States are discussed and presented in Table 3-9. More recent results using the PMF approach are included for Phoenix, AZ. Results obtained at a number of monitoring sites in the central and western United States by using the CMB model for PM_{10} are shown in Table 3-10. The sampling sites represent a variety of different source characteristics within different regions of Arizona, California, Colorado, Idaho, Illinois, Nevada, and Ohio. Definitions of source categories also vary from study to study. The results of the PM_{10} source apportionment studies were given in the 1996 PM AQCD and are presented here to allow easy comparison with results of $PM_{2.5}$ source apportionment studies. Chow and Watson (2002) present a detailed comparison of numerous studies using the CMB model performed mainly after 1995. Their comprehensive review includes the results of a number of studies that have been published as reports, such as the Northern Front Range Air Quality Study, in addition to those that have appeared in peer-reviewed journals.

There are several differences between the broadly defined source categories shown at the tops of Tables 3-9 and 3-10. These differences reflect the nature of sources that are important for producing the fine and coarse PM shown in Table 3-8. They also are related to improvements in the ability to distinguish between sources of similar nature (e.g., diesel and gasoline vehicles, meat cooking, and vegetation burning). The use of organic tracers allows motor vehicle emissions to be broken down into contributions from diesel and gasoline vehicles. In studies where this distinction cannot be made, the source type is listed as 'total motor vehicles' in the tables. The studies that were reported to be able to distinguish gasoline- from diesel-fueled vehicles reported mixed results for the contributions from gasoline and diesel

TABLE 3-9. RECEPTOR MODEL SOURCE CONTRIBUTIONS TO PM_{2.5}

Sampling Site	% Contribution												Total % Allocated
	Measured PM _{2.5} Concentration	Total Motor Vehicles	Diesel	Gasoline Vehicles	Road Dust, Soil	Vegetation Burning	Secondary Sulfate	Secondary Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	
Pasadena, CA 1982 ¹	28.2	—	18.8	5.7	12.4	9.6	20.9	7.4	5.3 ^a	9.2 ^b	8.5 ^c	1.1 ^d	98.9
Downtown LA, CA 1982 ¹	32.5	—	35.7	6.5	11.1	5.8	20.3	9.2	3.7 ^a	9.2 ^b	5.2 ^c	0.6 ^d	107.3
West LA, CA 1982 ¹	24.5	—	18	5.7	12.2	11	24.1	7.8	4.1 ^a	9.4 ^b	8.2 ^c	1.6 ^d	102.1
Rubidoux, CA 1982 ¹	42.1	—	12.8	0.7	13.1	1.2	13.8	24.7	4.5 ^a	12.1 ^b	4.5 ^c	0.5 ^d	87.9
Sacramento, CA Winter 1991-96 ²	39.5	24.5	—	—	1.2	18.1	4.5	36.6	—	—	—	—	84.9
Bakersfield, CA Winter 1996 ³	52	16	—	—	< 3	20	7	34	—	—	—	—	< 80
Fresno, CA Winter 1996 ³	63	13	—	—	< 3	19	5	32	—	—	—	—	< 85
Philadelphia, PA Summer 1982 ⁴	27	8.5	—	—	4.4	—	81.9 ^f	—	2.2 ^g	1.9 ^h	0.4 ⁱ	—	99.3
Camden, NJ Summer 1982 ⁴	28.3	9.2	—	—	3.2	—	81.3 ^f	0.4	2.5 ^g	2.5 ^h	0.7 ⁱ	—	99.8
Clarksboro, NJ Summer 1982 ⁴	26	5.8	—	—	2.7	—	84.6 ^f	—	0.8 ^g	1.5 ^h	0.4 ⁱ	—	95.8
Grover City, IL ENE ^j 1986-87 ⁵	—	—	—	—	2.3	—	83.2 ^f	—	9.7 ^k	3.0 ^l	1.2 ^g	—	99.4
Grover City, IL SSW ^j 1986-87 ⁵	—	—	—	—	—	—	59.0 ^f	—	11.6 ^k	11.9 ^l	4.1 ^g	4.6 ^m	91.2
Grover City, IL WNW ^j 1986-87 ⁵	—	2.4	—	—	5.1	—	88.5 ^f	—	2.8 ^k	—	—	—	98.8
Grover City, IL NNW ^j 1986-87 ⁵	—	—	—	—	3.1	—	86.6 ^f	—	3.4 ^l	3.0 ⁿ	—	—	96
Reno, NV Summer 1998 ⁶	7.8	68	—	—	14.5	4	11	2	0.6 ^g	—	—	—	100.1
Phoenix, AZ Summer 1995-98 ⁷	8.3	—	10.9	36.2	1.8	15	—	—	20.8 ⁿ	4.9 ^r	6.7 ^s	3.6 ^q	99.9
Phoenix, AZ Winter 1995-98 ⁷	13.8	—	14.5	38.9	1.1	8.9	—	—	9.5 ⁿ	4.5 ^r	18.7 ^s	4.1 ^q	100.2

¹Schauer et al. (1996)

²Motallebi (1999)

³Magliano et al. (1998)

⁴Dzubay et al (1988)

⁵Glover et al. (1991)

⁶Gillies et al. (2000)

⁷Ramadan et al. (2000)

^aSecondary and other organic compounds

^bSecondary ammonium

^cMeat cooking

^dVegetative detritus

^eValue represents sum of diesel and gasoline vehicle exhaust

^fIncluding associated cations and water

^gIncinerators

^hOil fly ash

ⁱFluidized catalyst cracker

^jWind direction

^kLead smelter

^lIron works

^mCopper smelter

ⁿCoal power plant

^oAs ammonium sulfate

^pAs ammonium nitrate

^qSea salt

^rWood burning

^sNonferrous smelting

TABLE 3-10. RECEPTOR MODEL SOURCE CONTRIBUTIONS TO PM₁₀

Sampling Site	Measured PM ₁₀ Concentration	% Contribution										Total % Allocated
		Primary Geological	Primary Construction	Primary Motor Vehicle Exhaust	Primary Vegetative Burning	Secondary Ammonium Sulfate	Secondary Ammonium Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	
Craycroft, AZ Winter 1989-1990 ¹	23.4	55.6	0	35.5	0.0	3	2.6	5.1 ^a	0.0	0	0.0	101.8
Hayden 1, AZ 1986 ¹	105.0	4.8	1.9 ^b	0.0	0.0	3.8	0.0	70.5 ^c	4.8 ^d	1.0 ^e	0.0	86.6
Hayden 2, AZ 1986 ¹	59.0	35.6	6.8 ^b	0.0	0.0	6.8	0.0	47.5 ^c	0.0	1.7	0.0	98.4
Rillito, AZ 1988 ²	79.5	53.7	17.4 ^b	1.5 ^f	0.0	0	0	14.6 ^e	0.0	0	0.0	87.2
Bakersfield, CA 1988-1989 ³	79.6	53.9	2	9.7	8.2	6.9	16	1.3 ^m	1.9 ⁿ	0.8 ^k	0.0	100.7
Crows Landing, CA 1988-1989 ³	52.5	61.3	0.0	4.2	6.5	5.3	12.4	1.0 ^m	1.9 ⁿ	2.3 ^k	0.0	94.9
Fellows, CA 1988-1989 ³	54.6	53.1	2.6	3.8	6.2	9.3	13.7	12.8 ^m	2.6 ⁿ	2.6 ^k	0.0	106.7
Fresno, CA 1988-1989 ³	71.5	44.5	0.0	9.5	7.1	5	14.5	0.4 ^m	1.9 ⁿ	0.1 ^k	0.0	83
Indio, CA ⁴	58.0	56.9	5.2	7.6	12.2	6.2	7.1	0.3 ^j	1.7 ^h	0.0	0.0	97.2
Kern Wildlife Refuge, CA 1988-1989 ³	47.8	31.6	4.2	4.6	8.4	6.9	3.1	1.0 ^m	3.1 ⁿ	1.5 ^k	0.0	64.4
Long Beach, CA 1986 ⁵	51.9	39.9	0.0	9.8 ⁱ	0.0	15.4	17.7	0.2 ^j	3.9 ^h	12.3 ^k	0.0	63.2
Long Beach, CA Summer 1987 ⁶	46.1	24.1	0.0	13.7	0.0	23.6	1.7	0.2 ^j	4.8 ^h	0.0	0.0	68.1
Long Beach, CA Fall 1987 ⁶	96.1	11.8	0.0	44.5	0.0	4	24.1	0.0 ^j	2.8 ^h	0.0	0.0	87.2
Riverside, CA 1988 ⁷	64	50.9	0.0	10.9	0.0	7.5	33.4	0.5 ^j	2.0 ^h	1.7 ^o	0.0	106.9
Rubidoux, CA 1986 ⁵	87.4	49.3	4.6	6.4 ⁱ	0.0	7.3	24.4	0.3 ^j	1.1 ^h	6.8 ^k	0	100.2
Rubidoux, CA Summer 1987 ⁶	114.8	30.4	3.9	15.1	0	8.3	23.9	0.0 ^j	4.4 ^h	0.0	0	86

TABLE 3-10 (cont'd). RECEPTOR MODEL SOURCE CONTRIBUTIONS TO PM₁₀

Sampling Site	Measured PM ₁₀ Concentration	% Contribution										Total % Allocated
		Primary Geological	Primary Construction	Primary Motor Vehicle Exhaust	Primary Vegetative Burning	Secondary Ammonium Sulfate	Secondary Ammonium Nitrate	Misc. Source 1	Misc. Source 2	Misc. Source 3	Misc. Source 4	
Rubidoux, CA Fall 1987 ⁶	112	17.1	14.4	27.1	0.0	1.9	28.2	0.0 ^j	1.0 ^h	0	0	89.7
Rubidoux, CA 1988 ⁷	87	55.2	0.0	11.7	0.0	6.1	24.9	0.6 ^j	1.7 ^h	6.6 ^o	0.0	106.8
San Nicolas Island, CA Summer 1987 ⁶	17.4	9.2	0	5.2	0.0	21.3	2.9	0.0 ^j	24.7 ^h	0	0	63.3
Stockton, CA 1989 ³	62.4	55.1	0.8	8.3	7.7	5	11.2	1.1 ^m	2.9 ⁿ	0.0 ^k	0	92.1
Pocatello, ID 1990 ⁸	100.0	8.3	7.5 ^a	0.1	0.0	0	0	0	0.0	84.1 ^r	0	100
S. Chicago, IL 1986 ⁹	80.1	34	3	3.5	0.0	19.2 ^s	—	18.9 ^t	2.7 ^u	0.0	0.0	81.3
S.E. Chicago, IL 1988 ¹⁰	41	35.9 ^v	0.0	2.2 ^f	0.0	18.8	—	2.0 ^t	0.7 ^h	2.7 ^w	18.8 ^g	81.1
Reno, NV 1986-87 ¹¹	30.0	49.7	0.0	33.3	6.3	4.3	2	0.0	0.0	0.0	0.0	95.6
Sparks, NV 1986-87 ¹¹	41	36.8	0.0	28.3	32.7	6.6	2.2	0.0	0.0	0.5 ^k	0.0	107.1
Follansbee, WV 1991 ¹²	66.0	15.2	0.0	53	0.0	24.2	—	14.1 ^l	0.0	0.0	0.0	106.5
Mingo, OH 1991 ¹²	60.0	20	0.0	23.3	6.8	25	—	5.7 ^t	18.3 ^x	0.0	0	99.1
Steubenville, OH 1991 ¹²	46	18	0.0	30.4	1.7	30.4	—	8.3 ^t	10.9 ^x	0.0	0.0	99.7

3-78

¹Chow et al. (1992a)

²Garfield; Ryan et al. (1988)

³Jail; Ryan et al. (1988)

⁴Thanukos et al. (1992)

⁵Chow et al. (1992b)

⁶Kim et al. (1992)

⁷Gray et al. (1988)

⁸Watson et al. (1994)

⁹Chow et al. (1992c)

¹⁰Houck et al. (1992)

¹¹Hopke et al. (1988)

¹²Vermette et al. (1992)

¹³Chow et al. (1988)

¹⁴Skidmore et al. (1992)

^aSmelter background aerosol

^bCement plant sources, including kiln stacks, gypsum pile, and kiln area

^cCopper ore

^dCopper tailings

^eCopper smelter building

^fHeavy-duty diesel exhaust emission

^gBackground aerosol

^hMarine aerosol, road salt, and sea salt plus sodium nitrate

ⁱMotor vehicle exhaust from diesel and leaded gasoline

^jResidual oil combustion

^kSecondary organic carbon

^lBiomass burning

^mPrimary crude oil

ⁿNaCl + NaNO₃

^oLime

^pRoad sanding material

^qAsphalt industry

^rPhosphorus/phosphate industry

^sRegional sulfate

^tSteel mills

^uRefuse incinerator

^vLocal road dust, coal yard road dust, and steel haul road dust

^wIncineration

^xUnexplained mass

vehicles. In the study in southern California, diesels made a larger contribution to $PM_{2.5}$ than did gasoline-fueled vehicles. On the other hand, the study in Phoenix, AZ, found the opposite result. Meat cooking is also distinguished from vegetation burning in more recent studies although both are considered to be part of biomass burning. Vegetation burning consists of contributions from residential fuel wood burning, wildfires, prescribed burning, and burning of agricultural and other biomass waste. Miscellaneous sources of fine particles include contributions from combustion sources, whereas miscellaneous sources of coarse particles consist of contributions from soil and sea spray and industrial processing of geological material (e.g., cement manufacturing). Although a large number of elements and chemical components are used to differentiate among source categories and although there can be a large number of source types affecting a given site, only a few broadly defined source types are needed to account for most of the mass of $PM_{2.5}$ and PM_{10} . At any given site, ≤ 5 source types account for $> 65\%$ of the mass of $PM_{2.5}$ (Table 3-9); and ≤ 5 source types account for $> 65\%$ of the mass of PM_{10} (Table 3-10).

Secondary sulfate is the dominant component of $PM_{2.5}$ samples collected in the studies of Dzubay et al. (1988) and Glover et al. (1991). Both studies found that sulfate at their monitoring site arose from regionally dispersed sources. Sulfate also represented the major component of $PM_{2.5}$ found in monitoring studies in the eastern United States shown in Appendix 6A of the 1996 PM AQCD. Primary and secondary organic components also make major contributions to $PM_{2.5}$ and in many areas may represent a larger component than sulfate. Contributions from road dust and soils are relatively minor, typically constituting $< 10\%$ of $PM_{2.5}$ in the studies shown in Table 3-9. Studies in the western United States shown in Table 3-9 have found larger contributions from motor vehicles, fugitive dust, and ammonium nitrate. The most notable difference in the relative importance of major source categories of $PM_{2.5}$ shown in Table 3-9 and PM_{10} shown in Table 3-10 involves crustal material, (e.g., soil, road dust), which represents about 40% on average of the total mass of PM_{10} in the studies shown in Table 3-10. The fraction is higher at sites located away from specific sources, such as sea spray or smelters. Emissions of crustal material are concentrated mainly in the $PM_{10-2.5}$ size range.

In Table 3-10, the magnitude of primary motor vehicle exhaust contributions is highly variable and ranges from 5 to 40% of average PM_{10} . Vehicle exhaust contributions are also variable at different sites within the same study area. The mean value and the variability of motor vehicle exhaust contributions reflects the proximity of sampling sites to roadways and

traffic conditions during the time of sampling. Many studies were conducted during the late 1980s when a portion of the vehicle fleet still used leaded gasoline. Pb and Br in motor vehicle emissions facilitated the distinction of motor vehicle contributions from other sources. Vehicles using leaded fuels have higher emission rates than vehicles using unleaded fuels. Pb also poisons automobile exhaust catalysts and produces adverse human health effects. As a result, Pb has been eliminated from vehicle fuels. Organic species such as *n*-pentacosane through *n*-nonacosane, cholestanes, ergostanes, sitostanes, and hopanes have replaced Pb as a source marker for motor vehicle emissions (e.g., Schauer and Cass, 2000). In their comprehensive review of CMB modeling studies undertaken since 1995, Chow and Watson (2002) note that, in 22 studies, fossil fuel combustion was found to be a large contributor to PM_{2.5} and PM₁₀ concentrations, with most of the contributions to primary PM originating from the exhaust of diesel and gasoline vehicles.

Marine aerosol is found, as expected, at coastal sites such as Long Beach (average 3.8% of total mass) and San Nicolas Island (25%). These contributions to PM₁₀ are relatively variable and are larger at the more remote sites. Individual values reflect proximity to local sources. Of great importance are the contributions from secondary ammonium sulfate in the eastern United States and ammonium nitrate in the western United States. Secondary ammonium sulfate is especially noticeable at sites in California's San Joaquin Valley (Bakersfield, Crows Landing, Fellows, Fresno, and Stockton) and in the Los Angeles area.

Because many source apportionment studies address problems in compliance with the National Ambient Air Quality Standards (NAAQS) and other air quality standards, samples selected for chemical analysis are often biased toward the highest PM₁₀ mass concentrations in the studies shown in Table 3-10. Thus, the average source contribution estimates shown in Table 3-10 are probably not representative of annual averages and may not be representative of a large spatial area for some source-dominated studies. For example, the study by Motallebi (1999) considered only days when the PM₁₀ concentration was greater than 40 µg/m³. Quoted uncertainties in the estimated contributions of the individual sources shown in Tables 3-9 and 3-10 range from 10 to 50%. Errors can be much higher when the chemical source profiles for different sources are highly uncertain or are too similar to distinguish one source from another.

Very few source apportionment studies using the CMB modeling technique have examined the spatial variability of source contributions at different sites within an urban area. As can be

seen from Table 3-9, Dzubay et al. (1988) found a uniform distribution of sulfate at the NE Airport in Philadelphia, PA, downtown Camden, NJ, and Clarksboro, NJ during the summer of 1982. The farthest distance between two monitoring sites (NE Airport and Clarksboro) was ~40 km. Magliano et al. (1998) examined the spatial variability of PM₁₀ source contributions at a number of sites in Fresno and Bakersfield, CA during the winter of 1995-1996 and reported values for 1 day (December 27, 1995). During that day, mobile sources contributed from 13.0 to 15.8 µg/m³, vegetation burning 5.1 to 11.1 µg/m³, ammonium sulfate 2.4 to 3.4 µg/m³, and ammonium nitrate 19.3 to 24.6 µg/m³ to PM₁₀ at the sites in Bakersfield. Mobile sources contributed 13.9 to 22.5 µg/m³, vegetation burning 8.2 to 15.7 µg/m³, ammonium sulfate 1.8 to 2.3 µg/m³, and ammonium nitrate 14.5 to 18.9 µg/m³ at the sites in Fresno. All of these components are expected to be found mainly in the PM_{2.5} size fraction. As can be seen, source contributions at different sites varied by factors of 1.2 to 2.2 in Bakersfield and by factors of 1.3 to 1.9 in Fresno on that day.

The receptor modeling methods outlined above do not explicitly include consideration of the distances between PM sources and the receptor site. Information about the relative importance of sources as a function of distance may be available from examination of data obtained by continuous monitoring methods. For example, concentration spikes are expected to be the result of transport from nearby sources because turbulent mixing in the atmosphere would not allow them to persist for very long. Short duration spikes in the time series of concentrations are assumed to result from emissions from local sources (0.1 to 1 km away) in this method. Contributions from sources located farther away are determined by comparisons between baselines measured at different sites. Details such as these are also lost in integrated 24-h samples. Watson and Chow (2001) used time series of black carbon (BC) obtained by aetholometers over 5-min intervals to estimate the contributions from sources located < 1 km away, 1 to 5 km away, and > 5 km away from a monitoring site in downtown Mexico City. They found that most of the BC was produced by sources scattered throughout the city and that sources located less than 1 km away from the site contributed only about 10% to BC concentrations even in the presence of local sources such as buses and trucks.

3.3.3 Background Concentrations of PM in the United States

This section contains information about the concentrations of “background” PM that are relevant for policy setting. For the purposes of this document (and consistent with the 1996 PM AQCD), policy relevant background (PRB) concentrations are those that would result in the United States from emissions from natural sources worldwide plus anthropogenic sources outside of North America. In other words, they are the concentrations that would exist if anthropogenic emissions in North America were zero. PRB concentrations of $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10} may be viewed as coming from two conceptually separate components: a reasonably consistent “baseline” component and an episodic component. The baseline component consists of the contribution from natural sources within North America and from transport of natural and anthropogenic sources outside of North America that is generally well characterized by a reasonably consistent distribution of daily values each year, although there is variability by region and season. The episodic component consists of infrequent, sporadic contributions from natural high-concentration events over shorter periods of time (e.g., hours to several days) both within North America (e.g., volcanic eruptions, large forest fires, dust storms) and outside of North America (e.g., transport related to dust storms from deserts in North Africa and China and storms at sea). These episodic natural events, as well as events like the uncontrolled biomass burning in Central America and southern Mexico, are essentially uncontrollable and do not necessarily occur in all years. The same processes that are responsible for transporting these emissions also transport anthropogenic emissions from outside North America.

It is impossible to reliably estimate PRB concentrations solely by examining mass measurements of $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10} . Rather, this section examines available monitoring data, source apportionment analysis, and information about various source contributions to inform a characterization of these two components of PRB PM concentrations and the associated uncertainties and limitations in estimating PRB concentrations. Appendix 3E presents data from relatively remote IMPROVE sites to provide some rough upper limits on PRB concentrations for the United States.

Annual average natural background concentrations of PM_{10} have been estimated to range from 4 to 8 $\mu\text{g}/\text{m}^3$ in the western United States and 5 to 11 $\mu\text{g}/\text{m}^3$ in the eastern United States. Corresponding $PM_{2.5}$ levels have been estimated to range from 1 to 4 $\mu\text{g}/\text{m}^3$ in the western United States and from 2 to 5 $\mu\text{g}/\text{m}^3$ in the eastern United States (U.S. Environmental Protection

Agency, 1996). The range of the natural background concentration of $PM_{10-2.5}$ can be roughly estimated by subtracting the minimum $PM_{2.5}$ value [$1 \mu\text{g}/\text{m}^3$ (western U.S.) and $2 \mu\text{g}/\text{m}^3$ (eastern U.S.)] from the maximum PM_{10} value [$8 \mu\text{g}/\text{m}^3$ (western U.S.) and $11 \mu\text{g}/\text{m}^3$ (eastern U.S.)] and by subtracting the maximum $PM_{2.5}$ value [$4 \mu\text{g}/\text{m}^3$ (western U.S.) and $5 \mu\text{g}/\text{m}^3$ (eastern U.S.)] from the minimum PM_{10} value [$4 \mu\text{g}/\text{m}^3$ (western U.S.) and $5 \mu\text{g}/\text{m}^3$ (eastern U.S.)]. These calculations indicate that the average $PM_{10-2.5}$ concentration is ≤ 1 to $7 \mu\text{g}/\text{m}^3$ in the western United States and ≤ 1 to $9 \mu\text{g}/\text{m}^3$ in the eastern United States. The estimated natural background concentrations given above do not include contributions from long-range transport from sources outside North America. [The range of annual mean $PM_{2.5}$ concentrations at IMPROVE network sites in the western United States (cf., Appendix E) is consistent with the range of values in the lowest 5th percentile annual mean $PM_{2.5}$ concentrations for specific sites in the AIRS data base ($2.8 \mu\text{g}/\text{m}^3$ to $6.9 \mu\text{g}/\text{m}^3$)]. However, $PM_{2.5}$ concentrations are much higher at sites in the eastern United States than at sites in the western United States. At most IMPROVE sites in the western United States, the annual mean concentration of $PM_{10-2.5}$ is higher than that of $PM_{2.5}$, and daily average $PM_{2.5}$ concentrations are moderately correlated ($r = 0.72$) with $PM_{10-2.5}$ concentrations. In contrast, $PM_{2.5}$ concentrations are higher than those of $PM_{10-2.5}$ at IMPROVE sites in the eastern United States, and $PM_{2.5}$ concentrations are only weakly correlated ($r = 0.26$) with those of $PM_{10-2.5}$. As discussed above, peak 24-h average concentrations due to natural sources may be substantially higher than the annual or seasonal average concentrations from natural sources, especially within areas affected by wildfires and dust storms.

Long-Range Transport from Outside North America

Windblown dust from dust storms in the Sahara desert has been observed in satellite images as plumes crossing the Atlantic Ocean and reaching the southeast coast of the United States (e.g., Ott et al., 1991). Dust transport from the deserts of Asia across the Pacific Ocean also occurs (Prospero, 1996). Most dust storms in the deserts of China occur in the spring following the passage of strong cold fronts after the snow has melted and before a surface vegetation cover has been established. Strong winds and unstable conditions result in the rapid transport of dust to altitudes of several kilometers, where it is transported by strong westerly winds out over the Pacific Ocean (Duce, 1995). Satellite images were used to track the progress

of a dust cloud from the Gobi desert to the northwestern United States during the spring of 1998 (Husar et al., 2000).

Satellite images obtained at visible wavelengths cannot track mineral dust across the continents because of a lack of contrast between the plume and the underlying surface. Other means must be used to track the spread of North African dust through the eastern United States. Perry et al. (1997) used two criteria ($PM_{2.5}$ soil concentration $> 3 \mu\text{g}/\text{m}^3$ and $\text{Al}:\text{Ca} > 3.8$) to distinguish between soil of local origin from soil originating in North Africa in characterizing the sources of PM in aerosol samples collected in the IMPROVE network. North African dust has been tracked as far north as Illinois (Gatz and Prospero, 1996) and Maine (Perry et al., 1997). The analysis of Perry et al. (1997) indicates that incursions of Saharan dust into the continental United States have occurred, on average, about three times per year from 1992 to 1995. These events persist for ~ 10 days on average, mainly during the summer. Large-scale dust events typically cover from 15 to 30% of the area of the continental United States and result in $\sim 8.7 \pm 2.3 \mu\text{g}/\text{m}^3$ increases of $PM_{2.5}$ levels throughout the affected areas during these events, with mean maximum dust contributions of $19.7 \pm 8.4 \mu\text{g}/\text{m}^3$ and a peak contribution of $32 \mu\text{g}/\text{m}^3$ to 24-h average $PM_{2.5}$ levels.

As can be expected, the frequency of dust events is highest in the southeastern United States. About half of these events are observed only within the state of Florida and are associated with dense hazes in Miami (Figure 3-22) during the summer (Prospero et al., 1987). North African dust is the dominant aerosol constituent in southern Florida during the summer, whereas soil dust constitutes only a minor fraction of PM during the remainder of the year (Prospero, 1999). Approximately 33 to 50% of the mass of the particles reaching southern Florida have aerodynamic diameters $< 2.5 \mu\text{m}$ (Prospero et al., 2001). During episodes when daily total dust concentrations ranged up to $100 \mu\text{g}/\text{m}^3$, it can be seen that daily $PM_{2.5}$ values of up to $50 \mu\text{g}/\text{m}^3$ could have resulted in Miami, FL.

Husar et al. (2001) documented the transport of dust from the Gobi and Taklimakan deserts to North America during April 1998. The PM_{10} concentration averaged over 150 stations in Washington, Oregon, California, Nevada, and Idaho reporting data to AIRS was $65 \mu\text{g}/\text{m}^3$ between April 26 and May 1, compared to about $20 \mu\text{g}/\text{m}^3$ during the rest of April and May. Data from several networks indicated that PM_{10} concentrations were over $100 \mu\text{g}/\text{m}^3$ in central British Columbia, Washington State, and Oregon. The highest PM concentrations observed




Figure 3-22. Monthly average Saharan dust components in all size fractions of the aerosol sampled in Miami, FL (from 1974 to 1996). Approximately one-third to one-half of Saharan dust is in the PM_{2.5} size range.

Source: Prospero (1999).

were 120 µg/m³ for PM₁₀ and 50 µg/m³ for PM_{2.5} at Chilliwack Airport in northwestern Washington State (Figure 3-23). Aircraft measurements made over the northwestern United States were consistent with a mass median diameter of the dust being between 2 and 3 µm.

Desert dust deposited over oceans provides nutrients to marine ecosystems (Savoie and Prospero, 1980), and such dust deposited on nutrient-depleted soils also provides nutrients to terrestrial ecosystems, e.g., Hawaiian rain forests (Chadwick et al., 1999). Microorganisms, including fungi and bacteria, have been found attached to African dust particles in the U.S. Virgin Islands (Griffin et al., 2001). The fungus, *Aspergillus sydowii*, which has been connected to the death of coral reefs, has been identified in air samples collected in the Caribbean during African dust transport events (Smith et al., 1996; Shinn et al., 2000). Measurements of the




Figure 3-23. PM_{2.5} and PM₁₀ concentrations measured at Chilliwack Airport, located in southwestern British Columbia, just before and during the Asian desert dust episode of April and May 1998.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

composition of Saharan dust in Miami indicate enhancements of nitrate, non-sea salt sulfate, ammonium, and trace metals above concentrations expected for clean marine air, suggesting pollution emitted in Europe and North Africa as sources (Prospero, 1999). It is likely that many other constituents will be found associated with dust from outside North America as more measurements are made. It should be noted that, as North African dust and associated material are transported northward through the United States during the summer, they are added to the mixture of primary and secondary PM generated domestically.

The transport of PM from uncontrolled biomass burning in Central America and southern Mexico resulted in anomalously high PM levels observed in southern Texas and generally elevated PM concentrations throughout the entire central and southeastern United States during

the spring and early summer of 1998. Biomass burning for agricultural purposes occurs normally during the spring of each year in Central America and southern Mexico. During the spring of 1998, fires burned uncontrollably because of abnormally hot and dry conditions associated with the intense El Niño of 1997-1998. PM₁₀ concentrations observed in the southern Rio Grande Valley were elevated substantially during the passage northward of the biomass burning plume produced by these fires, as shown in Figure 3-24. Elevated PM₁₀ concentrations also were found as far north as St. Louis, MO (Figure 3-25). As can be seen from Figure 3-24 and Figure 3-25, the elevations in PM concentrations were limited in duration. Uncontrolled wildfires occur in the United States every year, but their effects on air quality throughout the United States still need to be evaluated systematically. These fires can be widespread, and the frequency of their occurrence can vary markedly from year to year. For example, ~26,000 km² of forested land were consumed during 2000, but only a small fraction of this area was burnt during 2001 in the western United States. Wildfires also occur throughout the boreal forests of Canada. Wotawa and Trainer (2000) suggested that the plume from fires occurring in the Northwest Territories of Canada in early July 1995 may have extended throughout most of the eastern United States, resulting in elevated levels of CO and O₃. Simple scaling of their calculated excess CO concentrations because of the fires, by the ratio of emission factors of PM_{2.5} to CO, indicates that the excess PM_{2.5} concentrations in the plume may have ranged from ~5 µg/m³ in the Southeast and to close to 100 µg/m³ in the northern Plains States.

Sources Within North America

It is much more difficult to determine 24-h PRB concentrations in the absence of specific events such as those noted above, because contributions from anthropogenic sources located either nearby or elsewhere within North America can contribute substantially to observed values and perhaps overwhelm the contributions from PRB sources. Source-apportionment modeling techniques (described in Section 5.5 of the 1996 PM AQCD and updated in Section 3.3.2 of this document) can be helpful for this purpose. It is more likely that PRB concentrations can be more accurately determined in the West than in the East because contributions from pollution sources can overwhelm those from background sources in the East, and background source contributions can be lost within the errors of their source contributions. However, it should be noted that source apportionment techniques such as PMF and UNMIX are not able to distinguish




Figure 3-24. Time series of 24-h average PM₁₀ concentrations observed in the Rio Grande Valley during May 1998.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

between bioaerosol and organic PM from whatever source, mainly because of analytical limitations. In addition, background PM source contributions are contaminated by contributions from pollution sources during transport from source to receptor monitoring site. Thus, background source contributions derived by these methods should be regarded as upper limits on the true values.

Eastern United States

The use of data for PM_{2.5}, PM_{10-2.5}, and PM₁₀ from IMPROVE sites in the East will generally result in gross overestimates of PRB concentrations because of extensive contamination by anthropogenic sources. Song et al. (2001) derived source contributions to PM_{2.5} concentrations measured at Washington, DC, Brigantine, NJ, and Underhill, VT using PMF. They found that woodsmoke (consisting of contributions from wildfires and residential wood burning) could contribute $0.93 \pm 0.97 \mu\text{g}/\text{m}^3$; soil $0.11 \pm 0.22 \mu\text{g}/\text{m}^3$; and sea spray $0.90 \pm 1.0 \mu\text{g}/\text{m}^3$ on an annual basis at Brigantine, NJ throughout the 1989 to 1999 sampling period. They also derived contributions of $1.2 \pm 0.9 \mu\text{g}/\text{m}^3$ from woodsmoke; $0.32 \pm 0.61 \mu\text{g}/\text{m}^3$




Figure 3-25. PM₁₀ concentrations observed in St. Louis, MO, during May 1998.

Source: Aerometric Information Retrieval System (AIRS; U.S. Environmental Protection Agency, 2002b).

from soil; and $0.05 \pm 0.05 \mu\text{g}/\text{m}^3$ from sea spray at Underhill, VT from 1989 through 1999. The “background” sources contribute about 7% to annual average PM_{2.5} concentrations at Brigantine and about 12% at Underhill. The daily time series at the NJ and VT sites show striking variability in background components, characterized by spikes. Maximum daily values during these spikes are in the range of several $\mu\text{g}/\text{m}^3$. Song et al. concluded that such spikes in concentrations are likely caused by transient meteorological events, e.g., storms or transport of dust from northern Africa or other distant regions or by events such as wild fires. Contributions from all of these sources should be regarded as upper limits because of entrainment of pollutant emissions during transport from source to receptor of the background source emissions. Care should also be taken to ensure that distinctions are made between contributions from wildfires

versus those from residential wood burning on the basis of seasonality or observations of smoke plumes when using these results.

Pun et al. (2002) used a regional scale chemistry-transport model (CMAQ) to simulate O_3 and $PM_{2.5}$ concentrations arising from natural sources alone in model domains centered over Tennessee and over the mid-Atlantic region for several days in July 1995. These calculations were performed for meteorological conditions that resulted in high ambient ozone concentrations in the eastern United States. They found that natural sources contributed about $1.7 \mu\text{g}/\text{m}^3$ to Washington, DC, (ranging from about 0.6 to $3.1 \mu\text{g}/\text{m}^3$ in the mid-Atlantic domain) and about $1.2 \mu\text{g}/\text{m}^3$ to Nashville, TN, on a 24 h average basis. The formation of secondary organic PM from biogenic precursors may be expected to be maximized for these conditions; however, their contribution was estimated to be at most 15% of natural $PM_{2.5}$ or less than $1 \mu\text{g}/\text{m}^3$. The largest contributions in both cases came from natural $PM_{2.5}$ that was advected in from other regions of the United States. In addition to the sources considered above, contributions to both primary and secondary PM from events such as volcanic eruptions or geothermal activity are highly sporadic. The spatial and temporal distributions of secondary PM produced by background sources shown in Table 3-8 still remain to be investigated.

Western United States

As mentioned earlier, it is impossible to obtain reliable estimates of PRB concentrations solely on the basis of measurements of $PM_{2.5}$, $PM_{10-2.5}$, or PM_{10} . It is preferable to quantify contributions from both background and non-background sources by using compositional data in techniques such as source apportionment modeling. However, of those measured throughout the United States, the concentrations observed at several RRMS in the western United States probably come closest to what PRB concentrations might be in the West. In addition to looking at $PM_{2.5}$, PM_{10} , and $PM_{10-2.5}$, the $PM_{2.5}$ and PM_{10} mass with the sulfate component removed was examined to obtain better insight into non-anthropogenic background. However, it should be noted at the outset that an undetermined fraction of the sulfate will be PRB in nature as the result of long range transport from Asia, more local crustal sources or volcanic emissions, or from sea-spray (especially at coastal sites). Data from 19 IMPROVE sites are examined in Appendix 3E (including two eastern sites). The ranges of annual average non-sulfate $PM_{2.5}$ and non-sulfate PM_{10} as well as $PM_{10-2.5}$ (calculated as the difference between PM_{10} and $PM_{2.5}$) concentrations are

shown in Table 3-E1 and the corresponding ranges for the 90th percentile concentrations are summarized in Table 3-E2. The IMPROVE monitors did not measure coarse fraction sulfate. It must be recognized at the outset that these concentrations will only provide upper limits, and it is not clear over what spatial scale these concentrations can be extrapolated. However, at nine of the 16 sites in the western United States summarized in Appendix 3E, annual mean concentrations are within the range of PRB values estimated in the 1996 PM AQCD and given earlier in this section. At the other seven sites in the western United States, they are consistent with, although slightly above, the annual average values defined earlier in the 1996 PM AQCD. The annual mean sulfate concentrations at these western IMPROVE sites range significantly above the PRB values estimated in the 1996 PM AQCD.

Some screening should be performed to rule out transport from urban areas or other sources of anthropogenic PM on a day-by-day basis. It is important to note that there is much smaller relative variability in PM_{2.5} concentrations (from the minimum to the P₉₀ level) on a year-to-year basis at the western IMPROVE sites than at the eastern IMPROVE sites. This may be related to a lack of discrete transport events affecting most samples at the western sites compared to the eastern sites; but does not rule out the possibility that the western sites are receiving highly diluted contributions from many distant urban sources. Further inquiry is needed to address these issues. Likewise, data for PM_{10-2.5} concentrations at RRMS can be used for similar purposes.

Summary

It can be concluded from the above discussion that 24-h policy relevant background (PRB) concentrations are highly variable both spatially and temporally. Contributions to PRB concentrations from external sources (e.g., Asian and North African dust storms and Central American wildfires) can be significant on an episodic, but probably not on an annual basis. More local sources of primary PRB PM are also likely to be episodic, reflecting the occurrence of volcanic eruptions, wildfires, and storms that raise dust and sea spray. The influence from events such as these can be felt over thousands of square kilometers. Very little work has been done to quantify the magnitude and variability of contributions from the production of secondary PM. However, the one modeling study cited above found values < 1 µg/m³ as secondary PM. Perhaps the greatest possibility for estimating these concentrations comes from the application of

source apportionment techniques such as PMF (positive matrix factorization) to time series of species compositional data obtained at relatively remote monitoring sites to minimize interference from anthropogenic sources. In the absence of such results, some useful estimates may be obtained by examining the time series of PM concentrations at such sites with screening to eliminate days when concentrations are influenced by anthropogenic sources.

The above discussions do not explicitly consider the contributions from PBP. However, it should be noted there is very little data available for concentrations of PBP, and their contributions are likely to be highly spatially and temporally variable, especially on a seasonal basis. For example, one study conducted in Mainz, Germany (Mathias-Maser, 1998) found that PBP could contribute from about 17 to 20% of total aerosol volume and from 9 to 30% of total particle number in the size range from 0.35 μm to 50 μm , depending on season. Also, pollen can, at times, be a dominant contributor to ambient PM particles larger than 10 μm . However, rupture of pollen grains (especially during warm temperature and high humidity or rainfall conditions) can result in release of many smaller-size airborne cytoplasmic fragments ranging down to around 0.3 to 0.4 μm (see Appendix 7B of Chapter 7 for more information on pollen and other bioaerosols).

3.3.4 Emissions Estimates for Primary PM, and Precursors to Secondary PM (SO_2 , NO_x , VOCs, and NH_3) in the United States

In principle, source contributions to ambient PM also could be estimated on the basis of predictions made by chemistry-transport models (CTMs) or even on the basis of emissions inventories alone. Uncertainties in emissions inventories have arguably been regarded as representing the largest source of uncertainty in CTMs (Calvert et al., 1993). Apart from uncertainties in emission inventories, a number of other factors limit the ability of an emissions inventory-driven CTM to determine the effects of various sources on particle samples obtained at a particular location. CTM predictions represent averages over the area of a grid cell, which in the case of CMAQ (Community Model for Air Quality) and MAQSIP (Multiscale Air Quality Simulation Platform), ranges from 16 km^2 (4 $\text{km} \times 4 \text{ km}$) to 1296 km^2 (36 $\text{km} \times 36 \text{ km}$). CMAQ and MAQSIP constitute the CTMs within the overall Models3 framework, which also includes emissions processors, the meteorological model, and decision-support modules. The contributions of sources to pollutant concentrations at a monitoring site are controlled strongly

by local conditions that cannot be resolved by an Eulerian grid-cell model. Examples would be the downward mixing of tall-stack emissions and deviations from the mean flow caused by buildings. The effect of local sources at a particular point in the model domain may not be predicted accurately because their emissions would be smeared over the area of a grid cell or if the local wind fields at the sampling point deviated significantly from the mean wind fields calculated by the model. CTMs also have problems in predicting pollutant concentrations because of uncertainties in vertical mixing and in predicting concentrations of pollutants from stationary combustion sources resulting from uncertainties in estimates of plume rise. CTMs are an integral part of air quality management programs and are reviewed in the NARSTO *Fine Particle Assessment* (NARSTO, 2002).

Estimated emissions of primary PM_{2.5} from different sources in the United States are summarized in Table 3-11, and estimated emissions of precursors to secondary PM_{2.5} (SO₂, NO_x, VOCs, and NH₃) are summarized in Table 3-12. These estimates provide a rough overview of the relative importance of major PM sources in the United States. The emissions estimates are based on information presented in the EPA *National Air Pollutant Emission Trends Report, 1990-1999* (U.S. Environmental Protection Agency, 2001), to which the reader is referred for detailed tables showing trends in PM_{2.5} emissions from a number of source categories from 1990 to 1999. Detailed descriptions of the methodology for constructing emissions inventories for criteria pollutants, quality assurance procedures, and examples of calculations of emissions can be found in U.S. Environmental Protection Agency (1999). Although uncertainties associated with the estimates in the National Air Pollutant Emission Trends Report are not given therein, a discussion of uncertainties in emissions estimates is given in Section 3.3.5.

For the sake of completeness, an attempt was made to supplement the information given in the emissions tables in the Trends Report (which concentrates mainly on anthropogenic emissions) with information about emissions from natural sources. Details regarding the composition of the emissions of primary PM_{2.5} from the source categories shown in Table 3-11 are summarized in Appendix 3D, where available. Fugitive dust emissions are estimated to constitute over 50% of nationwide primary PM_{2.5} emissions according to Table 3-11. However, there are a number of issues concerning the methods for obtaining relevant emissions factor data for fugitive dust in field studies, as discussed in Section 3.3.5. An estimate of the production of PM_{2.5} from wind erosion on natural surfaces was not included in Table 3-11, because this

TABLE 3-11. EMISSIONS OF PRIMARY PM_{2.5} BY VARIOUS SOURCES IN 1999

Source	Emissions (10 ⁹ kg/y)	Major PM Components	Notes
On-road vehicle exhaust	0.21	Organic compounds, elemental carbon	Exhaust emissions from diesel (72%) and gasoline vehicles (28%).
Non-road vehicle exhaust	0.37	Organic compounds, elemental carbon	Exhaust emissions from off-road diesel (57%) and gasoline vehicles (20%); ships and boats (10%); aircraft (7%); railroads (6%).
Fossil fuel combustion	0.36	Crustal elements, trace metals	Fuel burning in stationary sources such as power plants (33%); industries (39%); businesses and institutions (25%); residences (3%).
Industrial processes	0.35	Metals, crustal material, organic compounds	Metals processing (29%); mineral products (27%); chemical mfg. (11%); other industries (33%).
Biomass burning	1.2	Organic compounds, elemental carbon	Managed burning (47%); residential wood burning (28%); agricultural burning (7%); wildfires (18%).
Waste disposal	0.48	Organic compounds, trace metals	Open burning (91%); incineration (9%).
Fugitive dust	3.3	Crustal elements	Dust raised by vehicles on paved (19%) and unpaved roads (40%); construction (15%), dust from raising crops (24%) and livestock (2%).
Windblown dust	NA ¹	Crustal elements	Dust raised by wind on bare land.
Other	0.02	Organic compounds, elemental carbon	Structural fires.
Total	6.2		

¹NA = not available.

Source: Adapted from U.S. Environmental Protection Agency (2001).

source is highly sporadic, occurs during periods of high winds, and the resulting emissions are too highly uncertain to be included here. As can be seen from a comparison of entries in Tables 3-11 and 3-12, estimates of emissions of potential precursors to secondary PM formation are considerably larger than those for estimates of primary PM_{2.5} emissions in the United States. The emissions of SO₂, NO_x, and NH₃ that are converted to PM should be multiplied by factors of 1.5, 1.35, and 1.07, respectively, to account for their chemical form in the aerosol phase. Estimating a factor for VOCs is somewhat less straightforward. Turpin and Lim (2001) recommend factors ranging from 1.4 to 2 to account for the conversion of OC to oxygen- and

TABLE 3-12. EMISSIONS OF PRECURSORS TO SECONDARY PM_{2.5} FORMATION BY VARIOUS SOURCES IN 1999

Precursor	Emissions (10 ⁹ kg/y)	Secondary PM Component	Notes
SO ₂	17	Sulfate	Exhaust from on-road (2%) and non-road (5%) engines and vehicles; fossil fuel combustion by electrical utilities, industries, other sources (85%); various industrial processes (7%); and other minor sources (1%).
NO _x ^{1,2}	26	Nitrate	Exhaust from on-road (34%) and non-road (22%) engines and vehicles; fossil fuel combustion by electrical utilities, industries, other sources (39%); lightning (4%); soils (4%); and other minor sources (5%).
Anthropogenic VOCs	16	Various mainly unidentified compounds of 'OC'	Evaporative and exhaust emissions from on-road (29%) and non-road (18%) vehicles; evaporation of solvents and surface coatings (27%); biomass burning (9%); storage and transport of petroleum and volatile compounds (7%); chemical and petroleum industrial processes (5%); other sources (5%).
Biogenic VOCs ¹	44	Various mainly unidentified compounds of 'OC'	Approximately 98% emitted by vegetation. Isoprene (35%); monoterpenes (25%); all other reactive and non-reactive compounds (40%).
NH ₃	45	Ammonium	Exhaust from on-road and non-road engines and vehicles (4%); chemical manufacturing (3%); waste disposal, recycling, and other minor sources (4%); livestock (73%); and fertilizer application (16%).

¹Includes estimates of natural sources from Guenther et al. (2000).

²Emissions expressed in terms of NO₂.

Source: Adapted from U.S. Environmental Protection Agency (2001).

nitrogen-containing compounds in the aerosol phase. There is some additional uncertainty in assigning a factor for the fraction of VOCs that are converted into OC in the aerosol phase. These factors are all greater than 1 and further underscore the potential importance of secondary PM precursor emissions relative to primary PM emissions. However, the emissions of precursors cannot be translated directly into rates of PM formation. Dry deposition and precipitation scavenging of some of these gaseous precursors and their intermediate oxidation products occur before they are converted to PM in the atmosphere, and most of the VOCs are oxidized to CO₂ rather than PM. In addition, some fraction of these gases are transported outside of the domain of the continental United States before being oxidized. Likewise, emissions of

these gases from areas outside the United States can result in the transport of their oxidation products into the United States.

As discussed in Section 3.3.1, the photochemical oxidation of SO₂ leads to the production of sulfate, whereas that of NO_x leads ultimately to particulate-phase nitrite and nitrate. Due to uncertainties it is difficult to calculate the rates of formation of SOPM from the emissions of VOC precursors. Smog chamber and laboratory studies discussed in Section 3.3.1 indicate that anthropogenic aromatic compounds and biogenic terpenoid compounds have the highest potential for forming SOPM; and, as can be seen from Table 3C-1, the dominant compounds tend to be those derived from these categories. Each of the source categories capable of emitting VOCs shown in Table 3-12 has components capable of forming SOPM, although in small yields (ranging typically up to several percent, cf., Section 3.3.1). The oxidation of lighter organic compounds leads ultimately to the formation of CO and CO₂. As discussed by Pandis et al. (1991) and in Section 3.3.1, soluble gas phase compounds, such as formaldehyde (CH₂O), other aldehydes, organic acids, etc., formed during the oxidation of a wide variety of hydrocarbons, can be incorporated into suspended particles. Although isoprene is a major component of biogenic emissions, its oxidation has not been found to result in the formation of new particles; whereas the oxidation of monoterpenes has. However, it should be remembered that soluble gas phase species such as CH₂O are formed during the oxidation of isoprene.

The emissions estimates shown in this section are based on averaged annual totals. However, annual averages do not reflect the variability of a number of emissions categories on shorter time scales. Residential wood burning in fireplaces and stoves, for example, is a seasonal practice that reaches its peak during cold weather. Cold weather also affects motor vehicle exhaust PM emissions, both in terms of chemical composition and emission rates (e.g., Watson et al., 1990b; Huang et al., 1994). Agricultural activities such as planting, fertilizing, and harvesting are also seasonal. Forest fires occur mainly during the local dry season and during periods of drought. Maximum dust production by wind erosion in the United States occurs during the spring, whereas the minimum occurs during the summer (Gillette and Hanson, 1989). Efforts are being made to account for the seasonal variations of emissions in the nationwide emissions inventories. Techniques for calculating emissions of criteria pollutants on a seasonal basis are given in U.S. Environmental Protection Agency (1999).

Trends in nationwide, annual average concentrations of PM₁₀ and precursor gases (SO₂, NO₂, and VOC) during the 10 years from 1992 to 2001 are shown in Table 3-13. As can be seen from Table 3-13, there have been substantial decreases in the ambient concentrations of PM₁₀, SO₂, and NO₂. Not enough data are available to define trends in VOC concentrations. There also have been substantial decreases in the emissions of all the species shown in Table 3-13, except for NO₂, although its average ambient concentration has decreased by 11%. These entries suggest that decreases in the average ambient concentration of PM₁₀ could have been produced by both decreases in emissions of primary PM₁₀ and in the formation of secondary PM₁₀. The large reductions in ambient SO₂ concentrations have resulted in reductions in sulfate formation that would have been manifest in PM_{2.5} concentrations on the regional scale in the East and Midwest, where sulfate has constituted a larger fraction of PM_{2.5} than in the West. Likewise, reductions in NO₂ concentrations would have had a more noticeable effect on PM_{2.5} concentrations in the West than in the East, because nitrate is a larger component of the aerosol in the West.

TABLE 3-13. NATIONWIDE CHANGES IN AMBIENT CONCENTRATIONS AND EMISSIONS OF PM₁₀ AND GASEOUS PRECURSORS TO SECONDARY PARTICULATE MATTER FROM 1992 TO 2001

	% Change 1992-2001	
	Ambient Concentration	Emissions
PM ₁₀	-14%	-13%*
PM _{2.5}	—	-10%*
SO ₂	-35% (SO ₂)	-24% (SO ₂)
NO _x	-11% (NO ₂)	-3% (NO _x)
VOC	—	-8%

*Includes only primary PM.

Source: U.S. Environmental Protection Agency (2002a).

Trends in aerosol components (i.e., nitrate, sulfate, carbon, etc.) are needed for a more quantitative assessment of the effects of changes in emissions of precursors. Aerosol nitrate and sulfate concentrations obtained at North Long Beach and Riverside, CA tracked downward trends in NO_x concentrations. SO_2 and sulfate concentrations have both decreased; however, the rate of decline of sulfate has been smaller than that of SO_2 , indicating the long range transport of sulfate from outside the airshed may be an important source in addition to the oxidation of locally generated SO_2 . There are a number of reasons why pollutant concentrations do not track well estimated reductions in emissions. Some of these reasons are related to atmospheric effects such as meteorological variability and secular changes in the rates of photochemical transformations and deposition (U.S. Environmental Protection Agency, 2000c). Other reasons are related to uncertainties in ambient measurements and in emissions inventories.

3.3.5 Uncertainties of Emissions Inventories

As described in the 1996 PM AQCD, it is difficult to quantitatively assign uncertainties to entries in emissions inventories. Methods that can be used to verify or place constraints on emissions inventories are sparse. In general, the overall uncertainty in the emissions of a given pollutant includes contributions from all of the terms used to calculate emissions (i.e., activity rates, emissions factors, and control device efficiencies). Additional uncertainties may arise during the compilation of an emissions inventory due to missing sources and computational errors. The variability of emissions can cause errors when annual average emissions are applied to applications involving shorter time scales.

Activity rates for well-defined point sources (e.g., power plants) should have the smallest uncertainty associated with their use, because emissions are monitored continuously in many cases and accurate production records need to be kept. On the other hand, activity rates for a number of very disperse fugitive sources are difficult to quantify. Emissions factors for easily measured fuel components that are released quantitatively during combustion (e.g., CO_2 , SO_2) should be the most reliable. Emissions of components formed during combustion are more difficult to characterize, as the emissions rates are dependent on factors specific to individual combustion units and on combustion stage (i.e., smoldering or active). Although the AP-42 emissions factors (U.S. Environmental Protection Agency, 1995) contain extensive information for a large number of source types, these data are very limited in the number of sources sampled.

The efficiency of control devices is determined by their design, age, maintenance history, and operating conditions. It is virtually impossible to assign uncertainties in control device performance because of these factors. It should be noted that the largest uncertainties occur for those devices that have the highest efficiencies (> 90%). This occurs because the efficiencies are subtracted from one, and small errors in assigning efficiencies can lead to large errors in emissions.

Ideally, an emissions inventory should include all major sources of a given pollutant. This may be an easy task for major point sources. However, area sources of both primary PM and precursors to secondary PM formation are more difficult to characterize than point sources; and, thus, they require special emphasis when preparing emission inventories. Further research is needed to better characterize the sources of pollutants to reduce this source of uncertainty. Errors also can arise from the misreporting of data, and arithmetic errors can occur in the course of compiling entries from thousands of individual sources. A quality assurance program is required to check for outliers and arithmetic errors. Because of the variability in emissions rates, there can be errors in the application of inventories developed on an annually averaged basis (as are the inventories shown in Tables 3-11 and 3-12) to episodes occurring on much shorter time scales. As an example, most modeling studies of air pollution episodes are carried out for periods of a few days.

Uncertainties in annual emissions were estimated to range from 4 to 9% for SO₂ and from 6 to 11% for NO_x in the 1985 NAPAP inventories for the United States (Placet et al., 1991). Uncertainties in these estimates increase as the emissions are disaggregated both spatially and temporally. The uncertainties quoted above are minimum estimates and refer only to random variability about the mean, assuming that the variability in emissions factors was adequately characterized and that extrapolation of emissions factors to sources other than those for which they were measured is valid. The estimates do not consider the effects of weather or variations in operating and maintenance procedures.

Fugitive dust sources, as mentioned above, are extremely difficult to quantify; and stated emission rates may represent only order-of-magnitude estimates. Although crustal dust emissions constitute about 50% of the total primary PM_{2.5} inventory, they constitute less than about 15% of the source strengths inferred from the receptor modeling studies shown in

Table 3-9. However, it should be remembered that secondary components (sulfate, nitrate, and some fractions of OC) often account for most of the mass of ambient PM_{2.5} samples.

Although mineral dust sources represent the major category in Table 3-11, their contributions are distributed much more widely than are those from combustion sources. Watson and Chow (2000) reexamined the methodology used to determine emissions of fugitive dust. The standard methods use data obtained by particle monitors stacked at several elevations from 1 to 2 m up to 7 to 10 m above the surface. However, small-scale turbulent motions and variable winds characterize atmospheric flow patterns immediately adjacent to the surface in this height range (Garratt, 1994). The depth of this turbulent layer is determined by surface roughness elements, and there is a high probability of particles being entrained in turbulent eddies and redepositing on the ground within a very short distance. In addition to the source-sampling problem referred to above, it should be remembered that dust often is raised in remote areas far removed from population centers. Precipitation or scavenging by cloud droplets and dry deposition removes particles during transport from the source area. In addition, gravitational settling can be an important loss mechanism for particles larger than a few micrometers in aerodynamic diameter.

As rough estimates, uncertainties in emissions estimates could be as low as 10% for the best characterized source categories; whereas emissions figures for windblown dust should be regarded as order-of-magnitude estimates. The application of emissions inventories to the estimation of source contributions at monitoring sites is also limited by the effects of local topography and meteorology. For example, Pinto et al. (1998) found that the contribution of power plants and residential space heating to PM_{2.5} concentrations in northwestern Bohemia are comparable on the basis of CMB receptor modeling. However, according to the emissions inventories, the contribution from power plants should have been roughly an order of magnitude larger than that from residential space heating. The difference between the two methods can be explained by noting that mixing of the emissions from the power plants downward to the surface is inhibited by strong surface inversions that develop during the winter season in this area.

There have been few field studies designed to test emissions inventories observationally. The most direct approach would be to use aircraft to obtain cross sections of pollutants upwind and downwind of major urban areas. The computed mass flux through a cross section of the urban plume can then be equated to emissions from the chosen city. This approach has been

attempted on a few occasions, but results have been ambiguous because of contributions from fugitive sources, variable wind flows, and logistic difficulties.

3.4 SUMMARY AND KEY CONCLUSIONS

The recently deployed PM_{2.5} FRM network has returned data for a large number of sites across the United States. Annual mean PM_{2.5} concentrations range from ~5 µg/m³ to ~30 µg/m³. In the eastern United States, the data from 1999 to 2001 indicate that highest quarterly mean concentrations and maximum concentrations most often occur during the summer. In the western United States, highest quarterly mean values and maximum values occur mainly during the winter at a number of sites, although there were exceptions to these general patterns. Sites affected strongly by sources of primary PM are expected to show winter maxima. Sources emitting precursors of secondary ammonium nitrate are also contributors to winter PM maxima. These findings are generally consistent with those based on longer term data sets such as these from the Metropolitan Acid Aerosol Characterization Study (MAAQS) in the eastern United States and the California Air Resources Board (CARB) network of dichotomous samplers in California. PM_{2.5} and PM₁₀ concentrations in a number of urban areas have generally declined over the past few decades, and they appear to have leveled off in the past few years.

Differences in annual mean PM_{2.5} concentrations between monitoring sites in urban areas examined are typically less than 6 or 7 µg/m³. However, on individual days, differences in 24-h average PM_{2.5} concentrations can be much larger. Some sites in metropolitan areas are highly correlated with each other but not with others, due to the presence of local sources, topographic barriers, etc. Although PM_{2.5} concentrations at sites within a MSA can be highly correlated, significant differences in their concentrations can occur on any given day. Consequently, additional measures should be used to characterize the spatial variability of PM_{2.5} concentrations. The degree of spatial uniformity in PM_{2.5} concentrations in urban areas varies across the country. These factors should be considered in using data obtained by the PM_{2.5} FRM network to estimate community-scale human exposure, and caution should be exercised in extrapolating conclusions obtained in one urban area to another. PM_{2.5} to PM₁₀ ratios were generally higher in the East than in the West, and values for this ratio are consistent with those found in numerous earlier studies presented in the 1996 PM AQCD.

Data for $PM_{10-2.5}$ are not as abundant as they are for $PM_{2.5}$, and their interpretation is complicated by the difference method used to determine their concentrations. The more sporadic nature of sources of $PM_{10-2.5}$ and its shorter atmospheric lifetime tend to result in lower spatial correlations for $PM_{10-2.5}$ than for $PM_{2.5}$ concentrations. Errors in the measurement of $PM_{2.5}$ and PM_{10} also result in calculations of lower spatial correlations of $PM_{10-2.5}$. Calculated concentrations of $PM_{10-2.5}$ are occasionally negative as reflected by $PM_{2.5}$ to PM_{10} ratios greater than one. Because analytical errors are generally larger for individual species than for total mass, similar problems arise in their determination in $PM_{10-2.5}$ samples by the difference approach. Some, but not all of these problems could be resolved by the use of dichotomous samplers that also provide a direct sample of $PM_{10-2.5}$ for compositional analyses.

Estimates of concentrations of individual species in $PM_{10-2.5}$ samples used in this chapter were limited to those obtained by dichotomous samplers. Generally, concentrations of most elements differ for $PM_{2.5}$ and $PM_{10-2.5}$. However, the available data suggest that concentrations of many metals are of the same order of magnitude in both size fractions. This is in marked contrast to the situation 20 years ago, when uncontrolled combustion sources were prevalent. At that time, concentrations of many metals, especially Pb, were much higher than today in fine-mode particles, and their concentrations were much higher in the fine mode than in the coarse mode. No substantive conclusions about contemporary concentrations and composition of ultrafine particles ($< 0.1 \mu m D_a$) can be drawn for the nation as a whole because of a lack of data.

Ambient PM contains both primary and secondary components. The results of ambient monitoring studies and receptor modeling studies indicate that $PM_{2.5}$ is dominated by secondary components in the eastern United States. General statements about the origin of organic carbon (OC) in ambient $PM_{2.5}$ samples cannot yet be made; and, so, the contribution of secondary components throughout the rest of the United States is still highly uncertain. Primary constituents represent smaller but still important components of $PM_{2.5}$. Crustal materials, which are primary constituents, constitute the largest measured fraction of $PM_{10-2.5}$ throughout the United States. Data for the concentration of bioaerosols in both the $PM_{2.5}$ and $PM_{10-2.5}$ size ranges are sparse. Data collected in several airsheds, including the Los Angeles Basin, Bakersfield and Fresno, CA and Philadelphia, PA airsheds, suggest that secondary PM

components are more uniformly distributed than are primary PM components. Compositional data obtained at multiple sites in other urban areas are sparse.

Because of the complexity of the composition of ambient $PM_{2.5}$ and $PM_{10-2.5}$, sources are best discussed in terms of individual constituents of both primary and secondary $PM_{2.5}$ and $PM_{10-2.5}$. Each of these constituents can have anthropogenic and natural sources as shown in Table 3-8. The distinction between natural and anthropogenic sources is not always obvious. Although windblown dust might seem to be the result of natural processes, highest emission rates are associated with agricultural activities in areas that are susceptible to periodic drought. Examples include the dust bowl region of the midwestern United States and the Sahel of Africa. There is also ongoing debate about characterizing wildfires as either natural or anthropogenic. Land management practices and other human actions affect the occurrence and scope of wildfires. Similarly, prescribed burning can be viewed as anthropogenic or as a substitute for wildfires that would otherwise eventually occur on the same land.

During the past decade, a significant amount of research has been carried out to improve our understanding of the atmospheric chemistry of secondary organic PM (SOPM) formation. Although additional sources of SOPM might still be identified, there appears to be a general consensus that biogenic compounds (monoterpenes, sesquiterpenes) and aromatic compounds (toluene, ethylbenzene) are the most significant SOPM precursors. A large number of compounds have been detected in biogenic and aromatic SOPM, although the chemical composition of these two categories has not been fully established, especially for aromatic SOPM. Transformations that occur during the aging of particles are still inadequately understood. There still exist large gaps in our current understanding of a number of key processes related to the partitioning of semivolatile compounds between the gas phase and ambient particles containing organic compounds, liquid water, and inorganic salts and acids. In addition, there is a general lack of reliable analytical methods for measuring multifunctional oxygenated compounds in the gas and aerosol phases.

The results of receptor modeling studies throughout the United States indicate that the combustion of fossil and biomass fuels is the major source of measured ambient $PM_{2.5}$. Fugitive dust, found mainly in the $PM_{10-2.5}$ range size, represents the largest source of measured ambient PM_{10} in many locations in the western United States. Quoted uncertainties in the source apportionment of constituents in ambient aerosol samples typically range from 10 to 50%. It is

apparent that a relatively small number of source categories, compared to the total number of chemical species that typically are measured in ambient monitoring-source receptor model studies, are needed to account for the majority of the observed mass of PM in these studies.

The application of any of the source apportionment techniques is still limited by the availability of source profile data as well as by the level of detail and the quality of ambient measurement data. Whereas the chemical mass balance (CMB) approach relies directly on source profile data, solutions from the positive matrix factorization (PMF) technique yield profiles for the factors that contribute to PM. However, there is some rotational ambiguity present in the solutions. Source profile data obtained by PMF must still be verified by comparison with observational data and these data can be used in techniques such as PMF to improve the solutions. Serious limitations still exist with regard to source profiles for organic compounds. The complexity of reactions involving organic compounds in particles adds to the difficulties of finding stable species that could be used as tracers.

As seen in Table 3-8, emissions of mineral dust, organic debris, and sea spray are concentrated mainly in the coarse fraction of PM₁₀ (> 2.5 μm D_a). A small fraction of this material is in the PM_{2.5} size range (< 2.5 μm D_a). Still, PM_{2.5} concentrations of crustal material can be appreciable, especially during dust events. It also should be remembered that from one-third to one-half of the Saharan dust reaching the United States is in the PM_{2.5} size range. Emissions from combustion sources (mobile and stationary sources and biomass burning) are also predominantly in the PM_{2.5} size range.

A number of sources contribute to policy relevant background (PRB) concentrations. Data obtained at relatively remote monitoring sites (RRMS) in the western United States could be used to place reasonable upper limits on PRB concentrations. More definitive results for both annual average and daily average concentrations could potentially be obtained from the application of source-receptor models and/or the application of large-scale chemistry transport models. Many areas in the East are affected by dust transported from northern Africa, and it has recently become apparent that many areas, especially, but not limited to the Northwest, are affected by dust transported from the deserts of Asia. In addition to crustal material, pollutants and primary biological aerosol particles (PBPs) are also transported during intercontinental transport events. Many areas are also affected by smoke from wildfires occurring within the United States or in Canada, Mexico, or Central America. Storms, in which the winds can

suspend material from the surface of the land or seas, also contribute soil, sea spray and PBP. Contributions of primary PM from natural sources and sources outside Northern America as given above are all episodic. Because concentrations of PBP are so poorly quantified, even though they can constitute significant portions of the organic fraction of the atmospheric aerosol, estimates of PRB concentrations will remain highly uncertain. Estimates of annually averaged PRB concentrations or their range have not changed from the 1996 PM AQCD.

REFERENCES

- Altshuller, A. P. (1983) Review: natural volatile organic substances and their effect on air quality in the United States. *Atmos. Environ.* 17: 2131-2165.
- Ansari, A. S.; Pandis, S. N. (1998) Response of inorganic PM to precursor concentrations. *Environ. Sci. Technol.* 32: 2706-2714.
- Atkinson, R. (1994) Gas-phase tropospheric chemistry of organic compounds. Washington, DC: American Chemical Society. (J. Phys. Chem. Ref. Data Monograph no. 2).
- Aumont, B.; Madronich, S.; Bey, I.; Tyndall, G. S. (2000) Contribution of secondary VOC to the composition of aqueous atmospheric particles: a modeling approach. *J. Atmos. Chem.* 35: 59-75.
- Bahadori, T.; Suh, H.; Koutrakis, P. (2000a) A cross-study analysis of ambient concentrations of fine mass. *J. Air Waste Manage. Assoc.*: in press.
- Bahadori, T.; Suh, H.; Koutrakis, P. (2000b) PM_{2.5} and PM₁₀ concentrations in four metropolitan cities in eastern United States. *J. Air Waste Manage. Assoc.*: in press.
- Blando, J. D.; Porcja, R. J.; Li, T.-H.; Bowman, D.; Liroy, P. J.; Turpin, B. J. (1998) Secondary formation and the Smoky Mountain organic aerosol: an examination of aerosol polarity and functional group composition during SEAVS. *Environ. Sci. Technol.* 32: 604-613.
- Cabada, J. C.; Pandis Spyros, N.; Robinson, A. L. (2002) Sources of atmospheric carbonaceous particulate matter in Pittsburgh, Pennsylvania. *J. Air Waste Manage. Assoc.* 52: 732-741.
- Cahill, T. A. (1996) [Data from IMPROVE data base.] Fort Collins, CO: Air Resource Specialists, Inc.
- California Air Resources Board. (1995) [Data base provided by John R. Holmes, Chief, Research Division.] Sacramento, CA: California Air Resources Board.
- Calvert, J. G.; Heywood, J. B.; Sawyer, R. F.; Seinfeld, J. H. (1993) Achieving acceptable air quality: some reflections on controlling vehicle emissions. *Science* (Washington, DC) 261: 37-45.
- Cass, G. R.; Hughes, L. A.; Bhave, P.; Kleeman, M. J.; Allen, J. O.; Salmon, L. G. (2000) The chemical composition of atmospheric ultrafine particles. *Philos. Trans. R. Soc. London Ser. A* 358: 2581-2592.
- Castro, L. M.; Pio, C. A.; Harrison, R. M.; Smith, D. J. T. (1999) Carbonaceous aerosol in urban and rural European atmospheres: estimation of secondary organic carbon concentrations. *Atmos. Environ.* 33: 2771-2781.
- Chadwick, O. A.; Derry, L. A.; Vitousek, P. M.; Huebert, B. J.; Hedin, L. O. (1999) Changing sources of nutrients during four million years of ecosystem development. *Nature* (London, UK) 397: 491-497.
- Chow, J. C.; Watson, J. G. (2002) Review of PM_{2.5} and PM₁₀ apportionment for fossil fuel combustion and other sources by the chemical mass balance receptor model. *Energy Fuels* 16: 222-260.
- Chow, J. C.; Watson, J. G.; Frazier, C. A.; Egami, R. T.; Goodrich, A.; Ralph, C. (1988) Spatial and temporal source contributions to PM₁₀ and PM_{2.5} in Reno, NV. In: Mathai, C. V.; Stonefield, D. H., eds. PM-10: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 438-457. (APCA transactions series TR-13).
- Chow, J. C.; Watson, J. G.; Richards, L. W.; Haase, D. L.; McDade, C.; Dietrich, D. L.; Moon, D.; Sloane, C. (1991) The 1989-90 Phoenix PM₁₀ study. Volume II: source apportionment. Final report. Phoenix, AZ: Arizona Department of Environmental Quality; Desert Research Institute document no. 8931.6F1.
- Chow, J. C.; Watson, J. G.; Lowenthal, D. H.; Frazier, C. A.; Hinsvark, B. A.; Prichett, L. C.; Neuroth, G. R. (1992a) Wintertime PM₁₀ and PM_{2.5} chemical compositions and source contributions in Tucson, Arizona. In: Chow, J. C.; Ono, D. M., eds. PM₁₀ standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 231-243. (A&WMA transactions series no. 22).
- Chow, J. C.; Watson, J. G.; Lowenthal, D. H.; Solomon, P. A.; Magliano, K. L.; Ziman, S. D.; Richards, L. W. (1992b) PM₁₀ source apportionment in California's San Joaquin Valley. *Atmos. Environ. Part A* 26: 3335-3354.
- Chow, J. C.; Liu, C. S.; Cassmassi, J.; Watson, J. G.; Lu, Z.; Pritchett, L. C. (1992c) A neighborhood-scale study of PM₁₀ source contributions in Rubidoux, California. *Atmos. Environ. Part A* 26: 693-706.
- Chow, J. C.; Watson, J. G.; Edgerton, S. A.; Vega, E.; Ortiz, E. (2002) Spatial differences in outdoor PM₁₀ mass and aerosol composition in Mexico City. *J. Air Waste Manage. Assoc.* 52: 423-434.
- Christoffersen, T. S.; Hjorth, J.; Horie, O.; Jensen, N. R.; Kotzias, D.; Molander, L. L.; Neeb, P.; Ruppert, L.; Winterhalter, R.; Virkkula, A.; Wirtz, K.; Larsen, B. R. (1998) cis-Pinic acid, a possible precursor for organic aerosol formation from ozonolysis of α -pinene. *Atmos. Environ.* 32: 1657-1661.
- Chung, A.; Herner, J. D.; Kleeman, M. J. (2001) Detection of alkaline ultrafine atmospheric particles at Bakersfield, California. *Environ. Sci. Technol.* 35: 2184-2190.

- Cocker, D. R., III; Clegg, S. L.; Flagan, R. C.; Seinfeld, J. H. (2001) The effect of water on gas-particle partitioning of secondary organic aerosol. Part I: α -pinene/ozone system. *Atmos. Environ.* 35: 6049-6072.
- Darlington, T. L.; Kahlbaum, D. F.; Heuss, J. M.; Wolff, G. T. (1997) Analysis of PM₁₀ trends in the United States from 1988 through 1995. *J. Air Waste Manage. Assoc.* 47: 1070-1078.
- Davis, A. P.; Shokouhian, M.; Ni, S. (2001) Loading estimates of lead, copper, cadmium, and zinc in urban runoff from specific sources. *Chemosphere* 44: 997-1009.
- Dolislager, L. J.; Motallebi, N. (1999) Characterization of particulate matter in California. *J. Air Waste Manage. Assoc.* 49: PM-45-56.
- Duce, R. A. (1995) Sources, distributions, and fluxes of mineral aerosols and their relationship to climate. In: Charlson, R. J.; Heintzenberg, J., eds. *Aerosol forcing of climate: report of the Dahlem workshop on aerosol forcing of climate; April 1994; Berlin, Federal Republic of Germany.* Chichester, United Kingdom: John Wiley & Sons, Ltd.; pp. 43-72.
- Dzubay, T. G.; Stevens, R. K.; Gordon, G. E.; Olmez, I.; Sheffield, A. E.; Courtney, W. J. (1988) A composite receptor method applied to Philadelphia aerosol. *Environ. Sci. Technol.* 22: 46-52.
- Edney, E. O.; Driscoll, D. J.; Weathers, W. S.; Kleindienst, T. E.; Conner, T. S.; McIver, C. D.; Li, W. (2001) Formation of polyketones in irradiated toluene/propylene/NO_x/air mixtures. *Aerosol Sci. Technol.* 35: 998-1008.
- Eldred, R. A.; Cahill, T. A. (1994) Trends in elemental concentrations of fine particles at remote sites in the United States of America. *Atmos. Environ.* 28: 1009-1019.
- Faust, B. C. (1994) Photochemistry of clouds, fogs, and aerosols. *Environ. Sci. Technol.* 28: 217A-222A.
- Fitz-Simons, T. S.; Mathias, S.; Rizzo, M. (2000) Analyses of 1999 PM data for the PM NAAQS review. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; November 17. Available: <http://www.epa.gov/oar/oaqps/pm25/analyses.html> [2 April, 2002].
- Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997a) Molecular speciation of secondary organic aerosol from photooxidation of the higher alkenes: 1-octene and 1-decene. *Atmos. Environ.* 31: 1953-1964.
- Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997b) Secondary organic aerosol from the photooxidation of aromatic hydrocarbons: molecular composition. *Environ. Sci. Technol.* 31: 1345-1358.
- Garratt, J. R. (1994) *The atmospheric boundary layer.* Cambridge, United Kingdom: Cambridge University Press. (Cambridge atmospheric and space science series).
- Gatz, D. F.; Prospero, J. M. (1996) A large silicon-aluminum aerosol plume in central Illinois: North African Desert dust? *Atmos. Environ.* 30: 3789-3799.
- Geron, C.; Rasmussen, R.; Arnts, R. R.; Guenther, A. (2000) A review and synthesis of monoterpene speciation from forests in the United States. *Atmos. Environ.* 34: 1761-1781.
- Geron, C. D.; Harley, P.; Guenther, A. (2001) Isoprene emission capacity for US tree species. *Atmos. Environ.* 35: 3341-3352.
- Gillette, D. A.; Hanson, K. J. (1989) Spatial and temporal variability of dust production caused by wind erosion in the United States. *J. Geophys. Res.* [Atmos.] 94: 2197-2206.
- Gillies, J. A.; Gertler, A. W.; Lowenthal, D.; Abu-Allaban, M.; Jennison, B.; Goodrich, A. (2000) Enhanced PM_{2.5} source apportionment using chemical mass balance receptor modeling and scanning electron microscopy. *Aerosol Sci. Technol.*: submitted.
- Glasius, M.; Lahaniati, M.; Calogirou, A.; Di Bella, D.; Jensen, N. R.; Hjorth, J.; Kotzias, D.; Larsen, B. R. (2000) Carboxylic acids in secondary aerosols from oxidation of cyclic monoterpenes by ozone. *Environ. Sci. Technol.* 34: 1001-1010.
- Glover, D. M.; Hopke, P. K.; Vermette, S. J.; Landsberger, S.; D'Auben, D. R. (1991) Source apportionment with site specific source profiles. *J. Air Waste Manage. Assoc.* 41: 294-305.
- Gone, J.-K.; Olmez, I.; Ames, M. R. (2000) Size distribution and probable sources of trace elements in submicron atmospheric particulate material. *J. Radioanal. Nucl. Chem.* 244: 133-139.
- Graedel, T. E.; Goldberg, K. I. (1983) Kinetic studies of raindrop chemistry: 1. Inorganic and organic processes. *J. Geophys. Res. C: Oceans Atmos.* 88: 10,865-10,882.
- Gray, H. A.; Landry, B.; Liu, C. S.; Henry, R. C.; Cooper, J. A.; Sherman, J. R. (1988) Receptor modeling for PM₁₀ source apportionment in the South Coast Air Basin of California. In: Mathai, C. V.; Stonefield, D. H., eds. *PM₁₀: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 399-418.* (APCA transactions series TR-13).
- Griffin, R. J.; Cocker, D. R., III; Seinfeld, J. H. (1999) Incremental aerosol reactivity: application to aromatic and biogenic hydrocarbons. *Environ. Sci. Technol.* 33: 2403-2408.

- Griffin, D. W.; Garrison, V. H.; Herman, J. R.; Shinn, E. A. (2001) African desert dust in the Caribbean atmosphere: microbiology and public health. *Aerobiologia* 17: 203-213.
- Grosjean, D. (1992) In situ organic aerosol formation during a smog episode: estimated production and chemical functionality. *Atmos. Environ. Part A* 26: 953-963.
- Grosjean, D.; Seinfeld, J. H. (1989) Parameterization of the formation potential of secondary organic aerosols. *Atmos. Environ.* 23: 1733-1747.
- Guenther, A.; Zimmerman, P.; Klinger, L.; Greenberg, J.; Ennis, C.; Davis, K.; Pollock, W.; Westberg, H.; Allwine, G.; Geron, C. (1996) Estimates of regional natural volatile organic compound fluxes from enclosure and ambient measurements. *J. Geophys. Res.* 101: 1345-1360.
- Guenther, A.; Geron, C.; Pierce, T.; Lamb, B.; Harley, P.; Fall, R. (2000) Natural emissions of non-methane volatile organic compounds, carbon monoxide, and oxides of nitrogen from North America. *Atmos. Environ.* 34: 2205-2230.
- Haagen-Smit, A. J. (1952) Chemistry and physiology of Los Angeles smog. *Ind. Eng. Chem.* 44: 1342-1346.
- Helmig, D.; Greenberg, J.; Guenther, A.; Zimmerman, P.; Geron, C. (1998) Volatile organic compounds and isoprene oxidation products at a temperate deciduous forest site. *J. Geophys. Res.* 103(D17): 22,397-22,414.
- Henry, R. C. (1997) History and fundamentals of multivariate air quality receptor models. *Chemom. Intell. Lab. Syst.* 37: 37-42.
- Hewitt, C. N. (2001) The atmospheric chemistry of sulphur and nitrogen in power station plumes. *Atmos. Environ.* 35: 1155-1170.
- Hidy, G. M. (1994) Atmospheric sulfur and nitrogen oxides: eastern North American source-receptor relationships. San Diego, CA: Academic Press, Inc.
- Hildemann, L. M.; Rogge, W. F.; Cass, G. R.; Mazurek, M. A.; Simoneit, B. R. T. (1996) Contribution of primary aerosol emissions from vegetation-derived sources to fine particle concentrations in Los Angeles. *J. Geophys. Res. [Atmos.]* 101: 19,541-19,549.
- Hoffmann, T.; Odum, J. R.; Bowman, F.; Collins, D.; Klockow, D.; Flagan, R. C.; Seinfeld, J. H. (1997) Formation of organic aerosols from the oxidation of biogenic hydrocarbons. *J. Atmos. Chem.* 26: 189-222.
- Hopke, P. K.; Wlaschin, W.; Landsberger, S.; Sweet, C.; Vermette, S. J. (1988) The source apportionment of PM₁₀ in south Chicago. In: Mathai, C. V.; Stonefield, D. H., eds. PM-10: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 484-494. (APCA transactions series TR-13).
- Houck, J. E.; Rau, J. A.; Body, S.; Chow, J. C. (1992) Source apportionment: Pocatello, Idaho PM₁₀ nonattainment area. In: Chow, J. C.; Ono, D. M., eds. PM₁₀ standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 219-230. (A&WMA transactions series no. 22).
- Huang, X.; Olmez, I.; Aras, N. K.; Gordon, G. E. (1994) Emissions of trace elements from motor vehicles: potential marker elements and source composition profile. *Atmos. Environ.* 28: 1385-1391.
- Husar, R. B.; Tratt, D. M.; Schichtel, B. A.; Falke, S. R.; Li, F.; Jaffe, D.; Gasso, S.; Gill, T.; Laulainen, N. S.; Lu, F.; Reheis, M. C.; Chun, Y.; Westphal, D.; Holben, B. N.; Gueymard, C.; McKendry, I.; Kuring, N.; Feldman, G. C.; McClain, C.; Frouin, R. J.; Merrill, J.; DuBois, D.; Vignola, F.; Murayama, T.; Nickovic, S.; Wilson, W. E.; Sassen, K.; Sugimoto, N.; Malm, W. C. (2001) Asian dust events of April 1998. *J. Geophys. Res. (Atmos.)* 106: 18,317-18,330.
- Inhalable Particulate Network. (1985) [Inhalable Particulate Network Database.] Research Triangle Park, NC: U.S. Environmental Protection Agency, National Exposure Research Laboratory; Jose M. Sune, contact.
- Izumi, K.; Fukuyama, T. (1990) Photochemical aerosol formation from aromatic hydrocarbons in the presence of NO_x. *Atmos. Environ. Part A* 24: 1433-1441.
- Jang, M.; Kamens, R. M. (1999) Newly characterized products and composition of secondary aerosols from the reaction of α -pinene with ozone. *Atmos. Environ.* 33: 459-474.
- Jang, M.; Kamens, R. M. (2001a) Atmospheric secondary aerosol formation by heterogeneous reactions of aldehydes in the presence of a sulfuric acid aerosol catalyst. *Environ. Sci. Technol.* 35: 4758-4766.
- Jang, M.; Kamens, R. M. (2001b) Characterization of secondary aerosol from the photooxidation of toluene in the presence of NO_x and 1-propene. *Environ. Sci. Technol.* 35: 3626-3639.
- Kamens, R.; Jang, M.; Chien, C.-J.; Leach, K. (1999) Aerosol formation from the reaction of α -pinene and ozone using a gas-phase kinetics-aerosol partitioning model. *Environ. Sci. Technol.* 33: 1430-1438.
- Kao, A. S.; Friedlander, S. K. (1995) Frequency distributions of PM₁₀ chemical components and their sources. *Environ. Sci. Technol.* 29: 19-28.
- Kavouras, I. G.; Mihalopoulos, N.; Stephanou, E. G. (1998) Formation of atmospheric particles from organic acids produced by forests. *Nature (London)* 395: 683-686.

- Kavouras, I. G.; Mihalopoulos, N.; Stephanou, E. G. (1999a) Formation and gas/particle partitioning of monoterpene photooxidation products over forests. *Geophys. Res. Lett.* 26: 55-58.
- Kavouras, I. G.; Mihalopoulos, N.; Stephanou, E. G. (1999b) Secondary organic aerosol formation vs primary organic aerosol emission: in situ evidence for the chemical coupling between monoterpene acidic photooxidation products and new particle formation over forests. *Environ. Sci. Technol.* 33: 1028-1037.
- Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. *Atmos. Environ.* 33: 2907-2913.
- Khalil, M. A. K.; Rasmussen, R. A. (1992) Forest hydrocarbon emissions: relationships between fluxes and ambient concentrations. *J. Air Waste Manage. Assoc.* 42: 810-813.
- Kim, B. M.; Zeldin, M. D.; Liu, C. S. (1992) Source apportionment study for state implementation plan development in the Coachella Valley. In: Chow, J. C.; Ono, D. M., eds. *PM₁₀ standards and nontraditional particulate source controls: an A&WMA/EPA international specialty conference, v. II*; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 979-991. (A&WMA transactions series no. 22).
- Kim, B. M.; Teffera, S.; Zeldin, M. D. (2000) Characterization of PM_{2.5} and PM₁₀ in the South Coast Air Basin of southern California: part 2—temporal variations. *J. Air Waste Manage. Assoc.* 50: 2045-2059.
- Kinney, P. L.; Aggarwal, M.; Northridge, M. E.; Janssen, N. A. H.; Shepard, P. (2000) Airborne concentrations of PM_{2.5} and diesel exhaust particles on Harlem sidewalks: a community-based pilot study. *Environ. Health Perspect.* 108: 213-218.
- Koch, S.; Winterhalter, R.; Uherek, E.; Kolloff, A.; Neeb, P.; Moortgat, G. K. (2000) Formation of new particles in the gas-phase ozonolysis of monoterpenes. *Atmos. Environ.* 34: 4031-4042.
- Koutrakis, P. (1995) [PM_{2.5} and PM₁₀ data for Philadelphia, 1992-1995]. Boston, MA: Harvard School of Public Health.
- Landis, M. S.; Norris, G. A.; Williams, R. W.; Weinstein, J. P. (2001) Personal exposures to PM_{2.5} mass and trace elements in Baltimore, MD, USA. *Atmos. Environ.* 35: 6511-6524.
- Larsen, B. R.; Di Bella, D.; Glasius, M.; Winterhalter, R.; Jensen, N. R.; Hjorth, J. (2001) Gas-phase OH oxidation of monoterpenes: gaseous and particulate products. *J. Atmos. Chem.* 38: 231-276.
- Leach, K. B.; Kamens, R. M.; Strommen, M. R.; Jang, M. (1999) Partitioning of semivolatile organic compounds in the presence of a secondary organic aerosol in a controlled atmosphere. *J. Atmos. Chem.* 33: 241-264.
- Leaderer, B. P.; Naeher, L.; Jankun, T.; Balenger, K.; Holford, T. R.; Toth, C.; Sullivan, J.; Wolfson, J. M.; Koutrakis, P. (1999) Indoor, outdoor, and regional summer and winter concentrations of PM₁₀, PM_{2.5}, SO₄²⁻, H⁺, NH₄⁺, NO₃⁻, NH₃, and nitrous acid in homes with and without kerosene space heaters. *Environ. Health Perspect.* 107: 223-231.
- Lin, J. J.; Noll, K. E.; Holsen, T. M. (1994) Dry deposition velocities as a function of particle size in the ambient atmosphere. *Aerosol Sci. Technol.* 20: 239-252.
- Lipfert, F. W. (1998) Trends in airborne particulate matter in the United States. *Appl. Occup. Environ. Hyg.* 13: 370-384.
- Magliano, K. L.; Ranzieri, A.; Solomon, P.; Watson, J. (1998) Chemical mass balance modeling of data from the 1995 integrated monitoring study. Sacramento, CA: California Air Resources Board; California Regional PM₁₀/PM_{2.5} Air Quality Study.
- Matthias-Maser, S. (1998) Primary biological aerosol particles: their significance, sources, sampling methods and size distribution in the atmosphere. In: Harrison, R. M.; Van Grieken, R., eds. *Atmospheric Particles*. New York, NY: John Wiley & Sons; pp. 349-368. (IUPAC series on Analytical and Physical Chemistry of Environmental Systems, v. 5).
- Mazurek, M.; Masonjones, M. C.; Masonjones, H. D.; Salmon, L. G.; Cass, G. R.; Hallock, K. A.; Leach, M. (1997) Visibility-reducing organic aerosols in the vicinity of Grand Canyon National Park: properties observed by high resolution gas chromatography. *J. Geophys. Res. [Atmos.]* 102: 3779-3793.
- McMurry, P. H.; Woo, K. S.; Weber, R.; Chen, D.-R.; Pui, D. Y. H. (2000) Size distributions of 3 to 10 nm atmospheric particles: implications for nucleation mechanisms. *Philos. Trans. R. Soc. London Ser. A* 358: 2625-2642.
- Monn, C. (2001) Exposure assessment of air pollutants: a review on spatial heterogeneity and indoor/outdoor/personal exposure to suspended particulate matter, nitrogen dioxide and ozone. *Atmos. Environ.* 35: 1-32.
- Motallebi, N. (1999) Wintertime PM_{2.5} and PM₁₀ source apportionment at Sacramento, California. *J. Air Waste Manage. Assoc.* 49: PM-25-34.
- NARSTO. (2002) Fine particle assessment (2nd review draft). Available: <http://www.cgenv.com/Narsto/> (13 February 2002).

- Neas, L. M. (1996) [Data provided for 1986 from Harvard six cities data base.] Boston, MA: Harvard School of Public Health.
- Norris, G. (1998) Air pollution and the exacerbation of asthma in an arid, western, US city [dissertation]. Seattle, WA: University of Washington. Available from: University Microfilms International, Ann Arbor, MI; publication no. 9836228.
- Norris, G.; Young-Pong, S. N.; Koenig, J. Q.; Larson, T. V.; Sheppard, L.; Stout, J. W. (1999) An association between fine particles and asthma emergency department visits for children in Seattle. *Environ. Health Perspect.* 107: 489-493.
- Odum, J. R.; Hoffmann, T.; Bowman, F.; Collins, D.; Flagan, R. C.; Seinfeld, J. H. (1996) Gas/particle partitioning and secondary organic aerosol yields. *Environ. Sci. Technol.* 30: 2580-2585.
- Odum, J. R.; Jungkamp, T. P. W.; Griffin, R. J.; Flagan, R. C.; Seinfeld, J. H. (1997a) The atmospheric aerosol-forming potential of whole gasoline vapor. *Science (Washington, DC)* 276: 96-99.
- Odum, J. R.; Jungkamp, T. P. W.; Griffin, R. J.; Forstner, H. J. L.; Flagan, R. C.; Seinfeld, J. H. (1997b) Aromatics, reformulated gasoline, and atmospheric organic aerosol formation. *Environ. Sci. Technol.* 31: 1890-1897.
- Ondov, J. M. (1996) Particulate tracers for source attribution: potential for application to California's San Joaquin Valley. *J. Aerosol Sci.* 27(suppl. 1): S687-S688.
- Ott, S.-T.; Ott, A.; Martin, D. W.; Young, J. A. (1991) Analysis of a trans-Atlantic Saharan dust outbreak based on satellite and GATE data. *Mon. Weather Rev.* 119: 1832-1850.
- Paatero, P.; Tapper, U. (1993) Analysis of different modes of factor analysis as least squares fit problems. *Chemom. Intelligent Lab. Syst.* 18: 183-194.
- Paatero, P.; Tapper, U. (1994) Positive matrix factorization: a non-negative factor model with optimal utilization of error estimates of data values. *Environmetrics* 5: 111-126.
- Pandis, S. N.; Paulson, S. E.; Seinfeld, J. H.; Flagan, R. C. (1991) Aerosol formation in the photooxidation of isoprene and β -pinene. *Atmos. Environ. Part A* 25: 997-1008.
- Pandis, S. N.; Harley, R. A.; Cass, G. R.; Seinfeld, J. H. (1992) Secondary organic aerosol formation and transport. *Atmos. Environ. Part A* 26: 2269-2282.
- Perry, K. D.; Cahill, T. A.; Eldred, R. A.; Dutcher, D. D.; Gill, T. E. (1997) Long-range transport of North African dust to the eastern United States. *J. Geophys. Res. [Atmos]* 102: 11,225-11,238.
- Pinto, J. P.; Pleasant, M.; Kellogg, R.; Torok, S.; Stiles, D.; Willis, R. D.; Koutrakis, P.; Allen, G. A. (1995) Chemical characterization of the ambient aerosol in Philadelphia. Presented at: Particulate matter: health and regulatory issues: an A&WMA international specialty conference; April; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association.
- Pinto, J. P.; Stevens, R. K.; Willis, R. D.; Kellogg, R.; Mamane, Y.; Novak, J.; Šantroch, J.; Beneš, I.; Leniček, J.; Bureš, V. (1998) Czech air quality monitoring and receptor modeling study. *Environ. Sci. Technol.* 32: 843-854.
- Pio, C.; Alves, C.; Duarte, A. (2001) Organic components of aerosols in a forested area of central Greece. *Atmos. Environ.* 35: 389-401.
- Placet, M.; Battye, R. E.; Fehsenfeld, F. C.; Bassett, G. W. (1991) Emissions involved in acidic deposition processes. In: Irving, P. M., ed. *Acidic deposition: state of science and technology, volume I: emissions, atmospheric processes, and deposition.* Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 1).
- Poore, M. W. (2000) Oxalic acid in PM_{2.5} particulate matter in California. *J. Air Waste Manage. Assoc.* 50: 1874-1875.
- Prospero, J. M. (1996) The atmospheric transport of particles to the ocean. In: Ittekkot, V.; Honjo, S.; Depetris, P. J., eds. *Particle flux in the ocean.* New York, NY: John Wiley & Sons, Ltd.; pp. 19-52. (SCOPE report no. 57).
- Prospero, J. M. (1999) Long-range transport of mineral dust in the global atmosphere: impact of African dust on the environment of the southeastern United States. *Proc. Natl. Acad. Sci. U.S.A.* 96: 3396-3403.
- Prospero, J. M.; Nees, R. T.; Uematsu, M. (1987) Deposition rate of particulate and dissolved aluminum derived from Saharan dust in precipitation at Miami, Florida. *J. Geophys. Res. [Atmos.]* 92: 14,723-14,731.
- Prospero, J. M.; Olmez, I.; Ames, M. (2001) Al and Fe in PM_{2.5} and PM₁₀ suspended particles in south-central Florida: the impact of the long range transport of African mineral dust. *J. Water Air Soil Pollut.* 125: 291-317.
- Pun, B. K.; Seigneur, C.; Grosjean, D.; Saxena, P. (2000) Gas-phase formation of water-soluble organic compounds in the atmosphere: a retrosynthetic analysis. *J. Atmos. Chem.* 35: 199-223.
- Pun, B. K.; Wu, S.-Y.; Seigneur, C. (2002) Contribution of biogenic emissions to the formation of ozone and particulate matter in the eastern United States. *Environ. Sci. Technol.* 36: 3586-3596.
- Ramadan, Z.; Song, X.-H.; Hopke, P. K. (2000) Identification of sources of Phoenix aerosol by positive matrix factorization. *J. Air Waste Manage. Assoc.* 50: 1308-1320.

- Rizzo, M.; Pinto, J. P. (2001) Initial characterization of fine particulate matter (PM_{2.5}) collected by the National Federal Reference Monitoring Network. Presented at: 94th annual conference & exhibition of the Air & Waste Management Association; June, Orlando, FL. Pittsburgh, PA: Air & Waste Management Association.
- Rodes, C. E.; Evans, E. G. (1985) Preliminary assessment of 10 µm particulate sampling at eight locations in the United States. *Atmos. Environ.* 19: 293-303.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993) Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. *Environ. Sci. Technol.* 27: 636-651.
- Ryan, W. M.; Badgett-West, C. R.; Holtz, D. R.; Peters, T. A.; Cooper, J. A.; Ono, D. (1988) Reconciliation of receptor and dispersion modeling impacts of PM₁₀ in Hayden, Arizona. In: Mathai, C. V.; Stonefield, D. H., eds. PM-10: implementation of standards, an APCA/EPA international specialty conference; February; San Francisco, CA. Pittsburgh, PA: Air Pollution Control Association; pp. 419-429. (APCA transactions series TR-13).
- Salma, I.; Maenhaut, W.; Záray, G. (2002) Comparative study of elemental mass size distributions in urban atmospheric aerosol. *J. Aerosol Sci.* 33: 339-356.
- Sarwar, G.; Corsi, R.; Kumura, Y.; Allen, D.; Weschler, C. J. (2002) Hydroxyl radicals in indoor environments. *Atmos. Environ.* 36: 3973-3988.
- Savoie, D. L.; Prospero, J. M. (1980) Water-soluble potassium, calcium, and magnesium in the aerosols over the tropical North Atlantic. *J. Geophys. Res.* 85: 385-392.
- Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. *J. Atmos. Chem.* 24: 57-109.
- Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environ. Sci. Technol.* 34: 1821-1832.
- Schauer, J. J.; Rogge, W. F.; Hildemann, L. M.; Mazurik, M. A.; Cass, G. R. (1996) Source apportionment of airborne particulate matter using organic compounds as tracers. *Atmos. Environ.* 30: 3837-3855.
- Schwartz, J.; Dockery, D. W.; Neas, L. M. (1996) Is daily mortality associated specifically with fine particles? *J. Air Waste Manage. Assoc.* 46: 927-939.
- Shinn, E. A.; Smith, G. W.; Prospero, J. M.; Betzer, P.; Hayes, M. L.; Garrison, V.; Barber, R. T. (2000) African dust and the demise of Caribbean coral reefs. *Geophys. Res. Lett.* 27: 3029-3032.
- Skidmore, L. W.; Chow, J. C.; Tucker, T. T. (1992) PM₁₀ air quality assessment for the Jefferson County, Ohio air quality control region. In: Chow, J. C.; Ono, D. M., eds. PM₁₀ standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. II; January; Scottsdale, AZ. Pittsburgh, PA: Air and Waste Management Association; pp. 1016-1031. (A&WMA transactions series no. 22).
- Smith, G. T.; Ives, L. D.; Nagelkerken, I. A.; Ritchie, K. B. (1996) Caribbean sea fan mortalities. *Nature (London)* 383: 487.
- Song, X.-H.; Polissar, A. V.; Hopke, P. K. (2001) Sources of fine particle composition in the northeastern US. *Atmos. Environ.* 35: 5277-5286.
- Spengler, J. D.; Briggs, S. L. K.; Özkaynak, H. (1986) Relationship between TSP measurements and size-fractionated particle mass measurements in six cities participating in the Harvard Air Pollution Health Study. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; EPA cooperative agreement no. CR-812667-01-0; December 5.
- Spengler, J. D.; Koutrakis, P.; Dockery, D. W.; Raizenne, M.; Speizer, F. E. (1996) Health effects of acid aerosols on North American children: air pollution exposures. *Environ. Health Perspect.* 104: 492-499.
- Strader, R.; Lurmann, F.; Pandis, S. N. (1999) Evaluation of secondary organic aerosol formation in winter. *Atmos. Environ.* 33: 4849-4863.
- Suarez, A. E.; Caffrey, P. F.; Ondov, J. M.; Thaug, K. C. (1996) Size distribution of Ir-tagged soot from diesel sanitation trucks. *J. Aerosol Sci.* 27(suppl. 1): S697-S698.
- Tanner, R. L.; Parkhurst, W. J. (2000) Chemical composition of fine particles in the Tennessee Valley region. *J. Air Waste Manage. Assoc.* 50: 1299-1307.
- Taylor, C. A., Jr.; Stover, C. A.; Westerdahl, F. D. (1998) Speciated fine particle (<2.5 µm aerodynamic diameter) and vapor-phase acid concentrations in southern California. Presented at: 91st annual meeting & exhibition of the Air & Waste Management Association; June; San Diego, CA. Pittsburgh, PA: Air & Waste Management Association; paper no. WA7401.
- Taylor, P. E.; Flagan, R. C.; Valenta, R.; Glovsky, M. M. (2002) Release of allergens as respirable aerosols: a link between grass pollen and asthma. *J. Allergy Clin. Immunol.* 109: 51-56.
- Thanukos, L. C.; Miller, T.; Mathai, C. V.; Reinhold, D.; Bennett, J. (1992) Intercomparison of PM₁₀ samplers and source apportionment of ambient PM₁₀ concentrations at Rillito, Arizona. In: Chow, J. C.; Ono, D. M., eds.

- PM₁₀ standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 244-261. (A&WMA transactions series no. 22).
- Tobias, H. J.; Ziemann, P. J. (2000) Thermal desorption mass spectrometric analysis of organic aerosol formed from reactions of 1-tetradecene and O₃ in the presence of alcohols and carboxylic acids. *Environ. Sci. Technol.* 34: 2105-2115.
- Turpin, B. J.; Huntzicker, J. J. (1991) Secondary formation of organic aerosol in the Los Angeles basin: a descriptive analysis of organic and elemental carbon concentrations. *Atmos. Environ. Part A* 25: 207-215.
- Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. *Atmos. Environ.* 29: 3527-3544.
- Turpin, B. J.; Lim, H.-J. (2001) Species contributions to PM_{2.5} mass concentrations: revisiting common assumptions for estimating organic mass. *Aerosol Sci. Technol.* 35: 602-610.
- Turpin, B. J.; Huntzicker, J. J.; Larson, S. M.; Cass, G. R. (1991) Los Angeles summer midday particulate carbon: primary and secondary aerosol. *Environ. Sci. Technol.* 25: 1788-1793.
- U.S. Environmental Protection Agency. (1995) Compilation of air pollutant emission factors. Volume I: stationary point and area sources. 5th ed. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. AP-42-ED-5. Available from: NTIS, Springfield, VA; PB95-196028.
- U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1999) Getting started: emissions inventory methods for PM-2.5. Research Triangle Park, NC: Office of Air Quality Planning and Standards, Emission Factor and Inventory Group; EIIP document series - volume IX. Available: <http://www.epa.gov/ttn/chief/eiip/techreport/volume09/index.html> [7 May 2003].
- U.S. Environmental Protection Agency. (2000a) Aerometric Information Retrieval System (AIRS). Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/airs/airsaq [7 May 2003].
- U.S. Environmental Protection Agency. (2000b) Air quality criteria for carbon monoxide. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/P-99/001F. Available: <http://www.epa.gov/ncea/pdfs/coaqcd.pdf> (7 May 2003).
- U.S. Environmental Protection Agency. (2000c) National air pollutant emission trends, 1900-1998. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA-454/R-00-002. Available: <http://www.epa.gov/ttn/chief/trends/trends98/> [7 May 2003].
- U.S. Environmental Protection Agency. (2001) National air quality and emissions trends report, 1999. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA 454/R-01-004. Available: <http://www.epa.gov/oar/aqtrnd99/> [7 May 2003].
- U.S. Environmental Protection Agency. (2002a) Latest findings on national air quality: 2001 status and trends. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-454/K-02-001.
- U.S. Environmental Protection Agency. (2002b) Aerometric Information Retrieval System (AIRS). Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/airs/airsaq [2 June 2003].
- Vermette, S. J.; Williams, A. L.; Landsberger, S. (1992) PM₁₀ source apportionment using local surface dust profiles: examples from Chicago. In: Chow, J. C.; Ono, D. M., eds. PM₁₀ standards and nontraditional particulate source controls, an A&WMA/EPA international specialty conference, v. I; January; Scottsdale, AZ. Pittsburgh, PA: Air & Waste Management Association; pp. 262-278. (A&WMA transactions series no. 22).
- Wainman, T.; Zhang, J.; Weschler, C. J.; Liroy, P. J. (2000) Ozone and limonene in indoor air: a source of submicron particle exposure. *Environ. Health Perspect.* 108: 1139-1145.
- Watson, J. G.; Chow, J. C. (2000) Reconciling urban fugitive dust emissions inventory and ambient source contribution estimates: summary of current knowledge and needed research. Reno, NV: Desert Research Institute; DRI document no. 6110.4F.
- Watson, J. G.; Chow, J. C. (2001) Estimating middle-, neighborhood-, and urban-scale contributions to elemental carbon in Mexico City with a rapid response aethalometer. *J. Air Waste Manage. Assoc.* 51: 1522-1528.
- Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990a) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. *Environ. Software* 5: 38-49.

- Watson, J. G.; Chow, J. C.; Pritchett, L. C.; Houck, J. A.; Ragazzi, R. A.; Burns, S. (1990b) Chemical source profiles for particulate motor vehicle exhaust under cold and high altitude operating conditions. *Sci. Total Environ.* 93: 183-190.
- Watson, J. G.; Chow, J. C.; Lu, Z.; Fujita, E. M.; Lowenthal, D. H.; Lawson, D. R.; Ashbaugh, L. L. (1994) Chemical mass balance source apportionment of PM₁₀ during the Southern California Air Quality Study. *Aerosol Sci. Technol.* 21: 1-36.
- West, J. J.; Ansari, A. S.; Pandis, S. N. (1999) Marginal PM_{2.5}: nonlinear aerosol mass response to sulfate reductions in the eastern United States. *J. Air Waste Manage. Assoc.* 49: 1415-1424.
- White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. *J. Air Waste Manage. Assoc.* 48: 454-458.
- Wilson, W. E.; Stockburger, L. (1990) Diurnal variations in aerosol composition and concentration. In: Masuda, S.; Takahashi, K., eds. *Aerosols: science, industry, health and environment: proceedings of the third international aerosol conference*, v. 2; September; Kyoto, Japan. Oxford, United Kingdom: Pergamon Press; pp. 962-965.
- Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. *J. Air Waste Manage. Assoc.* 47: 1238-1249.
- Winer, A. M.; Arey, J.; Atkinson, R.; Aschmann, S. M.; Long, W. D.; Morrison, C. L.; Olszyk, D. M. (1992) Emission rates of organics from vegetation in California's Central Valley. *Atmos. Environ. Part A* 26: 2647-2659.
- Winterhalter, R.; Neeb, P.; Grossmann, D.; Koloff, A.; Horie, O.; Moortgat, G. (2000) Products and mechanisms of the gas phase reaction of ozone with β -pinene. *J. Atmos. Chem.* 35: 165-197.
- Wongphatarakul, V.; Friedlander, S. K.; Pinto, J. P. (1998) A comparative study of PM_{2.5} ambient aerosol chemical databases. *Environ. Sci. Technol.* 32: 3926-3934.
- Woo, K.-S.; Chen, D.-R.; Pui, D. Y. H.; McMurry, P. H. (2001) Measurement of Atlanta aerosol size distributions: observations of ultrafine particle events. *Aerosol Sci. Technol.* 34: 75-87.
- Wotawa, G.; Trainer, M. (2000) The influence of Canadian forest fires on pollutant concentrations in the United States. *Science (Washington, DC)* 288: 324-328.
- Yu, J.; Cocker, D. R., III; Griffin, R. J.; Flagan, R. C.; Seinfeld, J. H. (1998) Gas-phase ozone oxidation of monoterpenes: gaseous and particulate products. *J. Atmos. Chem.* 34: 207-258.
- Yu, J.; Griffin, R. J.; Cocker, D. R., III; Flagan, R. C.; Seinfeld, J. H. (1999) Observation of gaseous and particulate products of monoterpene oxidation in forest atmospheres. *Geophys. Res. Lett.* 26: 1145-1148.
- Zhang, S.-H.; Shaw, M.; Seinfeld, J. H.; Flagan, R. C. (1992) Photochemical aerosol formation from α -pinene and β -pinene. *J. Geophys. Res. [Atmos.]* 97: 20,717-20,729.
- Zweidinger, R.; Purdue, L.; Fitzgerald, K.; Carmichael, L.; Kellog, R. (1998) NERL PM research monitoring platforms: Baltimore, Fresno and Phoenix, data report. Research Triangle Park, NC: U.S. Environmental Protection Agency, National Exposure Research Laboratory; report no. EPA/600/R-98/138. Available from: NTIS, Springfield, VA; PB99-105660.

APPENDIX 3A

Spatial and Temporal Variability of the Nationwide AIRS PM_{2.5} and PM_{10-2.5} Data Sets

Aspects of the spatial and temporal variability of 24-h average PM_{2.5} concentrations for 1999, 2000, and 2001 in a number of metropolitan statistical areas (MSAs) across the United States are presented in this Appendix. PM_{2.5} data for multiple sites in 27 urban areas have been obtained from the AIRS data base and analyzed for their seasonal variations, for their spatial correlations, and for their spatial uniformity. A number of aspects of the spatial and temporal variability of the PM_{2.5} data set from 1999 were presented in Rizzo and Pinto (2001) based in part on analyses given in Fitz-Simons et al. (2000).

Information about seasonal and spatial variability in PM_{2.5} concentrations within 27 MSAs across the United States are provided in the accompanying figures (Figures 3A-1 to 3A-27). Underneath the value for r , the 90th percentile (P_{90}) values of the absolute difference in PM_{2.5} concentrations (in $\mu\text{g}/\text{m}^3$) and the coefficient of divergence (COD) are provided in parentheses. Beneath these two measures of spatial variability, the number of observations used in the calculations of the statistics in part *c* of each figure is given.

Quality assured measurements from four monitoring sites for at least fifteen days during each calendar quarter for 1999, 2000, and 2001 (preferably) or for 2000 and 2001 at a minimum in a given MSA were required for their inclusion in the analyses presented in this appendix. The Columbia, SC and Baton Rouge, LA MSAs, which had only three sites meeting this criterion, are exceptions. Typically, at least 200 measurements were available for each monitoring site chosen. Monitoring sites were chosen without consideration of the land use type used to characterize their locations.

Because of changes in monitoring strategies, funding levels etc., there were year to year changes in monitoring sites meeting the above criteria in a number of MSAs. Data for the MSAs in Philadelphia, PA, Norfolk, VA, Pittsburgh, PA, Detroit, MI, Chicago, IL, Louisville, KY, St. Louis, MO, and the Dallas, TX have been analyzed only for 2000 and 2001 because of a lack of consistent coverage in 1999.

Information about seasonal and spatial variability in $PM_{10-2.5}$ concentrations within 17 MSAs across the United States are provided in the accompanying figures (Figures 3A-28 to 3A-44). Underneath the value for r , the 90th percentile values of the absolute difference in $PM_{2.5}$ concentrations (in $\mu\text{g}/\text{m}^3$) and the CODs are given in parentheses. Beneath these two measures of spatial variability, the number of observations used in the calculations of the statistics in part *c* of each figure is given. In order to maximize coverage, data were calculated for a number of sampling periods. Only Milwaukee, WI, and Salt Lake City, UT, had enough data for a 3 year average (1999 to 2001). Tampa, FL; Cleveland, OH; Steubenville, OH; Baton Rouge, LA; Portland, OR; and Riverside, CA had data for a 2 year average (2000 to 2001), as did Chicago, IL and Pittsburgh, PA (1999 to 2000). Other MSAs had only one year of data (for 2000 or 2001).

The COD was defined mathematically and used earlier in Chapter 3 as a measure of the degree of similarity between two data sets. A COD of zero implies that values in both data sets are identical, and a COD of one indicates that two data sets are completely different. Values of P_{90} provide a measure in absolute terms of differences in concentrations between sites, and CODs provide a relative measure of these differences. The maximum number of days of coincident data from paired sites were used to calculate correlation coefficients, values for P_{90} , and CODs. The correlation coefficients were also calculated by using only concurrent measurements obtained at all of the monitoring sites within urban areas meeting the above selection criteria. The correlation coefficients that were calculated differed only in the third significant figure between the two methods. Metrics used above for characterizing differences between separated monitors are applied to collocated monitors in Table 3A-1.

Figures 3A-28 to 3A-44 summarize information about the spatial and temporal variability of 24-h average $PM_{10-2.5}$ concentrations. Data are shown for a subset of MSAs included in the analyses for $PM_{2.5}$. Not all MSAs could be included, because of a lack of data for some. A schematic map showing locations of sampling sites within each MSA is given in part *a*, at the top of each figure. Also included in the map are major highways and a distance scale. A key giving the AIRS site ID numbers (#'s) is shown alongside each map. Box plots showing lowest, lower quartile, median, upper quartile and highest $PM_{2.5}$ concentrations for each calendar quarter are shown in part *b* of each figure. AIRS site ID #'s, annual mean concentrations, the number of observations, and the standard deviation of the data are shown above the box plots. Finally,

in part *c* of each figure, statistics characterizing the spatial variability in PM_{2.5} concentrations are given. For each site-pair, the Pearson correlation coefficient (*r*) is provided. Underneath each value for *r*, the 90th percentile of the absolute difference in PM_{10-2.5} concentrations, the COD, and number of observations is given. In some cases, because of negative concentration values, the COD may not be calculated. Dashes are shown for such cases.

It should be noted that the vertical axes for each MSA are drawn to different scales. Therefore, care should be taken in attempting to compare values in different MSAs. For PM_{2.5}, the maxima on the y-axis vary by up to a factor of four, from 40 µg/m³ to 150 µg/m³. The use of a single range of values would result in compression of the figures and loss of detail in a number of MSAs which have a lower range of values. Comparisons between MSAs can more easily be made by inspection of Figures 3-4a,b and 3-6a,b.



REFERENCES

- Fitz-Simons, T. S.; Mathias, S.; Rizzo, M. (2000) Analyses of 1999 PM data for the PM NAAQS review. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; November 17. Available: <http://www.epa.gov/oar/oaqps/pm25/analyses.html> [2 April, 2002].
- Rizzo, M.; Pinto, J. P. (2001) Initial characterization of fine particulate matter (PM_{2.5}) collected by the National Federal Reference Monitoring Network. Presented at: 94th annual conference & exhibition of the Air & Waste Management Association; June, Orlando, FL. Pittsburgh, PA: Air & Waste Management Association.

TABLE 3A-1. PERFORMANCE METRICS FOR PM_{2.5} FROM COLLOCATED SAMPLERS (Mean concentrations, the number of samples, the standard deviation at each site, and the Pearson correlation coefficient and the coefficient of divergence for the site pair are shown; concentrations are given in µg/m³).

Columbia, SC	Sampler 1	Sampler 2
	45-079-0019	45-079-0019
MEAN	15.3	15.5
N	226	211
SD	6.5	6.5
r		0.995
COD		0.022
Dallas, TX	Sampler 1	Sampler 2
	48-113-0069	48-113-0069
MEAN	12.7	13.1
N	687	116
SD	5.7	6
r		0.996
COD		0.032
Detroit, MI	Sampler 1	Sampler 2
	26-163-0001	26-163-0001
MEAN	16.5	16.2
N	637	111
SD	9.2	8.8
r		0.986
COD		0.041
Grand Rapids, MI	Sampler 1	Sampler 2
	26-081-0020	26-081-0020
MEAN	14	14.3
N	1050	181
SD	8.6	8.3
r		0.992
COD		0.059
Louisville, KY	Sampler 1	Sampler 2
	18-043-1004	18-043-1004
MEAN	15.8	16
N	196	104
SD	8.2	7.8
r		0.997
COD		0.027
Steubenville, OH	Sampler 1	Sampler 2
	54-029-0011	54-029-0011
Mean	16.5	16.6
Obs	342	325
SD	10.2	10.3
r		0.985
COD		0.087
Washington DC	Sampler 1	Sampler 2
	51805	51805
Mean	16.8	17.5
Obs	600	132
SD	9.7	11.1
r		0.953
COD		0.15

Philadelphia, PA MSA





c.

Site	A	B	C	D	E	F	G
A	1	0.91 (6.3, 0.14) 170	0.93 (5.2, 0.15) 167	0.87 (6.9, 0.19) 183	0.88 (5.0, 0.16) 176	0.94 (4.6, 0.15) 163	0.93 (5.1, 0.14) 166
B		1	0.84 (7.5, 0.19) 176	0.88 (7.4, 0.18) 194	0.83 (7.1, 0.18) 184	0.89 (7.4, 0.17) 169	0.85 (6.9, 0.18) 173
C			1	0.85 (7.5, 0.16) 199	0.88 (4.7, 0.13) 200	0.94 (4.9, 0.11) 177	0.89 (5.3, 0.13) 180
D				1	0.87 (6.1, 0.15) 208	0.94 (5.1, 0.11) 187	0.88 (4.0, 0.12) 193
E					1	0.90 (4.6, 0.11) 181	0.87 (4.5, 0.11) 185
F						1	0.96 (3.3, 0.08) 550
G							1

Figure 3A-1. Philadelphia, PA-NJ MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.

Washington, DC MSA





c.

Site	A	B	C	D	E	F
A	1	0.83 (5.1, 0.18) 784	0.91 (5.7, 0.15) 273	0.82 (7.4, 0.17) 691	0.84 (7.7, 0.19) 267	0.77 (8.6, 0.20) 262
B		1	0.84 (5.1, 0.16) 297	0.87 (6.2, 0.16) 731	0.79 (7.5, 0.20) 284	0.77 (7.6, 0.19) 281
C			1	0.96 (3.5, 0.08) 276	0.94 (4.9, 0.11) 312	0.85 (8.3, 0.16) 289
D				1	0.91 (4.5, 0.11) 266	0.82 (9.1, 0.18) 246
E					1	0.87 (8.5, 0.17) 284
F						1

Figure 3A-2. Washington, DC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Norfolk, VA MSA





c.

Site	A	B	C	D	E
A	1	0.94 (3.7, 0.09) 167	0.91 (5.0, 0.11) 180	0.96 (3.0, 0.08) 182	0.94 (3.5, 0.08) 184
B		1	0.96 (2.6, 0.09) 191	0.96 (3.1, 0.07) 194	0.93 (4.0, 0.09) 198
C			1	0.96 (3.6, 0.09) 206	0.92 (4.3, 0.11) 212
D				1	0.93 (3.6, 0.10) 217
E					1

Figure 3A-3. Norfolk, VA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Columbia, SC MSA



c.

Site	A	B	C
A	1	0.93 (3.3, 0.08) 316	0.95 (3.2, 0.07) 319
B		1	0.97 (2.8, 0.06) 313
C			1

Figure 3A-4. Columbia, SC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Atlanta, GA MSA




Figure 3A-5. Atlanta, GA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Birmingham, AL MSA






Figure 3A-6. Birmingham, AL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Tampa, FL MSA





c.

Site	A	B	C	D
A	1	0.79 (3.6, 0.10) 919	0.87 (4.0, 0.11) 920	0.87 (4.3, 0.12) 308
B		1	0.70 (4.6, 0.14) 939	0.71 (5.0, 0.13) 314
C			1	0.82 (3.1, 0.10) 325
D				1

Figure 3A-7. Tampa, FL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.

Cleveland, OH MSA



c.

Site	A	B	C	D	E	F	G	H
A	1	0.91 (7.1, 0.13) 320	0.96 (3.3, 0.12) 322	0.94 (5.4, 0.10) 314	0.92 (7.2, 0.16) 300	0.88 (9.0, 0.18) 308	0.89 (10.7, 0.21) 308	0.92 (8.1, 0.17) 265
B		1	0.92 (6.9, 0.14) 306	0.89 (9.4, 0.15) 296	0.85 (13.1, 0.21) 290	0.84 (12.9, 0.21) 304	0.84 (14.3, 0.23) 294	0.892 (11.2, 0.18) 256
C			1	0.93 (5.1, 0.14) 309	0.90 (8.15, 0.19) 300	0.87 (8.6, 0.20) 310	0.88 (10.8, 0.22) 307	0.90 (8.9, 0.18) 256
D				1	0.96 (4.7, 0.14) 295	0.91 (5.4, 0.16) 310	0.90 (7.8, 0.20) 306	0.91 (7.7, 0.18) 264
E					1	0.88 (4.9, 0.16) 304	0.91 (5.8, 0.15) 295	0.91 (5.8, 0.13) 247
F						1	0.89 (6.0, 0.18) 303	0.87 (5.3, 0.18) 261
G							1	0.90 (6.6, 0.15) 275
H								1

Figure 3A-8. Cleveland, OH MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Pittsburgh, PA MSA






Figure 3A-9. Pittsburgh, PA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Steubenville, OH MSA




c.


Site	A	B	C	D	E
A	1	0.85 (9.6, 0.16) 269	0.88 (8.0, 0.16) 294	0.86 (7.7, 0.16) 302	0.85 (7.9, 0.18) 290
B		1	0.87 (6.3, 0.16) 267	0.84 (8.6, 0.17) 277	0.79 (10, 0.20) 266
C			1	0.90 (7.4, 0.15) 320	0.89 (7.1, 0.16) 295
D				1	0.93 (6.2, 0.16) 307
E					1

Figure 3A-10. Steubenville, OH-Weirton, WV MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.

Detroit, MI MSA



AIRS Site ID	
Site A	26-099-0009
Site B	26-115-0005
Site C	26-125-0001
Site D	26-147-0005
Site E	26-163-0001
Site F	26-163-0015
Site G	26-163-0016
Site H	26-163-0025
Site I	26-163-0033
Site J	26-163-0036



c.

Site	A	B	C	D	E	F	G	H	I	J
A	1	0.89 (7.7, 0.16) 210	0.95 (5.0, 0.13) 172	0.95 (5.4, 0.14) 203	0.94 (6.7, 0.16) 202	0.92 (9.5, 0.19) 216	0.95 (6.8, 0.14) 202	0.93 (5.0, 0.13) 203	0.89 (12.8, 0.23) 205	0.90 (9.6, 0.18) 200
B		1	0.91 (6.5, 0.14) 169	0.83 (9.7, 0.20) 196	0.95 (5.6, 0.10) 200	0.91 (7.5, 0.15) 214	0.91 (6.8, 0.12) 202	0.91 (6.9, 0.13) 201	0.86 (11.9, 0.19) 202	0.90 (7.7, 0.13) 194
C			1	0.92 (6.4, 0.15) 161	0.96 (5.0, 0.12) 166	0.93 (7.3, 0.16) 174	0.96 (4.3, 0.10) 168	0.94 (5.0, 0.11) 164	0.91 (11.7, 0.19) 166	0.88 (8.9, 0.17) 162
D				1	0.88 (8.6, 0.19) 190	0.89 (10.1, 0.21) 203	0.92 (7.5, 0.17) 192	0.90 (6.5, 0.15) 190	0.84 (13.8, 0.25) 193	0.82 (11.8, 0.22) 190
E					1	0.96 (4.6, 0.09) 210	0.94 (4.6, 0.10) 580	0.94 (5.9, 0.13) 200	0.89 (10.8, 0.15) 196	0.93 (5.0, 0.10) 192
F						1	0.97 (4.8, 0.11) 210	0.90 (9.1, 0.17) 207	0.90 (8.7, 0.13) 210	0.91 (6.6, 0.12) 204
G							1	0.94 (6.4, 0.12) 204	0.92 (10.4, 0.16) 201	0.91 (7.1, 0.12) 192
H								1	0.86 (12.8, 0.21) 197	0.87 (9.7, 0.17) 191
I									1	0.90 (7.7, 0.13) 193
J										1

Figure 3A-11. Detroit MI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Grand Rapids, MI MSA




Figure 3A-12. Grand Rapids, MI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Milwaukee, WI MSA




Figure 3A-13. Milwaukee, WI MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Chicago, IL MSA






Figure 3A-14. Chicago, IL MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Gary, IN MSA





c.

Site	A	B	C	D
A	1	0.65 (9.1, 0.19) 823	0.93 (4.2, 0.13) 822	0.91 (5.0, 0.17) 277
B		1	0.59 (11.3, 0.20) 841	0.56 (10.9, 0.23) 277
C			1	0.92 (6.4, 0.17) 281
D				1

Figure 3A-15. Gary, IN MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.



Louisville, KY MSA



c.

Site	A	B	C	D	E
A	1	0.86 (5.8, 0.13) 279	0.90 (6.0, 0.12) 289	0.91 (3.9, 0.10) 315	0.91 (3.8, 0.08) 273
B		1	0.87 (6.0, 0.15) 260	0.89 (5.9, 0.14) 282	0.91 (5.3, 0.13) 250
C			1	0.93 (5.4, 0.12) 307	0.90 (5.6, 0.12) 256
D				1	0.93 (4.2, 0.11) 302
E					1

Figure 3A-16. Louisville, KY MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.





c.

Site	A	B	C	D	E	F	G	H	I	J	K
A	1	0.81 (9.8, 0.15) 207	0.73 (12.2, 0.20) 195	0.74 (11.9, 0.20) 207	0.67 (14.6, 0.22) 218	0.71 (14.0, 0.23) 215	0.68 (14.6, 0.23) 214	0.69 (15.2, 0.25) 217	0.69 (14.1, 0.21) 221	0.69 (15.0, 0.23) 207	0.68 (13.9, 0.20) 217
B		1	0.85 (6.8, 0.12) 195	0.86 (6.7, 0.12) 210	0.84 (7.3, 0.14) 217	0.84 (7.8, 0.16) 216	0.79 (7.5, 0.16) 215	0.84 (8.3, 0.17) 217	0.86 (6.7, 0.13) 226	0.85 (7.6, 0.15) 211	0.85 (6.5, 0.12) 221
C			1	0.86 (4.8, 0.13) 196	0.86 (5.9, 0.12) 209	0.94 (4.0, 0.11) 206	0.84 (6.3, 0.13) 207	0.90 (5.7, 0.13) 208	0.91 (5.2, 0.11) 228	0.91 (5.4, 0.12) 214	0.89 (5.0, 0.11) 224
D				1	0.83 (5.8, 0.14) 222	0.86 (5.7, 0.15) 221	0.81 (6.1, 0.15) 219	0.86 (6.1, 0.15) 222	0.86 (5.3, 0.14) 229	0.87 (5.5, 0.14) 216	0.84 (5.5, 0.14) 225
E					1	0.90 (5.3, 0.11) 233	0.90 (4.0, 0.11) 231	0.93 (4.1, 0.11) 234	0.95 (3.7, 0.08) 234	0.95 (3.8, 0.09) 218	0.93 (4.3, 0.09) 230
F						1	0.87 (4.8, 0.12) 230	0.94 (3.8, 0.09) 233	0.93 (5.1, 0.11) 233	0.93 (4.7, 0.10) 218	0.91 (5.4, 0.12) 229
G							1	0.92 (2.8, 0.10) 232	0.90 (3.7, 0.11) 231	0.90 (3.1, 0.10) 215	0.87 (5.3, 0.12) 227
H								1	0.95 (4.3, 0.10) 234	0.95 (2.9, 0.09) 218	0.94 (4.9, 0.12) 230
I									1	0.98 (2.8, 0.07) 671	0.97 (2.3, 0.06) 691
J										1	0.96 (3.3, 0.08) 652
K											1

Figure 3A-17. St. Louis, MO MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Baton Rouge, LA MSA





c.

Site	A	B	C
A	1	0.93 (2.7, 0.08) 326	0.93 (2.9, 0.09) 318
B		1	0.97 (2.5, 0.07) 1006
C			1

Figure 3A-18. Baton Rouge, LA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Kansas City, MO MSA



c.

Site	A	B	C	D	E	F
A	1	0.94 (1.9, 0.09) 320	0.92 (4.2, 0.12) 304	0.90 (4.3, 0.13) 318	0.93 (3.8, 0.10) 296	0.89 (5.9, 0.15) 320
B		1	0.90 (4.0, 0.13) 312	0.89 (4.1, 0.14) 322	0.90 (3.6, 0.12) 300	0.87 (6.2, 0.17) 326
C			1	0.96 (3.1, 0.09) 327	0.96 (2.8, 0.10) 300	0.90 (6.5, 0.16) 329
D				1	0.95 (2.9, 0.09) 940	0.95 (4.0, 0.11) 338
E					1	0.94 (5.1, 0.12) 314
F						1

Figure 3A-19. Kansas City, KS-MO MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Dallas, TX MSA




Figure 3A-20. Dallas, TX MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Boise, ID MSA






Figure 3A-21. Boise, ID MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.

Salt Lake City, UT MSA





c.

Site	A	B	C	D	E	F
A	1	0.89 (8.8, 0.22) 334	0.90 (9.9, 0.27) 331	0.94 (6.9, 0.19) 322	0.90 (8.2, 0.22) 315	0.94 (4.4, 0.15) 306
B		1	0.89 (7.0, 0.19) 327	0.92 (4.9, 0.17) 316	0.89 (6.1, 0.17) 313	0.88 (9.3, 0.20) 307
C			1	0.93 (7.1, 0.19) 314	0.96 (4.8, 0.17) 308	0.86 (11.4, 0.24) 302
D				1	0.94 (5.3, 0.16) 300	0.92 (8.7, 0.18) 297
E					1	0.89 (9.6, 0.20) 288
F						1

Figure 3A-22. Salt Lake City, UT MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Seattle, WA MSA





c.

Site	A	B	C	D	E
A	1	0.31 (15.0, 0.39) 303	0.28 (15.7, 0.43) 304	0.45 (8.8, 0.32) 291	0.37 (17.9, 0.39) 298
B		1	0.96 (3.6, 0.14) 1021	0.92 (6.8, 0.16) 774	0.81 (6.2, 0.17) 344
C			1	0.91 (7.5, 0.20) 779	0.79 (8.2, 0.20) 344
D				1	0.75 (8.5, 0.20) 327
E					1

Figure 3A-23. Seattle, WA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.

Portland, OR MSA



c.

Site	A	B	C	D
A	1	0.79 (6.5, 0.24) 433	0.90 (5.1, 0.20) 429	0.81 (4.5, 0.19) 427
B		1	0.77 (4.1, 0.14) 986	0.89 (4.3, 0.17) 477
C			1	0.83 (4.5, 0.19) 472
D				1

Figure 3A-24. Portland, OR MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Los Angeles, CA MSA






Figure 3A-25. Los Angeles-Long Beach, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P₉₀, COD), and number of measurements.

Riverside, CA MSA





c.

Site	A	B	C	D	E
A	1	0.94 (6.6, 0.10) 294	0.83 (14.3, 0.21) 289	0.93 (10.6, 0.13) 306	0.90 (10.5, 0.13) 278
B		1	0.81 (17.8, 0.23) 290	0.93 (13.3, 0.14) 313	0.91 (11.9, 0.13) 275
C			1	0.86 (11.8, 0.20) 302	0.78 (16.9, 0.22) 269
D				1	0.94 (8.9, 0.11) 290
E					1

Figure 3A-26. Riverside-San Bernadino, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.



San Diego, CA MSA



c.

Site	A	B	C	D
A	1	0.76 (10.0, 0.16) 270	0.73 (10.0, 0.19) 253	0.83 (7.6, 0.16) 255
B		1	0.85 (6.3, 0.13) 773	0.78 (9.7, 0.18) 769
C			1	0.73 (11.0, 0.20) 728
D				1

Figure 3A-27. San Diego, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000-2001; (c) Intersite correlation statistics for each data pair given as the correlation coefficient, (P_{90} , COD), and number of measurements.





c.

Site	A	B
A	1	0.70 (8.0, 0.37) 49
B		1

Figure 3A-28. Columbia, SC MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.



Tampa, FL MSA



c.

Site	A	B
A	1	0.81 (5.3, 0.17) 95
B		1

Figure 3A-29. Tampa, FL MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.





c.

Site	A	B	C	D	E	F
A	1	0.67 (23.2, 0.26) 182	0.67 (28.5, 0.28) 95	0.73 (17.9, 0.22) 97	0.62 (27.0, 0.31) 98	0.41 (40.0, 0.60) 94
B		1	0.65 (16.1, 0.22) 90	0.73 (11.7, 0.18) 93	0.69 (13.9, 0.62) 90	0.44 (24.9, 0.53) 89
C			1	0.66 (18.1, 0.23) 94	0.71 (10.6, 0.31) 102	0.49 (19.9, 0.50) 99
D				1	0.74 (15.4, 0.38) 93	0.31 (28.0, 0.59) 94
E					1	0.22 (20.4, 0.55) 99
F						1

Figure 3A-30. Cleveland, OH MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{10-2.5} concentrations (µg/m³) for 2000-2001; (c) Intersite correlation coefficients, (P₉₀, COD), and number of observations.

Steubenville, OH MSA





c.

Site	A	B	C	D
A	1	0.64 (10.9, 0.77) 83	0.69 (11.6, --) 91	0.68 (11.3, --) 100
B		1	0.54 (14.7, --) 166	0.48 (18.5, --) 88
C			1	0.69 (12.8, --) 97
D				1

Figure 3A-31. Steubenville, OH MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Detroit, MI MSA





c.

Site	A	B	C
A	1	0.58 (29.4, --) 53	0.54 (15.7, 0.54) 50
B		1	0.39 (34.9, 0.79) 51
C			1

Figure 3A-32. Detroit, MI MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2000; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Milwaukee, WI MSA



c.

Site	A	B
A	1	0.65 (9.2, 0.53) 150
B		1

Figure 3A-33. Milwaukee, WI MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 1999-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Chicago, IL MSA






Figure 3A-34. Chicago, IL MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 1999-2000; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Gary, IN MSA




c.


Site	A	B	C
A	1	0.79 (7.8, -) 49	0.63 (6.3, -) 49
B		1	0.60 (8.0, 0.83) 50
C			1

Figure 3A-35. Gary, IN MSA (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average PM_{10-2.5} concentrations (µg/m³) for 2001; (c) Intersite correlation coefficients, (P₉₀, COD), and number of observations.

Louisville, KY MSA





AIRS Site ID	
Site A	18-019-0005
Site B	21-029-0006



c.

Site	A	B
A	1	0.65 (5.5, 0.48) 46
B		1



Figure 3A-36. Louisville, KY MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu\text{g}/\text{m}^3$) for 2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.



c.

Site	A	B	C
A	1	0.70 (27.2, -) 51	0.73 (26.2, 0.76) 47
B		1	0.82 (13.0, 0.91) 50
C			1

Figure 3A-37. St. Louis, MO MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.





c.

Site	A	B
A	1	0.40 (22.4, 0.43) 96
B		1

Figure 3A-38. Baton Rouge, LA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Dallas, TX MSA





c.

Site	A	B	C	D
A	1	0.79 (4.5, 0.17) 54	0.71 (9.3, 0.22) 55	0.66 (16.5, 0.32) 54
B		1	0.69 (7.8, 0.18) 50	0.60 (13.2, 0.30) 50
C			1	0.69 (13.5, 0.24) 50
D				1

Figure 3A-39. Dallas, TX MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Salt Lake City, UT MSA





c.

Site	A	B	C
A	1	0.72 (28.7, --) 283	0.74 (9.8, --) 264
B		1	0.70 (27.6, 0.47) 274
C			1

Figure 3A-40. Salt Lake City, UT MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 1999-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Portland, OR MSA



c.

Site	A	B
A	1	0.69 (5.1, --) 107
B		1

Figure 3A-41. Portland, OR MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Los Angeles, CA MSA






Figure 3A-42. Los Angeles, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

Riverside, CA MSA



c.

Site	A	B	C	D
A	1	0.36 (42.6, 0.38) 184	0.32 (36.6, 0.39) 102	0.45 (39.0, 0.38) 107
B		1	0.62 (25.9, 0.32) 104	0.79 (18.2, 0.33) 108
C			1	0.80 (13.3, 0.28) 98
D				1

Figure 3A-43. Riverside, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2000-2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.




Figure 3A-44. San Diego, CA MSA. (a) Locations of sampling sites by AIRS ID#; (b) Quarterly distribution of 24-h average $PM_{10-2.5}$ concentrations ($\mu g/m^3$) for 2001; (c) Intersite correlation coefficients, (P_{90} , COD), and number of observations.

APPENDIX 3B

Aerosol Composition Data from the Speciation Network

The U.S. Environmental Protection Agency (EPA), working with state and local air quality agencies, began implementing an ambient air monitoring network in 1999/2000 to provide a consistent data set for the characterization and evaluation of trends in PM components (chemical species). The network was designed to include about 52 core trends sites across the United States and to provide a stable ongoing national perspective. In 1999, an initial thirteen sites were installed and operated to be used as a model for the deployment of the more comprehensive network consisting of the 52 core trends sites and roughly 200 additional locally relevant sites.

Data from the initial 13 sites were presented in the Third External Review Draft of this document, released by the EPA in 2001 for public comment and CASAC review. These sites were designed and operated to evaluate the suitability of various aerosol sampling devices for obtaining PM_{2.5} composition data, and data from those sites were summarized in Appendix 3B of the 2001 Third External Review Draft. Three types of collocated aerosol sampling devices were used to collect the data from February through July 2000, and data obtained from the three sampling devices were shown for each site. A complete description of the data, techniques used to analyze the filters, and the results of the evaluation of the performance of the sampling devices (including a number of caveats regarding the data) can be found in Coutant and Stetzer (2001) and the analyses of data in Coutant et al. (2001).

More recent measurements of ambient levels of PM_{2.5} constituents are presented in the tables provided below. Table 3B-1 presents the locations and general sampling schedules for 13 monitoring sites included for various reasons and representing a cross section of the country. Many of these sites were chosen because they are located in (or close to) MSAs in which risk assessments are to be performed. Sacramento, CA and Riverside, CA, for example, are located near San Francisco and Los Angeles, respectively. The data reported here are from the period October 2001 to September 2002. For this time period, a total of 51 sites (including both “Trends” and “non-Trends” sites) across the country have complete data (as defined by 50% of

observations available for every quarter for each of the major chemical species: sulfate, ammonium, nitrate, elemental carbon, organic carbon, and the five trace elements that go into the calculation of the crustal contribution to $PM_{2.5}$ (Al, Fe, Ca, Ti, and Si).

Summary statistics for concentrations of $PM_{2.5}$ and chemical species are given for each of the 13 sites in Table 3B-2. The number of samples (n) and the AIRS site code are given above each table. Entries in the tables provide the mean, minimum, and maximum component concentrations, and minimum detection limits for each component. Minimum detection levels (MDL) differ among the various sampling methods; these limits were estimated by Research Triangle Institute (RTI) in July of 2001 and are subject to review, revision, and reinterpretation. Anions and cations (ammonium, nitrate, sodium, potassium, and sulfate) were determined by ion chromatography; carbonaceous species were determined by the thermal optical transmittance method (NIOSH method); and trace elements (aluminum through zirconium) were determined by X-ray fluorescence spectrometry. The sulfate (calculated) entry is based on the XRF determination of sulfur. In general, relatively good agreement is found between the reconstructed mass and the $PM_{2.5}$ concentration measured by the collocated FRM monitor at each site. However, there are exceptions at several locations, as can be seen from inspection of Table 3B-2. Somewhat different sampling trains are used in the different sampling systems. These differences can result in differences in performance metrics and in differences in the entities that are measured. All samplers use a denuder in front of the filter for species to be analyzed by ion chromatography. In the RAAS and SASS samplers the denuder is followed by a nylon filter, whereas in the MASS samplers (Chicago, Houston, Seattle) a teflon filter is followed by a nylon filter. Particulate nitrate is collected on the teflon filter and is referred to as nonvolatile nitrate. However, there may be volatilization of nitrate containing compound from the teflon filter which is then adsorbed on the nylon filter. Nitrate extracted from the nylon filter is referred to as volatile nitrate.

Organic carbon (OC) concentrations are multiplied by a factor of 1.4 when calculating mass to account for the presence of H, N, and O in organic compounds on all samplers. Carbonate carbon has never been detected in any of the samples. Field blank corrections that could be applied to elemental carbon (EC) and OC concentrations are shown in Table 3B-3 for different samplers. Blank corrections for OC and EC shown in Table 3B-3 were applied to concentrations shown in Table 3B-2. However, subtracting blank corrections from OC concentrations results in negative values in several cases. A possible cause for results such as

these could have been that not enough blank samples were obtained to fully characterize the blank levels. Although the concentrations of 47 elements could be obtained by X-ray fluorescence spectrometry, the concentrations of many of these elements are beneath MDLs and are not shown. The same elements shown in Appendix 6A of the 1996 PM AQCD are shown here. The usual practice of denoting table entry values below MDL by (—) is followed here. Missing data for PM_{2.5} are also indicated by (—). Often, environmental data below MDL are represented by 0.5 x MDL for various purposes such as risk assessments and for mass closure. Unfortunately, speciation data are not available for the PM_{10-2.5} size fraction.

REFERENCES

- Coutant, B.; Stetzer, S. (2001) Evaluation of PM_{2.5} speciation sampler performance and related sample collection and stability issues: final report. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-454/R-01-008. Available: <http://www.epa.gov/ttn/amtic/pmspec.html> [5 April, 2002].
- Coutant, B.; Zhang, X.; Pivetz, T. (2001) Summary statistics and data displays for the speciation minitrends study: final report. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; contract no. 68-D-98-030.

TABLE 3B-1. PM_{2.5} SPECIATION SAMPLERS BY LOCATION: SITES SELECTED FOR SUMMARY IN 2004 PM AQCD

Sampler Type	Location_Name	State	Began Operation	AIRS_Code	POC	Schedule	Reason for Inclusion*
SASS	Burlington	VT	12/14/2000	000500070012	5	1-in-3	3
SASS	Philadelphia (AMS Laboratory)	PA	9/10/2001	000421010004	7	1-in-3	3
RAAS	Atlanta (South DeKalb)	GA	3/2/2001	000130890002	5	1-in-3	2
SASS	Detroit (Allen Park)	MI	12/14/2000	000261630001	5	1-in-3	1
MASS	Chicago (Com ED)	IL	5/22/2001	000170310076	5	Alt 1-in-3	3
RAAS	St. Louis (Blair Street)	MO	2/9/2000	000295100085	6	1-in-3	1
MASS	Houston (Deer Park)	TX	2/9/2000	000482011039	37778	1-in-3	3
SASS	Minneapolis (Philips)	MN	12/14/2000	000270530963	5	Alt 1-in-3	2
SASS	Boulder (Commerce City)	CO	2/18/2001	000080010006	5	Alt 1-in-3	3
SASS	Phoenix (Supersite)	AZ	2/21/2000	000040139997	7	1-in-3	1
MASS	Seattle (Beacon Hill)	WA	2/9/2000	000530330080	6	1-in-3	3
SASS	Sacramento (Del Paso Manor)	CA	2/9/2000	000060670006	5	1-in-3	2
SASS	Riverside-Rubidoux	CA	5/13/2001	000060658001	37746	1-in-3	1

* Reason for Inclusion:

1 = Completeness of geographic coverage

2 = PM₁₀ risk assessment city

3 = PM_{2.5} and PM₁₀ risk assessment city

**TABLE 3B-2a. BURLINGTON, VT SUMMARY DATA
(October 2001 to September 2002).**

All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 201; AIRS Site Code: 500070012

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	10.0	61.0	1.0	—
PM _{2.5} (Reconstructed Mass)	10.9	63.0	1.6	—
Sulfate (Calculated)	2.6	17.7	0.001	—
Sulfate (by I.C.)	2.8	18.8	0.17	0.012
Ammonium (by I.C.)	1.1	6.8	—	0.017
Sodium Ion (by I.C.)	0.22	2.1	—	0.030
Potassium (by I.C.)	0.035	0.38	—	0.014
Nitrate	1.3	8.8	—	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.26	0.84	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	2.5	31.4	-0.43	0.15
Aluminum	0.020	0.28	—	0.011
Arsenic	—	0.005	—	0.002
Barium	—	0.12	—	0.059
Bromine	0.002	0.007	—	0.002
Cadmium	—	0.014	—	0.011
Calcium	0.035	0.19	—	0.003
Chlorine	—	0.12	—	0.006
Chromium	0.002	0.060	—	0.002
Copper	0.002	0.014	0.001	0.001
Iron	0.041	0.390	0.007	0.002
Lead	—	0.016	—	0.006
Magnesium	—	0.19	—	0.018
Manganese	0.002	0.007	—	0.002
Molybdenum	—	0.012	—	0.005
Nickel	0.002	0.022	0.001	0.001
Phosphorous	—	0.092	—	0.006
Potassium	0.041	0.41	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	—	0.005	—	0.002
Silicon	0.069	0.54	—	0.008
Sodium	0.10	0.40	—	0.051
Strontium	—	0.004	—	0.003
Sulfur	0.88	5.9	—	0.007
Tin	—	0.033	—	0.018
Titanium	0.004	0.023	—	0.002
Vanadium	—	0.006	—	0.002
Zinc	0.008	0.057	0.007	0.002

**TABLE 3B-2b. PHILADELPHIA, PA SUMMARY DATA (October 2001 to
September 2002). All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 262;
AIRS Site Code: 421010004**

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	14.2	86.8	2.1	—
PM _{2.5} (Reconstructed Mass)	16.0	239.0	0.0	—
Sulfate (Calculated)	4.3	29.0	0.004	—
Sulfate (by I.C.)	4.4	30.5	0.020	0.012
Ammonium (by I.C.)	2.0	11.4	—	0.017
Sodium Ion (by I.C.)	0.21	1.6	—	0.030
Potassium (by I.C.)	0.042	0.83	—	0.014
Nitrate	2.1	8.7	0.050	0.008
Volatile Nitrate	0.84	4.0	0.030	—
Nonvolatile Nitrate	0.61	5.6	0.020	—
Elemental Carbon	0.66	2.3	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	3.1	50.5	-1.13	0.15
Aluminum	0.019	0.54	—	0.011
Arsenic	—	0.006	—	0.002
Barium	—	0.12	—	0.059
Bromine	0.004	0.013	—	0.002
Cadmium	—	0.017	—	0.011
Calcium	0.037	0.18	—	0.003
Chlorine	0.011	0.68	—	0.006
Chromium	0.002	0.018	—	0.002
Copper	0.004	0.025	0.003	0.001
Iron	0.084	0.53	—	0.002
Lead	0.005	0.025	—	0.005
Magnesium	—	0.30	—	0.018
Manganese	0.002	0.009	—	0.002
Molybdenum	—	0.012	—	0.005
Nickel	0.006	0.13	0.001	0.001
Phosphorous	—	0.039	—	0.006
Potassium	0.053	1.1	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	0.002	0.005	—	0.002
Silicon	0.086	1.1	—	0.008
Sodium	0.087	0.63	—	0.051
Strontium	—	0.016	—	0.003
Sulfur	1.45	9.6	—	0.007
Tin	—	0.035	—	0.018
Titanium	0.006	0.049	—	0.002
Vanadium	0.004	0.035	—	0.002
Zinc	0.015	0.095	0.007	0.001

TABLE 3B-2c. ATLANTA, GA SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 183; AIRS Site Code: 130890002

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	—	—	—	—
PM _{2.5} (Reconstructed Mass)	16.3	40.0	4.7	—
Sulfate (Calculated)	4.8	14.5	0.27	—
Sulfate (by I.C.)	4.8	15.2	0.88	0.011
Ammonium (by I.C.)	1.3	4.2	—	0.015
Sodium Ion (by I.C.)	0.27	1.7	—	0.028
Potassium (by I.C.)	0.044	0.31	—	0.013
Nitrate	0.70	3.5	0.16	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.90	3.5	—	0.13
Carbonate Carbon	—	—	—	0.13
Organic Carbon	4.3	11.2	0.8	0.13
Aluminum	0.028	0.65	—	0.004
Arsenic	0.001	0.014	—	0.001
Barium	—	0.054	—	0.024
Bromine	0.003	0.009	—	0.001
Cadmium	—	0.006	—	0.004
Calcium	0.037	0.28	0.006	0.001
Chlorine	0.003	0.049	—	0.002
Chromium	—	0.003	—	0.001
Copper	0.002	0.011	—	0.001
Gold	—	0.004	—	0.002
Iron	0.084	0.47	0.013	0.001
Lead	0.003	0.008	—	0.002
Magnesium	0.008	0.16	—	0.007
Manganese	0.002	0.011	—	0.001
Molybdenum	—	0.003	—	0.002
Nickel	—	0.002	—	0.001
Phosphorous	—	0.017	—	0.003
Potassium	0.060	0.36	0.015	0.001
Rubidium	0.005	0.002	—	0.001
Selenium	0.001	0.004	0.000	0.001
Silicon	0.11	1.4	0.010	0.003
Sodium	0.056	0.31	—	0.021
Strontium	0.001	0.005	0.001	0.001
Sulfur	1.6	4.8	0.089	0.003
Tin	—	0.019	—	0.007
Titanium	0.006	0.043	—	0.001
Vanadium	0.001	0.003	—	0.001
Zinc	0.008	0.034	0.001	0.001

TABLE 3B-2d. DETROIT, MI SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 189; AIRS Site Code: 261630001

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	16.6	53.1	2.6	
PM _{2.5} (Reconstructed Mass)	18.0	52.9	3.2	—
Sulfate (Calculated)	4.4	22.0	0.010	
Sulfate (by I.C.)	4.6	24.4	0.040	0.012
Ammonium (by I.C.)	2.2	9.3	0.009	0.017
Sodium Ion (by I.C.)	0.27	1.7	0.015	0.030
Potassium (by I.C.)	0.061	1.2	0.007	0.014
Nitrate	3.1	15.2	0.004	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.68	3.7	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	3.2	11.2	-0.46	0.15
Aluminum	0.025	0.53	—	0.011
Arsenic	0.002	0.010	—	0.002
Barium	—	0.11	—	0.059
Bromine	0.003	0.011	—	0.002
Cadmium	—	0.014	—	0.011
Calcium	0.069	0.33	—	0.003
Chlorine	0.017	0.65	—	0.006
Chromium	0.002	0.033	—	0.002
Copper	0.006	0.043	0.001	0.001
Iron	0.12	0.60	0.003	0.002
Lead	0.006	0.034	—	0.005
Magnesium	0.020	0.30	—	0.018
Manganese	0.004	0.025	—	0.002
Molybdenum	—	0.013	—	0.005
Nickel	0.002	0.022	0.001	0.001
Phosphorous	—	0.028	—	0.006
Potassium	0.078	1.3	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	0.002	0.011	—	0.002
Silicon	0.11	0.84	—	0.008
Sodium	0.098	0.53	—	0.051
Strontium	—	0.021	—	0.003
Sulfur	1.5	7.3	—	0.007
Tin	—	0.032	—	0.018
Titanium	0.007	0.079	—	0.002
Vanadium	0.002	0.012	—	0.002
Zinc	0.025	0.19	0.001	0.001

TABLE 3B-2e. CHICAGO, IL SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 139; AIRS Site Code: 170310076

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	15.7	42.7	3.6	—
PM _{2.5} (Reconstructed Mass)	15.0	44.5	3.4	—
Sulfate (Calculated)	4.3	22.8	0.64	—
Sulfate (by I.C.)	4.2	21.2	0.56	0.005
Ammonium (by I.C.)	1.9	8.8	0.17	0.007
Sodium Ion (by I.C.)	0.052	0.26	—	0.012
Potassium (by I.C.)	0.068	3.1	—	0.006
Nitrate	2.0	10.2	0.12	0.003
Volatile Nitrate	0.71	3.6	0.04	0.003
Nonvolatile Nitrate	1.3	9.9	0.03	0.003
Elemental Carbon	0.61	1.6	—	0.059
Carbonate Carbon	—	—	—	0.059
Organic Carbon	2.8	7.4	0.31	0.059
Aluminum	0.03	1.1	—	0.004
Arsenic	0.001	0.006	0.001	0.001
Barium	—	0.22	—	0.024
Bromine	0.003	0.011	—	0.001
Cadmium	—	0.007	—	0.004
Calcium	0.058	0.55	0.10	0.001
Chlorine	0.017	0.67	—	0.002
Chromium	0.001	0.005	—	0.001
Copper	0.004	0.056	—	0.001
Iron	0.091	0.53	0.014	0.001
Lead	0.006	0.040	—	0.002
Magnesium	0.017	0.69	—	0.007
Manganese	0.003	0.014	—	0.001
Molybdenum	—	0.006	—	0.002
Nickel	0.001	0.007	—	0.001
Phosphorous	0.002	0.015	—	0.003
Potassium	0.085	3.1	0.005	0.001
Rubidium	—	0.001	—	0.001
Selenium	0.001	0.004	0.001	0.001
Silicon	0.11	2.2	0.010	0.003
Sodium	0.048	0.58	—	0.021
Strontium	0.001	0.061	0.001	0.001
Sulfur	1.4	7.6	0.21	0.003
Tin	—	0.023	—	0.007
Titanium	0.004	0.070	—	0.001
Vanadium	0.001	0.002	—	0.001
Zinc	0.023	0.093	0.001	0.001

TABLE 3B-2f. ST. LOUIS, MO SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 324; AIRS Site Code: 295100085

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	15.6	52.1	3.8	—
PM _{2.5} (Reconstructed Mass)	16.4	53.2	2.8	—
Sulfate (Calculated)	4.3	22.5	0.61	—
Sulfate (by I.C.)	4.3	20.6	0.64	0.011
Ammonium (by I.C.)	1.9	7.2	0.14	0.015
Sodium Ion (by I.C.)	0.19	1.6	0.014	0.028
Potassium (by I.C.)	0.069	5.9	0.007	0.013
Nitrate	2.3	14.0	0.20	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.72	2.7	—	0.13
Carbonate Carbon	—	—	—	0.13
Organic Carbon	3.5	11.5	—	0.13
Aluminum	0.044	2.5	—	0.004
Arsenic	0.002	0.011	0.001	0.001
Barium	0.024	0.38	—	0.024
Bromine	0.004	0.16	—	0.001
Cadmium	—	0.010	—	0.004
Calcium	0.13	1.1	0.010	0.001
Chlorine	0.027	0.57	—	0.002
Chromium	0.002	0.034	—	0.001
Copper	0.018	0.72	—	0.001
Iron	0.24	1.3	0.012	0.001
Lead	0.014	0.09	—	0.002
Magnesium	0.011	0.79	—	0.007
Manganese	0.014	0.13	—	0.001
Molybdenum	—	0.007	—	0.002
Nickel	0.002	0.04	—	0.001
Phosphorous	0.004	0.095	—	0.003
Potassium	0.099	5.6	0.016	0.001
Rubidium	—	0.002	—	0.001
Selenium	0.001	0.006	—	0.001
Silicon	0.17	4.0	0.011	0.003
Sodium	0.045	0.43	—	0.021
Strontium	0.001	0.12	0.001	0.001
Sulfur	1.4	7.5	0.20	0.003
Tin	0.008	0.028	—	0.007
Titanium	0.008	0.14	—	0.001
Vanadium	0.001	0.011	—	0.001
Zinc	0.033	0.48	0.002	0.001

TABLE 3B-2g. HOUSTON, TX SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 229; AIRS Site Code: 482011039

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	12.4	21.9	5.9	—
PM _{2.5} (Reconstructed Mass)	11.0	48.0	0.0	—
Sulfate (Calculated)	3.4	24.5	0.004	—
Sulfate (by I.C.)	3.5	26.0	0.030	0.005
Ammonium (by I.C.)	1.2	8.8	—	0.007
Sodium Ion (by I.C.)	0.22	1.1	—	0.012
Potassium (by I.C.)	0.051	0.38	—	0.006
Nitrate	0.68	5.8	0.040	0.003
Volatile Nitrate	0.37	3.5	0.010	0.003
Nonvolatile Nitrate	0.32	4.2	0.020	0.003
Elemental Carbon	0.30	1.1	—	0.059
Carbonate Carbon	—	—	—	0.059
Organic Carbon	2.2	12.1	—	0.059
Aluminum	0.092	1.6	—	0.004
Arsenic	0.001	0.006	0.001	0.001
Barium	—	0.059	—	0.024
Bromine	0.003	0.022	—	0.001
Cadmium	—	0.008	—	0.004
Calcium	0.055	0.44	0.001	0.001
Chlorine	0.050	0.70	—	0.002
Chromium	0.001	0.009	—	0.001
Copper	0.003	0.024	—	0.001
Iron	0.073	0.83	—	0.001
Lead	0.002	0.007	—	0.002
Magnesium	0.020	0.38	—	0.007
Manganese	0.002	0.013	—	0.001
Molybdenum	0.002	0.022	—	0.002
Nickel	0.002	0.047	—	0.001
Phosphorous	0.002	0.021	—	0.003
Potassium	0.072	0.64	0.001	0.001
Rubidium	—	0.002	—	0.001
Selenium	0.001	0.001	0.001	0.001
Silicon	0.23	2.8	—	0.003
Sodium	0.17	1.1	—	0.021
Strontium	0.001	0.009	0.001	0.001
Sulfur	1.1	8.2	—	0.003
Tin	—	0.019	—	0.007
Titanium	0.007	0.091	—	0.001
Vanadium	0.003	0.011	—	0.001
Zinc	0.006	0.030	—	0.001

**TABLE 3B-2h. MINNEAPOLIS, MN SUMMARY DATA (October 2001 to
September 2002). All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 163;
AIRS Site Code: 270530963**

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	10.3	33.4	1.8	—
PM _{2.5} (Reconstructed Mass)	12.4	37.7	2.2	—
Sulfate (Calculated)	2.3	8.6	0.12	—
Sulfate (by I.C.)	2.4	8.9	0.24	0.012
Ammonium (by I.C.)	1.2	6.0	—	0.017
Sodium Ion (by I.C.)	0.27	3.7	—	0.030
Potassium (by I.C.)	0.059	1.7	—	0.014
Nitrate	2.3	16.4	0.098	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.39	1.8	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	2.6	9.9	—	0.15
Aluminum	0.028	0.67	—	0.011
Arsenic	0.002	0.012	—	0.002
Barium	—	0.15	—	0.059
Bromine	0.002	0.008	—	0.002
Cadmium	—	0.028	—	0.011
Calcium	0.071	0.46	—	0.003
Chlorine	0.008	0.17	—	0.006
Chromium	0.002	0.016	—	0.002
Copper	0.003	0.045	0.001	0.001
Iron	0.065	0.37	0.016	0.002
Lead	0.005	0.066	—	0.005
Magnesium	0.018	0.52	—	0.018
Manganese	0.002	0.009	—	0.002
Molybdenum	—	0.011	—	0.005
Nickel	0.002	0.014	0.001	0.001
Phosphorous	—	0.11	—	0.006
Potassium	0.068	1.8	0.005	0.003
Rubidium	—	0.002	—	0.002
Selenium	—	0.004	—	0.002
Silicon	0.12	1.3	—	0.008
Sodium	0.093	0.69	—	0.051
Strontium	—	0.026	—	0.003
Sulfur	0.78	2.9	0.040	0.007
Tin	—	0.034	—	0.018
Titanium	0.006	0.043	—	0.002
Vanadium	0.002	0.021	—	0.002
Zinc	0.008	0.039	0.001	0.001

TABLE 3B-2i. BOULDER, CO SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 161; AIRS Site Code: 080010006

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	9.5	26.9	2.1	—
PM _{2.5} (Reconstructed Mass)	11.3	28.2	3.7	—
Sulfate (Calculated)	1.5	5.3	0.31	—
Sulfate (by I.C.)	1.6	5.9	0.29	0.012
Ammonium (by I.C.)	0.79	4.3	0.009	0.017
Sodium Ion (by I.C.)	0.24	1.7	0.015	0.030
Potassium (by I.C.)	0.032	0.29	0.007	0.014
Nitrate	1.5	7.9	0.14	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	1.0	4.4	0.15	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	3.2	13.8	0.28	0.15
Aluminum	0.092	0.41	—	0.011
Arsenic	—	0.005	—	0.002
Barium	—	0.12	—	0.059
Bromine	0.002	0.011	—	0.002
Cadmium	—	0.016	—	0.011
Calcium	0.12	0.55	0.008	0.003
Chlorine	0.019	0.66	—	0.006
Chromium	0.002	0.009	—	0.002
Copper	0.004	0.013	0.001	0.001
Iron	0.13	0.47	0.016	0.002
Lead	0.005	0.036	—	0.005
Magnesium	0.020	0.15	—	0.018
Manganese	0.003	0.015	—	0.002
Molybdenum	—	0.010	—	0.005
Nickel	0.001	0.010	0.001	0.001
Phosphorous	0.006	0.096	—	0.006
Potassium	0.063	0.33	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	—	0.004	—	0.002
Silicon	0.28	0.97	0.027	0.008
Sodium	0.075	0.28	—	0.051
Strontium	—	0.005	—	0.003
Sulfur	0.49	1.8	0.10	0.007
Tin	—	0.033	—	0.018
Titanium	0.010	0.031	—	0.002
Vanadium	—	0.004	—	0.002
Zinc	0.023	0.53	0.001	0.001

TABLE 3B-2j. PHOENIX, AZ SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 275; AIRS Site Code: 040139997

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	9.9	32.1	3.3	—
PM _{2.5} (Reconstructed Mass)	12.0	158.0	0.8	—
Sulfate (Calculated)	1.2	2.9	0.001	—
Sulfate (by I.C.)	1.3	3.8	—	0.012
Ammonium (by I.C.)	0.5	2.6	—	0.017
Sodium Ion (by I.C.)	0.28	1.9	—	0.030
Potassium (by I.C.)	0.062	0.64	—	0.014
Nitrate	1.1	6.9	—	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.75	5.0	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	4.0	21.0	-0.66	0.15
Aluminum	0.12	0.87	—	0.011
Arsenic	0.002	0.009	—	0.002
Barium	—	0.13	—	0.059
Bromine	0.003	0.009	—	0.002
Cadmium	—	0.024	—	0.011
Calcium	0.15	0.82	0.005	0.003
Chlorine	0.047	0.40	—	0.006
Chromium	0.002	0.021	—	0.002
Copper	0.006	0.038	0.001	0.001
Iron	0.17	0.95	0.003	0.002
Lead	—	0.020	—	0.005
Magnesium	0.025	0.38	—	0.018
Manganese	0.004	0.027	—	0.002
Molybdenum	—	0.011	—	0.005
Nickel	0.003	0.13	—	0.001
Phosphorous	0.006	0.13	—	0.006
Potassium	0.11	0.69	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	—	0.005	—	0.002
Silicon	0.36	2.2	—	0.008
Sodium	0.10	0.52	—	0.051
Strontium	0.003	0.10	—	0.003
Sulfur	0.39	0.98	—	0.007
Tin	—	0.038	—	0.018
Titanium	0.013	0.058	—	0.002
Vanadium	—	0.005	—	0.002
Zinc	0.009	0.080	0.001	0.001

TABLE 3B-2k. SEATTLE, WA SUMMARY DATA (October 2001 to September 2002).
All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 314; AIRS Site Code: 530330080

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	8.2	29.1	1.9	—
PM _{2.5} (Reconstructed Mass)	8.0	28.5	1.5	—
Sulfate (Calculated)	1.4	4.9	0.057	—
Sulfate (by I.C.)	1.3	4.6	0.14	0.005
Ammonium (by I.C.)	0.47	2.0	—	0.007
Sodium Ion (by I.C.)	0.16	0.76	0.012	0.012
Potassium (by I.C.)	0.045	2.3	—	0.006
Nitrate	0.67	2.9	0.12	0.003
Volatile Nitrate	0.25	1.6	0.010	0.003
Nonvolatile Nitrate	0.41	2.5	0.020	0.003
Elemental Carbon	0.60	2.7	—	0.059
Carbonate Carbon	—	—	—	0.059
Organic Carbon	2.6	11.0	—	0.059
Aluminum	0.015	0.24	—	0.004
Arsenic	0.001	0.007	0.001	0.001
Barium	—	0.15	—	0.024
Bromine	0.002	0.014	—	0.001
Cadmium	—	0.007	—	0.004
Calcium	0.029	0.20	0.001	0.001
Chlorine	0.055	0.84	—	0.002
Chromium	0.002	0.016	—	0.001
Copper	0.003	0.045	—	0.001
Iron	0.053	0.29	0.001	0.001
Lead	0.004	0.078	—	0.002
Magnesium	0.014	0.19	—	0.007
Manganese	0.003	0.024	—	0.001
Molybdenum	0.002	0.011	—	0.002
Nickel	0.002	0.020	—	0.001
Phosphorous	—	0.028	—	0.003
Potassium	0.055	2.3	0.007	0.001
Rubidium	—	0.002	—	0.001
Selenium	0.001	0.008	0.001	0.001
Silicon	0.049	0.52	—	0.003
Sodium	0.15	0.84	—	0.021
Strontium	0.001	0.044	0.001	0.001
Sulfur	0.45	1.6	0.019	0.003
Tin	0.007	0.030	—	0.007
Titanium	0.003	0.026	—	0.001
Vanadium	0.003	0.029	—	0.001
Zinc	0.009	0.048	—	0.001

TABLE 3B-2I. SACRAMENTO, CA SUMMARY DATA (October 2001 to September 2002). All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 265; AIRS Site Code: 060670006

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	9.4	78.0	3.0	—
PM _{2.5} (Reconstructed Mass)	15.0	120.0	3.0	—
Sulfate (Calculated)	1.2	5.5	0.14	—
Sulfate (by I.C.)	1.3	4.1	0.06	0.012
Ammonium (by I.C.)	0.74	7.4	0.009	0.017
Sodium Ion (by I.C.)	0.38	3.5	—	0.030
Potassium (by I.C.)	0.077	3.6	—	0.014
Nitrate	2.3	23.4	—	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	0.66	8.4	—	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	5.2	54.1	0.18	0.15
Aluminum	0.040	0.52	—	0.011
Arsenic	0.002	0.006	—	0.002
Barium	—	0.26	—	0.059
Bromine	0.002	0.013	—	0.002
Cadmium	—	0.018	—	0.011
Calcium	0.043	0.56	0.005	0.003
Chlorine	0.052	1.8	—	0.006
Chromium	0.002	0.043	—	0.002
Copper	0.006	0.16	0.001	0.001
Iron	0.079	0.64	0.005	0.002
Lead	—	0.044	—	0.005
Magnesium	0.021	0.88	—	0.018
Manganese	0.002	0.013	—	0.002
Molybdenum	—	0.015	—	0.005
Nickel	0.010	0.61	0.001	0.001
Phosphorous	0.006	0.11	—	0.006
Potassium	0.10	3.3	—	0.003
Rubidium	—	0.003	—	0.002
Selenium	—	0.004	—	0.002
Silicon	0.12	1.1	0.011	0.008
Silver	—	0.015	—	0.010
Sodium	0.19	1.7	—	0.010
Strontium	—	0.077	—	0.051
Sulfur	0.41	1.8	0.045	0.003
Tin	—	0.035	0.009	0.007
Titanium	—	0.052	—	0.018
Vanadium	—	0.006	—	0.002
Zinc	0.005	0.11	—	0.002

TABLE 3B-2m. RIVERSIDE-RUBIDOUX, CA SUMMARY DATA (October 2001 to September 2002). All concentrations are given in $\mu\text{g}/\text{m}^3$; n = 161; AIRS Site Code: 060658001

Parameter	Mean	Max	Min	MDL
PM _{2.5} (FRM Mass)	28.6	78.2	2.6	—
PM _{2.5} (Reconstructed Mass)	30.5	79.2	3.1	—
Sulfate (Calculated)	3.6	9.6	0.23	—
Sulfate (by I.C.)	3.7	10.0	0.28	0.012
Ammonium (by I.C.)	4.8	16.8	—	0.017
Sodium Ion (by I.C.)	0.49	1.8	—	0.030
Potassium (by I.C.)	0.075	1.3	—	0.014
Nitrate	12.3	40.3	0.18	0.008
Volatile Nitrate	—	—	—	—
Nonvolatile Nitrate	—	—	—	—
Elemental Carbon	1.2	4.3	0.12	0.15
Carbonate Carbon	—	—	—	0.15
Organic Carbon	6.0	13.0	0.38	0.15
Aluminum	0.057	0.32	—	0.011
Arsenic	0.002	0.006	—	0.002
Barium	—	0.17	—	0.059
Bromine	0.006	0.016	—	0.002
Cadmium	—	0.017	—	0.011
Calcium	0.17	1.0	0.012	0.003
Chlorine	0.072	0.71	—	0.006
Chromium	0.003	0.024	—	0.002
Copper	0.006	0.037	0.001	0.001
Iron	0.17	0.58	0.019	0.002
Lead	0.006	0.026	—	0.005
Magnesium	0.032	0.31	—	0.018
Manganese	0.004	0.015	—	0.002
Molybdenum	—	0.024	—	0.005
Nickel	0.002	0.014	0.006	0.001
Phosphorous	0.007	0.067	—	0.006
Potassium	0.11	1.31	0.017	0.003
Rubidium	—	0.003	—	0.002
Selenium	—	—	—	0.006
Silicon	0.20	0.88	0.023	0.008
Silver	—	0.015	—	0.010
Sodium	0.20	0.94	—	0.051
Strontium	—	0.026	—	0.003
Sulfur	1.2	3.2	0.078	0.007
Tin	0.014	0.081	—	0.018
Titanium	0.012	0.041	—	0.002
Vanadium	0.006	0.017	—	0.002
Zinc	0.023	0.21	0.001	0.001

**TABLE 3B-3. BLANK CORRECTIONS FOR ELEMENTAL CARBON (EC),
ORGANIC CARBON (OC), AND TOTAL CARBON (TC)
IN THE SPECIATION NETWORK**

Sampler Type	Elemental Carbon		Organic Carbon		Total Carbon	
	EC Mass (µgC/filter)	EC Conc (µgC/m ³)	OC Mass (µgC/filter)	OC Conc (µgC/m ³)	TC Mass (µgC/filter)	TC Conc (µgC/m ³)
URG MASS	0.63	0.03	7.08	0.29	7.71	0.32
R & P 2300	3.21	0.22	12.93	0.9	16.13	1.12
Anderson RAAS	0.97	0.09	12.54	1.19	13.51	1.29
R & P 2025	1.67	0.07	18.42	0.77	19.91	0.83
MetOne SASS	1.03	0.11	13.75	1.42	14.78	1.53

APPENDIX 3C

Organic Composition of Particulate Matter

Although organic compounds typically constitute approximately 10 to 70% of the total dry fine particle mass in the atmosphere, organic PM concentrations, composition, and formation mechanisms are poorly understood. This is because organic particulate matter is an aggregate of hundreds of individual compounds spanning a wide range of chemical and thermodynamic properties (Saxena and Hildemann, 1996). The presence of multiphase or “semivolatile” compounds complicates collection of organic particulate matter. Furthermore, no single analytical technique currently is capable of analyzing the entire range of compounds present. Rigorous analytical methods frequently identify only 10 to 20% of the organic mass on the molecular level (Rogge et al., 1993). The data shown in Appendix 3C are meant to complement the data given for the inorganic components of particles in Appendix 6A of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996). Table 3C-1 lists a number of recent urban and some rural measurements of particulate organic and elemental carbon in μg of carbon/ m^3 ($\mu\text{g C}/\text{m}^3$). Emphasis is placed on measurements published after 1995. The analysis method and artifact correction procedure, if any, are indicated. Table 3C-2 presents information on recent (post-1990) studies concerning concentrations (in $\text{ng C}/\text{m}^3$) of particulate organic compounds found at selected U.S. sites.

**TABLE 3C-1. PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in $\mu\text{g C}/\text{m}^3$)
BASED ON STUDIES PUBLISHED AFTER 1995**

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
URBAN PM_{2.5}							
Offenberg and Baker (2000)	Chicago, IL	July 1994; Jan 1995			2.2 (3.8)	12 h	PM ₁₂ ; Imp; TOT
					1.7		PM _{1.4} ; Imp; TOT
Allen et al. (1999)	Uniontown, PA	July-Aug 1990	(0.8-8.4) ^a	(0.4-3.5) ^a 1.3 (3.1)		3 h 10 min	PM _{2.5} ; DQQ; TOR ^b Aeth
Pedersen et al. (1999)	Boston, MA	Jan-Dec 1995	5.8	1.7		24 h	PM _{2.0} ; Q; TOT
	Reading, MA (suburban)		4.0	0.7			
	Quabbin, MA (rural)		2.8	0.5			
	Rochester, NY (urban)		3.3	0.7			
	Brockport, NY (rural)		2.7	0.5			
IMPROVE (2000)	Washington, DC	1994-1998	3.4	1.1		24 h	PM _{2.5} ; QQ; TOR
	Seattle, WA		1.8	0.3			
Lewtas et al. (2001)	Seattle, WA	Apr-May 1999	8	1.4		23 h	PM _{2.5} ; DQA; EGA ^c
Khwaja (1995)	Schenectady, NY	Oct 24-26, 1991			23.2 (49.9)	6 h	PM _{1.0} ; Q; Th
Christoforou et al. (2000)	Azusa, CA	Jan-Dec 1993	9.4	1.3		24 h	PM _{2.1} ; Q; TOR
	Long Beach, CA		8.9	1.8			
	Central, LA		12.3	2.7			
	Rubidoux, LA		9.7	1.5			
	San Nicolas, LA		1.6	1.5			
Turpin and Huntzicker (1995)	Claremont, CA	Jun-Sept	na (29.4)	na (9.0)		2 h	PM _{2.5} ; Q+TQ; TOT ^d
	Long Beach, CA	Nov-Dec 1987	na (62.6)	na (24.6)		2-6 h	

**TABLE 3C-1 (cont'd). PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in $\mu\text{g C}/\text{m}^3$)
BASED ON STUDIES PUBLISHED AFTER 1995**

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
RURAL PM_{2.5}							
Klinedinst and Currie (1999)	Welby, CO Brighton, CO	Dec 1996-Jan 1997	5.6 (13.4) 3.6 (6.4)	3.3 (8.1) 1.9 (3.6)		6 h	PM _{2.5} ; Q; TOR
Andrews et al. (2000)	Look Rock, Smoky Mountains, TN	July-Aug 1995	2.2 2.7 1.2	0.4 0.1 0.2		12 h (day)	PM _{2.1} ; QQ; TOR ^e PM _{2.1} ; Q+TQ; TOR ^d PM _{1.8} ; Imp; TMO
Malm and Gebhart (1996)	Tahoma Wood, WA	June-Aug 1990	2.6 (7.4)	0.7 (2.2)		12 h	PM _{2.5} ; QQ; TOR ^f
IMPROVE (2000)	Three Sisters Wilderness, OR Rocky Mountains, CO Brigantine, NJ Acadia, MA Jefferson: James River Face Wilderness, VA Glacier, MT	1994-1998	0.9 1.0 2.0 1.2 3.8 2.4	0.2 0.2 0.5 0.2 0.7 0.4		24 h	PM _{2.5} ; QQ; TOR
Hegg et al. (1997)	150 km East of Mid-Atlantic Coast (0.02-4 km altitude)	July 1996			2.9 (5.4)		PM _{1.0} ; QQ ; EGA ^e
Cui et al. (1997)	Meadview, AZ	Aug 6-15, 1992			3	12 h	PM _{2.5} ; VDQA; EGA ^e
Chow et al. (1996)	Point Reyes, CA Altamont Pass, CA Pacheco Pass, CA Crows Landing, CA Academy, CA Button-Willow, CA Edison, CA Caliente, CA Sequoia, CA Yosemite, CA	July-Aug 1990	1.5 (2.7) 4.8 (7.2) 3.2 (6.1) 7.4 (12.7) 5.9 (8.7) 6.4 (10.6) 10.0 (12.8) 7.4 (10.7) 5.3 (7.0) 12.1 (25.8)	0.4 (0.6) 2.6 (3.9) 1.0 (1.3) 1.8 (2.5) 1.4 (2.4) 1.9 (2.7) 2.9 (4.1) 3.3 (4.4) 1.6 (3.0) 1.9 (3.5)		5-7 h	PM _{2.5} ; Q + TQ; TOR ^g

3C-3

**TABLE 3C-1 (cont'd). PARTICULATE ORGANIC AND ELEMENTAL CARBON CONCENTRATIONS (in $\mu\text{g C}/\text{m}^3$)
BASED ON STUDIES PUBLISHED AFTER 1995**

Reference	Location	Dates	OC Mean (Max)	EC Mean (Max)	TC Mean (Max)	Avg. Time	Notes
RURAL PM_{2.5}							
Malm and Day (2000)	Grand Canyon, AZ	July-Aug 1998	1.1 (1.6)	0.10 (0.3)		24 h	PM _{2.5} ; QQ; TOR ^f
PM₁₀							
Omar et al. (1999)	Bondville, IL	Jan-Dec 1994	2.6	0.2		24-48 h	PM ₁₀ ; Q; TOR
Gertler et al. (1995)	Bullhead City, AZ	Sept 1988-Oct 1989	6.0 (16.0)	1.9 (4.0)		24 h	PM ₁₀ ; Q; TOR
Chow et al. (1996)	Santa Barbara, CA (urban)	Jan-Dec 1989			8.8	24 h	PM ₁₀ ; Q; TOR
	Santa Maria, CA (urban)				4.6		
	Santa Ynez, CA (airport)				3.5		
	Gaviota, CA (rural SB)				3.4		
	Watt Road, CA (rural SB)				2.1		
	Anacapa Island, CA				3.1		
Lioy and Daisey (1987)	Newark, NJ	1982:					PM ₁₅ ; Q
		Summer	4.1	3.0			
		Winter	5.9	3.3			
	Elizabeth, NJ	Summer	2.1	1.7			
		Winter	7.1	2.3			
	Camden, NJ	Summer	2.2	1.3			
		Winter	5.2	2.0			

A limited amount of rural data is presented. In some cases, total carbon (TC = OC + EC) is reported. OC concentrations must be multiplied by the average molecular weight per carbon weight to convert to mass of particulate organic compounds. The location and dates over which sampling occurred are provided. Averaging time refers to the sampling duration. Sampling method: Q – quartz fiber filter; QQ – two quartz fiber filters in series; Q+TQ – a quartz fiber filter in one port and a Teflon followed by a quartz filter in a parallel port; Imp – cascade impactor; DQQ – denuder followed by two quartz fiber filters; DQA – denuder followed by quartz fiber filter and adsorbent; VDQA – virtual impactor inlet followed by denuder, quartz filter, and adsorbent. Analysis method is reported as follows: TOR – thermal optical reflectance; TOT – thermal optical transmittance; TMO – thermal MnO₂ oxidation; EGA – evolved gas analysis; Th – Thermal analysis; Aeth – Aethalometer. na – data not available.

^aRange is provided. It should be noted that samples were collected only during elevated pollution episodes and are not representative of average concentrations.

^bParticulate OC was considered to be the sum of front and back quartz fiber filters.

^cSum of adsorbent and filter after correction for inlet losses and denuder efficiency.

^dCorrected for adsorption by subtracting the Teflon-quartz back-up filter.

^eReported concentrations are corrected for adsorption by subtracting the quartz (TQ or QQ) back-up filter.

^fSampler contained two quartz fiber filters in series, but publication did not indicate whether the quartz back-up filter was subtracted to correct for adsorption.

^gCorrected for adsorption using Micro-Orifice Uniform Deposit Impactor (MOUDI) data from a collocated sampler.

TABLE 3C-2. PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.9}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
n-Alkanes								
n-tricosane	6.7	5.4			19.23 (57.7)			
n-tetracosane	6.4	4.7	42.3	12.7	6.04 (21.1)			
n-pentacosane	11.2	9.5	41.2	14.2	7.77 (21.3)			
n-hexacosane	8.2	4.3	29.9	10.7	2.08 (12.7)			
n-heptacosane	6.7	5.6	25.0	10.8	5.62 (15.1)			
n-octacosane	3.1	2.5	12.3	5.24	1.26 (9.0)			
n-nonacosane	7.1	4.7	33.8	23.6	7.70 (20.6)			
n-triacontane	2.7	2.5	7.39	4.27	0.76 (4.6)			
n-hentriacontane	12.6	9.6	16.1	9.66	5.24 (17.9)			
n-dotriacontane	1.5	1.5	2.61	3.50	0.41 (2.1)			
n-tritriacontane	2.1	2.3	5.02	3.31	1.49 (5.5)			
n-tetratriacontane	0.58	0.68						
Total n-alkanes	68.9	53.3	215.6	98.0	57.9			
n-Alkanoic Acids								
n-nonanoic acid	6.6	5.3						
n-decanoic acid	2.0	2.4	0.711	0.164				
n-undecanoic acid	2.8	6.0						
n-dodecanoic acid	5.3	7.0	0.905	0.803				
n-tridecanoic acid	4.3	4.9	6.17	1.78				
n-tetradecanoic acid	19.7	22.2	9.42	4.01				
n-pentadecanoic acid	5.3	6.1	33.7	5.63				
n-hexadecanoic acid (palmitic acid)	140.5	127.4	166	54.4				
n-heptadecanoic acid	4.7	5.2	13.6	3.77				
n-octadecanoic acid (stearic acid)	59.2	50.0	60.0	24.1				
n-nonadecanoic acid	1.1	1.1	10.7	2.58				
n-eicosanoic acid	5.1	6.1	41.2	10.4				
n-heneicosanoic acid	2.1	2.3	20.8	6.46				

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
n-Alkanoic Acids (cont'd)								
n-docosanoic acid	8.7	9.9	160	43.1				
n-tricosanoic acid	2.0	2.5	32.1	9.71				
n-tetracosanoic acid	11.8	16.5	205	78.0				
n-pentacosanoic acid	1.3	1.6	15.4	6.59				
n-hexacosanoic acid	5.6	9.3	174	81.3				
n-heptacosanoic acid	0.49	0.81	2.56	2.38				
n-octacosanoic acid	2.7	4.9	21.3	9.65				
n-nonacosanoic acid	0.33	0.57	1.46	2.11				
n-triacontanoic acid	1.0	2.2	4.32	5.79				
Total n-alkanoic acids	292.6	294.3	979.3	352.7				
n-Alkenoic Acids								
n-9-hexadecenoic acid			18.8	3.96				
n-9-octadecenoic acid	24.8	26.0	27.1	3.96				
n-9,12-octadecane-dienoic acid			13.6	1.83				
Total n-alkenoic acids	24.8	26.0	59.5	9.75				
n-Alkanals								
l-octanal					3.26 (14.4)			
n-nonanal	5.7	9.5	19.4	3.01	29.01 (62.8)			
n-decanal					23.58 (71.2)			
n-dodecanal					6.01 (16.4)			
n-tridecanal					6.50 (25.8)			
n-tetradecanal					9.62 (30.7)			
n-pentadecanal					12.47 (113.6)			
n-hexadecanal					17.45 (49.3)			
n-heptadecanal					24.09 (88.9)			
n-octadecanal					1.84 (11.7)			
Total n-alkanals	5.7	9.5	19.4	3.01	133.8			

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.0}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
n-Alkanols								
1-decanol					8.66 (64.1)			
1-dodecanol					21.29 (61.7)			
1-tetradecanol					13.59 (41.4)			
1-pentadecanol					4.50 (30.1)			
1-hexadecanol					27.42 (141.1)			
Total n-alkanols					75.5			
Aliphatic Dicarboxylic Acids								
oxalic acid (C ₂)						198 (360)		
malonic acid (propanedioic)	32.7	44.4				84 (107)		
methylmalonic acid (methylpropanedioic)			2.13	nd				
malonic acid (2-butenedioic)	0.66	1.3						
succinic acid (butanedioic)	66.5	51.2				102 (167)		
methylsuccinic acid (methylbutanedioic)	18	15.0	24.0	8.80				
glutaric acid (pentanedioic)	32.3	28.3	21.3	10.5				
methylglutaric acid (methylpentanedioic)	19.3	16.6						
hydroxybutanedioic acid	14.3	16.0						
adipic acid (hexanedioic)	14.1	14.1	3.39	3.07				
pimelic acid (heptanedioic)			2.22	1.03				
suberic acid (octanedioic)	3.4	4.1	4.41	13.4				
axelaic acid (nonanedioic)	29.0	22.8	19.9	8.22				
Total aliphatic dicarboxylic acids	230.3	213.8	77.4	45.0		384		

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.0}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
Ketocarboxylic Acids								
pyruvic acid (C ₃)						59 (103)		
glyoxylic acid (C ₂)						44 (68)		
<i>Total ketocarboxylic acids</i>						103		
Diterpenoid/Resin Acids								
dehydroabietic acid	23.6	22.6	98.5	8.01				
abietic acid			30.4	0.784				
13-isopropyl-5 α -podocarpa- 6,8,11,13-tetraen-16-oic acid	0.63	1.2						
8,15-pimaradien-18-oic acid	0.44	0.57	0.48	0.03				
pimaric acid	2.3	4.8	9.97	0.735				
isopimaric acid	1.3	2.3	127	7.95				
7-oxodehydroabietic acid	3.4	4.1	6.68	1.43				
abieta-6,8,11,13,15-pentaen- 18-oic acid			11.8	2.43				
abieta-8,11,13,15-tetraen- 18-oic acid			2.62	0.251				
sandaracopimaric acid	1.6	2.2	8.91	0.525				
<i>Total diterpenoid acids</i>	33.3	37.6	296.4	22.15				
Aromatic Polycarboxylic Acids								
1,2-benzene-dicarboxylic acid (phthalic acid)	60.0	55.7	9.16	6.78				
1,3-benzene-dicarboxylic acid	3.4	2.9	3.41	1.98				

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.0}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
Aromatic Polycarboxylic Acids (cont'd)								
1,4-benzene-dicarboxylic acid	2.8	1.5	5.16	4.48				
benzene tricarboxylic acids			14.4	8.77				
4-methyl-1,2-benzenedicarboxylic acid	27.8	28.8						
1,2,4-benzene-tricarboxylic acid (trimellitic acid)	0.52	0.84						
1,3,5-benzene-tricarboxylic acid (trimesic acid)	20.6	17.2						
1,2,4,5-benzene-tetracarboxylic acid (pyromellitic acid)	0.74	0.80						
Total aromatic polycarboxylic acids	115.9	107.7	32.1	22.0				
Polycyclic Aromatic Hydrocarbons								
retene	0.07	0.06	6.02	0.563				
fluoranthene	0.15	0.13	2.52	0.553				0.07 (0.26)
acephenanthrylene			0.834	0.302				0.02 (0.05)
pyrene	0.26	0.17	3.28	0.564				0.07 (0.26)
C ₁ -202 MW PAH			11.7	3.80				0.07 (0.36)
C ₂ -202 MW PAH								0.03 (0.32)
benz[<i>a</i>]anthracene	0.29	0.25	13.8	2.49				0.15 (1.09)
cyclopenta[<i>cd</i>]pyrene	0.23	0.41	1.90	0.496				0.14 (1.02)
benzo[<i>ghi</i>]-fluoranthene	0.39	0.30	6.05	1.25				0.20 (0.97)
C ₁ -226 MW PAH			10.1	1.48				0.14 (0.97)
chrysene/triphenylene	0.61	0.43	7.70	1.50				0.34 (1.62)

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.0}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
Polycyclic Aromatic Hydrocarbons (cont'd)								
C ₁ -228 MW PAH			17.6	5.35				0.34 (2.16)
C ₂ -228 MW PAH								0.09 (0.46)
benz[e]acephen-anthrylene								0.20 (1.00)
benzo[k]fluoranthene	1.15	1.20	8.69	2.13				0.22 (1.07)
benzo[b]fluoranthene	1.23	0.85	10.7	2.48				
benzo[j]fluoranthene			3.62	0.499				0.02 (0.10)
benzo[e]pyrene	0.97	0.93	7.20	1.98				0.22 (1.00)
benzo[a]pyrene	0.42	0.44	8.23	1.77				0.14 (0.80)
perylene			1.50	0.246				0.05 (0.51)
methyl-substituted 252 MW PAH								0.10 (0.88)
indeno[1,2,3- <i>cd</i>]-pyrene	0.37	0.42	6.84	2.56				0.29 (1.38)
indeno[1,2,3- <i>cd</i>]-fluoranthene	1.05	1.09	1.36	0.764				0.10 (0.46)
benzo[ghi]perylene	4.47	4.43	9.75	3.49				0.77 (4.23)
anthanthrene			0.180	0.131				
coronene								
Total polycyclic aromatic hydrocarbons	11.66	11.10	139.57	34.40				3.77
Oxygenated PAHs/ Polycyclic Aromatic Ketones/Quinones								
1,4-naphthoquinone								
1-acenaphthenone							0.26	
9-fluorenone							2.07	0.29 (1.04)
1,8-naphthalic anhydride							1.77	0.41 (1.65)
phenanthrenequinone							0.43	
phenalen-9-one								0.53 (2.23)
anthracene-9,10-dione								0.36 (1.14)
methylanthracene-9,10-dione								0.09 (0.24)
11H-benzo[a]fluoren-11-one							1.03	

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM ₁₀	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
Oxygenated PAHs/ Polycyclic Aromatic Ketones/Quinones (cont'd)								
7H-benzo[<i>c</i>]fluoren-7-one							0.37	
11H-benzo[<i>b</i>]fluoren-11-one							0.85	
1H-phenalen-1-one			7.96	0.588				
benzanthrone							1.18	
5,12-naphthacene-quinone							0.32	
7H-benz[<i>de</i>]-anthracen-7-one	0.81	0.84	7.80	1.48				
benz[<i>de</i>]anthracene-7-dione								0.20 (1.00)
benz[<i>a</i>]anthracene-7,12-dione	0.21	0.25						0.09 (0.31)
cyclopenta[<i>def</i>]phen-anthrone								0.05 (0.14)
benzo[<i>cd</i>]pyren-6-one	0.80	1.24						0.54 (2.47)
6H-benzo[<i>cd</i>]pyrene-6-one							1.34	
benzo[<i>a</i>]pyrene-6,12-dione							0.096	
Total polycyclic aromatic ketones/quinones	1.82	2.33	15.76	2.07			9.72	2.56
Steroids								
cholesterol	nd	1.9						
Substituted Phenols								
<i>p</i> -benzenediol			3.46	nd				
<i>m</i> -benzenediol			7.59	nd				
hydroxybenzaldehydes			2.64	0.604				
Total substituted phenols			13.69	0.604				

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut	Khwaja (1995) October 1991 (semiurban) no precut	Allen et al. (1997) Summer 1994 (urban) PM _{1.9}	Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO	Schenectady, NY	Kenmore Square, Boston, MA	Los Angeles Basin, CA
Guaiacol and Substituted Guaiacols								
guaiacol			0.889	0.832				
4-methylguaiacol			0.606	0.387				
trans-iso Eugenol			1.45	1.04				
vanillin			26.8	6.05				
acetovanillone			3.23	0.705				
guaiacyl acetone			10.8	4.29				
coniferyl aldehyde			47.0	nd				
Total guaiacol and substituted guaiacols			90.78	13.30				
Syringol and Substituted Syringols								
syringol			1.16	0.845				
4-methylsyringol			1.72	1.77				
4-ethylsyringol			2.28	2.39				
4-propylsyringol			0.871	nd				
4-propenylsyringol			4.38	1.40				
syringaldehyde			135	44.5				
acetosyringone			171	55.7				
acetonysyringol			406	68.1				
propionysyringol			32.1	16.2				
butyrylsyringol			15.3	6.18				
sinapyl aldehyde			15.9					
Total syringol and substituted syringols			785.7	197.1				
Sugars								
levoglucosan			7590	1100				
other sugars			1070	171				
Total sugars			8660	1271				

TABLE 3C-2 (cont'd). PARTICULATE ORGANIC COMPOUND CONCENTRATIONS (in ng C/m³) BASED ON STUDIES PUBLISHED AFTER 1990 AT SELECTED SITES

	Rogge et al. (1993) ^a Jan-Dec 1982 (annual average) PM _{2.1}		Schauer and Cass (2000) Dec 26-28, 1995 (pollution episode) PM _{2.5}		Veltkamp et al. (1996) July 24-Aug 4, 1989 no precut		Khwaja (1995) October 1991 (semiurban) no precut		Allen et al. (1997) Summer 1994 (urban) PM _{1.9}		Fraser et al. (1998) Sept 8-9, 1993 (urban)
	Los Angeles, CA	Pasadena, CA	Fresno, CA	Bakersfield, CA	Niwot Ridge, CO		Schenectady, NY		Kenmore Square, Boston, MA		Los Angeles Basin, CA
Other Compounds											
divanillyl			19.4	3.18							
divanillyl methane			2.39	nd							
vanillylmethylguaiacol			3.24	0.568							
Total other			25.0	3.75							
N-Containing Compounds											
3-methoxypyridine	0.86	1.4									
isoquinoline	1.1	1.1									
1-methoxypyridine	0.27	0.24									
1,2-dimethoxy-4-nitro-benzene	1.8	3.9									
dihydroxynitrobenzene											1.62 (10.52)
Total N-containing compounds	4.03	6.64									1.62
Total Quantified Organic Compound Mass	789	764	11410	2075	267		487		10		8
Total Organic Compound Mass			55700	18700							
Percent of Organic Mass Quantified	8-15% (a)	8-15% (a)	20%	11%			<3%				
Percent of Organic Mass Extractable and Elutable	45-60% (a)	45-60% (a)	30%	21%							

Mean values are provided with maximum concentrations in parentheses.

^a Rogge et al. (1993) summarized these percentages for all four Los Angeles Basin sampling sites (West LA, Downtown LA, Pasadena, and Rubidoux). Only Downtown LA and Pasadena data are shown here.

REFERENCES

- Allen, J. O.; Dookeran, N. M.; Taghizadeh, K.; Lafleur, A. L.; Smith, K. A.; Sarofim, A. F. (1997) Measurement of oxygenated polycyclic aromatic hydrocarbons associated with a size-segregated urban aerosol. *Environ. Sci. Technol.* 31: 2064-2070.
- Allen, G. A.; Lawrence, J.; Koutrakis, P. (1999) Field validation of a semi-continuous method for aerosol black carbon (aethalometer) and temporal patterns of summertime hourly black carbon measurements in southwestern PA. *Atmos. Environ.* 33: 817-823.
- Andrews, E.; Saxena, P.; Musarra, S.; Hildemann, L. M.; Koutrakis, P.; McMurry, P. H.; Olmez, I.; White, W. H. (2000) Concentration and composition of atmospheric aerosols from the 1995 SEAVS experiment and a review of the closure between chemical and gravimetric measurements. *J. Air Waste Manage. Assoc.* 50: 648-664.
- Chow, J. C.; Watson, J. G.; Lu, Z.; Lowenthal, D. H.; Frazier, C. A.; Solomon, P. A.; Thuillier, R. H.; Magliano, K. (1996) Descriptive analysis of PM_{2.5} and PM₁₀ at regionally representative locations during SJVAQS/AUSPEX. In: Parrish, D.; Trainer, M.; Rao, S. T.; Solomon, P. A., eds. A&WMA international specialty conference on regional photochemical measurements and modeling, part 2; November 1993; San Diego, CA. *Atmos. Environ.* 30: 2079-2112.
- Christoforou, C. S.; Salmon, L. G.; Hannigan, M. P.; Solomon, P. A.; Cass, G. R. (2000) Trends in fine particle concentration and chemical composition in southern California. *J. Air Waste Manage. Assoc.* 50: 43-53.
- Cui, W.; Machir, J.; Lewis, L.; Eatough, D. J.; Eatough, N. L. (1997) Fine particulate organic material at Meadview during the project MOHAVE summer intensive study. *J. Air Waste Manage. Assoc.* 47: 357-369.
- Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T.; Rasmussen, R. A. (1998) Air quality model evaluation data for organics. 5. C₆-C₂₂ nonpolar and semipolar aromatic compounds. *Environ. Sci. Technol.* 32: 1760-1770.
- Gertler, A. W.; Lowenthal, D. A.; Coulombe, W. G. (1995) PM₁₀ source apportionment study in Bullhead City, Arizona. *J. Air Waste Manage. Assoc.* 45: 75-82.
- Hegg, D. A.; Livingston, J.; Hobbs, P. V.; Novakov, T.; Russell, P. (1997) Chemical apportionment of aerosol column optical depth off the mid-Atlantic coast of the United States. *J. Geophys. Res.* 102: 25,293-25,303.
- IMPROVE: interagency monitoring of protected visual environments [database]. (2000) [Data on particulate organic and elemental carbon concentrations after 1995]. Fort Collins, CO: National Park Service (NPS); Cooperative Institute for Research in the Atmosphere (CIRA). Available at: <http://vista.cira.colostate.edu/improve/> [2001, January 26].
- Khwaja, H. (1995) Atmospheric concentrations of carboxylic acids and related compounds at a semiurban site. *Atmos. Environ.* 29: 127-139.
- Klinedinst, D. B.; Currie, L. A. (1999) Direct quantification of PM_{2.5} fossil and biomass carbon within the northern front range air quality study's domain. *Environ. Sci. Technol.* 33: 4146-4154.
- Lewtas, J.; Pang, Y.; Booth, D.; Reimer, S.; Eatough, D. J.; Gundel, L. A. (2001) Comparison of sampling methods for semi-volatile organic carbon associated with PM_{2.5}. *Aerosol. Sci. Technol.* 34: 9-22.
- Lioy, P. J.; Daisey, J. M. (1987) Toxic air pollution: a comprehensive study of non-criteria air pollutants. Chelsea, MI: Lewis Publishers.
- Malm, W. C.; Day, D. E. (2000) Optical properties of aerosols at Grand Canyon National Park. *Atmos. Environ.* 34: 3373-3391.
- Malm, W. C.; Gebhart, K. A. (1996) Source apportionment of organic and light-absorbing carbon using receptor modeling techniques. *Atmos. Environ.* 30: 843-855.
- Offenberg, J. H.; Baker, J. E. (2000) Aerosol size distributions of elemental and organic carbon in urban and over-water atmospheres. *Atmos. Environ.* 34: 1509-1517.
- Omar, A. H.; Biegalski, S.; Larson, S. M.; Landsberger, S. (1999) Particulate contributions to light extinction and local forcing at a rural Illinois site. *Atmos. Environ.* 33: 2637-2646.
- Pedersen, D. U.; Durant, J. L.; Penman, B. W.; Crespi, C. L.; Hemond, H. F.; Lafleur, A. L.; Cass, G. R. (1999) Seasonal and spatial variations in human cell mutagenicity of respirable airborne particles in the northeastern United States. *Environ. Sci. Technol.* 33: 4407-4415.
- Rogge, W. F.; Mazurek, M. A.; Hildemann, L. M.; Cass, G. R.; Simoneit, B. R. T. (1993) Quantification of urban organic aerosols at a molecular level: identification, abundance and seasonal variation. *Atmos. Environ. Part A* 27: 1309-1330.
- Saxena, P.; Hildemann, L. M. (1996) Water-soluble organics in atmospheric particles: a critical review of the literature and applications of thermodynamics to identify candidate compounds. *J. Atmos. Chem.* 24: 57-109.

- Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environ. Sci. Technol.* 34: 1821-1832.
- Turpin, B. J.; Huntzicker, J. J. (1995) Identification of secondary organic aerosol episodes and quantitation of primary and secondary organic aerosol concentrations during SCAQS. *Atmos. Environ.* 29: 3527-3544.
- U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- Veltkamp, P. R.; Hansen, K. J.; Barkley, R. M.; Sievers, R. E. (1996) Principal component analysis of summertime organic aerosols at Niwot Ridge, Colorado. *J. Geophys. Res. [Atmos.]* 101: 19,495-19,504.

APPENDIX 3D

Composition of Particulate Matter Source Emissions

This appendix includes discussions of the elemental composition of emissions from various source categories discussed in Table 3-8. Discussions in this appendix incorporate material dealing with the inorganic components of source emissions from Chapter 5 of the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996), provides updates to that material, and adds material describing the composition of organic components in source emissions. Primary emphasis is placed in the discussions on the composition of PM_{2.5} sources.

Soil and Fugitive Dust

The compositions of soils and average crustal material are shown in Table 3D-1 (adapted from Warneck, 1988). Two entries are shown as representations of average crustal material. Differences from the mean soil composition shown can result from local geology and climate. Major elements in both soil and crustal profiles are Si, Al, and Fe, which are found in the form of various minerals. In addition, organic matter constitutes a few percent, on average, of soils. In general, the soil profile is similar to the crustal profiles, except for the depletion of soluble elements such as Ca, Mg, Na, and K. It should be noted that the composition of soils from specific locations can vary considerably from these global averages, especially for elements like Ca, Mg, Na, and K.

Fugitive dust emissions arise from paved and unpaved roads, building construction and demolition, parking lots, mining operations, storage piles, feed lots, grain handling, and agricultural tilling in addition to wind erosion. Figure 3D-1 shows examples of size distributions in dust from paved and unpaved roads, agricultural soil, sand and gravel, and alkaline lake bed sediments, which were measured in a laboratory resuspension chamber as part of a study in California (Chow et al., 1994). This figure shows substantial variation in particle size among some of these fugitive dust sources. The PM_{1.0} abundance (6.9%) in the total suspended PM (TSP) from alkaline lake bed dust is twice its abundance in paved and unpaved road dust. Approximately 10% of the TSP is in the PM_{2.5} fraction and ~50% of TSP is in the PM₁₀ fraction.

TABLE 3D-1. AVERAGE ABUNDANCES OF MAJOR ELEMENTS IN SOIL AND CRUSTAL ROCK

Elemental Abundances (ppm)			
Element	Soil (1)	Crustal Rock	
		(2)	(3)
Si	330,000	277,200	311,000
Al	71,300	81,300	77,400
Fe	38,000	50,000	34,300
Ca	13,700	36,300	25,700
Mg	6,300	20,900	33,000
Na	6,300	28,300	31,900
K	13,600	25,900	29,500
Ti	4,600	4,400	4,400
Mn	850	950	670
Cr	200	100	48
V	100	135	98
Co	8	25	12

Source: (1) Vinogradov (1959); (2) Mason (1966); (3) Turekian (1971), Model A; as quoted in Warneck (1988).




Figure 3D-1. Size distribution of particles generated in a laboratory resuspension chamber.

Source: Chow et al. (1994).

The sand/gravel dust sample shows that 65% of the mass is in particles larger than the PM₁₀ fraction. The PM_{2.5} fraction of TSP is ~30 to 40% higher in alkaline lake beds and sand/gravel than in the other soil types. The tests were performed after sieving and with a short (< 1 min) waiting period prior to sampling. It is expected that the fraction of PM_{1.0} and PM_{2.5} would increase with distance from a fugitive dust emitter, as the larger particles deposit to the surface faster than do the smaller particles.

The size distribution of samples of paved road dust obtained from a source characterization study in California is shown in Figure 3D-2. As might be expected, most of the emissions are in the coarse size mode. The chemical composition of paved road dust obtained in Denver, CO during the winter of 1987-1988 is shown in Figure 3D-3. The chemical composition of paved road dust consists of a complex mixture of PM from a wide variety of sources. Hopke et al. (1980) found that the inorganic composition of urban roadway dust in samples from Urbana, IL could be described in terms of contributions from natural soil, automobile exhaust, rust, tire wear, and salt. Automobile contributions arose from: exhaust emissions enriched in Pb; Fe as rust; tire wear particles enriched in Zn; brake linings enriched in Cr, Ba, and Mn; and cement particles derived from roadways by abrasion. In addition to organic compounds from combustion and secondary sources, road dust also contains biological material such as pollen and fungal spores.

Very limited data exist for characterizing the composition in organic compounds found in resuspended paved road dust and soil dust. The only reported measurements are from Rogge et al. (1993a) and Schauer and Cass (2000), which consist of data for the fine particle fraction. The resuspended road dust sample analyzed by Rogge et al. (1993a) was collected in Pasadena, CA during May of 1988. The sample analyzed by Schauer and Cass (2000) is a composite sample collected at several sites in the Central Valley of California in 1995. In both cases, road dust samples were resuspended in the laboratory. Samples were drawn through a PM_{2.0} cyclone upstream of the collection substrate to remove particles with aerodynamic diameters > 2.0 μm. It is unclear if these samples are representative of road dust in other locations of the United States. Table 3D-2 summarizes the organic compounds measured in these road dust samples.




Figure 3D-2. Size distribution of California source emissions, 1986.

Source: Houck et al. (1989, 1990).




Figure 3D-3. Chemical abundances for PM_{2.5} emissions from paved road dust in Denver, CO. Solid bars represent fractional abundances, and the error bars represent variability in species abundances. Error bars represent detection limits when there are no solid bars.

Source: Watson and Chow (1994).

TABLE 3D-2. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS PRESENT IN FINE PARTICLE ROAD DUST SAMPLE

Source	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Pasadena Road Dust (Rogge et al., 1993a)	n-Alkanes	0.13	C ₁₇ , C ₁₉ , C ₂₁
	n-Alkanoic acids	0.37	Palmitic acid and stearic acid
	n-Alkenoic acids	0.028	Oleic acid and linoleic Acid
	Petroleum biomarkers	0.017	Hopanes and steranes
	PAH	0.0059	No dominant compounds
	n-Alkanals	0.046	Octacosanol and triacontanal
	n-Alkanols	0.021	Hexacosanol and octacosanol
San Joaquin Valley Road Dust (Schauer and Cass, 2000)	n-Alkanes	0.023	No dominant compounds
	n-Alkanoic acids	0.23	Palmitic acid and stearic acid
	n-Alkenoic acids	0.095	Oleic acid, linoleic acid, and hexadecenoic acid

Stationary Sources

The elemental composition of primary PM emitted in the fine fraction from a variety of power plants and industries in the Philadelphia area is shown in Table 3D-3 as a representative example of emissions from stationary fossil-fuel combustion sources (Olmez et al., 1988). Entries for the coal fired power plant show that Si and Al, followed by sulfate, are the major primary constituents produced by coal combustion, whereas fractional abundances of elemental carbon (EC) were much lower and organic carbon (OC) species were not detected. Sulfate is the major particulate constituent released by the oil-fired power plants examined in this study; and, again, EC and OC are not among the major species emitted. Olmez et al. (1988) also compared their results to a number of similar studies and concluded that their data should have much wider applicability to receptor model studies in other areas with some of the same source types. The high temperature of combustion in power plants results in the almost complete oxidation of the carbon in the fuel to CO₂ and very small amounts of CO. Combustion conditions in smaller boilers and furnaces allow the emission of unburned carbon and sulfur in more reduced forms such as thiophenes and inorganic sulfides. A number of trace elements are greatly enriched over crustal abundances in different fuels, such as Se in coal and V, Zn, and Ni in oil. In fact, the higher V content of the fuel oil than in coal could help account for the higher sulfate seen in the profiles from the oil-fired power plant compared to the coal-fired power plant, because V at combustion temperatures found in power plants is known to catalyze the oxidation of reduced sulfur species. During combustion at lower temperatures, the emission of reduced sulfur species also occurs. For example, Huffman et al. (2000) identified sulfur species emitted by the combustion of several residual fuel oils (RFO) in a fire tube package boiler that is meant to simulate conditions in small institutional and industrial boilers. They found that sulfur was emitted not only as sulfate (26 to 84%), but as thiophenes (13 to 39%) with smaller amounts of sulfides and elemental S. They also found that Ni, V, Fe, Cu, Zn, and Pb are present mainly as sulfates in emissions. Linak et al. (2000) found, when burning RFO, that the fire tube package boiler produced particles with a bimodal size distribution in which ~0.2% of the mass was associated with particles < 0.1 μm AD, with the rest of the mass lying between 0.5 and 100 μm. Miller et al. (1998) found that larger particles consisted mainly of cenospheric carbon; whereas trace metals and sulfates were found concentrated in the smaller particles in a fire tube package boiler. In contrast, when RFO was burning in a refractory-lined combustor that is meant to

**TABLE 3D-3. COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES
IN THE PHILADELPHIA AREA**

Species (Units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary Al Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
C-v (%)	ND		2.7 ± 1.2	3	0.75 ± 0.63	4	1.6 ± 1.5	2	ND		0.57 ± 0.26	4
C-e (%)	0.89 ± 0.12	3	7.7 ± 1.5	3	0.22 ± 0.17	4	0.18 ± 0.10	2	0.16 ± 0.05	3	3.5 ± 0.2	4
NH ₄ (%)	1.89 ± 0.19	3	3.5 ± 1.6	3	3.7 ± 1.7	4	2.2 ± 0.9	2	0.43 ± 0.22	3	0.36 ± 0.07	4
Na (%)	0.31 ± 0.03	3	3.0 ± 0.8	3	3.3 ± 0.8	3	16.3 ± 0.8	1	0.38 ± 0.05	3	6.6 ± 3.5	3
Al (%)	14 ± 2	3	0.45 ± 0.09	3	0.94 ± 0.08	3	1.74 ± 0.09	1	6.8 ± 1.2	3	0.25 ± 0.10	3
Si (%)	21.8 ± 1.6	9	1.9 ± 0.6	9	2.6 ± 0.4	11	3.1 ± 2.2	2	9.8 ± 20.0	9	1.7 ± 0.3	10
P (%)	0.62 ± 0.10	9	1.5 ± 0.4	9	1.0 ± 0.2	11	0.45 ± 0.27	2	ND		0.63 ± 0.12	10
S (%)	3.4 ± 0.6	9	11 ± 2	9	13 ± 1	11	3 ± 4	2	4.2 ± 12.6	9	2.9 ± 0.8	10
SO ₄ (%)	11.9 ± 1.2	3	40 ± 4	3	45 ± 7	4	5.9 ± 2	2	38 ± 4	3	6.8 ± 2.3	4
Cl (%)	0.022 ± 0.11	3	0.019 ± 0.009	2	ND		21 ± 4	1	ND		29 ± 5	3
K (%)	1.20 ± 0.09	9	0.16 ± 0.05	9	0.21 ± 0.03	11	10.9 ± 1.5	2	0.031 ± 0.005	9	7.6 ± 2.3	10
Ca (%)	1.4 ± 0.5	3	3.6 ± 1.0	3	2.3 ± 1.0	3	0.12 ± 0.09	2	0.030 ± 0.004	9	0.23 ± 0.10	10
Sc (ppm)	42 ± 2	3	0.17 ± 0.02	3	0.47 ± 0.02	3	0.092 ± 0.039	1	2.7 ± 0.4	3	0.11 ± 0.02	1
Ti (%)	1.1 ± 0.2	3	0.040 ± 0.044	9	0.12 ± 0.02	11	0.024 ± 0.003	2	0.38 ± 0.1	3	0.030 ± 0.015	10
V (ppm)	550 ± 170	3	11,500 ± 3000	3	20,000 ± 3000	3	36 ± 7	1	250 ± 70	3	8.6 ± 5.3	2
Cr (ppm)	390 ± 120	3	235 ± 10	3	230 ± 70	3	410 ± 20	1	59 ± 8	3	99 ± 31	3
Mn (ppm)	290 ± 15	3	380 ± 40	3	210 ± 50	3	120 ± 15	1	14 ± 3	3	165 ± 40	3
Fe (%)	7.6 ± 0.4	3	1.6 ± 0.2	3	1.7 ± 0.4	3	0.31 ± 0.02	1	0.20 ± 0.03	9	0.22 ± 0.05	3
Co (ppm)	93 ± 10	3	790 ± 150	3	1100 ± 200	3	13 ± 2	1	15 ± 2	3	3.7 ± 0.8	3
Ni (ppm)	380 ± 50	9	15,000 ± 5000	9	19,000 ± 2000	11	300 ± 100	2	220 ± 30	9	290 ± 40	10

3D-7

TABLE 3D-3 (cont'd). COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES IN THE PHILADELPHIA AREA

Species (units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary Al Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
Cu (ppm)	290 ± 20	9	980 ± 320	9	1100 ± 500	11	450 ± 200	2	14 ± 8	9	1300 ± 500	3
Zn (%)	0.041 ± 0.005	3	1.3 ± 0.3	3	0.78 ± 0.30	3	0.079 ± 0.006	1	0.0026 ± 0.0007	3	10.4 ± 0.5	3
As (ppm)	640 ± 80	3	33 ± 6	1	50 ± 16	3	15 ± 6	1	ND		64 ± 34	3
Se (ppm)	250 ± 20	3	26 ± 9	3	23 ± 7	3	66 ± 3	1	15 ± 1	3	42 ± 16	3
Br (ppm)	35 ± 8	3	90 ± 60	9	45 ± 17	11	630 ± 70	2	5.6 ± 1.8	9	2300 ± 800	10
Rb (ppm)	190 ± 80	1	ND		ND		97 ± 38	1	ND		230 ± 50	2
Sr (ppm)	1290 ± 60	9	160 ± 50	9	280 ± 70	11	ND		36 ± 6	9	87 ± 14	10
Zr (ppm)	490 ± 190	9	140 ± 180	9	100 ± 120	11	ND		130 ± 50	2	ND	
Mo (ppm)	170 ± 60	2	930 ± 210	3	1500 ± 300	3	ND		ND		240 ± 130	10
Ag (ppm)	ND		ND		ND		ND		ND		71 ± 15	3
Cd (ppm)	ND		ND		ND		ND		ND		1200 ± 700	3
In (ppm)	0.71 ± 0.04	2	ND		ND		ND		ND		4.9 ± 1.4	3
Sn (ppm)	ND		320 ± 230	9	200 ± 80	11	550 ± 540	2	ND		6700 ± 1900	10
Sb (ppm)	^a		370 ± 410	3	1020 ± 90	3	6100 ± 300	1	7.7 ± 1.5	3	1300 ± 1000	3
Cs (ppm)	9.2 ± 0.9	2	ND		ND		ND		ND		5.9 ± 3.0	3
Ba (ppm)	ND		1960 ± 100	3	2000 ± 500	3	ND		290 ± 90	2	ND	
La (ppm)	120 ± 10	3	130 ± 30	3	450 ± 30	3	19 ± 2	1	3300 ± 500	3	1.1 ± 0.5	1
Ce (ppm)	180 ± 10	2	89 ± 23	3	360 ± 20	3	ND		2700 ± 400	3	ND	
Nd (ppm)	80 ± 26	3	28 ± 5	2	230 ± 20	3	ND		1800 ± 250	3	ND	
Sm (ppm)	23 ± 2	3	3.7 ± 0.7	3	20.5 ± 1.5	3	ND		170 ± 20	3	ND	

3D-8

TABLE 3D-3 (cont'd). COMPOSITION OF FINE PARTICLES RELEASED BY VARIOUS STATIONARY SOURCES IN THE PHILADELPHIA AREA

Species (units)	Eddystone Coal-Fired Power Plant	N	Oil-Fired Power Plants				Secondary AI Plant	N	Fluid Cat. Cracker	N	Municipal Incinerator	N
			Eddystone	N	Schuylkill	N						
Eu (ppm)	5.1 ± 0.5	3	ND		0.65 ± 0.23	3	ND		4.9 ± 0.7	3	ND	
Gd (ppm)	ND		ND		ND		ND		71 ± 10	3	ND	
Tb (ppm)	3.3 ± 0.3	3	ND		0.90 ± 0.29	3	ND		8.9 ± 1.3	3	ND	
Yb (ppm)	10.3 ± 0.5	1	ND		ND		ND		3.7 ± 0.4	3	ND	
Lu (ppm)	ND		ND		ND		ND		0.59 ± 0.17	3	ND	
Hf (ppm)	5.8 ± 0.8	3	0.39 ± 0.07	1	ND		ND		0.99 ± 0.08	3	ND	
Ta (ppm)	ND		ND		ND		ND		0.56 ± 0.10	3	ND	
W (ppm)	20 ± 8	1	60 ± 5	2	ND		ND		ND		ND	
Au (ppm)	ND		0.054 ± 0.017	2	ND		ND		ND		0.56 ± 0.27	3
Pb (%)	0.041 ± 0.004	9	1.8 ± 0.6	9	1.0 ± 0.2	11	0.081 ± 0.014	2	0.0091 ± 0.0021	9	5.8 ± 1.2	10
Th (ppm)	24 ± 2	3	1.9 ± 0.5	2	ND		ND		6.2 ± 0.7	3	ND	
% mass	24 ± 2	6	93.5 ± 2.5	6	96 ± 2	6	81 ± 10	2	97 ± 2	7	89 ± 2	7

^a Omitted because of sample contamination.

N = Number of samples.

ND = Not detected.

The “% mass” entries give the average percentage of the total emitted mass found in the fine fraction.

1 ppm = 10⁻⁴%

Source: Adapted from Olmez et al. (1988).

3D-9

simulate combustion conditions in a large utility residual oil fired boiler, Linak et al. (2000) found that particles were distributed essentially unimodally, with a mean diameter of about 0.1 μm .

Apart from emissions in the combustion of fossil fuels, trace elements are emitted as the result of various industrial processes such as steel and iron manufacturing and nonferrous metal production (e.g., for Pb, Cu, Ni, Zn, and Cd). As may be expected, emissions factors for the various trace elements are highly source-specific (Nriagu and Pacyna, 1988). Inspection of Table 3D-3 reveals that the emissions from the catalytic cracker and the oil-fired power plant are greatly enriched in rare-earth elements (e.g., La) compared to other sources.

Emissions from municipal waste incinerators are heavily enriched in Cl, arising mainly from the combustion of plastics and metals that form volatile chlorides. The metals can originate from cans or other metallic objects, and some metals such as Zn and Cd are also used as additives in plastics or rubber. Many elements such as S, Cl, Zn, Br, Ag, Cd, Sn, In, Sb, and Pb are enormously enriched compared to their crustal abundances. A comparison of the trace elemental composition of incinerator emissions in Philadelphia, PA (shown in Table 3D-3) with those in Washington, DC and Chicago, IL (Olmez et al., 1988) shows agreement for most constituents to within less than a factor of two.

Very limited data exist for characterizing the chemical composition of organic compounds present in particulate emissions from industrial-scale stationary fuel combustion. Oros and Simoneit (2000) have reported on the abundance and distribution of organic constituents in coal smokes burned under laboratory conditions. Their work provides the basis for further investigation addressing the emissions of coal fired boilers.

Rogge et al. (1997a) measured the composition of the organic constituents in the PM emissions from a 50 billion kJ/h boiler that was operating at 60% capacity and was burning Number 2 distillate fuel oil. The fine carbon PM emissions from this boiler across five tests were composed of an average of 14% OC and 86% EC (Hildemann et al., 1991). Significant variability in the distribution of organic compounds present in the emissions from two separate tests was observed. Most of the identified organic mass consisted of n-alkanonic acids, aromatic acids, n-alkanes, PAH, oxygenated PAH, and chlorinated compounds. It is unclear if these emissions are representative of typical fuel oil combustion units in the United States. Rogge

et al. (1997b) measured the composition of hot asphalt roofing tar pots, and Rogge et al. (1993b) measured the composition of emissions from home appliances that use natural gas.

Motor Vehicles

Particulate emissions in exhaust from gasoline- and diesel-powered vehicles have changed significantly over the past 25 years (Sawyer and Johnson, 1995; Cadle et al., 1999). These changes have resulted from reformulation of fuels, the wide application of exhaust-gas treatment in gasoline-powered motor vehicles, and changes in engine design and operation. Because of these evolving tailpipe emissions, along with the wide variability of emissions between vehicles of the same class (Hildemann et al., 1991; Cadle et al., 1997; Sagebiel et al., 1997; Yanowitz et al., 2000), well-defined average emissions profiles for the major classes of motor vehicles have not been established. Two sampling strategies have been employed to obtain motor vehicle emissions profiles: (1) the measurement of exhaust emissions from vehicles operating on dynamometers and (2) the measurement of integrated emissions of motor vehicles driving through roadway tunnels. Dynamometer testing can be used to measure vehicle emissions operating over an integrated driving cycle and allows the measurement of emissions from individual vehicles. However, dynamometer testing requires considerable resources and usually precludes testing a very large number of vehicles. In contrast, a large number of vehicles can be readily sampled in tunnels; however, vehicles driving through tunnels operate over limited driving conditions, and the measurements represent contributions from a large number of vehicle types. As a result, except in a few cases, tunnel tests have not been effective at developing chemically speciated PM emissions profiles for individual motor vehicle classes. Rather, several studies have measured the contribution of both OC and EC to the PM emissions from different classes of motor vehicles operating on chassis dynamometers.

The principal PM components emitted by diesel- and gasoline-fueled vehicles are OC and EC as shown in Tables 3D-4a and 4b. As can be seen, the variability among entries for an individual fuel type is large and overlaps that found between different fuel types. On average, the abundance of EC is larger than that of OC in the exhaust of diesel vehicles, whereas OC is the dominant species in the exhaust of gasoline-fueled vehicles. Per vehicle mile, total carbon emissions from light and heavy duty diesel vehicles can range from 1 to 2 orders of magnitude higher than those from gasoline vehicles.

TABLE 3D-4a. ORGANIC AND ELEMENTAL CARBON FRACTIONS OF DIESEL AND GASOLINE ENGINE PARTICULATE MATTER EXHAUST

Engine Type	Organic Carbon	Elemental Carbon
Heavy-duty diesel ^a	19 ± 8%	75 ± 10%
Heavy-duty diesel (SPECIATE) ^b	21 - 36%	52 - 54%
Light-duty diesel ^c	30 ± 9%	61 ± 16%
Light-duty diesel (SPECIATE) ^b	22 - 43%	51 - 64%
Gasoline (hot stabilized) ^a	56 ± 11%	25 ± 15%
Gasoline (“smoker” and “high emitter”) ^{a,c}	76 ± 10%	7 ± 6%
Gasoline (cold start) ^a	46 ± 14%	42 ± 14%

^a Fujita et al. (1998) and Watson et al. (1998).

^b SPECIATE database (U.S. Environmental Protection Agency, 1999).

^c Norbeck et al. (1998).

Source: U.S. Environmental Protection Agency (2002).

As might be expected, most of the PM emitted by motor vehicles is in the PM_{2.5} size range. Particles in diesel exhaust are typically trimodal (consisting of a nuclei mode, an accumulation mode, and a coarse mode) and are log-normal in form (Kittelson, 1998). More than 90% of the total number of particles are in the nuclei mode, which contains only about 1 to 20% of the particle mass with a mass median diameter of about 0.02 μm, whereas the accumulation mode (with a mass median diameter of about 0.25 μm) contains most of the mass with a smaller fraction (5 to 20%) contained in the coarse mode. Kerminin et al. (1997), Bagley et al. (1998), and Kleeman et al. (2000) also have shown that gasoline- and diesel-fueled vehicles produce particles that are mostly < 2.0 μm in diameter. Cadle et al. (1999) found that 91% of PM emitted by in-use gasoline vehicles in the Denver area was in the PM_{2.5} size range, which increased to 97% for “smokers” (i.e., light-duty vehicles with visible smoke emitted from their tailpipes) and 98% for light-duty diesels. Durbin et al. (1999) found that about 92% of the PM was < 2.5 μm for smokers and diesels. The mass median diameter of the PM emitted by the gasoline vehicles sampled by Cadle et al. (1999) was ~0.12 μm and increased to ~0.18 μm for smokers and diesels. Corresponding average emissions rates of PM_{2.5} found by Cadle et al. (1999) were 552 mg/mile for diesels; 222 mg/mile for gasoline smokers; and 38 mg/mile for other gasoline

TABLE 3D-4b. CONTRIBUTION OF ORGANIC CARBON TO PARTICULATE MATTER CARBON EMISSIONS IN MOTOR VEHICLE EXHAUST COLLECTED FROM VEHICLES OPERATED ON CHASSIS DYNAMOMETERS

	Year of Tests	Test Cycle	Number of Vehicles	OC % of Total Carbon	Notes
GASOLINE POWERED VEHICLES					
Light-duty vehicles	1996-97	FTP	195 ^a	70	A
High-CO/VOC-emitting smokers	1994	IM-240	7	91	B
High-CO/VOC-emitting nonsmokers	1994	IM-240	15	76	B
Catalyst-equipped vehicles	Mid-1980s	FTP	7	69	C
Noncatalyst vehicles	Mid-1980s	FTP	6	89	C
DIESEL VEHICLES					
Light-duty diesel vehicles	1996-1997	FTP	195 ^a	40	A
Medium-duty diesel vehicles	1996	FTP	2	50 ^b	D
Heavy-duty diesel vehicles	1992	^c	6	42	E
Heavy-duty diesel vehicles	Mid-1980s	^c	2	45	C

Notes:

- A. From Cadle et al. (1999). Average of summer and winter cold start emissions.
- B. From Sagebiel et al. (1997). Hot start testing of vehicles identified as either high emitters of carbon monoxide or volatile organic compounds (VOCs).
- C. From Hildemann et al. (1991). Cold start tests.
- D. From Schauer et al. (1999). Hot start tests of medium duty vehicles operating on an FTP cycle.
- E. From Lowenthal et al. (1994). Only includes measurement of vehicles powered by diesel fuel operated without an exhaust particulate trap.

^a A total of 195 light-duty vehicles were tested that include both gasoline- and diesel-powered vehicles.

^b Fraction of PM consisting of OC was measured with and without an organics denuder upstream of particulate filter. Results reported here represent measurement without an organics denuder for consistency with other measurements. Using an organics denuder, the OC comprised 39% of the PM carbon.

^c Driving cycle comprised of multiple idle, steady acceleration, constant speed, deceleration steps (see reference for more details).

vehicles. The values for gasoline smokers and for diesels appear to be somewhat lower than those given in Table 3D-5, whereas the value for other gasoline vehicles falls in the range given for low and medium gasoline vehicle emissions.

Examples of data for the trace elemental composition of the emissions from a number of vehicle classes obtained December 1997 in Colorado, as part of the North Frontal Range Air

TABLE 3D-5. EMISSION RATES (mg/mi) FOR CONSTITUENTS OF PARTICULATE MATTER FROM GASOLINE AND DIESEL VEHICLES

	Gasoline Vehicles				Diesel Vehicles	
	Low	Medium	High	Smoker	Light Duty	Heavy Duty
TC	9.07 ± 0.75	41.30 ± 1.68	207.44 ± 7.29	456.38 ± 16.80	373.43 ± 13.75	1570.69 ± 58.24
OC	6.35 ± 0.54	26.02 ± 1.31	95.25 ± 4.28	350.24 ± 15.27	132.01 ± 5.82	253.94 ± 16.12
EC	2.72 ± 0.52	15.28 ± 0.99	112.19 ± 5.82	106.14 ± 5.42	241.42 ± 12.11	1316.75 ± 55.33
NO ₃ ⁻	0.039 ± 0.027	0.057 ± 0.028	0.141 ± 0.031	0.964 ± 0.051	1.474 ± 0.071	1.833 ± 1.285
SO ₄ ²⁻	0.158 ± 0.036	0.518 ± 0.043	0.651 ± 0.052	2.160 ± 0.137	2.902 ± 0.165	3.830 ± 1.286
Na	0.060 ± 0.063	0.023 ± 0.111	0.052 ± 0.092	0.000 ± 0.000	0.000 ± 0.000	1.288 ± 2.160
Mg	0.036 ± 0.022	0.068 ± 0.027	0.041 ± 0.033	0.000 ± 0.000	0.000 ± 0.000	1.061 ± 0.729
Al	0.083 ± 0.016	0.078 ± 0.016	0.057 ± 0.014	0.000 ± 0.000	0.000 ± 0.000	0.321 ± 0.543
Si	0.066 ± 0.008	0.279 ± 0.011	0.714 ± 0.012	0.000 ± 0.000	0.000 ± 0.000	8.018 ± 0.221
P	0.035 ± 0.004	0.152 ± 0.007	0.113 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.407 ± 0.136
S	0.085 ± 0.006	0.442 ± 0.009	0.822 ± 0.022	2.515 ± 0.116	2.458 ± 0.124	3.717 ± 0.111
Cl	0.024 ± 0.012	0.038 ± 0.012	0.081 ± 0.020	0.140 ± 0.117	0.228 ± 0.114	0.881 ± 0.221
K	0.010 ± 0.009	0.019 ± 0.009	0.031 ± 0.035	0.033 ± 0.386	0.000 ± 0.426	0.064 ± 0.248
Ca	0.060 ± 0.010	0.212 ± 0.011	0.210 ± 0.030	0.362 ± 0.250	0.150 ± 0.304	0.716 ± 0.107
Fe	0.143 ± 0.004	0.756 ± 0.005	1.047 ± 0.010	2.438 ± 0.054	0.515 ± 0.057	0.376 ± 0.055
Ni	0.001 ± 0.004	0.005 ± 0.004	0.011 ± 0.005	0.008 ± 0.017	0.014 ± 0.018	0.002 ± 0.057
Cu	0.002 ± 0.004	0.016 ± 0.003	0.021 ± 0.005	0.071 ± 0.018	0.024 ± 0.021	0.001 ± 0.062
Zn	0.048 ± 0.003	0.251 ± 0.004	0.265 ± 0.023	0.188 ± 0.272	0.000 ± 0.299	0.707 ± 0.032
Br	0.001 ± 0.002	0.016 ± 0.002	0.079 ± 0.003	0.047 ± 0.012	0.003 ± 0.014	0.012 ± 0.050
Ba	0.013 ± 0.136	0.009 ± 0.138	0.011 ± 0.299	0.380 ± 2.175	0.428 ± 2.390	0.493 ± 3.108
Pb	0.007 ± 0.006	0.085 ± 0.005	0.255 ± 0.008	0.345 ± 0.032	0.153 ± 0.033	0.008 ± 0.154

Source: Lawson and Smith (1998).

Quality Study (NFRAQS), are shown in Table 3D-5. As can be seen from Table 3D-5, emissions of total carbon (TC), which is equal to the sum of organic carbon (OC) and elemental carbon (EC), from gasoline vehicles are highly variable. Gillies and Gertler (2000) point out that there is greater variability in the concentrations of trace elements and ionic species than for OC

and EC among different source profiles, e.g., SPECIATE (U.S. Environmental Protection Agency, 1999; Lawson and Smith, 1998; Norbeck et al., 1998). They suggest that this may arise because emissions of trace elements are not related only to the combustion process, but also to their abundances in different fuels and lubricants and to wear and tear during vehicle operation. Emissions from gasoline smokers are comparable to those from light-duty diesel vehicles. Thus, older, poorly maintained gasoline vehicles could be significant sources of PM_{2.5} (Sagebiel et al., 1997; Lawson and Smith, 1998), in addition to being significant sources of gaseous pollutants (e.g., Calvert et al., 1993). Durbin et al. (1999) point out that although smokers constitute only 1.1 to 1.7% of the light-duty fleet in the South Coast Air Quality Management District in California, they contribute roughly 20% of the total PM emissions from the light-duty fleet. In general, motor vehicles that are high emitters of hydrocarbons and CO also will tend to be high emitters of PM (Sagebiel et al., 1997; Cadle et al., 1997). Particle emission rates, even in newer vehicles, also are correlated with vehicle acceleration; and emissions occur predominantly during periods of heavy acceleration (Maricq et al., 1999).

Although the data shown in Table 3D-5 indicate that sulfur (mainly in the form of sulfate) is a minor component of PM_{2.5} emissions, sulfur (S) may be the major component of the ultrafine particles emitted by either diesel or internal combustion engines (Gertler et al., 2000). It is not clear what the source of the small amount of Pb seen in the auto exhaust profile is. It is very difficult to find suitable tracers for automotive exhaust, because Pb has been removed from gasoline.

Several tunnel studies have measured the distribution of OC and EC in the integrated exhaust of motor vehicle fleets comprising several classes of motor vehicles (Pierson and Brachaczek, 1983; Weingartner et al., 1997a; Fraser et al., 1998a). The study by Fraser et al. (1998a) found that OC constituted 46% of the carbonaceous PM emissions from the vehicles operating in the Van Nuys tunnel in Southern California in the Summer of 1993. Although diesel vehicles constituted only 2.8% of the vehicles measured by Fraser et al. (1998a), the contribution of OC to the total particulate carbon emissions obtained in the Van Nuys tunnels is in reasonable agreement with the dynamometer measurements shown in Table 3D-4b.

Very few studies have reported comprehensive analyses of the organic composition of motor vehicle exhaust. The measurements by Rogge et al. (1993c) are the most comprehensive but are not expected to be the best representation of current motor vehicle emissions, because

these measurements were made in the mid-1980s. Measurements reported by Fraser et al. (1999) were made in a tunnel study conducted in 1993 and represent integrated gasoline and diesel powered vehicle emissions. In addition, exhaust emissions from two medium-duty diesel vehicles operating over an FTP cycle were analyzed by Schauer et al. (1999). A unique feature of both the measurements by Fraser et al. (1999) and Schauer et al. (1999) is that they include the quantification of unresolved complex mixture (UCM), which comprises aliphatic and cyclic hydrocarbons that cannot be resolved by gas chromatography (GC) (Schauer et al., 1999). Schauer et al. (1999) have shown that all of the organic compound mass in their diesel exhaust samples could be extracted and eluted by CG/MS techniques even though not all of the organic compound mass can be identified on a single compound basis. Table 3D-6 summarizes the composition of motor vehicle exhaust measured by Fraser et al. (1999) and Schauer et al. (1999).

Several studies have measured the distribution of polycyclic aromatic hydrocarbons (PAHs) in motor vehicle exhaust from on-road vehicles (Westerholm et al., 1991; Lowenthal et al., 1994; Venkataraman et al., 1994; Westerholm and Egeback, 1994; Reilly et al., 1998; Cadle et al., 1999, Weingartner et al., 1997b; Marr et al., 1999). Cadle et al. (1999) found that high molecular weight PAHs (PAHs with molecular weights greater than or equal to 202 g/mole) constitute 0.1 to 7.0% of the PM emissions from gasoline- and diesel-powered light-duty vehicles. It is important to note, however, that PAHs with molecular weights of 202 (fluoranthene, acephenanthrylene, and pyrene), 226 (benzo[ghi]fluoranthene and cyclopenta[cd]pyrene), and 228 (benz[a]anthracene, chrysene, and triphenylene) exist in both the gas-phase and particle-phase at atmospheric conditions (Fraser et al., 1998b), although those with molecular weight of 228 are predominantly associated with particles, with only traces in the gas-phase (Arey et al., 1987). Excluding these semivolatile PAHs, the contribution of nonvolatile PAHs to the PM emitted from the light-duty vehicles sampled by Cadle et al. (1999) ranges from 0.013 to 0.18%. These measurements are in good agreement with the tunnel study conducted by Fraser et al. (1999) and the heavy-duty diesel truck and bus exhaust measurements by Lowenthal et al. (1994), except that the nonvolatile PAH emissions from the heavy-duty diesel vehicles tested by Lowenthal et al. (1994) were moderately higher, making up ~0.30% of the PM mass emissions.

**TABLE 3D-6. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS
EMITTED FROM MOTOR VEHICLES**

Source	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Gasoline and diesel-powered vehicles driving through the Van Nuys Tunnel (Fraser et al., 1999) ^a	n-Alkanes	0.009	C ₂₁ through C ₂₉
	Petroleum biomarkers	0.078	Hopanes and steranes
	PAHs	0.38	No dominant compound
	Aromatic acids	0.29	Benzenedicarboxylic acids
	Aliphatic acids	0.21	Palmitic and stearic acids
	Substituted aromatic	0.042	No dominant compound
	UCM ^b	23.0	
Medium-duty diesel vehicles operated over an FTP Cycle (Schauer et al., 1999)	n-Alkanes	0.22	C ₂₀ through C ₂₈
	Petroleum biomarkers	0.027	Hopanes and steranes
	PAHs	0.54	No dominant compound
	Aliphatic acids	0.24	n-Octadecanoic acid
	Aromatic acids	0.014	Methylbenzoic acid
	Saturated cycloalkanes	0.037	C ₂₁ through C ₂₅
	UCM ^b	22.2	

^a Includes emissions of brake wear, tire wear, and resuspension of road dust associated with motor vehicle traffic.

^b Unresolved complex mixture.

Biomass Burning

In contrast to the mobile and stationary sources discussed earlier, emissions from biomass burning in wood stoves and forest fires are strongly seasonal and can be highly episodic within their peak emissions seasons. The burning of fuelwood is confined mainly to the winter months and is acknowledged to be a major source of ambient air PM in the northwestern United States during the heating season. Forest fires occur primarily during the driest seasons of the year in different areas of the country and are especially prevalent during prolonged droughts.

Particulate matter produced by biomass burning outside the United States (e.g., in Central America during the spring of 1988) also can affect ambient air quality in the United States.

An example of the composition of fine particles (PM_{2.5}) produced by wood stoves is shown in Figure 3D-4. These data were obtained in Denver during the winter of 1987-1988 (Watson and Chow, 1994). As was the case for motor vehicle emissions, OC and EC are the major components of particulate emissions from wood burning. It should be remembered that the relative amounts shown for OC and EC vary with the type of stove, the stage of combustion, and the type and condition of the fuelwood. Fine particles are dominant in smoke studies of wood burning emissions. For instance, the mass median diameter of wood particles was found to be ~0.17 μm in a study of the emissions from burning hardwood, softwood, and synthetic logs (Dasch, 1982).




Figure 3D-4. Chemical abundances for PM_{2.5} emissions from wood burning in Denver, CO. Solid bars represent fractional abundances, and the error bars represent variability in species abundances. Error bars represent detection limits when there are no solid bars.

Source: Watson and Chow (1994).

Kleeman et al. (1999) showed that the particles emitted by the combustion of wood in fireplaces are predominately < 1.0 µm in diameter, such that the composition of fine PM (PM_{2.5}) emitted from fireplace combustion of wood is representative of the total PM emissions from this source. Hildemann et al. (1991) and McDonald et al. (2000) reported that fireplace and wood stove smoke consists of 48 to 71% OC and 2.9 to 15% EC. Average OC and EC contents for these measurements are shown in Table 3D-7. It should be noted that the two methods used for the measurements shown in Table 3D-7 have been reported to produce different relative amounts of OC and EC for wood smoke samples but show good agreement for total carbon (OC + EC) measurements (Chow et al., 1993).

TABLE 3D-7. MASS EMISSIONS, ORGANIC CARBON, AND ELEMENTAL CARBON EMISSIONS FROM RESIDENTIAL COMBUSTION OF WOOD

Wood Type	Combustion Type	Average Mass Emission Rate (g/kg of wood burned)	Number of Tests	Percent Organic Carbon ^a	Percent Elemental Carbon ^a	References
Softwood	Fireplace	13.0	2	48.4	5.2	Hildemann et al. (1991)
Softwood	Fireplace	5.14	5	58.5	15.0	McDonald et al. (2000)
Hardwood	Fireplace	5.28	3	48.4	2.9	Hildemann et al. (1991)
Hardwood	Fireplace	5.66	5	63.2	7.0	McDonald et al. (2000)
Hardwood	Wood Stove	3.96	8	71.2	9.0	McDonald et al. (2000)

^a Hildemann et al. (1991) used the method described by Birch and Cary (1996) to measure EC; and McDonald et al. (2000) used the method reported by Chow et al. (1993) to measure OC.

Hawthorne et al. (1988) and Hawthorne et al. (1989) measured gas-phase and particle-phase derivatives of guaiacol (2-methoxyphenol), syringol (2,6-dimethoxyphenol), phenol, and catechol (1,2-benzenediol) in the downwind plume of 28 residential wood stoves and fireplaces. Rogge et al. (1998) reported a broad range of particle-phase organic compounds in the wood

smoke samples collected by Hildemann et al. (1991), which include n-alkanes, n-alkanoic acids, n-alkenoic acids, dicarboxylic acids, resin acids, phytosterols, PAHs, and the compounds reported by Hawthorne et al. (1989). Supplementing these measurements, McDonald et al. (2000) reported the combined gas-phase and particle-phase emissions of PAHs and the compounds quantified by Hawthorne et al. (1989). The measurements by Rogge et al. (1998), which represent a comprehensive data set of the organic compounds present in wood smoke aerosol, are summarized in Table 3D-8. It should be noted, however, that these nearly 200 compounds account for only ~15 to 25% of the OC particle mass emitted from the residential combustion of wood. Simoneit et al. (1999) have shown that levoglucosan constitutes a noticeable portion of the organic compound mass not identified by Rogge et al. (1998). In addition, Elias et al. (1999) used high-temperature gas chromatography/mass spectrometry (HTGC-MS) to measure high-molecular-weight organic compounds in smoke from South American leaf and stem litter biomass burning. These compounds cannot be measured by the analytical techniques employed by Rogge et al. (1998) and, therefore, are strong candidates to make up some of the unidentified organic mass in the wood smoke samples analyzed by Rogge et al. (1998). These compounds, which include triterpenyl fatty acid esters, wax esters, triglycerides, and high-molecular-weight n-alkan-2-ones, are expected to be present in North American biomass smoke originating from agricultural burning, forest fires, grassland fires, and wood stove/fireplace emissions.

Measurements of aerosol composition, size distributions, and aerosol emissions factors have been made in biomass burning plumes, either on towers (Susott et al., 1991) or aloft on fixed-wing aircraft (e.g., Radke et al., 1991) or helicopters (e.g., Cofer et al., 1988). As was found for wood stove emissions, the composition of biomass burning emissions is strongly dependent on the stage of combustion (i.e., flaming, smoldering, or mixed), and the type of vegetation (e.g., forest, grassland, scrub). Over 90% of the dry mass in particulate biomass burning emissions is composed of OC (Mazurek et al., 1991). Ratios of OC to EC are highly variable, ranging from 10:1 to 95:1, with the highest ratio found for smoldering conditions and the lowest for flaming conditions. Emissions factors for total particulate emissions increase by factors of two to four in going from flaming to smoldering stages in the individual fires studied by Susott et al. (1991).

**TABLE 3D-8. SUMMARY OF PARTICLE-PHASE ORGANIC COMPOUNDS
EMITTED FROM THE COMBUSTION OF WOOD IN FIREPLACES***

Biomass Type	Compound Class	Contribution to Particulate Mass (%)	Dominant Contributors to Emissions of Compound Class
Fireplace combustion of softwood	n-Alkanes	0.039	C ₂₁ through C ₃₁
	n-Alkanoic acids	0.45	C ₁₆ , C ₁₈ , C ₂₀ , C ₂₁ , C ₂₂ , C ₂₄
	n-Alkenoic acids	0.12	Oleic and linoleic acid
	Dicarboxylic acids	0.36	Malonic acid
	Resin acids	1.28	Abietic, dehydroabietic, isopimaric, pimaric, and sandaracopimaric acids
	Substituted phenols	3.30	Benzenediols and guaiacols
	Phytosterols	0.37	β-Sitosterol
	PAHs	0.092	Fluoranthene and pyrene
	Oxygenated PAH	0.019	1H-phenalen-1-one
Fireplace combustion of hardwood	n-Alkanes	0.044	C ₂₁ through C ₂₉
	n-Alkanoic acids	1.33	C ₁₆ , C ₂₂ , C ₂₄ , C ₂₆
	n-Alkenoic acids	0.049	Oleic and linoleic acid
	Dicarboxylic acids	0.42	Succinic acid
	Resin acids	0.11	Dehydroabietic acid
	Substituted phenols	8.23	Benzenediols, guaiacols, and syringols
	Phytosterols	0.21	β-sitosterol
	PAHs	0.13	No dominant compounds
	Oxygenated PAHs	0.020	1H-phenalen-1-one

*Note: Measurements were made using a dilution sampler and no semivolatile organic compound sorbent.

Source: Rogge et al. (1998).

Particles in biomass burning plumes from a number of different fires were found to have three distinguishable size modes: (1) a nucleation mode, (2) an accumulation mode, and (3) a coarse mode (Radke et al., 1991). Based on an average of 81 samples, ~70% of the mass was found in particles < 3.5 μm in aerodynamic diameter. The fine particle composition was found to be dominated by tarlike, condensed hydrocarbons; and the particles were usually

spherical in shape. Figure 3D-2 shows additional information for the size distribution of particles produced by vegetation burning.

An example of ambient data for the composition of PM_{2.5} collected at a tropical site that was heavily affected by biomass burning is shown in Table 3D-9. The samples were collected during November 1997 on the campus of Sriwijaya University, located in a rural setting on the island of Sumatra in Indonesia (Pinto et al., 1998). The site was subjected routinely to levels of PM_{2.5} well in excess of the U.S. NAAQS as a result of biomass fires in Indonesia from summer 1997 through spring 1998. As can be seen from a comparison of the data shown in Table 3D-9 with those shown in Figure 3D-4, there are a number of similarities and differences (especially with regard to the heavy metal content) in the abundances of many species. The abundances of some crustal elements (e.g., Si, Fe) are higher in Table 3D-9 than in Figure 3D-4, perhaps reflecting additional contributions of entrained soil dust.

TABLE 3D-9. MEAN AEROSOL COMPOSITION AT TROPICAL SITE (SRIWIJAYA UNIVERSITY, SUMATRA, INDONESIA) AFFECTED HEAVILY BY BIOMASS BURNING EMISSIONS^a

Component	Abundance (%)	Component	Abundance (%)
OC	76	Cr	BD ^b
EC	1.2	Mn	BD ^b
SO ₄ ²⁻	11	Fe	3.9×10^{-2}
Al	BD ^b	Ni	$< 3.8 \times 10^{-5}$
Si	9.3×10^{-2}	Cu	4.8×10^{-4}
Cl	4.4	Zn	3.1×10^{-3}
K	0.7	As	6.4×10^{-4}
Ca	4.5×10^{-2}	Se	2.8×10^{-4}
Ti	4.2×10^{-3}	Br	3.6×10^{-2}
V	BD ^b	Pb	3.1×10^{-3}

^aThe mean PM_{2.5} concentration during the sampling period (November 5 through 11, 1997) was 264 µg/m³.

^bBeneath detection limit.

Source: Pinto et al. (1998).

Limited emissions data that include organic compound speciation information have been reported for agricultural burning (Jenkins et al., 1996), forest fires (Simoneit, 1985), and grassland burning (Standley and Simoneit, 1987). Jenkins et al. (1996) present PAH emissions factors for the combustion of cereals (barley, corn, rice, and wheat), along with PAH emissions factors for wood burning. Profiles of organic compounds in emissions from meat cooking (Rogge et al., 1991) and cigarette smoke (Rogge et al., 1994) also have been obtained.

Natural Sources

Although sea-salt aerosol production is confined to salt water bodies, it is included here because many marine aerosols can exert a strong influence on the composition of the ambient aerosol in coastal areas. In some respects, the production of sea-salt aerosols is like that of windblown dust, in that both are produced by wind agitation of the surface. The difference between the two categories arises because sea-salt particles are produced from the bursting of air bubbles rising to the sea surface. Air bubbles are formed by the entrainment of air into the water by breaking waves. The surface energy of a collapsing bubble is converted to kinetic energy in the form of a jet of water that can eject drops above the sea surface. The mean diameter of the jet drops is about 15% of the bubble diameter (Wu, 1979). Bubbles in breaking waves range in size from a few μm to several mm in diameter. Field measurements by Johnson and Cooke (1979) of bubble size spectra show maxima in diameters at around 100 μm , with the bubble size distribution varying as (d/d_0) with $d_0 = 100 \mu\text{m}$.

Because sea-salt particles receive water from the surface layer, which is enriched in organic compounds, aerosol drops are composed of this organic material in addition to sea salt (about 3.5% by weight in seawater). The major ionic species by mass in seawater are: Na^+ (30.7%), Cl^- (55.0%), SO_4^{2-} (7.7%), Mg^{2+} (3.6%), Ca^{2+} (1.2%), K^+ (1.1%), HCO_3^- (0.4%), and Br^- (0.2%) (Wilson, 1975). The composition of the marine aerosol also reflects the occurrence of displacement reactions that enrich sea-salt particles in SO_4^{2-} and NO_3^- while depleting them of Cl^- and Br^- .

Sea salt is concentrated in the coarse mode, with a mass median diameter of $\sim 7 \mu\text{m}$ for samples collected in Florida, the Canary Islands, and Barbados (Savoie and Prospero, 1982). The size distribution of sulfate is distinctly bimodal. Sulfate in the coarse mode is derived from sea water, but sulfate in the submicron aerosol arises from the oxidation of dimethyl sulfide

(CH₃SCH₃) or DMS. DMS is produced during the decomposition of marine micro-organisms. DMS is oxidized to methane sulfonic acid (MSA), a large fraction of which is oxidized to sulfate (e.g., Hertel et al., 1994).

Apart from sea spray, other natural sources of particles include the suspension of organic debris and volcanism. Profiles of organic compounds in vegetative detritus have been obtained by Rogge et al. (1993d). Particles released from plants in the form of seeds, pollen, spores, leaf waxes, and resins, range in size from ~1 to 250 μm (Warneck, 1988). Fungal spores and animal debris, such as insect fragments, also are to be found in ambient aerosol samples in this size range. Although material from all the foregoing categories may exist as individual particles, bacteria usually are found attached to other dust particles (Warneck, 1988). Smaller bioaerosol particles include viruses, individual bacteria, protozoa, and algae (Matthias-Maser and Jaenicke, 1994). In addition to natural sources, other sources of bioaerosols include industry (e.g., textile mills), agriculture, and municipal waste disposal (Spendlove, 1974). Size distributions for bioaerosols have not been as well characterized as for other categories of airborne particles.

Trace metals are emitted to the atmosphere from a variety of sources such as sea spray, wind-blown dust, volcanoes, wildfires, and biotic sources (Nriagu, 1989). Biologically mediated volatilization processes (e.g., biomethylation) are estimated to account for 30 to 50% of the worldwide total Hg, As, and Se emitted annually; whereas other metals are derived principally from pollens, spores, waxes, plant fragments, fungi, and algae. It is not clear, however, how much of the biomethylated species are remobilized from anthropogenic inputs. Median ratios of the natural contribution to globally averaged total sources for trace metals are estimated to be 0.39 (As), 0.15 (Cd), 0.59 (Cr), 0.44 (Cu), 0.41 (Hg), 0.35 (Ni), 0.04 (Pb), 0.41 (Sb), 0.58 (Se), 0.25 (V), and 0.34 (Zn), suggesting a significant natural source for many trace elements. It should be noted, however, that these estimates are based on emissions estimates that have uncertainty ranges of an order of magnitude.

REFERENCES

- Arey, J.; Zielinska, B.; Atkinson, R.; Winer, A. M. (1987) Polycyclic aromatic hydrocarbon and nitroarene concentrations in ambient air during a wintertime high-NO_x episode in the Los Angeles basin. *Atmos. Environ.* 21: 1437-1444.
- Bagley, S. T.; Gratz, L. D.; Johnson, J. H.; McDonald J. F. (1998) Effects of an oxidation catalytic converter and a biodiesel fuel on the chemical, mutagenic, and particle size characteristics of emissions from a diesel engine. *Environ. Sci. Technol.* 32: 1183-1191.
- Birch, M. E.; Cary, R. A. (1996) Elemental carbon-based method for occupational monitoring of particulate diesel exhaust: methodology and exposure issues. *Analyst* 121: 1183-1190.
- Cadle, S. H.; Mulawa, P. H.; Ball, J.; Donase, C.; Weibel, A.; Sagebiel, J. C.; Knapp, K. T.; Snow, R. (1997) Particulate emission rates from in-use high-emitting vehicles recruited in Orange county, California. *Environ. Sci. Technol.* 31: 3405-3412.
- Cadle, S. H.; Mulawa, P.; Hunsanger, E. C.; Nelson, K.; Ragazzi, R. A.; Barrett, R.; Gallagher, G. L.; Lawson, D. R.; Knapp, K. T.; Snow, R. (1999) Light-duty motor vehicle exhaust particulate matter measurement in the Denver, Colorado, area. *J. Air Waste Manage. Assoc.* 49: PM-164-174.
- Calvert, J. G.; Heywood, J. B.; Sawyer, R. F.; Seinfeld, J. H. (1993) Achieving acceptable air quality: some reflections on controlling vehicle emissions. *Science* (Washington, DC) 261: 37-45.
- Chow, J. C.; Watson, J. G.; Pritchett, L. C.; Pierson, W. R.; Frazier, C. A.; Purcell, R. G. (1993) The DRI thermal/optical reflectance carbon analysis system: description, evaluation and applications in U.S. air quality studies. In: Lodge, J. P., Jr. Fourth international conference on carbonaceous particles in the atmosphere; April 1991; Vienna, Austria. *Atmos. Environ. Part A* 27: 1185-1201.
- Chow, J. C.; Watson, J. G.; Houck, J. E.; Pritchett, L. C.; Rogers, C. F.; Frazier, C. A.; Egami, R. T.; Ball, B. M. (1994) A laboratory resuspension chamber to measure fugitive dust size distributions and chemical compositions. *Atmos. Environ.* 28: 3463-3481.
- Cofer, W. R., III; Levine, J. S.; Sebacher, D. I.; Winstead, E. L.; Riggan, P. J.; Brass, J. A.; Ambrosia, V. G. (1988) Particulate emissions from a mid-latitude prescribed chaparral fire. *J. Geophys. Res. [Atmos.]* 93: 5207-5212.
- Dasch, J. M. (1982) Particulate and gaseous emissions from wood-burning fireplaces. *Environ. Sci. Technol.* 16: 639-645.
- Durbin, T. D.; Smith, M. R.; Norbeck, J. M.; Truex, T. J. (1999) Population density, particulate emission characterization, and impact on the particulate inventory of smoking vehicles in the South Coast Air Quality Management District. *J. Air Waste Manage. Assoc.* 49: 28-38.
- Elias, V. O.; Simoneit, B. R. T.; Pereira, A. S.; Cabral, J. A.; Cardoso, J. N. (1999) Detection of high molecular weight organic tracers in vegetation smoke samples by high-temperature gas chromatography-mass spectrometry. *Environ. Sci. Technol.* 33: 2369-2376.
- Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T. (1998a) Gas-phase and particle-phase organic compounds emitted from motor vehicle traffic in a Los Angeles roadway tunnel. *Environ. Sci. Technol.* 32: 2051-2060.
- Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T.; Rasmussen, R. A. (1998b) Air quality model evaluation data for organics. 5. C6-C22 nonpolar and semipolar aromatic compounds. *Environ. Sci. Technol.* 32: 1760-1770.
- Fraser, M. P.; Cass, G. R.; Simoneit, B. R. T. (1999) Particulate organic compounds emitted from motor vehicle exhaust and in the urban atmosphere. *Atmos. Environ.* 33: 2715-2724.
- Fujita, E.; Watson, J. G.; Chow, J. C.; Robinson, N. F.; Richards, L. W.; Kumar, N. (1998) Northern front range air quality study, volume C: source apportionment and simulation methods and evaluation. Reno, NV: Desert Research Institute; prepared for Colorado State University, Cooperative Institute for Research in the Atmosphere.
- Gertler, A. W.; Abu-Allaban, M.; Coulombe, W.; Gilles, J. A.; Pierson, W. R.; Rogers, C. F.; Sagebiel, J. C.; Tarnay, L. (2000) Preliminary results of a tunnel study to characterize mobile source particulate emissions. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.
- Gillies, J. A.; Gertler, A. W. (2000) Comparison and evaluation of chemically speciated mobile source PM_{2.5} particulate matter profiles. *J. Air Waste Manage. Assoc.* 50: 1459-1480.
- Hawthorne, S. B.; Miller, D. J.; Barkley, R. M.; Krieger, M. S. (1988) Identification of methoxylated phenols as candidate tracers for atmospheric wood smoke pollution. *Environ. Sci. Technol.* 22: 1191-1196.

- Hawthorne, S. B.; Krieger, M. S.; Miller, D. J.; Mathiason, M. B. (1989) Collection and quantitation of methoxylated phenol tracers for atmospheric pollution from residential wood stoves. *Environ. Sci. Technol.* 23: 470-475.
- Hertel, O.; Christensen, J.; Hov, Ø. (1994) Modelling of the end products of the chemical decomposition of DMS in the marine boundary layer. *Atmos. Environ.* 28: 2431-2449.
- Hildemann, L. M.; Markowski, G. R.; Jones, M. C.; Cass, G. R. (1991) Submicrometer aerosol mass distributions of emissions from boilers, fireplaces, automobiles, diesel trucks, and meat-cooking operations. *Aerosol Sci. Technol.* 14: 138-152.
- Hopke, P. K.; Lamb, R. E.; Natusch, D. F. S. (1980) Multielemental characterization of urban roadway dust. *Environ. Sci. Technol.* 14: 164-172.
- Houck, J. E.; Chow, J. C.; Watson, J. G.; Simons, C. A.; Pritchett, L. C.; Goulet, J. M.; Frazier, C. A. (1989) Determination of particle size distribution and chemical composition of particulate matter from selected sources in California: volume I and executive summary (final report). Sacramento, CA: California Air Resources Board.
- Houck, J. E.; Goulet, J. M.; Chow, J. C.; Watson, J. G.; Pritchett, L. C. (1990) Chemical characterization of emission sources contributing to light extinction. In: Mathai, C. V., ed. *Visibility and fine particles: an A&WMA/EPA international specialty conference; October 1989; Estes Park, CO. Pittsburgh, PA: Air & Waste Management Association; pp. 437-446. (A&WMA transactions series no. TR-17).*
- Huffman, G. P.; Huggins, F. E.; Shah, N.; Huggins, R.; Linak, W. P.; Miller, C. A.; Pugmire, R. J.; Meuzelaar, H. L. C.; Seehra, M. S.; Manivannan, A. (2000) Characterization of fine particulate matter produced by combustion of residual fuel oil. *J. Air Waste Manage. Assoc.* 50: 1106-1114.
- Jenkins, B. M.; Jones, A. D.; Turn, S. Q.; Williams, R. B. (1996) Particle concentrations, gas-particle partitioning, and species intercorrelations for polycyclic aromatic hydrocarbons (PAH) emitted during biomass burning. *Atmos. Environ.* 30: 3825-3835.
- Johnson, B. D.; Cooke, R. C. (1979) Bubble populations and spectra in coastal waters: a photographic approach. *J. Geophys. Res. C: Oceans Atmos.* 84: 3761-3766.
- Kerminen, V. M.; Makela, T. E.; Ojanen, C. H.; Hillamo, R. E.; Vilhunen, J. K.; Rantanen, L.; Havers, N.; Von Bohlen, A.; Klockow, D. (1997) Characterization of the particulate phase in the exhaust from a diesel car. *Environ. Sci. Technol.* 31: 1883-1889.
- Kittelson, D. B. (1998) Engines and nanoparticles: a review. *J. Aerosol Sci.* 29: 575-588.
- Kleeman, M. J.; Schauer, J. J.; Cass, G. R. (1999) Size and composition distribution of fine particulate matter emitted from wood burning, meat charbroiling, and cigarettes. *Environ. Sci. Technol.* 33: 3516-3523.
- Kleeman, M. J.; Schauer, J. J.; Cass, G. R. (2000) Size and composition distribution of fine particulate matter emitted from motor vehicles. *Environ. Sci. Technol.* 34: 1132-1142.
- Lawson, D. R.; Smith, R. E. (1998) The northern front range air quality study: a report to the Governor and General Assembly. Fort Collins, CO: Colorado State University; December.
- Linak, W. P.; Miller, C. A.; Wendt, J. O. L. (2000) Comparison of particle size distributions and elemental partitioning from the combustion of pulverized coal and residual fuel oil. *J. Air Waste Manage. Assoc.* 50: 1532-1544.
- Lowenthal, D. H.; Zielinska, B.; Chow, J. C.; Watson, J. G.; Gautam, M.; Ferguson, D. H.; Neuroth, G. R.; Stevens, K. D. (1994) Characterization of heavy-duty diesel vehicle emissions. *Atmos. Environ.* 28: 731-743.
- Maricq, M. M.; Podsiadlik, D. H.; Chase, R. E. (1999) Gasoline vehicle particle size distributions: comparison of steady state, FTP, and US06 measurements. *Environ. Sci. Technol.* 33: 2007-2015.
- Marr, L. C.; Kirchstetter, T. W.; Harley, R. A.; Miguel, A. H.; Hering, S. V.; Hammond, S. K. (1999) Characterization of polycyclic aromatic hydrocarbons in motor vehicle fuels and exhaust emissions. *Environ. Sci. Technol.* 33: 3091-3099.
- Mason, B. (1966) *Principles of geochemistry*. 3rd ed. New York, NY: John Wiley & Sons, Inc.
- Matthias-Maser, S.; Jaenicke, R. (1994) Examination of atmospheric bioaerosol particles with radii > 0.2 µm. *J. Aerosol Sci.* 25: 1605-1613.
- Mazurek, M. A.; Cofer, W. R., III; Levine, J. S. (1991) Carbonaceous aerosols from prescribed burning of a boreal forest ecosystem. In: Levine, J. S., ed. *Global biomass burning: atmospheric, climatic, and biospheric implications*. Cambridge, MA: MIT Press; pp. 258-263.
- McDonald, J. D.; Zielinska, B.; Fujita, E. M.; Sagebiel, J. C.; Chow, J. C.; Watson, J. G. (2000) Fine particle and gaseous emission rates from residential wood combustion. *Environ. Sci. Technol.* 34: 2080-2091.
- Miller, C. A.; Linak, W. P.; King, C.; Wendt, J. O. L. (1998) Fine particle emissions from heavy fuel oil combustion in a firetube package boiler. *Combust. Sci. Technol.* 134: 477-502.

- Norbeck, J. M.; Durbin, T. D.; Truex, T. J. (1998) Measurement of primary particulate matter emissions from light-duty motor vehicles. Riverside, CA: University of California, College of Engineering, Center for Environmental Research and Technology; prepared for Coordinating Research Council, Inc. and South Coast Air Quality Management District.
- Nriagu, J. O. (1989) A global assessment of natural sources of atmospheric trace metals. *Nature (London)* 338: 47-49.
- Nriagu, J. O.; Pacyna, J. M. (1988) Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature (London)* 333: 134-139.
- Olmez, I.; Sheffield, A. E.; Gordon, G. E.; Houck, J. E.; Pritchett, L. C.; Cooper, J. A.; Dzubay, T. G.; Bennett, R. L. (1988) Compositions of particles from selected sources in Philadelphia for receptor modeling applications. *JAPCA* 38: 1392-1402.
- Oros, D. R.; Simoneit, B. R. T. (2000) Identification and emission rates of molecular tracers in coal smoke particulate matter. *Fuel* 79: 515-536.
- Pierson, W. R.; Brachaczek, W. W. (1983) Particulate matter associated with vehicles on the road. II. *Aerosol Sci. Technol.* 2: 1-40.
- Pinto, J. P.; Grant, L. D.; Hartlage, T. A. (1998) Report on U.S. EPA air monitoring of haze from S.E. Asia biomass fires. Research Triangle Park, NC: U.S. Environmental Protection Agency, National Center for Environmental Assessment; report no. EPA/600/R-98/071.
- Radke, L. F.; Hegg, D. A.; Hobbs, P. V.; Nance, J. D.; Lyons, J. H.; Laursen, K. K.; Weiss, R. E.; Riggan, P. J.; Ward, D. E. (1991) Particulate and trace gas emissions from large biomass fires in North America. In: Levine, J. S., ed. *Global biomass burning: atmospheric, climatic, and biospheric implications*. Cambridge, MA: MIT Press; pp. 209-224.
- Reilly, P. T. A.; Gieray, R. A.; Whitten, W. B.; Ramsey, J. M. (1998) Real-time characterization of the organic composition and size of individual diesel engine smoke particles. *Environ. Sci. Technol.* 32: 2672-2679.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1991) Sources of fine organic aerosol. 1. Charbroilers and meat cooking operations. *Environ. Sci. Technol.* 25: 1112-1125.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993a) Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust: roads as sources and sinks. *Environ. Sci. Technol.* 27: 1892-1904.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993b) Sources of fine organic aerosol. 5. Natural gas home appliances. *Environ. Sci. Technol.* 27: 2736-2744.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993c) Sources of fine organic aerosol. 2. Noncatalyst and catalyst-equipped automobiles and heavy-duty diesel trucks. *Environ. Sci. Technol.* 27: 636-651.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993d) Sources of fine organic aerosol. 4. Particulate abrasion products from leaf surfaces of urban plants. *Environ. Sci. Technol.* 27: 2700-2711.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1994) Sources of fine organic aerosol. 6. Cigarette smoke in the urban atmosphere. *Environ. Sci. Technol.* 28: 1375-1388.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1997a) Sources of fine organic aerosol. 8. Boilers burning no. 2 distillate fuel oil. *Environ. Sci. Technol.* 31: 2731-2737.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1997b) Sources of fine organic aerosol. 7. Hot asphalt roofing tar pot fumes. *Environ. Sci. Technol.* 31: 2726-2730.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1998) Sources of fine organic aerosol. 9. Pine, oak, and synthetic log combustion in residential fireplaces. *Environ. Sci. Technol.* 32: 13-22.
- Sagebiel, J. C.; Zielinska, B.; Walsh, P. A.; Chow, J. C.; Cadle, S. H.; Mulawa, P. A.; Knapp, K. T.; Zweidinger, R. B.; Snow, R. (1997) PM-10 exhaust samples collected during IM-240 dynamometer tests of in-service vehicles in Nevada. *Environ. Sci. Technol.* 31: 75-83.
- Savoie, D. L.; Prospero, J. M. (1982) Particle size distribution of nitrate and sulfate in the marine atmosphere. *Geophys. Res. Lett.* 9: 1207-1210.
- Sawyer, R. F.; Johnson, J. H. (1995) Diesel emissions and control technology. Diesel exhaust: a critical analysis of emissions, exposure, and health effects. Cambridge, MA: Health Effects Institute.
- Schauer, J. J.; Cass, G. R. (2000) Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environ. Sci. Technol.* 34: 1821-1832.

- Schauer, J. J.; Kleeman, M. J.; Cass, G. R.; Simoneit, B. R. T. (1999) Measurement of emissions from air pollution sources. 2. C₁ through C₃₀ organic compounds from medium duty diesel trucks. *Environ. Sci. Technol.* 33: 1578-1587.
- Simoneit, B. R. T. (1985) Application of molecular marker analysis to vehicular exhaust for source reconciliations. *Int. J. Environ. Anal. Chem.* 22: 203-233.
- Simoneit, B. R. T.; Schauer, J. J.; Nolte, C. G.; Oros, D. R.; Elias, V. O.; Fraser, M. P.; Rogge, W. F.; Cass, G. R. (1999) Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles. *Atmos. Environ.* 33: 173-182.
- Spendlove, J. C. (1974) Industrial, agricultural, and municipal microbial aerosol problems. In: *Current problems in aerobiology: proceedings of the thirtieth general meeting of the Society for Industrial Microbiology*; August 1973; Evanston, IL. *Dev. Ind. Microbiol.* 15: 20-27.
- Standley, L. J.; Simoneit, B. R. T. (1987) Characterization of extractable plant wax, resin, and thermally matured components in smoke particles from prescribed burns. *Environ. Sci. Technol.* 21: 163-169.
- Susott, R. A.; Ward, D. E.; Babbitt, R. E.; Latham, D. J. (1991) The measurement of trace emissions and combustion characteristics for a mass fire. In: Levine, J. S., ed. *Global biomass burning: atmospheric, climatic, and biospheric implications*. Cambridge, MA: MIT Press; pp, 245-257.
- Turekian, K. K. (1971) Elements, geochemical distribution of. In: *Encyclopedia of science and technology*, v. 4. New York, NY: McGraw-Hill Book Company; pp. 627-630.
- U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1999) SPECIATE. Washington, DC: Technology Transfer Network-Clearinghouse for Inventories & Emissions Factors. Available: <http://www.epa.gov/ttn/chief/software/speciate/index.html> (2 June 2003).
- U.S. Environmental Protection Agency. (2002) Health assessment document for diesel engine exhaust. Washington, DC: Office of Research and Development; report no. EPA/600/8-90/057.
- Venkataraman, C.; Lyons, J. M.; Friedlander, S. K. (1994) Size distributions of polycyclic aromatic hydrocarbons and elemental carbon. 1. Sampling, measurement methods, and source characterization. *Environ. Sci. Technol.* 28: 555-562.
- Vinogradov, A. P. (1959) *The geochemistry of rare and dispersed chemical elements in soils*. 2nd ed. New York, NY: Consultants Bureau, Inc.
- Warneck, P. (1988) *Chemistry of the natural atmosphere*. New York, NY: Academic Press, Inc.
- Watson, J. G.; Chow, J. C. (1994) Clear sky visibility as a challenge for society. *Annu. Rev. Energy Environ.* 19: 241-266.
- Watson, J. G.; Fujita, E. M.; Chow, J. C.; Zielinska, B.; Richards, L. W.; Neff, W.; Dietrich, D. (1998) Northern front range air quality study final report. Reno, NV: Desert Research Institute; prepared for Colorado State University, Cooperative Institute for Research in the Atmosphere.
- Weingartner, E.; Burtscher, H.; Baltensperger, U. (1997a) Hygroscopic properties of carbon and diesel soot particles. *Atmos. Environ.* 31: 2311-2327.
- Weingartner, E.; Keller, C.; Stahel, W. A.; Burtscher, H.; Baltensperger, U. (1997b) Aerosol emission in a road tunnel. *Atmos. Environ.* 31: 451-462.
- Westerholm, R.; Egeback, K.-E. (1994) Exhaust emissions from light- and heavy-duty vehicles: chemical composition, impact of exhaust after treatment, and fuel parameters. In: *Symposium of risk assessment of urban air: emissions, risk identification, and risk quantitation*; May-June 1992; Stockholm, Sweden. *Environ. Health Perspect.* 102(suppl. 4) 13-23.
- Westerholm, R. N.; Almén, J.; Li, H.; Rannug, J. U.; Egeback, K.-E.; Grägg, K. (1991) Chemical and biological characterization of particulate-, semivolatile-, and gas-phase-associated compounds in diluted heavy-duty diesel exhausts: a comparison of three different semivolatile-phase samplers. *Environ. Sci. Technol.* 25: 332-338.
- Wilson, T. R. S. (1975) Salinity and the major elements of sea water. In: Riley, J. P.; Skirrow, G., eds. *Chemical oceanography*: v. 1. 2nd ed. London, United Kingdom: Academic Press Inc.; pp. 365-413.
- Wu, J. (1979) Spray in the atmospheric surface layer: review and analysis of laboratory and oceanic results. *J. Geophys. Res. C: Oceans Atmos.* 84: 1693-1704.

APPENDIX 3E

Characterization of PM_{2.5}, PM₁₀, and PM_{10-2.5} Concentrations at IMPROVE Sites

In this appendix, concentrations of PM_{2.5}, PM_{10-2.5} and PM₁₀ are presented based on daily values in the PM data set obtained at 18 sites in the IMPROVE network. The purpose of these analyses is to provide summaries of concentrations of PM_{2.5} and PM_{10-2.5} that could be used for characterizing human and ecosystem exposures to PM under the least polluted conditions that are found in the United States and to provide upper limits on policy relevant background (PRB) PM concentrations (cf., Section 3.3.3). This is accomplished by summarizing nonsulfate PM_{2.5} and PM_{10-2.5} concentrations and their variability at relatively remote monitoring sites (RRMS) that are not obviously influenced by local and/or regional anthropogenic pollution sources in the United States. The rationale for looking at nonsulfate PM_{2.5} is that the sulfate component of PM_{2.5} is almost all anthropogenic in origin and, therefore, nonsulfate PM_{2.5} provides a better estimate of upper limits on PRB PM concentrations.

The potential for receiving contributions from anthropogenic sources exists at every monitoring site. Thus, sites were chosen for this analysis where it was judged that anthropogenic sources do not contribute extensively to the monitored concentrations. These sites are characterized by lower interannual variability and smaller ranges of values across the percentile distribution in concentrations and by lower probabilities of being affected by transport from regional pollution sources than other sites. Such sites are almost all in the western United States. This analysis includes only two sites in the eastern United States and two areas in the midwest.

The 24-h average data available at the selected 18 IMPROVE sites from 1990 through 2002 were summarized on an annual basis; and annual mean values and annual 90th percentiles of the 24-h concentrations were calculated. The data capture requirement for these summaries required that each site have least 11 observations in each calendar quarter, and only years in which this requirement was satisfied for all four quarters were used. The monitoring site in Yellowstone National Park was relocated in 1996; thus, the designations for the two locations are Yellowstone National Park 1 and Yellowstone National Park 2. The Voyageurs National

Park site was also relocated. The designations for the two locations are analogous to those used for Yellowstone National Park.

Both $PM_{2.5}$ and PM_{10} mass were examined with the sulfate component removed to obtain better insight into non-anthropogenic background. Daily fine sulfate concentrations were calculated by $4.125 * S$, where S is the measured mass of fine sulfur (Malm, et al., 1994). Since sulfate is not a significant component of coarse fraction particles and measurements are not available, we calculated an estimate of the coarse PM ($PM_{10-2.5}$) by the difference between PM_{10} and $PM_{2.5}$.

Information about the range of annual mean nonsulfate (and total) $PM_{2.5}$ and PM_{10} mass concentrations, and $PM_{10-2.5}$ mass concentrations, is summarized in Table 3E-1, and the range of 90th percentile concentrations is summarized in Table 3E-2. The 90th percentile was chosen to reduce the impact of infrequent, episodic events that can result in exceptionally high daily values. As can be seen from Table 3E-1, the lowest annual average nonsulfate $PM_{2.5}$ concentrations among all sites examined were recorded at Denali National Park (AK), with concentrations there ranging from an annual mean of 0.7 to 2.4 $\mu\text{g}/\text{m}^3$. The site with the next lowest observed annual mean (among the 18 sites analyzed) was Bridger Wilderness (WY), where annual means ranged from 1.5 to 2.2 $\mu\text{g}/\text{m}^3$. The highest annual mean nonsulfate $PM_{2.5}$ concentrations at the sites examined were observed in Glacier National Park (MT), where values ranged from 3.8 to 5.5 $\mu\text{g}/\text{m}^3$. Annual means of coarse PM ranged from a low of 1.1 to 5.6 $\mu\text{g}/\text{m}^3$ at Denali National Park to a high of 5.6 to 10.8 $\mu\text{g}/\text{m}^3$ at Big Bend National Park, TX. Over the period of the observational record, a substantial range in annual mean concentrations in all three size ranges is observed, indicating the importance of examining a number of years in attempting to characterize long-term averages. Inspection of Table 3E-1 shows that the range in annual average $PM_{10-2.5}$ concentrations was generally larger than for $PM_{2.5}$. However, it should be noted that $PM_{10-2.5}$ concentrations are obtained by a difference technique and, so, are more strongly affected by errors in the determination of both $PM_{2.5}$ and PM_{10} .

As can be seen in Table 3E-2, 90th percentile 24-h nonsulfate $PM_{2.5}$ levels ranged from 1.5 to 10.5 $\mu\text{g}/\text{m}^3$. The 90th percentiles of 24-h coarse PM ranged from 1.9 to 27.6 $\mu\text{g}/\text{m}^3$. As can be seen from Table 3E-2, concentrations of PM in all three size fractions at the 90th percentile level are often several times higher than the mean concentrations at all sites. In addition, the range of values at the 90th percentile level is much greater than that at lower

**TABLE 3E-1. RANGES OF ANNUAL MEAN PM CONCENTRATIONS AT IMPROVE
MONITORING SITES ($\mu\text{g}/\text{m}^3$)**

Site	PM _{2.5}		PM ₁₀		Coarse PM
	Nonsulfate	(Total)	Nonsulfate	(Total)	
Acadia National Park, ME	2.6-4.7	(4.9-8.2)	4.6-11.3	(7.3-15.0)	1.8-6.0
Big Bend National Park, TX	2.7-4.9	(5.0-7.8)	8.8-15.7	(11.3-18.6)	5.6-10.8
Boundary Waters Canoe Area, MN	2.6-3.9	(4.4-5.8)	5.0-10.2	(7.0-12.0)	2.3-7.3
Bryce Canyon National Park, UT	1.7-2.4	(2.6-3.4)	4.4-7.6	(5.3-8.5)	2.5-5.6
Bridger Wilderness, WY	1.5-2.2	(2.1-2.9)	3.7-6.5	(4.3-7.3)	1.9-4.7
Canyonlands National Park, UT	1.9-3.2	(2.8-4.0)	5.1-10.5	(6.3-11.7)	3.2-8.0
Denali National Park, AK	0.7-2.4	(1.1-3.2)	2.0-7.5	(2.4-8.3)	1.1-5.6
Gila Wilderness, NM	2.4-3.4	(3.4-4.5)	4.9-7.9	(6.0-9.2)	2.5-5.0
Glacier National Park, MT	3.8-5.5	(4.8-6.5)	7.6-14.2	(8.5-15.2)	3.7-9.6
Lassen Volcanic National Park, CA	1.7-4.5	(2.1-5.1)	4.0-8.1	(4.6-8.5)	1.8-6.4
Lone Peak Wilderness, UT	3.1-5.3	(4.1-6.9)	7.1-10.9	(8.1-12.5)	3.7-6.0
Lye Brook Wilderness, VT	2.3-4.8	(4.5-8.8)	4.2-9.7	(7.0-13.6)	1.6-4.8
Redwood National Park, CA	2.8-4.6	(3.6-5.4)	6.0-10.6	(7.2-11.7)	3.3-6.5
Three Sisters Wilderness, OR	2.0-5.4	(2.7-6.5)	4.0-8.1	(4.6-9.1)	1.9-4.4
Voyageurs National Park 1, MN	3.2-3.5	(5.1-5.9)	5.7-11.2	(8.1-13.1)	2.8-7.8
Voyageurs National Park 2, MN	2.6-5.4	(4.1-7.2)	5.2-10.8	(7.0-12.5)	2.6-5.3
Yellowstone National Park 1, WY	2.0-3.0	(2.6-3.6)	6.0-9.2	(6.6-9.9)	3.8-7.0
Yellowstone National Park 2, WY	1.7-4.1	(2.3-4.7)	3.6-9.0	(4.2-9.6)	1.9-5.0

TABLE 3E-2. RANGES OF ANNUAL 90th PERCENTILE VALUES OF DAILY PM CONCENTRATIONS AT IMPROVE MONITORING SITES ($\mu\text{g}/\text{m}^3$)

Site	PM _{2.5}		PM ₁₀		Coarse PM
	Nonsulfate	(Total)	Nonsulfate	(Total)	
Acadia National Park, ME	4.8-10.5	(8.8-16.5)	9.0-20.2	(12.7-26.0)	3.6-12.0
Big Bend National Park, TX	4.7-9.5	(8.7-12.8)	15.9-29.0	(18.5-34.5)	10.4-18.7
Boundary Waters Canoe Area, MN	5.5-8.0	(8.1-12.0)	9.4-13.8	(13.6-17.1)	4.8-7.5
Bryce Canyon National Park, UT	2.9-5.0	(4.4-6.1)	7.4-13.8	(8.9-14.8)	4.7-12.0
Bridger Wilderness, WY	2.8-4.6	(3.9-5.8)	7.3-14.5	(8.3-15.1)	3.5-12.3
Canyonlands National Park, UT	3.2-7.3	(4.5-7.9)	9.5-16.1	(10.9-16.9)	6.3-14.7
Denali National Park, AK	1.5-5.1	(2.2-7.4)	3.6-17.2	(4.3-17.9)	1.9-11.7
Gila Wilderness, NM	4.1-6.3	(5.6-8.0)	8.3-14.4	(10.2-15.9)	4.2-9.6
Glacier National Park, MT	7.2-10.0	(8.0-11.0)	16.2-36.5	(16.5-36.8)	8.1-27.6
Lassen Volcanic National Park, CA	3.7-7.7	(4.0-9.1)	7.6-15.6	(8.5-16.0)	3.7-14.7
Lone Peak Wilderness, UT	5.4-9.8	(6.7-11.1)	11.7-19.6	(13.4-21.5)	6.6-10.9
Lye Brook Wilderness, VT	5.4-9.1	(11.4-18.2)	8.5-15.8	(14.4-22.9)	3.3-8.8
Redwood National Park, CA	5.2-9.4	(6.3-9.9)	11.8-21.8	(13.2-23.7)	5.9-13.0
Three Sisters Wilderness, OR	4.5-8.0	(5.8-10.0)	7.3-16.2	(8.5-18.0)	4.0-9.6
Voyageurs National Park 1, MN	5.7-6.4	(7.7-11.3)	13.9-22.0	(17.9-23.1)	9.0-17.4
Voyageurs National Park 2, MN	5.1-5.5	(7.4-8.1)	9.6-11.9	(12.0-15.5)	5.4-6.0
Yellowstone National Park 1, WY	3.9-5.7	(4.7-6.6)	11.7-20.6	(12.3-21.3)	8.8-16.5
Yellowstone National Park 2, WY	3.4-4.9	(4.1-6.0)	6.7-10.3	(7.4-10.8)	4.2-5.8

3E-4

levels. These findings indicate that extreme events are important for shaping the frequency distribution of PM that is observed. Wildfires play a major role in defining the year-to-year variability at specific sites, especially in the West. Notable examples include the massive forest fire in 1988 in Yellowstone National Park and those that occurred in 2000 in many western states. Dust storms also play a role in arid climates. The variability in annual average $PM_{10-2.5}$ concentrations is again related to several factors. Variability in factors that govern the production of soil dust from the surface and the production of primary biological aerosol particles may be largely responsible. Wildfires also produce PM in this size range (cf., Appendix 3D).

Examples of the interannual variability in the percentile distributions of $PM_{2.5}$ concentrations for 1997 through 2001 are shown in Figure 3E-1(a,b) for two sites. As can be seen, at concentrations less than the 90th percentile (P_{90}) level, year-to-year differences in $PM_{2.5}$ can be less than $1 \mu\text{g}/\text{m}^3$ at sites such as Bridger, WY. Differences at the P_{90} level are much larger, both on an absolute and on a relative basis at eastern sites.

Percentile distributions by season are shown in Figure 3E-2(a,b) for $PM_{2.5}$ and Figure 3E-3(a,b) for $PM_{10-2.5}$. For most sites, the variability in concentrations on a quarterly basis increases substantially beyond the P_{90} level, as can be seen from inspection of Figure 3E-2(a,b) for $PM_{2.5}$ and Figure 3E-3(a,b) for $PM_{10-2.5}$. Perhaps the most striking features seen in Figure 3E-2(a,b) and Figure 3E-3(a,b) are the concentration changes associated with the 95th (P_{95}) and 99th (P_{99}) percentile events, which represent extreme value events. Most of these events occur in the third calendar quarter; however, high concentrations also do occur at some sites during the second and fourth calendar quarters. If locally derived climatologic seasons were used instead of calendar quarters, a more accurate depiction of the seasonal variability of these events may have been obtained. In most cases, there is consistency in the behavior of $PM_{2.5}$ and PM_{10} , which suggests that the episodes of higher concentrations could be associated with sources that produce mainly $PM_{2.5}$ such as wildfires and/or anthropogenic combustion sources. Wildfires are limited to hotter and drier times of the year, but anthropogenic sources can contribute to high concentrations during other seasons. Additional factors which would tend to produce a third quarter maximum include: the enhanced production of secondary PM from anthropogenic and biogenic precursors during summer months; wildfires that are located in the East or elsewhere during summer; and surface dust produced locally and/or in northern Africa.

REFERENCES

Malm, W. C.; Sisler, J. F.; Huffman, D.; Eldred, R. A.; Cahill, T. A. (1994) Spatial and seasonal trends in particle concentration and optical extinction in the United States. *J. Geophys. Res. [Atmos.]* 99: 1347-1370.





Figure 3E-1a,b. Interannual variability in 24-h average PM_{2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY.




Figure 3E-2a,b. Seasonal variability in 24-h average PM_{2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY.





Figure 3E-3a,b. Seasonal variability in 24-h average PM_{10-2.5} concentrations observed at selected IMPROVE sites: (a) Bridger Wilderness, WY; (b) Yellowstone National Park, WY.

4. ENVIRONMENTAL EFFECTS OF AIRBORNE PARTICULATE MATTER

4.1 INTRODUCTION

This chapter assesses information providing inputs to U.S. EPA decision making on secondary National Ambient Air Quality Standards (NAAQS) aimed at protecting against welfare effects of ambient airborne particulate matter (PM). Specifically, it assesses the effects of atmospheric PM on the environment, including: (a) direct and indirect effects on vegetation and natural ecosystem integrity, (b) effects on visibility, and (c) effects on man-made materials, as well as (d) relationships of atmospheric PM to climate change processes.

4.2 EFFECTS OF AMBIENT AIRBORNE PARTICULATE MATTER ON VEGETATION AND NATURAL ECOSYSTEMS

Introduction

The effects of airborne particles are manifested via physical and chemical effects exerted at the individual plant level. However, plants are key members of ecosystems, which are structurally complex communities comprised of populations of plants, animals (including humans), insects, and microorganisms that interact with one another and with their non-living (abiotic) chemical and physical environment in which they exist (Odum, 1989; U.S. Environmental Protection Agency, 1993). All life on Earth is dependent on chemical energy in the form of carbon compounds to sustain their life processes. Terrestrial vegetation, via the process of photosynthesis, provides approximately half of the carbon that annually cycles between the Earth and the atmosphere (Chapin and Ruesch, 2001).

Ecosystems respond to stresses through their constituent organisms. The responses of plant species and populations to environmental perturbations (such as those caused by atmospheric PM) depend on their genetic constitution (genotype), their life cycles, and their microhabitats. Stresses that produce changes in their physical and chemical environment apply selection pressures on individual organisms (Treshow, 1980). The changes that occur within populations and plant communities reflect these new and different pressures. A common

response in a community under stress is the elimination of the more sensitive populations and an increase in abundance of species that tolerate or are favored by stress (Woodwell, 1970, Guderian et al., 1985).

The present section is organized to discuss: (1) factors affecting deposition of airborne PM on plants and ecosystems and then (2) the effects of PM deposition on individual plants, plant populations, forest trees and terrestrial ecosystems. As such, the section is organized to follow, in rough outline, the *Framework for Assessing and Reporting on Ecological Condition* recommended in a report by the Ecological Processes and Effects Committee (EPEC) of the EPA's Science Advisory Board (Science Advisory Board, 2002), which states:

“The purpose of this report is to provide the Agency with a sample framework that may serve as a guide for designing a system to assess, and then report on, ecological condition at local, regional, or national scale. The sample framework is intended as an organizing tool that may help the Agency decide what ecological attributes to measure and how to aggregate those measurements into an understandable picture of ecological integrity.” This framework is not actually a risk assessment per se, but it can be used to “construct a report of ecological condition” that characterizes the ecological integrity of an ecosystem based on “the relationship between common anthropogenic stressors and one or more of the six Essential Ecological Attributes.”

The EPEC report provides a useful approach for organizing discussions of stressor effects on ecosystem components at successive levels of complexity.

4.2.1 Ecological Attributes

The EPEC Framework provides a checklist of generic ecological attributes that should be considered when evaluating the integrity of ecological systems (see Table 4-1). The six generic ecological attributes, termed Essential Ecological Attributes (EEAs), represent groups of related ecological characteristics (Science Advisory Board, 2002; Harwell et al., 1999) and include: landscape conditions, biotic conditions, chemical and physical characteristics, ecological processes, hydrology and geomorphology, and natural disturbance regimes. All of the EEAs are interrelated (i.e., changes in one EEA may directly or indirectly affect other EEAs).

The first three ecological attributes listed in Table 4-1 are primarily “patterns,” whereas the last three are “processes.” Ecological science has used “patterns” and “processes” as terms to describe features of ecological systems for many years (e.g., Bormann and Likens, 1979).

TABLE 4-1. ESSENTIAL ECOLOGICAL ATTRIBUTES AND REPORTING CATEGORIES

<p>Landscape Condition</p> <ul style="list-style-type: none"> • Extent of Ecological System/Habitat Types • Landscape Composition • Landscape Pattern and Structure <p>Biotic Condition</p> <ul style="list-style-type: none"> • Ecosystems and Communities <ul style="list-style-type: none"> – Community Extent – Community Composition – Trophic Structure – Community Dynamics – Physical Structure • Species and Populations <ul style="list-style-type: none"> – Population Size – Genetic Diversity – Population Structure – Population Dynamics – Habitat Suitability • Organism Condition <ul style="list-style-type: none"> – Physiological Status – Symptoms of Disease or Trauma – Signs of Disease <p>Chemical and Physical Characteristics (Water, Air, Soil, and Sediment)</p> <ul style="list-style-type: none"> • Nutrient Concentrations <ul style="list-style-type: none"> – Nitrogen – Phosphorus – Other Nutrients • Trace Inorganic and Organic Chemicals <ul style="list-style-type: none"> – Metals – Other Trace Elements – Organic Compounds • Other Chemical Parameters <ul style="list-style-type: none"> – pH – Dissolved Oxygen – Salinity – Organic Matter – Other • Physical Parameters 	<p>Ecological Processes</p> <ul style="list-style-type: none"> • Energy Flow <ul style="list-style-type: none"> – Primary Production – Net Ecosystem Production – Growth Efficiency • Material Flow <ul style="list-style-type: none"> – Organic Carbon Cycling – Nitrogen and Phosphorus Cycling – Other Nutrient Cycling <p>Hydrology and Geomorphology</p> <ul style="list-style-type: none"> • Surface and Groundwater flows <ul style="list-style-type: none"> – Pattern of Surface flows – Hydrodynamics – Pattern of Groundwater flows – Salinity Patterns – Water Storage • Dynamic Structural Characteristics <ul style="list-style-type: none"> – Channel/Shoreline Morphology, Complexity – Extent/Distribution of Connected Floodplain – Aquatic Physical Habitat Complexity • Sediment and Material Transport <ul style="list-style-type: none"> – Sediment Supply/Movement – Particle Size Distribution Patterns – Other Material Flux <p>Natural Disturbance Regimes</p> <ul style="list-style-type: none"> • Frequency • Intensity • Extent • Duration
--	--

Source: Science Advisory Board (2002).

Of main concern in this chapter are relationships between a certain class of diverse airborne stressors from anthropogenic sources (i.e., PM) and one or more of the EEAs. Changes in patterns resulting from responses of vegetation and ecosystems to the effects of fine and coarse

PM deposition, along with known or possible effects on ecological processes associated with changes in the patterns, are discussed in the subsections that follow.

The reader is also referred to several other sources for more detailed discussions of several topics only briefly alluded to or addressed here. For example, an extensive discussion of various types of effects of acidic deposition is presented in the U.S. National Acid Precipitation Assessment Program (NAPAP) Biennial Report to Congress: An Integrated Assessment Program (National Science and Technology Council, 1998). Additionally, ecological effects of acidic precipitation and nitrate deposition on aquatic systems are discussed in the EPA document *Air Quality Criteria Document for Nitrogen Oxides* (U.S. Environmental Protection Agency, 1993); and sulfate deposition and effects, as related to wetlands and aquatic habitats, are discussed in U.S. Environmental Protection Agency (1982). Effects of lead on crops, vegetation, and ecosystems are assessed in the EPA document *Air Quality Criteria for Lead* (U.S. Environmental Protection Agency, 1986). Lastly, effects of “certain pesticides, metal compounds, chlorinated organic compounds, and nitrogen compounds” are discussed in *Deposition of Air Pollutants to the Great Waters, Third Report to Congress* (U.S. Environmental Protection Agency, 2000a).

4.2.2 Ecosystem Exposures – Particle Deposition

Airborne particles, their precursors, and their transformation products are removed from the atmosphere by wet and dry deposition processes. This atmospheric cleansing process fortunately lowers the long-term buildup of lethal concentrations of these pollutants in the air and moderates the potential for direct human health effects caused by their inhalation. Unfortunately, these deposition processes also mediate the transfer of PM pollutants to other environmental media where they can and do alter the structure, function, diversity, and sustainability of complex ecosystems.

The potential effects of PM deposition on vegetation and ecosystems encompass the full range, scales, and properties of biological organization listed under Biotic Condition, Table 4-1. Exposure to a given mass concentration of airborne PM, however, may lead to widely differing responses, depending on the particular mix of deposited particles. Particulate matter is not a single pollutant, but rather a heterogeneous mixture of particles differing in size, origin, and chemical composition. This heterogeneity exists across individual particles within samples from

individual sites and, to an even greater extent, between samples from different sites. Thus far, atmospheric PM has been defined, for regulatory purposes, mainly by size fractions and less clearly so in terms of chemical nature, structure, or source. While size is related to the mode and magnitude of deposition to vegetated landscapes and may be a useful surrogate for chemical constitution, PM size classes do not necessarily have specific differential relevance for vegetation effects (Whitby, 1978; U.S. Environmental Protection Agency, 1996a); i.e., both fine- and coarse-mode particles may affect plants. Much of the burden of sulfates (SO_4^{2-}), nitrates (NO_3^-), ammonium salts (NH_4^+), and hydrogen ions (H^+) resides in the atmosphere either dissolved in fog water or as liquid or solid aerosols. Therefore, assessment of atmospheric PM deposition and effects on vegetation unavoidably include discussion of NO_3^- and SO_4^{2-} and associated compounds involved in acidic and acidifying deposition. Other important issues relate to trace elements and heavy metals often found in ambient airborne PM.

4.2.2.1 Fine and Coarse Particulate Matter

Fine and coarse PM have a number of contrasting properties that affect their impact on vegetated systems (see Chapter 2, Table 2-1 of this document). The model results of Wiman and Agren (1985) and the measurements of Lovett and Lindberg (1993) addressing the complexity of deposition processes in patchy forested landscapes and vertical stratification within canopies reveal clear distinctions between the deposition behavior of fine and coarse particles. For one, coarse particles settle nearer their site of formation than do fine particles. Also, the chemical constitution of individual particles is strongly correlated with size (i.e., most S and much N is present on fine particles, whereas much of the base cation and heavy metal burden is present on coarse particles) and influences the predicted landscape loading of specific elements. Atmospheric PM may also act as a carrier for other directly phytotoxic materials (e.g., herbicides). Fine PM dominates the surface area of particles suspended in the atmosphere, while coarse PM dominates the mass of such airborne particles. Surface area may become more central to ecological impact assessment as recognition of the oxidizing capacity of fine particles, their interactions with other pollutants such as ozone (O_3), and their adsorption of phytoactive organic compounds such as herbicides become more fully appreciated. Fine and coarse particles respond to changes in atmospheric humidity, precipitation, and wind through different mechanisms, differentially altering their deposition characteristics.

Fine PM may exhibit similar mass concentrations at different sites (Figure 4-1) and yet be composed of very different constituents. In eastern North America, sulfate is typically the major component of this fraction, in contrast to the West where nitrate is a key component (Figure 4-1; cf., the eastern Appalachian site and the western California site). On the other hand, in the urban center of Mexico City (Hidy et al., 2000), an environment more similar to the western than eastern U.S., concentrations of fine PM of about 300 $\mu\text{g}/\text{m}^3$ are found, and sulfate concentrations are three times that of nitrate. In contrast to sulfur and nitrogen, the contributions of organic and elemental carbon to the eastern and western U.S. sites were similar (Figure 4-1), although soil-derived geologic material was greater at the more arid western site.




Figure 4-1. The diversity of fine PM from sites in the western and eastern United States.

Source: Modified from Sisler and Malm (2000).

Fine PM is typically more diverse than coarse PM and is secondary in nature, having condensed from the vapor phase or been formed by chemical reaction from gaseous precursors

in the atmosphere (see Chapter 2). Fine PM derives from atmospheric gas-to-particle conversion reactions involving nucleation, condensation, and coagulation, and from evaporation of water from contaminated fog and cloud droplets. Sulfur and nitrogen oxides (SO_x and NO_x) are often oxidized to their respective acids and neutralized with ammonium cations as particulate salts. Fine PM may also contain condensates of volatile organic compounds, volatilized metals, and products of incomplete combustion (see Chapter 3).

Coarse PM, in contrast, is mainly primary in nature, having been emitted from area, point, or line sources as fully formed particles derived from abrasion and crushing processes, soil disturbances, desiccation of marine aerosol from bursting bubbles, hygroscopic fine PM expanding with humidity to coarse mode, and/or gas condensation directly onto preexisting coarse particles (see Chapters 2 and 3). Suspended primary coarse PM may contain iron, silica, aluminum, and base cations from soil, plant and insect fragments, pollen, fungal spores, bacteria, and viruses, as well as fly ash, brake linings, debris, and automobile tire fragments. Coarse-mode particles can be altered by chemical reactions and/or physical interactions with gaseous or liquid contaminants.

The coating of coarse particles with semivolatile materials can substantially affect their deposition and potential for biological effects. For example, nitrogen exhibits a strongly bimodal size distribution: the peak above $1\ \mu\text{m}$ can be attributed to nitric acid (HNO_3) adsorption onto coarse alkaline particles; and that below $1\ \mu\text{m}$ can be attributed to gas phase condensation of ammonia with either sulfuric or nitric acid yielding either $(\text{NH}_4)_2\text{SO}_4$ or NH_4NO_3 aerosol. HNO_3 has an extremely high deposition velocity, nearly independent of the physiography of the surface. Therefore, formation of ammonium nitrate reduces local nitrogen deposition, because the deposition velocity of these particles is much less than that of HNO_3 gas.

Similarly, anthropogenic emissions of sulfur are mostly as sulfur dioxide (SO_2), which is hydrophilic, rapidly hydrated, and subsequently oxidized to sulfate (SO_4^{2-}), which is about 30-fold less phytotoxic than SO_2 . The ratio of $\text{SO}_4^{2-}:\text{SO}_2$ increases with aging of the air mass and, therefore, with distance from the source. Sulfate is thus a widespread regional/global pollutant and is sufficiently hygroscopic that, in humid air, it exists significantly in the coarse PM fraction. It is unusual for injurious levels of particulate sulfate to be deposited upon vegetation in the vicinity of emission sources, while direct injury due to SO_2 is commonly observed near uncontrolled point sources. In this case, gas-to-particle conversion is of benefit

to vegetation. The chemical composition of gaseous precursors of PM and the formation of sulfates and nitrates is discussed in Section 2.1.3 of Chapter 2.

Since enactment of the 1990 Clean Air Act Amendments, the atmospheric mix of PM precursors in the United States has changed substantially. That is, as emissions of SO₂ have declined, emissions of oxides of nitrogen (NO_x) have remained about the same, but emissions of cations have decreased (Hedin et al., 1994; Likens et al., 1996). This is almost certainly due to increased suspension of wind-borne geologic material from exposed soils.

For characterization of tropospheric chemistry, the deposition of O₃, NO_x, peroxides, and ammonia are first-order problems, followed by deposition of organics, SO₂, and particulate sulfate and nitrate (Wesely and Hicks, 2000). For impact on vegetation, however, the order may be different and may include different species — notably SO₂, fluoride (where it still exists as a problem), particulate heavy metals, base cations, sulfate and nitrate. In spite of the current regulatory focus on non-speciated PM, exposure to a given mass concentration of PM may lead to widely differing phytotoxic outcomes depending upon the particular mix of PM constituents involved. This variability has not been characterized adequately. Though effects of specific chemical fractions of PM have been described, there has been relatively little research aimed at defining the effects of unspciated PM on plants or ecosystems.

4.2.2.2 Diversity of Deposition Modes

Atmospheric deposition of particles to ecosystems takes place via both wet and dry processes, through three major routes: (1) wet, by precipitation scavenging in which particles are deposited in rain and snow; (2) much slower dry deposition; and (3) occult deposition (so named because it was hidden from measurements which determined the previous two types of deposition) by fog, cloud-water, and mist interception (Table 4-2). Unlike gaseous dry deposition, neither the solubility of the particle nor the physiographical activity of the surface are likely to be of first order importance in determining particulate dry deposition velocity (V_d). Factors that contribute to surface wetness or stickiness may be critical determinants of deposition efficiency. Available tabulations of deposition velocity are highly variable and suspect. Recent evidence indicates that all three modes of deposition (wet, occult, and dry) must be considered in determining inputs to watersheds or ecosystems, because each may dominate over specific

**TABLE 4-2. TYPES AND DETERMINANTS OF PARTICULATE DEPOSITION
AND IMPACT TO VEGETATION**

Type of Deposition	Determinant of Deposition	Quantifiable Factors
<i>Dry deposition</i>	Ambient Concentration	Distance from Source Emission Strength
	Atmospheric Conditions	Wind Speed Stability Mixing Height Temperature Humidity Dew Formation
	Aerosol Properties	Chemical Reactivity Particle Solubility Aerodynamic Diameter Biological Availability Hygroscopicity
	Surface Roughness	Terrain Discontinuity Leaf Pubescence Leaf Shape Plant Density Branch Spacing Tissue Flexibility
	Vegetation Condition	Surface Wetness Salt Exudates Organic Exudates Insect Excreta
<i>Wet deposition</i>	Ambient Concentration	Distance from Source Emission Strength
	Atmospheric Conditions	Mixing Height Timing of Precipitation Intensity of Precipitation Duration of Precipitation
	Aerosol Properties	Chemical Reactivity Particle Solubility Biological Availability
	Surface Roughness	Terrain Discontinuity Leaf Pubescence Leaf Area Index Nature of Exposed Bark and Stem
<i>Occult deposition</i>	As Above	Combination of Above Factors

intervals of time or space and ultimately, by interception and impaction, onto vegetation or other rough elements in the landscape.

The distribution of deposition between wet, dry, and occult modes varies substantially between locations for both nitrogen and sulfur (Figure 4-2). Clearly, rainfall and snowfall will determine the magnitude of wet deposition. Precipitation events clean the air so that dry deposition is eliminated or reduced during subsequent dry periods. Occult deposition depends upon landscape interception of the cloud base (Cape, 1993). This may occur at high elevation sites, in coastal areas subject to onshore advection, or in low-lying interior areas subject to radiation fogs. Thus, ecosystem exposure is determined by the mode, and to some extent the magnitude, of deposition. Total deposition among mountain sites is related to the magnitude of the occult deposition, particularly for nitrogen (Figure 4-2). Topography and vegetation characteristics influence the deposition modes differently (Table 4-3). In general, dry deposition is the most sensitive to features of the vegetated surface, and wet deposition is the least sensitive.

Comparison of micrometeorological and other methods for estimating particle deposition velocity (Erisman et al., 1997) suggests that there is little discrepancy between contrasting methodologies for estimating particle deposition and that this conclusion holds for both anions and cations, with the exception of nitrogenous species, which appear to interact with foliage in more complex ways. These comprehensive studies in the Speulder forest in the Netherlands indicated that aerosol deposition represents a considerable fraction of total deposition to the landscape. At this location, occult deposition was relatively insignificant, but dry deposition accounted for about one-fourth of alkaline-earth cation deposition.

Wet Deposition

Wet deposition results from the incorporation of atmospheric particles and gases into cloud droplets and their subsequent precipitation as rain or snow, or from the scavenging of particles and gases by raindrops or snowflakes as they fall (Lovett, 1994). Precipitation scavenging, in which particles are incorporated in hydrometers and deposited in the resulting rain and snow, includes rainout (within-cloud incorporation by nucleation) and washout (below-cloud scavenging by impaction). Wet deposition generally is confounded by fewer factors than dry or occult deposition and has been easier to quantify. Total inputs from wet deposition to vegetative canopies can be significant (Table 4-3), although not all wet deposition involves particle




Figure 4-2. Relative importance of three modes of deposition of nitrate (A) and sulfate (B) at high elevation sites (Unsworth and Wilshaw, 1989; Fowler et al., 1989; Mueller, 1988; Aneja and Murthy, 1994).

TABLE 4-3. RELATIVE IMPORTANCE OF WET, DRY, PARTICULATE, AND TOTAL DEPOSITION TO THREE FOREST SITES ^a

Site	Deposition							
	Total Nitrogen ^b				Total Sulfur ^c			
	Wet (%)	Dry (%)	Particle (%)	Total (kg/ha)	Wet (%)	Dry (%)	Particle (%)	Total (kg/ha)
Duke Forest	75	25	0.11	9.87	64	33	2.7	17.20
Cary Forest	71	20	0.94	5.80	76	20	4.2	7.60
Austin Forest	71	29	0.58	6.57	83	13	4.3	7.79

^a Data from Allen et al. (1994). Sampling was by triple filter pack so that fine-mode particles could be sampled preferentially. An average particle deposition velocity of 0.9 cm/s was derived as in Hicks et al. (1987).

^b Wet nitrogen consists of NO₃⁻ and NH₄⁺; dry nitrogen consists of vapor phase HNO₃ and NO₂; and particulate nitrogen consists of NO₃⁻.

^c Wet sulfur consists of SO₄²⁻, dry sulfur consists of vapor phase SO₂, and particulate sulfur consists of pSO₄⁻².

scavenging because gaseous pollutants also dissolve in raindrops during precipitation events (Lovett, 1994). This contribution is obscured during measurements, because wet deposition is measured simply by chemical analysis of total precipitation collected in clean, nonreactive buckets. Exclusion of dry deposited material (as opposed to dissolved gaseous species) requires closure or covering of the vessels except during precipitation.

Wet deposition is largely a function of precipitation amount and ambient pollutant concentrations. Surface properties (Table 4-2) have little effect on wet deposition, compared to dry or occult deposition, although leaves (depending on their surface properties of wettability, exposure, and roughness) retain liquid and solubilized PM. Extensive vegetative canopies typically develop leaf area indices (LAI; ratio of projected leaf area to ground area) much greater than 1. Thus, any material deposited via precipitation to the upper stratum of foliage is likely to be intercepted by several foliar surfaces before reaching the soil.

Landscape characteristics may also affect wet deposition. Forested hillsides often receive four- to six-fold greater inputs of wet deposition than short vegetation in nearby valleys. This effect is due to a variety of orographic effects (Unsworth and Wilshaw, 1989) and the closer aerodynamic coupling to the atmosphere of tall forest canopies than of the shorter canopies in the valleys. This leads to more rapid foliar drying in forest canopies, which reduces the

residence time but concentrates the solubilized particulate materials available for foliar uptake on the cuticular surface more quickly; and concentration increases the thermodynamic driving force for foliar uptake (Fowler et al., 1991; Unsworth, 1984; Schönherr and Huber, 1977). Following wet deposition, humidity and temperature conditions strongly influence the extent of biological effects, because of the competing effects of drying versus concentrating the solutions, which influence the rate of metabolic uptake of surface solutes (Swietlik and Faust, 1984). The net consequence of these factors on direct physical effects of wet deposited PM on leaves is not known.

Rainfall introduces new wet deposition and redistributes throughout the canopy previously dry-deposited PM, particularly coarse particles that were preferentially deposited in the upper foliage (Peters and Eiden, 1992). Both effects scale the likelihood of foliar contact and potential direct PM effects on vegetation nearly linearly with canopy leaf area. The concentrations of suspended and dissolved materials are typically highest at the onset and decline with duration of individual precipitation events (Lindberg and McLaughlin, 1986; Hansen et al., 1994). Sustained rainfall removes much of the accumulation of dry-deposited PM from foliar surfaces, reducing direct foliar effects and combining the associated chemical burden with the wet-deposited material (Lovett and Lindberg, 1984; Lovett, 1994) for transfer to the soil. Intense rainfall may contribute substantial total particulate inputs to vegetated land surfaces, mostly via the soil, but is less effective as a source of directly bioavailable or injurious pollutants to foliar surfaces. This washing effect, combined with differential foliar uptake and foliar leaching of different chemical constituents of PM, alters the composition of the rainwater that reaches the soil. Low intensity precipitation events, in contrast, may deposit significantly more particulate pollutants to foliar-surfaces than high intensity precipitation events. Because of the short duration and limited atmospheric cleansing, the concentration of PM in the final precipitation that remains in contact with foliar surfaces may be high. Additionally, low-intensity events may enhance foliar uptake through the hydrating of some previously dry-deposited particles. The combination of dry deposition to foliage and subsequent wet removal increases the potential for PM to exert effects via soil pathways: first by enhancing dry deposition relative to adjacent unvegetated surfaces; and second by accelerating passage along with wet deposited PM material by throughfall and stemflow to the soil. Once, in the soil, PM may affect the important ecosystem-level biogeochemical cycles of major, minor, and trace elements.

Dry Deposition

Dry particulate deposition is a complex, poorly characterized process controlled primarily by atmospheric stability, macro- and micro-surface roughness, particle diameter, and surface characteristics (Table 4-2; Hosker and Lindberg, 1982). The range of particle sizes, the diversity of canopy surfaces, and the variety of chemical constituents in airborne PM have slowed progress in both prediction and measurement of dry particulate deposition. Deposition of particles suspended regionally and throughout the full depth of the planetary boundary layer (PBL) is controlled by different mechanisms within the three distinct atmospheric transport zones above the surface. In the lower atmosphere, fine particles are transported by turbulent eddies of mechanical and convective origin. In the relatively unstirred, laminar boundary layer surrounding individual surface elements, Brownian diffusion dominates. Near the surface, actual deposition and contact with the surface is mediated by impaction (El-Shobokshy, 1985).

Dry deposition of atmospheric particles to plant and soil is a much slower process than wet or occult deposition, but it occurs nearly continuously and affects all exposed surfaces (Hicks, 1986). In dry deposition, particles at the large end of the spectrum (i.e., $> 5 \mu\text{m}$ diameter) are deposited mainly by gravitational sedimentation and inertial impaction. Smaller particles, especially those with diameters between ≈ 0.2 and $2 \mu\text{m}$, are not readily dry-deposited and tend to travel long distances in the atmosphere until their eventual deposition, most often via precipitation. This long-distance transport of fine aerosols is largely responsible for the regional nature of acid deposition (Lovett, 1994). A major conclusion from atmospheric deposition research is the realization that dry deposition is usually a significant and, in some cases, a dominant portion of total atmospheric deposition to an ecosystem (Lovett, 1994). Plant parts of all types, including those not currently physiologically active, along with exposed soil and water surfaces, receive steady deposits of dry dusts, elemental carbon encrustations, grease films, tarry acidic coatings, and heterogeneous secondary particles formed from gaseous precursors (U.S. Environmental Protection Agency, 1982).

Deposition fluxes may be calculated from measurements, estimates, or modeled values of mass concentration, C , at a specified measurement height and the total conductance or deposition velocity, V_d , from this height to the surface (equation 4-1; Hicks et al., 1987). These modeling techniques are closely allied with the micrometeorological techniques used to measure such fluxes. The flux, F , may be inferred as:

$$F = V_d * (C_z - C_0), \quad (4-1)$$

where F is flux to the surface, C_z is the particle concentration at measurement height z , C_0 is the particle concentration at receptor sites in the canopy (usually assumed equal to 0), and V_d is the overall deposition velocity. The flux is controlled by V_d and C_z .

Vertical transport of particles through the lower atmosphere to the vicinity of the vegetation elements is by turbulence and sedimentation, such that:

$$V_d = V_t + V_s, \quad (4-2)$$

in which V_t (inner, left-hand pathway of Figure 4-3) is a turbulent diffusion term, and V_s is a sedimentation term that dominates deposition of very coarse particles (Figure 4-4) and increases with particle size (Figure 4-5; dotted line). Sedimentation may be considered a pathway parallel to turbulent transport (Figure 4-4), but this is an oversimplification. V_s affects the concentration of particles near the surface where eddy transport may occur and also governs the redeposition of some fraction of the particles that were lost to resuspension or rebound following deposition by impaction. For this reason, V_s is included (Figure 4-3) in the composite surface resistance term ($R_a R_{cp} V_s$) as well as in the parallel sedimentation term.

For submicron particles for which sedimentation is negligible (Hicks et al., 1987; Monteith and Unsworth, 1990; Wesely, 1989), the resistance catena Ohm's Law Analogy analogous to the law used to describe transport of heat, momentum, or gases may be adequate, as:

$$V_d = V_t = [r_a + r_b + r_c]^{-1}, \quad (4-3)$$

where V_t is the deposition velocity due to turbulent transport of particles or other entities through the atmosphere; r_a is aerodynamic resistance (inverse of conductance or velocity) associated with the efficiency of turbulent transport above the canopy; r_b is the boundary layer resistance associated with diffusional transport through the still air layer immediately adjacent to canopy elements; and r_c is canopy resistance associated with physiological control of leaf porosity via stomata in the leaf surface. Significant departures from the analogy arise near the surface




Figure 4-3. A simplified resistance catena representing the factors controlling deposition of particles to the surface. Vegetation-specific parameters are not explicitly considered.

Source: Modified after Hicks et al. (1987).

(Chamberlain, 1975; Sehmel, 1980), where particles that were transported efficiently by turbulent eddies are slowed substantially in the laminar boundary layer, reducing the efficiency of impaction. The preservation of momentum in this zone declines with decreasing diameter; however, this is offset by an increase in Brownian diffusivity with decreasing diameter (Figure 4-3). Aerodynamic streamlines are parallel to the surface of each roughness element, so that deposition ultimately depends on diffusion to the surface. The transition from impaction to diffusion is likely blurred in the presence of leaf pubescence extending beyond the boundary layer. These conflicting trends lead to a broad range over which empirical measurements of V_d and particle size are relatively independent (Figure 4-3), further demonstrating the importance of the quasilaminar boundary layer (Lamaud et al., 1994; Shinn, 1978).




Figure 4-4. The relationship between deposition velocity of selected particulate materials and the distribution of the material between the coarse- and fine-aerosol fractions.

Source: Data from Foltescu et al. (1994). Ranges for Mn and Fe are from Davidson and Wu (1989).

The aerodynamic term (r_a) decreases with increasing wind speed, turbulence, and friction velocity and increases with measurement height and atmospheric stability. It describes the capacity of turbulent eddies to transport material, momentum, and heat between the measurement height and the roughness height of the surface. Coarse particles may not be carried efficiently by the high frequency eddies near the surface and may fall more rapidly than they diffuse by either Brownian or turbulent processes. Thus, the relevance of r_a breaks down as V_s increases. Indeed because V_s (equation 4-2) is independent of a concentration gradient, the electrical analogy is a theoretically flawed approximate approach (Venkatram and Pleim, 1999).




Figure 4-5. The relationship between particle diameter and deposition velocity for particles. Values measured in wind tunnels by Little and Wiffen (1977) over short grass with wind speed of 2.5 m/s closely approximate the theoretical distribution determined by Peters and Eiden (1992) for a tall spruce forest. These distributions reflect the interaction of Brownian diffusivity (descending dashed line), which decreases with particle size and sedimentation velocity (ascending dotted line from Stokes Law), which increases with particle size. Intermediate-sized particles (≈ 0.1 to $1.0 \mu\text{m}$) are influenced strongly by both particle size and sedimentation velocity, and deposition is independent of size.

Deposition Velocity

Because the final stage of deposition for particles involves either impaction following deceleration through a quasilaminar boundary layer or diffusion through this boundary layer, its effective depth is a critical determinant of V_d (Wiman et al., 1985; Peters and Eiden, 1992). The term corresponding to the boundary layer resistance for gases (r_b ; equation 4-3) incorporates the absence of form drag for gases. This parameter decreases with increasing turbulence and particle diffusivity; it is poorly characterized for gases (depending critically on canopy morphology, vertical wind profiles, and gust penetration) and is of extremely limited usefulness for particles.

Once delivered by turbulent transport or sedimentation to the vicinity of vegetative surface elements, a variety of particle size-dependent mechanisms come into play, some differing substantially from those governing gaseous deposition. The concepts of r_b (the still air or boundary layer resistance) and r_c (the canopy or surface resistance) are not generally applicable to deposition of polydisperse particles. Because of the roles of momentum and bounce-off and the complication by reentrainment back into the airstream following deposition of a particle to the surface, the factors determining the effective r_b and r_c for particle deposition are not as independent as for gases. They are replaced in some resistance formulations (e.g., Hicks et al., 1987) by the term, r_{cp} , that combines near-surface and surface effects and by a mathematically derived composite term, $R_a R_{cp} V_s$, that combines atmospheric, surface, and sedimentation effects (Figure 4-4). This latter term was insignificant for the submicron sulfate component considered originally in its derivation (Hicks et al., 1987); however, it scales with the square of particle diameter, so that its general applicability to polydisperse particles is unclear. In general, transport between the turbulent air column and the leaf surface through the laminar boundary layer remains difficult to describe (Lindberg and McLaughlin, 1986).

Current estimates of regional particulate dry deposition (e.g., Edgerton et al., 1992; Brook et al., 1999) infer fluxes from the product of variable and uncertain measured or modeled particulate concentrations and even more variable and uncertain measured or modeled estimates of dry deposition velocity parameterized for a variety of specific surfaces (e.g., Brook et al., 1999). However, even for specific sites and well defined particles, uncertainties in F are largest in the values of V_d , which are typically characterized by the large ranges and variances described in other sources (e.g., Bytnerowicz et al., 1987a,b and Hanson and Lindberg, 1991, for nitrogen-containing particles; McMahon and Denison, 1979 and Hicks et al., 1987, for general treatment). The nature of the vegetative cover to which particulate deposition occurs has a moderate to substantial effect on the components of V_d . The surface resistance (Hicks et al., 1987) is a significant and highly site-specific component of total resistance that is difficult to predict along with site, seasonal, and diurnal effects on the atmospheric components of total resistance.

Early models of dry particulate deposition to vegetation (e.g., U.S. Environmental Protection Agency, 1982; Chamberlain, 1975; Davidson and Friedlander, 1978; Garland, 1978; Little and Wiffen, 1977; McMahon and Denison, 1979; Sehmel, 1980; Sehmel and Hodgson, 1976; Slinn, 1977, 1978) used this paradigm (e.g., equation 4-3) to deal with transport to the

near-surface regime explicitly including conventional micrometeorological and particle size considerations. Alternative modeling treatments have attempted to parameterize the geometry of vegetative receptor surfaces and within-canopy micrometeorology (Wiman and Ågren, 1985; Peters and Eiden, 1992). Chemical reactivity, particle shape and density, rates of physiological sequestration, and reentrainment by gusts of wind remain to be addressed. Modeling the deposition of particles to vegetation is at a relatively early stage of development, and it is not currently possible to identify a best or most generally applicable modeling approach. These approaches have been further elaborated with canopy-specific choices among the available models and with specific incorporation of capture efficiencies by Brook et al. (1999).

Methods of Measuring Dry Deposition

Methods of measuring dry deposition of particles are more restricted than for gaseous species and fall into two major categories: surface analysis methods, which include all types of measurements that examine contaminant accumulation on surfaces of interest, and atmospheric flux methods, which measure contaminants in the atmosphere from which one may estimate the flux (Davidson and Wu, 1990). Surface extraction or washing methods characterize the accumulation of particles on natural receptor surfaces of interest or on experimental surrogate surfaces. These techniques rely on methods designed specifically to remove only surface-deposited material (Lindberg and Lovett, 1985). Total surface rinsate may be equated to accumulated deposition or to the difference in concentrations in rinsate between exposed and control (sheltered) surfaces and may be used to refine estimates of deposition (John et al., 1985; Dasch, 1987). In either case, foliar extraction techniques may underestimate deposition to leaves because of uptake and translocation processes that remove pollutants from the leaf surface (Taylor et al., 1988; Garten and Hanson, 1990). Foliar extraction methods also cannot distinguish sources of chemicals (e.g., N) deposited as gases from those deposited as particles (e.g., HNO_3 or NO_3^- from nitrogen dioxide [NO_2], or NH_3 from NH_4^+ [Bytnerowicz et al., 1987a,b; Dasch, 1987; Lindberg and Lovett, 1985; Van Aalst, 1982]). Despite these limitations, these methods are often used in the development of in-canopy deposition models (McCartney and Aylor, 1987).

Deposition of pollutants by wet deposition is relatively straightforward to determine through analysis of precipitation samples. Dry deposition of pollutants, on the other hand, is

more difficult to measure. A significant limitation on current capacity to estimate regional impacts of PM is inadequate knowledge of the mechanism and factors governing particle dry deposition to diverse surfaces. This has hindered efforts to develop robust measurement techniques for particle deposition (distinct from atmospheric concentration) and has compromised efforts to develop generally applicable deposition models for particles. The National Dry Deposition Network (NDDN) was established in 1986 to document the magnitude, spatial variability, and trends in dry deposition across the United States. Currently, the network operates as a component of the Clean Air Status and Trends Network (CASTNet) (Clarke et al., 1997).

Dry deposition is not measured directly, but is determined by an inferential approach (i.e., fluxes are calculated as the product of measured ambient concentration and a modeled deposition velocity). This method is appealing and widely used because atmospheric concentrations are relatively easy to measure when compared to dry deposition fluxes, and models have been developed to calculate deposition velocities (Lovett, 1994). Ambient pollutant concentrations, meteorological conditions, and land use data required for the inferential model are routinely collected at CASTNet dry deposition sites. Monitored chemical species include ozone, sulfate, nitrate, ammonium, sulfur dioxide, and nitric acid. The temporal resolution for the ambient concentration measurements and dry deposition flux calculations is hourly for ozone and weekly for the other chemical substances (Clarke et al., 1997). Isotopic labeling of dry deposited PM (e.g., sulfate with ^{35}S) prior to experimental surface exposures and extractions (Garten et al., 1988) can provide more precise differentiation between the deposition rates of related chemical species (e.g., SO_4^{2-} from SO_2).

At the whole-canopy level, natural surface washing by rainfall may be used to estimate dry deposition of PM and gases during the preceding dry period (Cape et al., 1992; Davidson and Wu, 1990; Draaijers and Erisman, 1993; Erisman, 1993; Fahey et al., 1988; Lindberg and Lovett, 1992; Lovett and Lindberg, 1993; Reiners and Olson, 1984; Sievering, 1987). Collection and analysis of stem flow and throughfall provides useful estimates of particulate deposition when compared to directly sampled precipitation. The method is most precise for strictly PM deposition when gaseous deposition is a small component of the total dry deposition and when leaching or uptake of compounds of interest out of or into the foliage (i.e., N, S, base cations) is not a significant fraction of the total depositional flux (Davidson and Wu, 1990; Draaijers and

Erisman, 1993; Lindberg and Lovett, 1992; Lovett and Lindberg, 1993). Throughfall sampling of sulfate deposition (Garten et al., 1988; Lindberg and Garten, 1988; Lindberg et al., 1990) often suggests substantial foliar exchange. Other throughfall studies (e.g., Erisman, 1993; Fahey et al., 1988) may lack sufficient specificity for evaluating dry particle deposition.

Careful timing of throughfall and stemflow measurements after prolonged dry periods, with simultaneous direct measurement of unchanged precipitation, can be used to determine the magnitude of dry deposition between precipitation events. Indeed, this foliar washing technique, whether using subsequent precipitation or experimental lavage, is one of the best available methods to determine dry deposition of PM to vegetated ecosystems. Major limitations include the extreme site specificity of the measurements, the substantial labor requirement that normally precludes regional coverage, and the restriction to elements that are conserved within the vegetative system (thereby excluding semivolatile organics, ammonium salts, and gases such as ozone).

Surrogate surfaces have not been found that can adequately replicate essential features of natural surfaces; and therefore the surfaces currently used do not produce reliable estimates of particle deposition to the landscape. Deposition to surrogate surfaces deployed in extensive plant canopies may provide a measure of particle deposition to the surrounding foliage or soil surfaces. For example, a uniform population of 0.8 μm gold colloid particles deposited with similar coverage to leaves of *Phaseolus vulgaris* and to upward-facing inert surfaces (Klepper and Craig, 1975). However, dry deposition of particles to foliage and to flat inert surrogate surfaces (polycarbonate Petri dishes) in a deciduous forest resulted in greater accumulation on the inert surfaces, although both surfaces accumulated particles of a similar range of sizes (Lindberg and Lovett, 1985). These differences in deposition/accumulation remain to be fully characterized and hinder the surrogate techniques in providing quantitative deposition estimates.

Micrometeorological methods employ an eddy covariance, eddy accumulation, or flux gradient protocol in contrast to washing or extracting of receptor surfaces for quantifying dry deposition. These techniques require measurements of PM concentrations and of atmospheric transport processes. They are currently well developed for ideal conditions of flat, homogeneous, and extensive landscapes and for chemical species for which accurate and rapid sensors are available. Additional studies have expanded the variety of such species and extended

these techniques to more complex terrain (McMillen, 1988; Hicks et al., 1984; Wesely and Hicks, 1977).

The eddy covariance technique measures vertical fluxes of gases and fine particles directly from calculations of the mean covariance between the vertical component of wind velocity and pollutant concentration (Wesely et al., 1982). This technique is particularly limited by its requirement for sensors capable of acquiring concentration data at 5 to 20 Hz. For the flux-gradient or profile techniques, vertical fluxes are calculated from a concentration difference and an eddy exchange coefficient determined at discrete heights (Erisman et al., 1988; Huebert et al., 1988). Businger (1986), Baldocchi (1988), and Wesely and Hicks (1977) evaluated the benefits and pitfalls of these micrometeorological flux measurements for gases. Most measurements of eddy transport of PM have used chemical sensors (rather than mass or particle counting) to focus on specific PM components. These techniques have not been well developed for generalized particles and, for the same reasons that limit mathematical description of particle deposition, may be less suitable for coarse particles that are transported efficiently in high frequency eddies (Gallagher et al., 1988).

Factors Affecting Dry Deposition

Ambient Concentration. The ambient concentration of particles (C_z ; equation 4-1), the parameter for which there is the most data (for example, see Chapter 3, this document), is at best an indicator of exposure. However, it is the amount of PM actually entering the immediate plant environment that determines the biological effect. The linkage between ambient concentration and delivery to vegetation is the deposition velocity (V_d ; equation 4-1). C_z is determined by regional and local emission sources, regional circulation, and weather. It may be locally sensitive to removal from the atmosphere by deposition, but this effect is generally small. Average annual concentrations for NO_3^- exhibit much more variability than SO_4^{2-} and a definite pattern of higher concentrations in the Midwest than elsewhere. The highest concentrations are observed (i.e., $> 2 \mu\text{g m}^{-3}$) in the agricultural areas of the Midwest, while the lowest are seen at forested sites in New England and the southern Appalachian Mountains. Annual average concentrations SO_4^{2-} of $5.0 \mu\text{g m}^{-3}$ are observed over nearly the entire eastern United States from New York and Michigan to northern Mississippi and Alabama (Edgerton et al., 1992).

Deposition increases linearly with the concentration of many materials over a broad range. This allows atmospheric cleansing to take place and accounts for the greater surface impact of pollutant particles during pollution episodes. A serious limitation of the V_d formulation used to infer deposition of specific chemical species that exist in a range of particle sizes is an appropriate specification of their concentration. Most sulfur emissions are readily oxidized to sulfite, bisulfite, and SO_4^{2-} . In the presence of atmospheric ammonia, particulate ammonium sulfate is formed. However, this material is hygroscopic and will increase in mass and diameter in the presence of high humidity, altering its deposition behavior. Similarly, coalescence of small particles into larger aggregates and adsorption of gaseous pollutants onto existing coarse particles complicate the association of particle size with concentration of individual chemical species.

Distance from the source, and the resulting residence time in the atmosphere, control the relative concentrations of surface-reactive materials (NO , SO_2) of secondary particles that take some time to form in the atmosphere and of coarse particles that exhibit high rates of deposition by sedimentation near the source. These interacting processes affect the time required for the formation of secondary particles by gas-to-particle conversion reactions, resulting in a greater ratio of dry to wet deposition near emission sources where gaseous sulfur dioxide (gSO_2) deposition predominates than at greater distances where rainout of particulate SO_4^{2-} (pSO_4^{2-}) may dominate (Barrie et al., 1984) and where dry deposition of pSO_4^{2-} may be greater than that of gSO_2 . The effect of gas-to-particle conversion on dry deposition of a specific chemical species can be substantial because V_d for SO_2 is approximately 0.33 ± 0.17 cm/s; whereas it is approximately 0.16 ± 0.08 cm/s for SO_4^{2-} . These phase conversions affect both C_z and the effective V_d which together control dry depositional fluxes (equation 4-1). The neutralization of acidic gaseous and particulate species by alkaline coarse particles has been described in arid regions, but it may be more prevalent in urban New York where coarse particles are observed to be neutral because alkaline cations approximately balance gaseous acidic species (Lovett et al., 2000). The deposition of the acidic materials in the urban environment is likely enhanced by incorporation into these previously formed coarse particles.

Similarly, the ratio of coarse to fine particle concentrations determines the effective V_d for chemically speciated particles (Figure 4-4). This ratio reflects the size-dependent deposition processes that govern delivery of PM to receptor surfaces (Figure 4-5). For example, SO_4^{2-} was

found predominantly on fine submicron particles; whereas potassium (K^+), calcium (Ca^{2+}), and nitrate (NO_3^-) ions were associated most often with coarse particles larger than $2\ \mu m$ (Lindberg and Lovett, 1985). However, concentrations of particulate S and K^+ within a coniferous canopy were strongly correlated (Wiman and Lannefors, 1985), suggesting a primary source of coarse-mode sulfur particles. Many researchers reported bimodal mass-size distributions of NO_3^- with one peak in the fine mode and the other peak in the coarse mode (Wu and Okada, 1994). The behavior of NO_3^- in ambient aerosols depends not only on the concentrations of gaseous nitric acid and ammonia, but also on the chemical composition of the particles and atmospheric conditions (Tang, 1980). When sea-salt particles were transported atmospherically from the Pacific Ocean in the Nagoya, Japan area, the amount of nitrate in the coarse particle size range increased. Coarse particle formation on sea salt under these conditions becomes a major pathway for nitrate. The heterogeneous reaction of NaCl with gaseous HNO_3 is considered to be the dominating pathway for nitrate formation (Wu and Okada, 1994). As a result, marine and continental particle size spectra for both N and S differ substantially: a peak in the coarse mode is generally apparent near marine sources (Milford and Davidson, 1987). The issue for NO_3^- is further confounded by uncertain discrimination between gaseous and particulate N species by current sampling methods. The substantial effect of particle size on V_d (Figure 4-5) implies a need for size resolution as well as chemically speciated ambient PM concentrations, even within the PM_{10} fraction.

Particle Effects on V_d . Particle size is a key determinant of V_d as noted above; but, unfortunately, the size spectra may be quite complex. The particles in the study of Lindberg and Lovett (1985) at Walker Branch Watershed had median diameters ranging from 3 to $5\ \mu m$; but approximately 25% of the particles had diameters $< 1\ \mu m$ (0.2 to $0.3\ \mu m$), and 5 to 20% of the particles were much larger aggregates. The aggregated particles are significant in that chemically they reflect their fine particle origins, but physically they behave like large particles and deposit by sedimentation. Direct observations with SEM demonstrate that particle morphology can be highly variable. Many submicron particles can be observed on trichomes (leaf hairs), although most particles are in the 5 to $50\ \mu m$ diameter range. Large aggregated particles in excess of $100\ \mu m$ also are seen; larger carbonaceous aggregate particles are

especially common (Smith, 1990a). Trichomes are efficient particle receptors; however, they are reduced in size by “weathering” and occasionally break off during the growing season.

In the size range around approximately 0.1 to 1.0 μm , where V_d is relatively independent of particle diameter (Figure 4-5), deposition is controlled by macroscopic roughness properties of the surface and by the stability and turbulence of the atmospheric surface layer. The resistance catena (Figure 4-3) is less useful in this size range and, in some treatments, has been abandoned entirely (e.g., Erisman et al., 1994; equation 4-4). Impaction and interception dominate over diffusion, and the V_d is considerably (up to two orders of magnitude; Figure 4-2) lower than for particles either smaller or larger (Shinn, 1978). The deposition velocity may be parameterized in this size range as a function of friction velocity,

$$V_d = (a/b)u^*, \quad (4-4)$$

where a depends on atmospheric stability and b depends on surface roughness (Wesely et al., 1985; Erisman et al., 1994). Similar formulations have been presented in terms of turbulence (standard deviation of wind direction) and wind speed (e.g., Wesely et al., 1983), both determinants of u^* .

Deposition of particles between 1 and 10 μm in diameter, including the coarse mode of PM_{10} , is strongly dependent on particle size (Shinn, 1978). Larger particles within this size range are collected more efficiently at typical wind speeds than are smaller particles (Clough, 1975), suggesting the importance of impaction. Impaction is related to wind speed, the square of particle diameter, and the inverse of receptor diameter as a depositing particle fails to follow the streamlines of the air in which it is suspended around the receptor. When particle trajectory favors a collision, increasing either wind speed or the ratio of particle size to receptor cross section makes collision nearly certain. Similarly, as these parameters become very small, the probability of collision becomes negligible. However, the shape parameter for the more common range of situations between these extremes remains poorly characterized (Peters and Eiden, 1992; Wiman and Ågren, 1985).

As particle size increases above 1 μm , deposition is governed increasingly by sedimentation (Figure 4-5) and decreasingly by turbulence and impaction. Particles between ~ 10 and 24 μm (Gallagher et al., 1988) are both small enough to be transported efficiently by

turbulent eddies to the surface and large enough to impact with sufficient momentum to overcome boundary layer effects. These particles deposit highly efficiently and relatively independently of particle size. Deposition of the largest suspended particles (e.g., $> 50 \mu\text{m}$) is governed, through sedimentation and the corresponding terminal settling velocity, V_s , almost entirely by size. These particles are not transported efficiently by small-scale eddies near the surface.

Theoretically based models for predicting particle deposition velocities have been published by Bache (1979a,b), Davidson et al. (1982), Noll and Fang (1989), Slinn (1982), and Wiman (1985). These models deal primarily with low canopies or individual elements of canopy surfaces. Wiman and Ågren (1985) developed an aerosol deposition model that specifically treats the problem of particle deposition to forests where turbulence plays a particularly important role, especially at roughness transitions such as forest edges. They found that deposition of supermicron particles is controlled by complex interactions among particle size and concentration, forest structure, and aerodynamics; whereas deposition of fine (submicron) particles is controlled by particle concentration and forest structure.

Empirical measurements of V_d for fine particles under wind tunnel and field conditions are often several-fold greater than predicted by available theory (Unsworth and Wilshaw, 1989). A large number of transport phenomena, including streamlining of foliar obstacles, turbulence structure near surfaces, and various phoretic transport mechanisms remain poorly parameterized in current models. The discrepancy between measured and predicted values of V_d may reflect such model limitations or experimental limitations in the specification of the effective size and number of receptor obstacles, as suggested by Slinn (1982).

Available reviews (Davidson and Wu, 1990; McMahon and Denison, 1979; Nicholson, 1988; Sehmel, 1980; Slinn, 1982; U.S. Environmental Protection Agency, 1982, 1996a) suggest the following generalizations: (1) particles $> 10 \mu\text{m}$ exhibit a variable V_d between ≈ 0.5 and 1.1 cm/s depending on friction velocities; whereas a minimum particle V_d of 0.03 cm/s exists for particles in the size range 0.1 to $1.0 \mu\text{m}$; (2) the V_d of particles is approximately a linear function of friction velocity; and (3) deposition of particles from the atmosphere to a forest canopy is from 2 to 16 times greater than deposition in adjacent open terrain (i.e., grasslands or other vegetation of low stature).

Leaf Surface Effects on V_d . The term r_c (Equation 4-3) reflects the chemical, physical, or physiographical characteristics of the surface that govern its ability to capture, denature, or otherwise remove particulate material from the atmospheric surface layer. For gases, relevant surface properties involve the physiological state of the vegetation, including stomatal opening and mesophyll antioxidant activity, as well as the chemical reactivity of the exposed surface with the specific gas. For particles, relevant surface properties involve stickiness, microscale roughness, and cross-sectional area. These properties determine the probability of impaction and bounce (e.g., Shinn, 1978). The chemical composition of PM is not usually a primary determinant of deposition velocity. At the microscopic scale where Van der Waals forces may determine particle bounce and reentrainment, the chemical properties of both surface and particle may be significant but remain poorly understood.

Stickiness may itself depend on previous deposition of deliquescent particles that prolong leaf wetness, on the wettability of foliar surfaces, and on the presence of sticky residues such as honeydew deposited by aphids. These factors increase deposition by decreasing bounce-off, wind reentrainment, and, to some extent, wash-off by precipitation.

The distribution of particles on, and the efficiency of deposition to, vegetation also varies with leaf shape and plant part. Particles are more prevalent on the adaxial (upper surface facing twig) surface than on the abaxial (lower surface away from the twig). Peripheral leaf areas tend to be the cleanest, with most particles accumulating in the midvein, central portion of leaves. The rough area surrounding the stomatal pores was not found to be a preferential site for particle deposition or retention (Smith and Staskawicz, 1977). Most particles were located near veins with smaller particles localized on the trichomes. The greatest particle loading on dicotyledonous leaves is frequently on the adaxial (upper) surface at the base of the blade, just above the petiole junction. Precipitation washing probably plays an important role in this distribution pattern. Lead particles accumulated to a greater extent on older than younger needles and twigs of white pine, indicating that wind and rain were insufficient to fully wash the foliage. Fungal mycelia (derived from windborne spores) were frequently observed in intimate contact with other particles on other leaves, which may reflect minimal reentrainment of the spore due to shelter by the particles, mycelia development near sources of soluble nutrients provided by the particles, or simply codeposition (Smith and Staskawicz, 1977). This pattern is significant and could yield further insight into deposition mechanisms.

Leaves with complex shapes collect more particles than those with regular shapes, but conifer needles are more effective than broad leaves in accumulating particles. The edge to area ratio (Woodcock, 1953) is also a key determinant of salt deposition to individual artificial leaves. A strong negative correlation was observed under wind tunnel conditions between the area of individual leaves and the deposition of coarse particles (Little, 1977). Small twigs and branches were more effective particle collectors than were large branches and trunks of trees (Smith, 1984). Lead particles accumulated 20-fold more on woody stems than on needles of white pine (*Pinus strobus*), even though needles displayed a 10-fold greater total surface area (Heichel and Hankin, 1976). Deposition was heaviest at the tips of individual leaves.

Rough, pubescent broadleaf discs collected coarse (5.0 μm) particles up to 7-fold more efficiently than glabrous leaf discs (Little, 1977). Laminae, petioles, and stems all differed in collection efficiency. Pubescent leaves of sunflower (*Helianthus annuus*) collected coarse particles nearly an order of magnitude more efficiently than the glabrous leaves of tulip poplar (*Liriodendron tulipifera*) under wind tunnel conditions (Wedding et al., 1975). The rough pubescent leaves of nettle (*Urtica dioica*) were more effective in capturing coarse PM_{10} than were the densely tomentose leaves of poplar (*Populus alba*) or smooth leaves of beech (*Fagus sylvatica*).

Conifer needles are more efficient than broad leaves in collecting particles by impaction. This reflects the small cross section of the needles relative to the larger leaf laminae of broadleaves and the greater penetration of wind into conifer than broadleaf canopies. Conifers were more effective in removing coarse ($\approx 20 \mu\text{m}$) particles of ragweed pollen from the atmosphere than were broadleaf trees (Steubing and Klee, 1970) and in intercepting even coarser rain particles (Smith, 1984). Conifers were also more effective in retaining and accumulating particles against reentrainment by wind and removal by rain, particularly on foliar surfaces where they are likely to be most biologically active. Seedlings of white pine (*Pinus strobus*) and oak (*Quercus rubra*) initially retained between a quarter (pine) and a third (oak) of very coarse (88 to 175 μm) ^{134}Cs -tagged quartz particles applied under field conditions (Witherspoon and Taylor, 1971). After 1 h, the pine retained over 20% of the ^{134}Cs -tagged particles; whereas the oak retained only approximately 3%. Long-term retention of the particles was concentrated at the base of the fascicles in pine and near the surface roughness caused by the vascular system on leaves of oak. The sheltered locations available in the conifer foliage contribute substantially to

greater retention of particles. For similar reasons, grasses also are efficient particle collectors (Smith and Staskawicz, 1977) with long-term retention mostly in the ligule and leaf sheath.

Because of the strong relationship between particle size and deposition, the sharply increasing humidity gradient near transpiring foliar surfaces may cause hygroscopic particles to behave, at the immediate surface, as larger particles than reflected in ambient measurements at reference heights. This needs to be further considered (Wesely and Hicks, 2000). Recent deposition models (Ruijgrok et al., 1997; Zhang et al., 2001) have introduced the role of ambient humidity but lack sufficient emphasis on the role of vegetation itself in modifying humidity near the surface. This may be significant, as the size of a dry, 0.5 μm particle of ammonium sulfate may increase to about 3.5 μm at saturated humidity (Ruijgrok et al., 1997). Kinetic analyses have suggested that full (95 to 100%) equilibration to the new diameter will occur during the deposition process in ambient humidity.

Wind tunnel studies also demonstrated equivalent deposition properties for 3.4 μm particles of dense lead chloride and the 6.8 μm particles of the less dense uranine dye. These particles were shown to be aerodynamically equivalent, substantiating the use of aerodynamic diameter as a classification parameter for particle deposition.

Canopy Surface Effects on V_d . In general, surface roughness increases particulate deposition. As a result, V_d is typically greater for a forest than for a field or nonwoody wetland and greater for a field than for a water surface. The contrasting transport properties and deposition velocities of different size particles lead to predictable patterns of deposition. For coarse particles, the upwind leading edges of forests, hedge rows, and individual plants, as well as of individual leaves, are primary sites of deposition. Impaction at high wind speed and the sedimentation that follows the reduction in wind speed and carrying capacity of the air in these areas lead to preferential deposition of larger particles.

Air movement is slowed in proximity to vegetated surfaces. Resulting log profiles of wind and pollutant concentrations in the near-surface turbulent boundary layer above canopies reflect surface characteristics of roughness length, friction velocity, and displacement height. Plasticity, streamlining, and oscillations of foliar elements also alter the aerodynamic roughness and the level of within-canopy turbulence. Canopies of uneven age or with a diversity of species are typically aerodynamically rougher and receive larger inputs of pollutants than do smooth, low, or

monoculture vegetation (Garner et al., 1989; Wiman and Ågren, 1985). Canopies on slopes facing the prevailing winds and individual plants on the windward edges of discontinuities in vegetative cover over which roughness increases receive larger inputs of pollutants than more sheltered, interior canopy regions. For example, some 80% of coarse particulate sea salt was deposited on the upwind edge of a hedgerow (Edwards and Claxton, 1964), and the ragweed pollen concentration was reduced by 80% within 100 m of the upwind edge of a forest (Neuberger et al., 1967).

Beier et al. (1992) and Beier (1991) discussed two methods for estimating the dry deposition of base cations to forest edges: (1) a difference method between measured precipitation and throughfall concentrations of base cations, and (2) a calculation method based on known ratios of Na^+ deposition in wet and dry forms (Ulrich, 1983). A combination of these two approaches has produced the best estimates of SO_4^{2-} , Ca^{2+} , Mg^{2+} , and K^+ particle deposition. Using these methods, particulate SO_4^{2-} (Beier, 1991) and particulate Ca^{2+} , Mg^{2+} , and K^+ (Beier et al., 1992) were found to decrease by an order of magnitude from the forest edge to the forest interior. A number of authors also have shown that particle deposition is elevated at forest edges when compared to a uniform forest canopy (Draaijers et al., 1988; Grennfelt, 1987; Lindberg and Owens, 1993), and Draaijers et al. (1992) reported that differences are likely to exist between forest types because of variable canopy structure. Draaijers et al. (1988) further emphasized that enhanced particle deposition at or near forest edges is strongly dependent on the velocity and wind direction during observations.

The factors leading to horizontal gradients are confounded by time- and distance-related sedimentation. For example, geologic dust (mostly around 7 μm aerodynamic diameter) collects on stems of wild oats (*Avena* spp.; Davidson and Friedlander, 1978) and on eastern white pine (Heichel and Hankin, 1972; Smith 1973) downwind of roadways. Rapid sedimentation of coarse crustal particles suggests that potential direct effects may be restricted to roadway margins, forest edges, and, because of the density of unpaved roads in agricultural areas, crop plants.

Simulated deposition to an ecologically complex, mixed canopy was considerably higher than to a pure spruce stand in which most of the leaf area was concentrated in regions of low wind speed. Limitations to the application of these models to predict deposition over large regions include an incomplete understanding both of the nature of microscopic particle-surface

interactions and of the effects of complex terrain and species composition on macroscopic transport processes.

Macroscopic turbulent transport processes, related to r_a at successive layers through the canopy, can be separated from microscopic processes related to r_b and r_c (or r_{cp}) at each deposition surface (e.g., Peters and Eiden, 1992; Wiman and Lannefors, 1985). The macroscopic approach deals with deposition as the product of a turbulent diffusion coefficient and a given concentration (C_z) at each canopy layer, both of which vary with particle size and with height (z) in the canopy. The microscale parameters involve those factors that determine absorption of a particle at each surface as captured imperfectly by r_c . Shelter effects caused by the crowding of foliar elements within the canopy can be ignored if the wind speed within each canopy layer is specified. This approach requires knowledge of the vertical distribution of particle concentration and foliage density in the canopy airspace along with profiles of wind speed or turbulence.

Once introduced into a forest canopy, elements associated with coarse particles tend to decrease markedly with canopy depth; whereas elements associated with fine particles do not (Lovett and Lindberg, 1992). Trace elements and alkaline-earth elements are enriched below the canopies of both southern (Lindberg et al., 1986) and northern (Eaton et al., 1973) hardwood forests. Vertical gradients in concentration of coarse particles and of elements associated with coarse particles were observed in a mixed conifer/birch forest canopy (Wiman and Lannefors, 1985; Wiman et al., 1985) and in a mixed oak forest (e.g., Ca^{2+} , Figure 4-6a; Lovett and Lindberg, 1992). The highly reactive gas, HNO_3 , also exhibited a vertical gradient but with a steep decline at the top of the canopy (Figure 4-6b). Lovett and Lindberg (1992) studied concentration profiles of various gases and particles within an closed canopy forest and concluded that coarse particle concentrations associated with elements like Ca^{2+} would decrease markedly with depth in the canopy, but they found only minor reductions with depth in the concentrations of fine aerosols containing SO_4^{2-} , NH_4^+ , and H^+ . These data suggest that all foliar surfaces within a forest canopy are not equally exposed to particle deposition: upper canopy foliage tends to receive maximum exposure to coarse and fine particles, but foliage within the canopy tends to receive primarily fine aerosol exposures. Fine-mode particles (e.g., sulfate, Figure 4-6c) and unreactive gases typically do not exhibit such vertical profiles, suggesting that their uptake is smaller in magnitude and more evenly distributed throughout the canopy.




Figure 4-6. Vertical stratification of diverse, chemically speciated particles in a mixed oak forest. Nitric acid vapor, as a highly reactive, nonparticulate species, is shown for comparison. The horizontal dotted line indicates canopy height, and Δ indicates the percent depletion from above to below the canopy.

Source: Based on data from Lovett and Lindberg (1992).

In multilayer canopies, simultaneous reentrainment and deposition may effectively uncouple deposition from local concentration. Polydisperse size distributions of many chemical species effectively prevent the use of a single estimate of V_d for any element if highly accurate results are required.

Although gradients (Figures 4-6 and 4-7) may be related to local V_d within the canopy (Bennett and Hill, 1975), the absence of a gradient may reflect either low or very high rates of




Figure 4-7. The relationship between particle size and concentration below a spruce canopy with wind velocity at a height of 16.8 m equaling 5 m/s.

Source: Modified from Peters and Eiden (1992).

deposition relative to turbulent replenishment from above the canopy (Tanner, 1981). Below- or within-canopy emissions may confound interpretation of vertical gradients. Linear gradients of the gaseous pollutants hydrogen fluoride (HF) and O_3 reflected large uptake rates; whereas small gradients in NO suggested little uptake by foliage (Bennett and Hill, 1973, 1975). However, soil efflux of NO could have complicated the latter interpretation. The lack of a vertical gradient and a peak near the top of the active canopy in particulate K^+ (Figure 4-6d) was interpreted as evidence for a biogenic source within the deciduous forest canopy with moderate rates of deposition (Lindberg et al., 1986; Lovett and Lindberg, 1992).

The size dependence of this vertical stratification of particulate concentration (see Figure 4-7) is reflected in current simulation models (Wiman et al., 1985; Peters and Eiden, 1992). The model of Wiman and Ågren (1985) predicts a uniform vertical distribution of fine-mode particles and a pronounced vertical gradient of coarse-mode particles which is in agreement with observations (Lovett and Lindberg, 1992).

Simulation of the horizontal deposition patterns at the windward edge of a spruce forest downwind of an open field with the canopy between 1 and 25 m above the ground indicated that deposition was maximal at the forest edge, where wind speed and impaction were greatest. Simulation of the vertical deposition pattern was more complex. Deposition was not greatest at the top of the canopy where wind speed was highest, but at $z = 0.75 h$, where the balance between leaf area (obstacles for impaction) and wind speed (momentum for impaction) was optimal, although neither parameter alone was maximal. Simulated deposition in this spruce forest increased considerably with increasing LAI at the forest edge, where wind speed was insensitive to LAI but the number of obstacles increased. Inside the forest, where both wind speed and impaction increasingly were attenuated by increasing LAI, deposition increased only marginally in spite of the increase in obstacle frequency.

To scale surface-specific measurements of particle deposition to forest or crop canopies, conversions of the following type have been suggested:

$$V_{d,\text{canopy}} = V_{d,\text{surface}} * \text{scaling factor}, \quad (4-5)$$

with empirical scaling factors proposed by Lindberg et al. (1988).

To appropriately scale surface-specific measurements of particle deposition to landscapes, one must consider the complexity of grassland, crop, and forest canopies in order to avoid serious over- or underestimates of particle deposition. Individual species exposed to similar ambient concentrations may receive a range of particulate loading that is more closely related to foliar damage than the ambient concentration (Vora and Bhatnagar, 1987).

Both uptake and release of specific constituents of PM may occur within a single canopy (e.g., K^+ ; Lovett and Lindberg, 1992). The leaf cuticular surface is a region of dynamic exchange processes occurring through leaching and uptake. Exchange occurs with epiphytic microorganisms and bark and through solubilization and erosion of previously deposited PM. Vegetation emits a variety of particles and particulate precursor materials. Terpenes and isoprenoids predominate and, on oxidation, become condensation nuclei for heterogeneous particle formation. Salts and exudates on leaves and other plant parts are continually abraded and suspended as particles, as are plant constituents from living and dead foliage (Rogge et al., 1993a). Soil minerals, including radioactive strontium, nutrient cations and anions, and trace

metals are transferred to the active upper foliage and then to the atmosphere in this way. Although not representing a net addition to an ecosystem, particle release from vegetation is a mechanism for redistributing chemical pollutants derived from the soil or prior deposition within a canopy, potentially enhancing direct effects and confounding estimates of V_d .

Range of Deposition Velocity

As noted in an earlier criteria document (U.S. Environmental Protection Agency, 1982) and in McMahon and Denison (1979), estimates of V_d for PM_{10} particles to vegetation are variable and suggest a minimum between 0.1 and 1.0 μm , as predicted using first principles (Monteith and Unsworth, 1990; Sehmel, 1980). Determinations in wind tunnels with passive collectors and micrometeorological methods tend to converge in this range. The range of V_d for sulfate from passive collectors was found to be from 0.147 to 0.356 cm/s; and, from eddy covariance techniques, a mean V_d of 0.27 cm/s was observed (Dolske and Gatz, 1984). Micrometeorological techniques over grass (Wesely et al., 1985); indirect, inert collector techniques within an oak forest (Lindberg and Lovett, 1985); and many other empirical determinations (e.g., McMahon and Denison, 1979; Table 4-4) generally support this range. Over aerodynamically smooth snow (Duan et al., 1988; Table 4-5), measurements of V_d were an order of magnitude smaller. Very coarse particles, often non-size-specified primary geologic material, frequently exhibit V_d greater than 1.0 cm/s (e.g., Clough, 1975). The increase in V_d with decreasing size below 0.1 μm is probably hidden in most empirical determinations of V_d because the total mass in this fraction is very small despite the large number of individual particles. Table 4-6 shows published estimates of V_d with variability estimates for fine particles of specified aerodynamic diameters dominated by a range of chemical species.

Ibrahim et al. (1983) evaluated the deposition of ammonium sulfate particles to a range of surfaces and found that particles having a mean diameter of 0.7 μm had deposition velocities ranging from 0.039 to 0.096 cm/s. Larger particles (having mean diameters of 7 μm) had greater deposition velocities (between 0.096 and 0.16 cm/s). The authors further concluded that the hygroscopic nature of the sulfate particle could increase its size and enhance deposition near sources of water (e.g., snow). Using eddy correlation approaches, Hicks et al. (1989) found a mean daily V_d for sulfur-containing PM to be 0.16 cm/s. However, they suggested that the V_d value could be as high as 1 cm/s during the day and near zero at night.

TABLE 4-4. REPORTED MEAN DEPOSITION VELOCITIES (V_d) FOR SULFATE, CHLORINE, NITRATE, AND AMMONIUM-ION-CONTAINING PARTICLES

Chemical Species/ Surface	V_d (cm/s) ^a	Method	Reference
SO₄²⁻			
Inert plates	0.13	Extraction	Lindberg and Lovett (1985)
Inert plates	≈0.14	Extraction	Lindberg et al. (1990)
Inert plates	0.14	Extraction	Davidson and Wu (1990) ^a
Inert bucket	0.51	Extraction	Davidson and Wu (1990) ^a
Foliage	0.29	Extraction	Davidson and Wu (1990) ^a
Chaparral	0.15	Extraction	Bytnerowicz et al. (1987a)
Grass canopy	0.10	Gradient	Allen et al. (1991)
Grass canopy	0.07	Gradient	Nicholson and Davies (1987)
Pine foliage	0.07	Extraction	Wiman (1981)
Plant canopies	0.50	Gradient	Davidson and Wu (1990) ^a
Grass canopy	0.22	Eddy covariance	Weseley et al. (1985)
Cl⁻			
Inert bucket	3.1	Extraction	Dasch and Cadle (1985)
Inert bucket	5	Extraction	Dasch and Cadle (1986)
Beech canopy	1	Throughfall	Höfken et al. (1983)
Spruce canopy	1.9	Throughfall	Höfken et al. (1983)
NO₃⁻			
Inert plates	0.4-2	Extraction	Lindberg et al. (1990)
<i>Ceanothus</i>	0.4	Extraction	Bytnerowicz et al. (1987a)
<i>Glycine max</i>	0.24	Extraction	Dolske (1988)
<i>Ligustrum</i>	0.1-0.5	Extraction	John et al. (1985)
<i>Quercus</i>	0.7-1.1	Extraction	Dasch (1987)
<i>Quercus</i> summer	0.55	Throughfall	Lovett and Lindberg (1984)
<i>Quercus</i> winter	0.7	Throughfall	Lovett and Lindberg (1984)
<i>Quercus</i> summer	0.3	—	Lovett and Lindberg (1986)
<i>Quercus</i> winter	0.1	—	Lovett and Lindberg (1986)
<i>Pinus</i>	0.5-1.3	Extraction	Dasch (1987)
<i>Pasture</i>	0.7-0.8	Gradient	Huebert et al. (1988)
<i>Ulmus</i>	1.1	Extraction	Dasch (1987)
NH₄⁺			
<i>Calluna/Molina</i>	0.18	Gradient	Duyzer et al. (1987)
<i>Ceanothus</i>	0.4	Extraction	Bytnerowicz et al. (1987a)
<i>Kalmia</i>	0.03-0.14	Extraction	Tjepkema et al. (1981)
<i>Pinus</i>	0.01-0.06	Extraction	Dasch (1987)

^a These data represent the mean of data by measurement technique as reported in the cited reference. The reader is referred to the referenced articles for information on the specific cations contributing to the means.

TABLE 4-5. REPRESENTATIVE EMPIRICAL MEASUREMENTS OF DEPOSITION VELOCITY (V_d) FOR PARTICULATE DEPOSITION

V_d $x \pm SE$ (cm/s)	Particle Size (μm)	Method	Reference
0.034 \pm 0.014 0.021 \pm 0.005	0.15-0.30 0.5-1.0	Eddy covariance with optical counter, flat snow surface	Duan et al. (1988)
0.1 \pm 0.03	37987	Profile, fine SO_4^{2-} , short grass	Allen et al. (1991)
0.22 \pm 0.06	0.1-2.0	Eddy covariance with flame photometer plus denuder, 40-cm grass, fine SO_4^{2-}	Wesely et al. (1985)
0.13 \pm 0.02 0.75 \pm 0.24 1.1 \pm 0.1	(SO_4^{2-}) (K^+) (Ca^{2+})	Inert surface collectors (petri dish) in oak forest	Lindberg and Lovett (1985)
0.9 2.5 9.4	2.75 5.0 8.5	Wind tunnel to pine shoots; polystyrene beads; within-canopy wind speed, 2.5 m/s	Chamberlain and Little (1981)

TABLE 4-6. REPORTED MEAN DEPOSITION VELOCITIES FOR POTASSIUM (K^+), SODIUM (Na^+), CALCIUM (Ca^{2+}), AND MAGNESIUM (Mg^{2+}) BASE CATION CONTAINING PARTICLES

Chemical Species/Surface	V_d (cm/s)	Method	Reference
K^+			
Inert plates	0.75	Extraction	Lindberg and Lovett (1985)
Inert bucket	0.51-2.4	Extraction	Dasch and Cadle (1985)
Na^+			
Inert bucket	1.7-2.9	Extraction	Dasch and Cadle (1985)
Inert plate	0.8-8.2	Extraction	
Ca^{2+}			
Inert plates	1.1	Extraction	Lindberg and Lovett (1985)
Inert plates	\approx 2	Extraction	Lindberg et al. (1990)
Inert bucket	1.7-3.2	Extraction	McDonald et al. (1982)
Mg^{2+}			
Inert bucket	1.1-2.7	Extraction	Dasch and Cadle (1985)

Lindberg et al. 1990 found a wide discrepancy between deposition velocities for NO_3^- between study sites in Oak Ridge, TN (≈ 2 cm/s) and Göttingen, Germany (≈ 0.4 cm/s). They suggest that the increased V_d at Oak Ridge could be explained by the primary occurrence of NO_3^- in coarse particles that exhibit greater V_d than fine particles (Davidson et al., 1982). Large values of V_d for base-cation-containing particles (> 1 cm/s) suggest their occurrence in coarse particles (Lindberg and Lovett, 1985).

The several attempts to estimate V_d for SO_4^{2-} , NO_3^- , and NH_4^+ with the throughfall mass balance approach (Davidson and Wu, 1990; Gravenhorst et al., 1983; Höfken and Gravenhorst, 1982) have produced higher V_d values that are considered suspect. They have not been included in Tables 4-4 and 4-6. Overestimates of V_d for SO_4^{2-} and NO_3^- particles derived from throughfall mass balance approaches may be the result of gaseous SO_2 and HNO_3 gaseous deposition to foliar surfaces (Lindberg and Lovett, 1985). A similar contribution of NH_3 deposition may lead to erroneously high V_d values for NH_4^+ when the throughfall method is attempted in areas of high NH_3 concentrations. Dolske's (1988) reported V_d values for NO_3^- deposition to soybean ranged from 0.4 to 31 with a mean of 0.24 cm/s. However, because Dolske's leaf extraction measurements included a component of HNO_3 vapor, the V_d values may represent more than deposition caused by aerosol nitrate alone.

The quantitative importance of dry particulate deposition depends upon the chemical species, topography, precipitation regime, and surface characteristics, including vegetation properties. Across the diverse landscapes of the Integrated Forest Study (Johnson and Lindberg, 1992a), the relative contribution of dry deposition for Ca^{2+} ranged from about 0% to nearly 90% (Figure 4-8). In contrast, for S the total range was from just over 0% to about 30%. An average for these forest systems demonstrates that deposition of (usually coarse) base cations was nearly 50% by dry particulate deposition (Figure 4-9). Both N and S were around 15%. While the relative significance of dry particle deposition varies from site to site, it cannot be excluded from the analysis at any site.

In a recent review, Wesely and Hicks (2000) concluded that a comprehensive understanding of particulate deposition remains a distant goal. In general, there is only modest confidence in available particulate deposition parameterizations at this time, although recent experimental and theoretical efforts to improve this situation have been made (e.g., Erisman et al., 1997, and companion articles; Zhang et al., 2001; Kim et al., 2000).




Figure 4-8. Range of percent of total deposition delivered in the dry particulate fraction, across the sites of the Integrated Forest Study.

Source: Johnson and Lindberg (1992a).




Figure 4-9. Contribution of particulate deposition to total deposition of nitrogen, sulfur, and base cations.

Source: Modified from Johnson and Lindberg (1992a); Lovett (1992); Lovett and Lindberg (1993); Lindberg et al. (1990); Kelly and Meagher (1986).

The successful treatment of dry deposition of gaseous pollutants (e.g., SO₂ and O₃) linking turbulent deposition to surface physiological properties has allowed incorporation of these models into landscape and regional scale models. This has allowed gaseous deposition to be

scaled up for purposes of atmospheric chemistry and vegetation damage assessment. These advances have not been matched by progress in deposition of PM. A serious remaining impediment is the lack of suitable techniques for measuring deposition of heterogeneous and polydisperse particles, such as the gradient and eddy covariance techniques that are used for gaseous pollutant species. As with gaseous pollutants, parameterization of particle deposition in hilly terrain, to patchy surfaces (small agricultural fields, forest edges), and under extremely windy conditions, remains to be fully developed. These limitations must be addressed before a full accounting of regional PM effects on vegetation in natural and managed ecosystems can be achieved.

Occult Deposition

Gaseous pollutant species may dissolve in the suspended water droplets of fog and clouds. The stability of the atmosphere and persistence of the droplets often allow a gas/liquid phase equilibrium to develop. This permits the use of air mass history or ambient concentrations of specific pollutants to estimate fog or cloud water concentrations. Further estimates of the deposition velocity of the polluted droplets allows calculation of depositional fluxes. Unfortunately, interception of fog or cloud droplets by plant parts or other receptor surfaces remains difficult to predict and to measure. Fog formation influences the total atmospheric burden and deposition of particulate matter (Pandis and Seinfeld, 1989) by accreting and removing particles from the air, by facilitating particle growth through aqueous oxidation reactions, and by enhancing deposition as noted. Aqueous condensation may occur onto preexisting fine particles, and such particles may coalesce or dissolve in fog or cloud droplets. Material transported in fog and cloud water and intercepted by vegetation escapes detection by measurement techniques designed to quantify either dry or wet deposition; hence it is hidden (i.e., “occult”) from the traditional measurements.

Occult deposition of fog and cloud droplets is by impaction and gravitational settling in concert with the instantaneous particle (droplet) size. This mode of deposition may be significant at high elevation sites, particularly near the base of orographic clouds. While coastal fog may be generally more pristine than high-elevation continental fogs, this is not true in areas subject to inland radiation fog (e.g., the increasingly polluted San Joaquin Valley of Central California) and in areas where the marine layer is advected through a highly polluted urban area

(e.g., the Los Angeles basin of Southern California). Fog water in these areas may be at least as contaminated as that occurring at higher elevations.

Low elevation radiation fog has different formation and deposition characteristics than high elevation cloud or coastal fog water droplets. A one-dimensional deposition model has recently been described for a radiation fog episode (Von Glasow and Bott, 1999). A substantially greater concentration of key polluting species (e.g., NO_3^- , SO_4^{2-} , organics) may be observed in smaller than in larger droplets in fog (Collett et al., 1999). Acidity differences exceeding 1 pH unit were also observed in the San Joaquin Valley winter radiation fog, with smaller particles being more acidic. This has implications for aqueous phase oxidation of sulfur and nitrogen compounds, in particular, while sulfur oxidation by ozone (the dominant reaction in this environment even during winter) is well known in typically acidic fog droplets. However, the alkaline larger droplets in the San Joaquin Valley could lead to greater nitrate production through aqueous ozonation reactions (Collett et al., 1999). The size class distinctions have substantial implications for deposition of particulate pollutant species in the fog droplets due to the larger V_d for impaction and occult deposition of the larger fog particles.

Acidic cloud water deposition has been associated with forest decline in industrialized areas of the world (Anderson et al., 1999). Clouds can contain high concentrations of acids and other ions. Cloud water typically is 5 to 20 times more acid than rain water. This can increase pollutant deposition and increases the exposure of vegetation and soils at high-elevation sites by more than 50% when compared with rainfall and dry deposition.

The widespread injury to mountain forests documented in West Germany since the 1970s and other parts of Europe, and more recently in the Appalachian Mountains, has been attributed to cloud water exposure reducing the cold tolerance of red spruce. Forest injury also has been attributed to increased leaching of cations and amino acids and increased deposition of nitrogen and aluminum toxicity resulting from acidic deposition and the combined effect of acidic precipitation, acid fog, oxidants, and heavy metals (Anderson et al., 1999). The Mountain Acid Deposition Program (MADPro) was initiated in 1993 as part of the Clean Air Status and Trends Network (CASTnet). MADPro monitoring efforts focused on the design and implementation of an automated cloud water collection system in combination with continuous measurement of cloud liquid water content (LWC) and meteorological parameters relevant to the cloud deposition process.

Using the MADPro automated cloud-water collectors at three selected mountain sites (Whiteface Mt., NY; Whitetop Mt., Va; and Clingman's Dome, TN), samples taken hourly from nonprecipitating clouds during nonfreezing seasons of the year from 1994 to 1997 and promptly analyzed for pH, conductivity, and the concentrations of dissolved ions gave an indication of exposures at each of the three sites. Cloud LWC was measured at each site. The mean cloud water frequencies and LWC were higher at Whiteface Mountain, NY than in the southern Appalachians. The four most prevalent ions found in cloud water samples, in order of decreasing concentrations, were usually SO_4^{2-} , H^+ , NH_4^+ , and NO_3^- . The concentrations of these ions tended to co-vary within cloud events and typically there was an inverse relationship between LWC of the cloud and ionic concentration of the cloudwater. Highest ionic concentrations were seen in mid-summer. Ionic concentrations of samples from southern sites were significantly higher than samples from Whiteface Mountain, but further analysis indicated that this was at least partly due to north to south differences in the LWC of clouds (Anderson et al., 1999).

Several factors make occult deposition particularly effective for the delivery of dissolved and suspended materials to vegetation. Concentrations of particulate-derived materials are often many-fold higher in cloud or fog water than in precipitation or ambient air in the same area due to orographic effects and from gas-liquid partitioning coefficients of specific chemical species. Fog and cloud water deliver PM in a hydrated (and, therefore, bioavailable) form to foliar surfaces. Previously dry-deposited PM may also become hydrated through delinquency or by dissolution in the film of liquid water from fog deposition. The presence of fog itself maintains conditions of high relative humidity and low radiation, thus reducing evaporation and contributing to the persistence of these hydrated particles on leaf surfaces. Deposition of fog water is very efficient (Fowler et al., 1991) with a V_d (fog 10 to 24 μm ; Gallagher et al., 1988) essentially equal to the aerodynamic conductance for momentum transfer $(r_a)^{-1}$. This greatly enhances deposition by sedimentation and impaction of submicron aerosol particles that exhibit very low V_d prior to fog droplet formation (Fowler et al., 1989). The near equivalence of V_d and $(r_a)^{-1}$ simplifies calculation of fog water deposition and reflects the absence of vegetative physiological control over surface resistance. Fog particles outside this size range may exhibit V_d below $(r_a)^{-1}$. For smaller particles, this decline reflects the increasing influence of still air and boundary layer effects on impaction as particle size and momentum decline. For larger particles,

momentum is sufficient to overcome these near surface limitations, but V_d may decline as turbulent eddy transport to the surface becomes inefficient with increasing inertia (Gallagher et al., 1988). The deposition to vegetation for PM in fog droplets is directly proportional to wind speed, droplet size, concentration, and fog density (liquid water content per volume air) although the latter two may be inversely related. In some areas, typically along foggy coastlines or at high elevations, occult deposition represents a substantial fraction of total deposition to foliar surfaces (Fowler et al., 1991; Figure 4-2).

4.2.2.3 Magnitude of Deposition

Dry deposition of PM is most effective for coarse particles including primary geologic material and for elements such as iron and manganese. Wet deposition is most effective for fine particles of atmospheric (secondary) origin (e.g., nitrogen and sulfur, Table 4-6) and elements such as cadmium, chromium, lead, nickel, and vanadium (Reisinger, 1990; Smith, 1990a,b,c; Wiman and Lannefors, 1985). The occurrence of occult deposition is more restricted. The relative magnitudes of the different deposition modes varies with ecosystem type, location, elevation, and chemical burden of the atmosphere. For the Walker Branch Watershed, a deciduous forest in rural eastern Tennessee, dry deposition constituted a major fraction of the total annual atmospheric input of cadmium and zinc ($\approx 20\%$), lead ($\approx 55\%$), and manganese ($\approx 90\%$). Whereas wet deposition fluxes during precipitation events exceeded dry deposition fluxes by one to four orders of magnitude (Lindberg and Harriss, 1981), dry deposition was nearly continuous. Immersion of high-elevation forests in cloudwater may occur for 10% or more of the year, significantly enhancing transfer of PM and dissolved gases to the canopy. Occult deposition in the Hawaiian Islands dominated total inputs of inorganic N (Heath and Huebert, 1999). Much of this N was volcanically derived during the generation of volcanic fog in part through reactions with seawater. In this humid climate, the dominance of occult rather than wet deposition is notable.

High-elevation forests receive larger particulate deposition loadings than equivalent low elevation sites. Higher wind speeds enhance the rate of aerosol impaction. Orographic effects enhance rainfall intensity and composition and increase the duration of occult deposition. Coniferous species in these areas with needle-shaped leaves also enhance impaction and retention of PM delivered by all three deposition modes (Lovett, 1984).

In more arid regions, such as the western United States, the importance of dry deposition may be larger. In the San Gabriel Mountains of southern California, for example, while annual deposition of SO_4^{2-} (partly of marine origin) was dominated by wet deposition (Fenn and Kiefer, 1999), deposition of NO_3^- was dominated by dry deposition, as was that of NH_4^+ at two of three sites. Similarly, at a series of low elevation sites in southern California (Padgett et al., 1999), dry deposition of NO_3^- was dominated by dry deposition. In both cases, however, the contribution of gaseous HNO_3 was probably substantial.

Nitrates, Sulfates, and Cations

Much particulate sulfate and nitrate is found on particles in the 0.1- to 1.0- μm size range (U.S. Environmental Protection Agency, 1982). However, most sulfate and nitrate, base cation and heavy metal inputs to forested ecosystems result from the deposition of larger particles (Chapter 2) (Lindberg and Lovett, 1985; Lindberg et al., 1982). The influence of aerodynamic diameter is particularly critical for nitrogen species, because they exist as a wide range of particle sizes in the atmosphere (Milford and Davidson, 1987). For example, at many sites in North America, NO_3^- is characterized by a bimodal size distribution with modes above and below 1 μm . The supermicron particles are often the result of reactions between HNO_3 and coarse alkaline aerosols (Wolff, 1984) as, for example, in the San Joaquin Valley of California (Lindberg et al., 1990). Although the annual deposition of NH_4^+ is distributed similarly among the fine and coarse particles, particulate NO_3^- is found predominantly in the coarse-particle fraction (Table 4-7). Similar to the pattern for NH_4^+ , the estimated annual deposition of SO_4^{2-} particles occurs in both the fine- and coarse-particulate fractions (Table 4-8), while base cation deposition is virtually restricted to contributions from coarse particles (Table 4-9).

Although the annual chemical inputs to ecosystems from particle deposition is significant by itself, it is important to compare it with the total chemical inputs from all sources of atmospheric deposition (i.e., precipitation, particles, and gaseous dry deposition). Figure 4-10 shows the mean percentage contribution of NO_3^- and NH_4^+ , SO_4^{2-} , and base cation-containing particles to the total nitrogen, sulfur, and base cation deposition load to forest ecosystems (derived from Tables 4-7 through 4-9). Although the mean contribution of particulate deposition to cumulative nitrogen and sulfur deposition is typically less than 20% of annual inputs from all

TABLE 4-7. MEAN ANNUAL NITROGEN DEPOSITION (equivalents/ha/year) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL NITROGEN DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

Region/Forest Type	Location	Dry Particle Deposition						Annual Total Nitrogen ^a	References ^b
		Fine			Coarse				
		NO ₃ ⁻	NH ₄ ⁺	NO ₃ ⁻	NH ₄ ⁺	NO ₃ ⁻	NH ₄ ⁺		
North America									
Douglas fir	Washington	2	9	58	27	96	345	1,2	
Loblolly pine	Georgia	2	27	36	37	102	647	1,2	
Loblolly pine	North Carolina	3	35	89	66	193	997	1,2	
Loblolly pine	Tennessee	0.8	18	27	7	53	699	1,2	
Loblolly pine	Tennessee	0.2	14	8	1	23	410 ^c	3	
Slash pine	Florida	6	16	105	4	131	431	1,2	
White pine	North Carolina	1	22	23	23	69	510	1,2	
Red spruce	Maine	2	14	64	102	182	545	1,2	
Red spruce	North Carolina	3	74	133	43	253	1,939	1,2	
Red spruce	New York	1	9	5	2	17	1,136	1,2	
Alder	Washington	1	5	58	27	91	339	1,2	
Maple/Beech	New York	0.3	5	37	13	55	567	1,2	
Oak	Tennessee					307	857	4	
Oak	Tennessee	1	36	83	8	128	720	5	
Europe									
Norway spruce	Norway	8	21	52	29	110	775	1,2	
Norway spruce	Germany	21	62	56	4	143	1,250 ^c	3	

^aIncludes deposition from precipitation, gases, and particles.

^b1 = Johnson and Lindberg (1992a), 2 = Lovett (1992) and Lovett and Lindberg (1993), 3 = Lindberg et al. (1990), 4 = Kelly and Meagher (1986), 5 = Lindberg et al. (1986).

^cIncludes only the growing season from April to October 1987.

TABLE 4-8. MEAN ANNUAL SULFATE DEPOSITION (equivalents/ha/year) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL SULFUR DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

Region/Forest Type	Location	Dry Particle Deposition			Annual Sulfur Deposition ^a	References ^b
		Fine	Coarse	Total		
North America						
Douglas fir	Washington	12	62	74	320	1,2
Loblolly pine	Georgia	47	74	121	776	1,2
Loblolly pine	North Carolina	57	59	116	1,050	1,2
Loblolly pine	Tennessee	27	69	96	941	1,2
Loblolly pine	Tennessee	22	44	66	570 ^c	3
Slash pine	Florida	25	129	154	514	1,2
White pine	North Carolina	33	55	88	552	1,2
Red spruce	Maine	34	115	149	585	1,2
Red spruce	North Carolina	135	161	296	2,214	1,2
Red spruce	New York	10	4	14	1,096	1,2
Alder	Washington	6	62	68	325	1,2
Maple/Beech	New York	7	32	39	488	1,2
Oak forest	Tennessee	70	190	260	1,600	4
Europe						
Norway spruce	Germany	58	97	155	1,100 ^c	3
Norway spruce	Norway	19	79	98	663	12

^aIncludes deposition from precipitation, gases, and particles.

^b1 = Johnson and Lindberg (1992a), 2 = Lindberg (1992) and Lindberg and Lovett (1992), 3 = Lindberg et al. (1990); 4 = Lindberg et al. (1986).

^cIncludes only the growing season from April to October 1987.

TABLE 4-9. MEAN ANNUAL BASE CATION DEPOSITION (equivalents/ha/year) FROM FINE AND COARSE PARTICLES COMPARED TO TOTAL BASE CATION DEPOSITION FROM ALL SOURCES TO A VARIETY OF FOREST ECOSYSTEMS

Region/Forest Type	Location	Dry Base Cation Particles Deposition ^a			Total Annual Deposition ^b	References ^c
		Fine	Coarse	Total		
North America						
Douglas fir	Washington	9	180	189	670	1,2
Loblolly pine	Georgia	17	130	147	300	1,2
Loblolly pine	North Carolina	30	340	370	780	1,2
Loblolly pine	Tennessee	6.1	220	226	410	1,2
Mixed Hardwood	Tennessee	9.5 ^d	121 ^d	131 ^d	899 ^d	3
Slash pine	Florida	17	600	617	1,200	1,2
White pine	North Carolina	4	150	154	280	1,2
Red spruce	Maine	5	150	155	240	1,2
Red spruce	North Carolina	25	440	465	1,500	1,2
Red spruce	New York	2	36	38	230	1,2
Alder	Washington	4	180	184	660	1,2
Maple/Beech	New York	1	84	85	200	1,2
Oak forest	Tennessee	11	312	323	452	3
Europe						
Spruce	Germany ^d					3
Spruce	Norway	6	180	186	390	1,2

^aIncludes K⁺, Na⁺, Ca²⁺, and Mg²⁺.

^bIncludes deposition from precipitation, gases, and particles.

^c1 = Ragsdale et al. (1992), 2 = Johnson and Lindberg (1992a), 3 = Lindberg et al. (1986).

^dIncludes H⁺, K⁺, Na⁺, Ca²⁺, Mg²⁺ for the growing season from April to October 1987.

^eIncludes only Ca²⁺ and K⁺.




Figure 4-10. Mean (\pm SE) percent of total nitrogen, sulfur, or base cation deposition contributed by fine plus coarse particles. Data are means from Tables 4-7 through 4-9.

atmospheric sources, particulate inputs of base cations average half the total base cations entering forest ecosystems from the atmosphere.

An extensive comparison of particle to total chemical deposition is provided by the Integrated Forest Study (IFS; Johnson and Lindberg, 1992a; Lovett, 1994; Lovett and Lindberg, 1993; Lindberg and Lovett, 1992; Ragsdale et al., 1992). Other similar data sets are available (Kelly and Meagher, 1986; Miller et al., 1993; Lindberg et al., 1986, 1990). These data in (Tables 4-7 through 4-9) clearly indicate that the contribution of coarse and fine aerosols to deposition to forest ecosystems is strongly dependent on the chemical species.

Dry deposition is an important flux of sulfur and nitrogen compounds at all of the IFS sites and ranges from 9 to 59% of total (wet + dry + cloud) deposition for sulfur, 25 to 70% for NO_3^- and 2 to 33% for NH_4^+ . Only for NH_4^+ is wet deposition consistently greater than dry deposition (Lovett, 1994).

After emission from their sources, air pollutants are transformed and transported by atmospheric processes until deposited from the atmosphere to an aquatic or terrestrial ecosystem.

As a result, ground-level concentrations of an air pollutant depend on the proximity to the sources, prevailing meteorology, and nature and extent of atmospheric reactions between the source and the receptor (Holland et al., 1999). A more direct relationship exists between source strength and downwind ambient concentrations for primary air pollutants (e.g., SO₂) than for secondary pollutants (e.g., SO₄²⁻). Interaction of the chemical and physical atmospheric processes and source locations for all of the pollutants have a tendency to produce data patterns that show large spatial and temporal variability.

Holland et al. (1999) analyzed CASTNet monitoring data and using generative additive models (GAM) estimated the form and magnitude of trends of airborne concentrations of SO₂, SO₄²⁻, and nitrogen from 1989 to 1995 at 34 rural long-term CASTNet monitoring sites in the eastern United States. These models provide a highly flexible method for describing potential nonlinear relationships between concentrations, meteorology, seasonality, and time (e.g., how weekly SO₂ varies as a function of temperature). For most of the 34 sites in the eastern United State, estimates of change in SO₂ concentrations showed a decreasing functional form in 1989 to 1990, followed by a relatively stable period during 1991 to 1992), then a sharper decline beginning in 1994 (Holland et al., 1999).

Regional trends of seasonal and annual wet deposition and precipitation-weighted concentrations (PWCs) of sulfate in the United States over the period 1980 to 1995 were developed by Shannon (1999) from monitoring data and scaled to a mean of unity. In order to reduce some effects of year to year climatological variability, the unitless regional deposition and PWC trends were averaged (hereafter referred to a CONCDEP). During the 16-year period examined in the study, estimated aggregate emissions of SO₂ in the United States and Canada fell ~12% from about 1980 to 1982, it remained roughly level for a decade and then fell another ~15% from 1992 to 1995 — for an overall decrease of about 18%. Eastern regional trends of sulfate concentrations and deposition and their average CONCDEP, also exhibited patterns of initial decrease, near steady state, and final decrease with year to-to-year variability. The overall relative changed in CONCDEPs are greater than the changes in SO₂ emissions.

Concentrations and calculated deposition (concentration times amount of water) of SO₄²⁻ at the Hubbard Brook Experimental Forest (HBEF) in the White Mountains of central New Hampshire have been measured since June of 1964 (Likens et al., 2001). These measurements represent the longest continuous record of precipitation chemistry in North America. The

long-term measurements generally concur with those of Shannon (1999) discussed above. Major declines in emissions of SO₂ have been observed during recent decades in the eastern United States and have been correlated with significant decreases in SO₄²⁻ concentrations in precipitation (Shannon, 1999).

Deposition of sulfates and nitrates are clearly linked to emissions. Reduction in emissions must occur before concentrations can be reduced below current levels (Likens et al., 2001). Deposition is the key variable as sensitive ecosystems in the eastern North America have not yet shown improvement in response to decreased emissions of SO₂ (Driscoll et al., 1989; Likens et al., 1996). Clearly, additions of other chemicals, such as nitric acid and base cations, must be considered in addition to sulfur when attempting to resolve the acid rain problem (Likens et al., 1996, 1998). The effects of sulfur and nitrogen deposition on ecosystems are discussed in Section 4.2.2.2.

The long-term record indicates that a reduction in the deposition of basic cations (Ca²⁺, Mg⁺², K⁺, Na⁺) in bulk precipitation was associated with significant declines in SO₄²⁻ deposition cited above for the HBEF region (Driscoll et al., 1989). Decreases in streamwater concentrations of basic cations have decreased simultaneously, suggesting that streamwater concentrations of basic cations are relatively responsive to changes in atmospheric inputs. Regardless of the cause, the decline in atmospheric influx of basic cations could have important effects on nutrient availability as well as on the acid/base status of soil and drainage water (Driscoll et al., 1989).

Trace Elements

Deposition velocities for fine particles to forest surfaces have been reported in the range of 1 to 15 cm/s (Smith, 1990a). For example, total, annual heavy metal deposition amounts are highly variable depending on specific forest location and upwind source strength (Table 4-10). Lindberg et al. (1982) quantified the dry deposition of heavy metals to inert surfaces and to leaves of an upland oak forest. As noted for other chemical species, V_d was highly dependent on particle size and chemical species (Table 4-11) with the larger particles depositing more efficiently.

The preferential association of heavy metals with fine particles results in reduced control in emission control systems. Metal removal efficiencies for baghouse filters are typically 95 to

TABLE 4-10. MEAN (\pm SE) PARTICLE SIZE, DEPOSITION RATES, AND DERIVED DEPOSITION VELOCITIES (V_d) FOR HEAVY METAL DEPOSITION TO THE UPPER CANOPY (INERT PLATES OR LEAVES) OF AN UPLAND OAK FOREST

Metal	Particle Size (μm)	Deposition Rate	V_d
		(pg/cm/h)	(cm/s)
Manganese	3.4 ± 0.7	91 ± 23	6.4 ± 3.6
Cadmium	1.5 ± 0.7	0.3 ± 0.1	0.37 ± 0.18
Zinc	0.9 ± 0.2	6 ± 1	0.38 ± 0.1
Lead	0.5	23 ± 8	0.06 ± 0.01

Source: Lindberg et al. (1982).

TABLE 4-11. TOTAL HEAVY METAL DEPOSITION TO TEMPERATE LATITUDE FORESTS

Heavy Metal	Forest Deposition kg ha/yr (Range)
Cadmium	0.002-0.02
Copper	0.016-0.24
Lead	0.099-1000
Nickel	0.014-0.15
Zinc	0.012-0.178

Source: Smith (1990c).

99% for all but mercury, but fine particle capture is much less efficient. Wet scrubber efficiency varies with design and pressure drop, typically 50 to 90% (McGowan et al., 1993). Fine particles also have the longest atmospheric residence times and, therefore, can be carried long distances. Depending on climate conditions and topography, fine particles may remain airborne for days to months and may be transported 1,000 to 10,000 km or more from their source. This long-distance transport and subsequent deposition qualify heavy metals as regional- and global-

scale air pollutants. Ecosystems immediately downwind of major emissions sources (such as power generating, industrial, or urban complexes) may receive locally heavy inputs. Mass balance budgets (inputs and outputs) of seven heavy metals (cadmium, copper, iron, lead, manganese, nickel, and zinc) have been determined at the Hubbard Brook Experimental Forest (White Mountain National Forest) in New Hampshire. This forest is about 120 km northwest of Boston and relatively distant from major sources of heavy metal emissions. However, continental air masses that have passed over centers of industrial and urban activity also frequently follow storm tracks over northern New England. Resulting annual inputs for seven heavy metals at Hubbard Brook for 1975 to 1991 are presented in Figure 4-11. Note that the 44-fold decrease in Pb deposition is correlated with removal of Pb from motor vehicle fuels.




Figure 4-11. Annual total deposition of heavy metals to Hubbard Brook Experimental Forest, NH.

Source: Smith (1990a).

Trace element investigations conducted in roadside, industrial, and urban environments have shown that impressive burdens of particulate heavy metals accumulate on vegetative surfaces. Lead deposition to roadside vegetation (prior to its removal from fuel) was 5 to 20, 50 to 200, and 100 to 200 times the Pb deposition to agricultural crops, grasses, and trees, respectively, in non-roadside environments. In an urban setting, it has been estimated that the leaves and twigs of a 30 cm (12 in.) diameter sugar maple remove 60, 140, 5800, and 820 mg of Cd, Cr, Pb, and Ni, respectively, during a single growing season (Smith, 1973).

Semivolatile Organics

Organic compounds partition between gas and particle phases, and particulate deposition depends largely on the particle sizes available for adsorption (Pankow, 1987; Smith and Jones, 2000). Dry deposition of organic materials (e.g., dioxins, dibenzofurans, polycyclic aromatics) is often dominated by the coarse fraction, even though mass loading in this size fraction may be small (Lin et al., 1993) relative to the fine PM fraction. For example, measurements in Bavaria in both summer and winter revealed that > 80% of organics were in the fine (< 1.35 μm) fraction (Kaupp and McLachlan, 1999). Nevertheless, in most cases, calculated values of dry deposition were dominated by the material adsorbed to coarse particles. Wet deposition, in contrast, was dominated by the much larger amount of material associated with fine particles. In this Bavarian environment (where monthly precipitation is about 50 mm in winter and summer), wet deposition dominated, with dry deposition accounting for only 14 to 25% of total deposition (Kaupp and McLachlan, 1999). Lower relative contents of more volatile species in summer than winter (Kaupp and McLachlan, 1999) indicate the critical importance of gas-particle phase interconversions in determining deposition.

Effective deposition of PM is required before biological effects on plants or ecosystems can occur. It is clear that substantially improved techniques for monitoring and predicting deposition will be required to characterize with certainty quantitative relationships between rates and quantities of PM deposition associated with such effects.

4.2.3 Assessment of Atmospheric PM Deposition Effects

Introduction

The discussion in the pages that follow assesses and characterizes the overall ecological condition or integrity of the ecosystems within the United States affected by the deposition of anthropogenic stressors associated with PM and indicates their status. The six Essential Ecological Attributes (EEAs) – the landscape condition, biotic condition, and chemical/ physical characteristics, ecological processes, hydrology/geomorphology, and natural disturbance regimes (Table 4-1) – provide a hierarchical framework for assessing ecosystem status. Measurable characteristics related to structure, composition, or functioning of ecological systems may be determined by the use of endpoints or ecological indicators of condition that are significant either ecologically or to society (Harwell et al., 1999).

The relationships among the EEAs are complex because all are interrelated (i.e., changes in one EEA may affect, directly or indirectly, every other EEA). The ecological processes create and maintain patterns composed of the elements in the system and their arrangement; in turn, the patterns affect how the processes are expressed (Science Advisory Board, 2002). Changes in patterns or processes result in changes in the status and functioning of an ecosystem. The information in the sections that follow discusses changes in landscape and biotic patterns and in ecological and chemical/physical processes resulting from the stressors in PM deposition (Figure 4-12).

The elements of biotic condition are organized in a nested hierarchy with several levels. These include the structural and composition aspects (patterns) of the biota within landscape, ecosystem or ecological community, species/population, organism, and genetic/molecular levels (Science Advisory Board, 2002). Within these biological levels of organization, changes in the biodiversity, composition, and structural elements relate directly to functional integrity (such as trophic status or structural integrity within habitats). Changes in biodiversity are of particular significance in altering the functioning of ecosystems.

As previously stated, ecosystems are dynamic, self-adjusting, self-maintaining, complex, adaptive systems, in which the patterns of the higher levels of biotic organization emerge from the interactions and selection processes at localized levels (Levin, 1998). Ecosystem components must have an adequate supply of energy, mineral nutrients, and water to maintain themselves and function properly. During the ecological processes of energy and material flow,




Figure 4-12. Sample stressors and the essential ecological attributes they affect (after Science Advisory Board, 2002).

the energy obtained by plants (the producers) from sunlight during photosynthesis (primary production) and chemical nutrients (e.g., nitrogen, phosphorus, sulfur) taken up from the soil are transferred to other species (the consumers) within the ecosystem through food webs. The movement of chemical nutrients (materials) through an ecosystem is cyclic, as the nutrients are used or stored and eventually returned to the soil by decomposer organisms. Energy, on the

other hand, is transferred from organism to organism through an ecosystem in food webs and, finally, is dissipated into the environment as heat (Odum, 1993).

Ecosystem and community patterns are characterized by the interaction of their component species, the ecosystems processes of energy flow, nutrient flux, water and material flow, and by the effects their activities have on the physical and chemical environment. The flows of energy and nutrient cycling provide the interconnectedness among the elements of the biotic hierarchy and transform the community from a random collection of numerous species into an integrated whole, an ecosystem. Elucidating these interactions across scales is fundamental to understanding the relationships between biodiversity and ecosystem functioning (Levin, 1998).

Human existence on this planet depends on the life-support services provided by the interaction of the different EEAs. Both ecosystem structure (biotic conditions) and function (ecological processes) play essential roles in providing goods (products) and services (Table 4-12; Daily, 1997). Ecosystem processes maintain clean water, clean air, a vegetated Earth, and a balance of organisms — the functions that enable humans to survive. The benefits they impart include the absorption and breakdown of pollutants, cycling of nutrients, binding of soil, degradation of organic waste, maintenance of a balance of gases in the air, regulation of radiation balance and climate, and the fixation of solar energy (World Resources Institute, 2000; Westman, 1977; Daily, 1997). The economic benefits and values associated with ecosystem functions, goods, and services as well as the need to preserve them because of their value to human life are discussed by Costanza et al. (1997) and Pimentel et al. (1997).

Goods such as food crops, timber, livestock, fish, and drinking water usually have market value, while ecosystem services such as flood-control benefits, wildlife habitat, cycling of nutrients, and removal of air pollutants, usually are very difficult to measure (Goulder and Kennedy, 1997).

Attempts have been made to calculate the value of biodiversity and the world's ecosystem services and natural capital (Goulder and Kennedy 1997; Pimentel et al., 1997; Constanza et al., 1997). These have been controversial because of a lack of agreement on the measurement and philosophical basis for placing value on ecosystem services (Goulder and Kennedy, 1997; Heal, 2000; Moomaw, 2002). Constanza et al. (1997) state that it may never be possible to make a precise estimate of the services provided by ecosystems; however, their estimates indicate the relative importance of services, not their true values, considering that the loss of ecosystems can

TABLE 4-12. PRIMARY GOODS AND SERVICES PROVIDED BY ECOSYSTEMS

Ecosystem	Goods	Services
<i>Agroecosystems</i>	Food crops Fiber crops Crop genetic resources	Maintain limited watershed functions (infiltration, flow control, and partial soil protection) Provide habitat for birds, pollinators, and soil organisms important to agriculture Sequester atmospheric carbon Provide employment
<i>Coastal Ecosystems</i>	Fish and shellfish Fishmeal (animal feed) Seaweeds (for food and industrial use) Salt Genetic resources	Moderate storm impacts (mangroves, barrier islands) Provide wildlife (marine and terrestrial) habitat and breeding areas/hatcheries/nurseries Maintain biodiversity Dilute and treat wastes Provide harbors and transportation routes Provide human and wildlife habitat Provide employment Contribute aesthetic beauty and provide recreation
<i>Forest Ecosystems</i>	Timber Fuelwood Drinking and irrigation water Fodder Nontimber products (vines, bamboos, leaves, etc.) Food (honey, mushrooms fruit, and other edible plants; game) Genetic resources	Remove air pollutants, emit oxygen Cycle nutrients Maintain array of watershed functions (infiltration, purification, flow control, soil stabilization) Maintain biodiversity Sequester atmospheric carbon Moderate weather extremes and impacts Generate soil Provide employment Provide human and wildlife habitat Contribute aesthetic beauty and provide recreation
<i>Freshwater</i>	Drinking and irrigation water Fish Hydroelectricity Genetic resources	Buffer water flow (control timing and volume) Dilute and carry away wastes Cycle nutrients Maintain biodiversity Provide aquatic habitat Provide transportation corridor Provide employment Contribute aesthetic beauty and provide recreation
<i>Grassland Ecosystems</i>	Livestock (food, game, hides, and fiber) Drinking and irrigation water Genetic resources	Maintain array of watershed functions (infiltration, purification, flow control, and soil stabilization) Cycle nutrients Remove air pollutants and emit oxygen Maintain biodiversity Generate soil Sequester atmospheric carbon Provide human and wildlife habitat Provide employment Contribute aesthetic beauty and provide recreation

Source: World Resources Institute (2000).

affect human existence. Constanza et al. (1997) refer to the loss of “total values” for all portions of all ecosystem services, whereas the relevant economic analysis for the PM standard setting is for marginal changes in ecosystem goods and services due to varying levels of air pollution effects, generally not total loss of services.

The prevailing economic approach to ascertaining value discussed above is benefit-cost analysis. Benefit-cost analysis implicitly adopts the utilitarian basis for value. That is, the value of a given living thing is the amount of human benefit or satisfaction the thing provides (Goulder and Kennedy, 1997). Benefit-cost analysis determines the environmental and public health damage and then calculates an appropriate level of emissions reduction required to lower concentrations to a standard-setting level. As the costs and benefits extend into the future at differing rates, the question arises of what discount rate to use in determining the net present value of the costs and benefits. A degraded ecosystem continues to provide fewer services for the indefinite future (Moomaw, 2002).

The approach of Harwell et al. (1999) and the report of the Ecological Processes and Effects Committee of the SAB (Science Advisory Board, 2002) also point out the need to understand human effects and their costs on ecosystems so that management can define what ecological conditions are to be desired. Further, they state that the establishment of ecological goals involves a close linkage between scientists and decision makers in which science informs decision makers and the public by characterizing the ecological conditions that are achievable under particular management regimes. Decision makers then can make choices that reflect societal values including issues of economics, politics, and culture. For management to achieve their goals, the general public, scientific community, resource managers, and decision makers need to be routinely apprised of the condition or integrity of ecosystems so that ecological goals may be established (Harwell et al., 1999).

Biodiversity, the variety of life, encompasses all levels of biological organization, including species, individuals, populations, and ecosystems (Wilson, 1997). Human-induced changes in biotic diversity and alterations in the structure and functioning of ecosystems are the two most dramatic ecological trends of the past century (Vitousek et al., 1997). There are few ecosystems on Earth today that are not influenced by humans (Freudenburg and Alario, 1999; Vitousek et al., 1997; Matson et al., 1997; Noble and Dirzo, 1997). The scientific literature is filled with discussions of the importance of ecosystem structure (patterns) and function

(processes). The deposition of PM from the atmosphere has the potential to alter ecosystem structure and function by altering nutrient cycling and changing biodiversity. We must, therefore, determine how ecosystems respond to both natural and anthropogenic stresses so as to know whether anthropogenic stresses are affecting ecosystem services and products (Table 4-12).

Concern has risen in recent years regarding the consequences of changing the biological diversity of ecosystems (Tilman, 2000; Ayensu et al., 1999; Wall, 1999; Hooper and Vitousek, 1997; Chapin et al., 1998). The concerns arise because human activities are creating disturbances that decrease biodiversity, alter the complexity and stability of ecosystems, and produce changes in ecological processes and the structure, composition and function of ecosystems (Figure 4-13; Pimm, 1984; Levin, 1998; Chapin et al., 1998; Peterson et al., 1998; Tilman, 1996; Tilman and Downing, 1994; Wall, 1999; Daily and Ehrlich, 1999). The above changes can affect ecosystem services vital to human life.




Figure 4-13. Linkages among various ecosystem goods and services (food, water, biodiversity, forest products) and other driving forces (climate change).

Source: Modified from Ayensu et al. (1999).

4.2.3.1 Effects on Vegetation and Ecosystems

Exposure to a given mass concentration of airborne PM may lead to widely differing phytotoxic responses, depending on the particular mix of deposited particles. Effects of particulate deposition on individual plants or ecosystems are difficult to characterize because of the complex interactions among biological, physicochemical, and climatic factors. Most direct effects, other than regional effects associated with global changes, occur in the severely polluted areas surrounding industrial point sources, such as limestone quarries, cement kilns, and metal smelting facilities. Fine particles are more widely distributed from their sources than are coarse particles. Experimental applications of PM constituents to foliage typically elicit little response at the more common ambient concentrations. The diverse chemistry and size characteristics of ambient PM and the lack of clear distinction between effects attributed to phytotoxic particles and to other air pollutants further confound understanding of the direct effects on foliar surfaces. The majority of the documented toxic effects of particles on vegetation reflect their chemical content (e.g., acid/base, trace metal, nutrient), surface properties, or salinity. Studies of the direct effects of particles on vegetation have not yet advanced to the stage of reproducible exposure experiments. The difficulties of experimental application of ambient particles to vegetation have been discussed by Olszyk et al. (1989). Studies indicate many phytotoxic gases are deposited more readily, assimilated more rapidly, and lead to greater direct injury of vegetation than do most common particulate materials (Guderian, 1986). The dose-specific responses (dose-response curves) obtained in early experiments following the exposure of plants to phytotoxic gases generally have not been observed following the application of particles.

Unlike gaseous dry deposition, neither the solubility of the particles nor the physiographical activity of the surface is likely to be of first order importance in determining deposition velocity (V_d). Factors that contribute to surface wetness and stickiness may be critical determinants of deposition efficiency. Available tabulation of deposition velocities are highly variable and suspect. However, high-elevation forests receive larger particle deposition loadings than equivalent lower elevations sites because of higher wind speeds and enhanced rates of aerosol impaction, orographic effects on rainfall intensity and composition, increased duration of occult deposition, and, in many areas, the dominance of coniferous species with needle-shaped leaves (Lovett, 1984), (see Section 4.2.2.3 for a discussion of deposition efficacy). Recent evidence indicates that all three modes of deposition (wet, occult, and dry)

must be considered in determining inputs to ecosystems or watersheds, because each may dominate over specific intervals of space.

Atmospheric PM may affect vegetation directly following deposition on foliar surfaces or indirectly by changing the soil chemistry or by changing the amount of radiation reaching the Earth's surface through PM-induced climate change processes. Indirect effects, however, are usually the most significant because they can alter nutrient cycling and inhibit plant nutrient uptake.

Direct Effects of Particulate Matter Deposition

Coarse and fine PM deposition affects both patterns in the elements of EEA categories subsumed under biotic condition and processes subsumed under chemical/physical and ecological processes in the EPEC framework described earlier. Measurable responses have been observed as reductions in photosynthesis, changes in soil salinity, and foliar effects resulting from nitrate, sulfate, and acidic and heavy metal deposition.

Particles transferred from the atmosphere to foliar surfaces may reside on the leaf, twig, or bark surface for extended periods; be taken up through the leaf surface; or be removed from the plant via resuspension to the atmosphere, washing by rainfall, or litter-fall with subsequent transfer to the soil. Any PM deposited on above-ground plant parts may exert physical or chemical effects. The effects of "inert" PM are mainly physical; whereas those of toxic particles are both chemical and physical. The effects of dust deposited on plant surfaces or soil are more likely to be associated with their chemistry, rather than their mass and chemical effects may be more important than any physical effects (Farmer, 1993). Nevertheless, vegetative surfaces represent filtration and reaction/exchange sites (Tong, 1991; Youngs et al., 1993).

Direct Effects of Coarse Particles

Coarse particles, ranging in size from 2.5 to 100 μm , are chemically diverse, arise predominantly from local sources, and are typically deposited near their sources because of their sedimentation velocities. Airborne coarse particles are derived from the following sources: road, cement kiln, and foundry dust; fly ash; tire particles and brake linings; soot and cooking oil droplets; biogenic materials (e.g, plant pollen, fungal spores, bacteria and viruses); abraded plant fragments; sea salt; and hydrated deliquescent particles of otherwise fine aerosol. In many rural

areas and some urban areas, the majority of the mass in the coarse particle mode derives from the elements silicon, aluminum, calcium, and iron, suggesting a crustal origin as fugitive dust from disturbed land, roadways, agriculture tillage, or construction activities. Rapid sedimentation of coarse particles tends to restrict their direct effects on vegetation largely to roadsides and forest edges.

Physical Effects — Radiation. Dust can cause physical and chemical effects. Deposition of inert PM on above-ground plant organs sufficient to coat them with a layer of dust may result in changes in radiation received, a rise in leaf temperature, and the blockage of stomata. Increased leaf temperature and heat stress, reduced net photosynthesis, and leaf chlorosis, necrosis, and abscission were reported by Guderian (1986). Road dust decreased the leaf temperature on *Rhododendron catawbiense* by ~4 °C (Eller, 1977); whereas foundry dust caused an 8.7 °C increase in leaf temperature of black poplar (*Populus nigra*) under the conditions of the experiment (Guderian, 1986). Deciduous (broad) leaves exhibited larger temperature increases because of particle loading than did conifer (needle) leaves, a function of poorer coupling to the atmosphere. Inert road dust caused a three- to four-fold increase in the absorption coefficient of leaves of English ivy (Eller, 1977; Guderian, 1986) for near infrared radiation (NIR; 750 to 1350 nm). Little change in absorption occurred for photosynthetically active radiation (PAR; 400 to 700 nm). The increase in NIR absorption may be accounted for by a decrease in reflectance and transmission in these wavelengths. The amount of energy entering the leaf increased by ~30% in the dust-affected leaves. Deposition of coarse particles increased leaf temperature and contributed to heat stress, reduced net photosynthesis, and caused leaf chlorosis, necrosis, and abscission (Dässler et al., 1972; Parish, 1910; Guderian, 1986; Spinka, 1971).

Starch storage in dust-affected leaves increased with dust loading when under high (possibly excessive) radiation, but decreased following dust loading when radiation was limiting. These modifications of the radiation environment had a large effect on single-leaf utilization of light. The boundary layer properties, determined by leaf morphology and environmental conditions, strongly influenced the direct effects of particle deposition on radiation heating (Eller, 1977; Guderian, 1986) and on gas exchange. Brandt and Rhoades (1973) attributed the reduction in the growth of trees to crust formation from limestone dust on the leaves. Crust

formation reduced photosynthesis and the formation of carbohydrates needed for normal growth, induced premature leaf-fall, damaged leaf tissues, inhibited growth of new tissue, and reduced starch storage. Dust may decrease photosynthesis, respiration, and transpiration; and it may allow penetration of phytotoxic gaseous pollutants, thereby causing visible injury symptoms and decreased productivity. Permeability of leaves to ammonia increased with increasing dust concentrations and decreasing particle size (Farmer, 1993).

Dust also has been reported to physically block stomata (Krajíčková and Mejstřík, 1984). Stomatal clogging by PM from automobiles, stone quarries, and cement plants was also studied by Abdullah and Iqbal (1991). The percentage of clogging was low in young leaves when compared with old, mature leaves and the amount of clogging varied with species and locality. The maximum clogging of stomata observed was about 25%. The authors cited no evidence that stomatal clogging inhibited plant function. The heaviest deposit of dust usually occurs on the upper surface of broad-leaved plants; whereas the majority of the stomata are on the lower surface where stomatal clogging would be less likely.

Chemical Effects. The chemical composition of PM is usually the key phytotoxic factor leading to plant injury. On hydration, cement-kiln dust liberates calcium hydroxide, which can penetrate the epidermis and enter the mesophyll; in some cases, this has caused the leaf surface alkalinity to reach a pH of 12. Lipid hydrolysis, coagulation of the protein compounds, and ultimately plasmolysis of the leaf tissue reduce the growth and quality of plants (Guderian, 1986). In experimental studies, applications of cement-kiln dust of known composition for 2 to 3 days yielded dose-response curves between net photosynthetic inhibition or foliar injury and dust application rate (Darley, 1966). Lerman and Darley (1975) determined that leaves must be misted regularly to produce large effects. Alkalinity was probably the essential phytotoxic property of the applied dusts.

Salinity. Particulate matter enters the atmosphere from oceans following the mixing of air into the water column and the subsequent bursting of bubbles at the surface. The effervescence of bubbles on the surface of the ocean forcefully ejects droplets of sea water into the air. These droplets, concentrated by evaporation, are carried inland by wind and deposited on the seaward side of coastal plants (Boyce, 1954). This occurs largely at the surf line (i.e., near land and

potentially sensitive terrestrial receptors). This process can be a significant source of sulfate, sodium, chloride, and trace elements (as well as living material) in the atmospheric aerosol that impacts coastal vegetation. Sea-spray particles (Taback et al., 1979) are approximately 24% greater in size than 10 μm , and 54% are between 3 and 10 μm . Thus, only about 20% are fine (0 to 2.5 μm) particles; and deposition by sedimentation and impaction is concentrated near the coast, whereas the particle size distribution shifts toward the fine fraction over longer inland transport distances. Airborne concentrations of this marine PM decrease quickly with distance inland from the surf line both by deposition and dilution within the atmospheric mixed layer (McKay et al., 1994; Nelis et al., 1994). Near-shore sediments, with associated pollutants present in coastal runoff, may be suspended in the surf and reentrained into the air. This can be a substantial source of microorganisms and of radionuclides to coastal vegetation (Nelis et al., 1994; McKay et al., 1994).

Sea-salt particles can serve as nuclei for the absorption and subsequent reaction of other gaseous and particulate air pollutants. Both nitrate and sulfate from the atmosphere have been found to associate with coarse and fine sea-salt particles (Wu and Okada, 1994). Direct effects on vegetation reflect these inputs, as well as the classical salt injury caused by the sodium and chloride that constitute the bulk of these particles. Foliar accumulation of airborne salt particles may lead to foliar injury, thusly affecting the species composition in coastal environments (Smith, 1984).

The effects of winds and sea spray on coastal vegetation has been reported in the literature since the early 1800s (Boyce, 1954). However, there has been some controversy about whether the injury to coastal vegetation resulted from windblown aerial salts or from mechanical injury (i.e., sand blasting) due to wind alone. Though the significance of sea water dashed on fore dunes and rocky coasts had been recognized by several authors, Wells and Shunk (1937, 1938) and Wells (1939) were the first to recognize the importance of salt spray in coastal ecology. Wells and Shunk (1937) reported that salt spray carried over dunes was the most important factor influencing growth form, zonation, and succession in coastal dunes. Salt spray injury was recorded 1.25 miles inland on the North Carolina coast. On the basis of observations in the Cape Fear area, they determined that the shape of coastal "wind form" shrubs were the result of sea spray carried by high winds. They found injury on shrubs only near the coast, while those at

greater distances inland showed no injury whatsoever after a strong southeast wind that persisted for a period of 19 hours during cloudy weather and abundant soil moisture.

To determine the cause of injury, injured and uninjured shoots were titrated for chlorides. A marked difference was observed between the injured and uninjured shoots (Wells and Shunk, 1937, 1938). Experimental spraying of shoots of woody plants with seawater resulted in a pattern of injury similar to the injury observed on seaside shrubs. The absence of the more inland species, such as persimmon (*Diosporos virginiana*), turkey oak (*Quercus laevis*), longleaf pine (*Pinus palustris* Mill., *P. australis*), and wire grass (*Aristida stricta*), was explained on the basis of intolerance of these species to salt spray. The dominance of live oak (*Quercus virginiana*), as a practically pure stand on Smith Island (also known as Bald Head Island), NC and along the eastern and southern NC coast, was determined by Wells (1939) to be due to the tree's tolerance to salt spray. He termed the long term stabilization of the live oak stand as a new type of climax, the "salt spray climax." The later work of Oosting and Billings (1942) near Beaufort, NC corroborated the findings of Wells and Shunk (1937, 1938).

The report by Boyce (1954) is probably the most extensive on salt-spray communities. Dune sands in many coastal areas have been shown to have extremely low concentrations of dissolved salts. Studies have indicated that the salt content of the coastal dunes of Virginia, Massachusetts, and California did not exceed the maximum occurring in ordinary cultivated soils. Oosting and Billings (1942) found no correlation between soil salinity and plant distribution on the North Carolina coast. Although surface crusts of sand dunes have been shown to have high concentrations of chlorides which could be attributed to sea spray, concentrations of chlorides in underlying layers was low. The salt content of the surface layer varied with exposure of the dunes to oceanic winds (Boyce, 1954).

Boyce (1954), Wells (1939), and Wells and Shunk (1938) concluded on the basis of their studies that necrosis and death of plant tissues results from the high deposition of salt spray and high accumulation of the chloride ion (Cl^-) in the plant tissues. Very little salt is taken up by plant roots; most enters through the aerial organs. Leaves of plants exposed to salt spray show a distinct pattern of injury (Wells and Shunk, 1938). Necrotic areas first appear at the leaf tips and upper margins and then progress slowly in an inverted "V" toward the petiole. This leaf injury pattern was verified experimentally. Mechanical injury resulting from leaves and twigs beating against each other in the wind causes the formation of small lesions through which salt

can enter. After entry into the plant, the chloride ion is rapidly translocated to the apices of the leaves and twigs where it accumulates to injurious concentrations and results in the death of only a portion of the plant. The differential deposition and translocation of the chloride ion results in the death of the seaward leaves and twigs. The result is the continued growth of the uninjured branches in an inland direction. As a result, the canopy angle varies with the intensity of the spray (Boyce, 1954).

Little or no mineral ions are available in the silicate sands of the coastal dunes. Consequently, plants obtain mineral ions needed for growth from the salt spray. Seawater contains all of the mineral ions required for plant growth except nitrogen and phosphorus. The amount of nitrogen and phosphorus in seawater varies over a wide range (Boyce, 1954). Experiments indicated that available nitrogen in sea spray was a conditioning factor. Low nitrogen availability increased the tolerance of dune species to salt spray. Increasing the availability of nitrogen resulted in a different pattern of plant zonation and distribution. Dicotyledonous species were restricted to areas of lower spray intensity. The severity of chloride injury was associated more with the amount of available potassium than with the concentration of chlorides within the limits of 280 to 360 mg Cl/liter (Boyce, 1954).

Other sources of phytotoxic saline PM include aerosols from cooling towers and roadway deicing salt. Cooling towers used to dissipate waste heat from steam-electric power generating facilities may emit salt if brackish salt water is used as a coolant (McCune et al., 1977; Talbot, 1979). Foliar injury is related to salt droplets deposited by sedimentation or impaction from cooling tower drift. The distance of the salt drift determines the amount of deposition and location of injury. The environmental conditions most conducive to injury were absence of precipitation, which can wash salt off leaves, and high relative humidity (Talbot, 1979). Increased injury is associated with wind speed and salt concentrations. Typical toxicity symptoms from acute exposures include marginal foliar necrosis and lesions, shoot-tip dieback, leaf curl, and interveinal necrosis (McCune et al., 1977). Based on experimental data, Grattan et al. (1981) observed that, to cause injury, salt deposited on leaf surfaces must dissolve and be absorbed into leaf tissue. Their work also indicated the importance of RH in foliar uptake. If RH remained below 70%, even heavy deposition of salt did not induce injury in peppers, soybeans, or tomatoes.

Injury to vegetation from the application of deicing salt was related to salt spray blown or drifting from the highways (Hofstra and Hall, 1971; Viskari and Kärenlampi, 2000). The most severe injury was observed nearest the highways. The results presented in these studies agree with that of Wyttenbach et al. (1989), who observed that conifers planted near roadway margins in the eastern United States often exhibit foliar injury due to toxic levels of saline aerosols deposited from deicing solutions. Piatt and Krause (1974) demonstrated that road and site factors influence the spread of deicing salt into forested areas. The slope away from the road influenced the distance from the road where injury was observed and the percent slope was correlated with distance.

Effects of Fine Particles

Fine PM in rural areas is generally secondary in nature, having condensed from the vapor phase or been formed by chemical reaction from gaseous precursors in the atmosphere, and is generally smaller than 1 to 2.5 μm . Nitrogen and sulfur oxides, volatile organic gases, condensates of volatilized metals, and products of incomplete combustion are common precursors for fine PM. Many of these materials react with an oxidizing atmosphere to contribute to high secondary PM concentrations during summer months in many U.S. areas or during late fall and winter in areas with high nitrate concentrations. The conclusion that sufficient data were not available for adequate quantification of dose-response functions for direct effects of fine aerosols on vegetation, reached in the 1982 PM/SO_x AQCD (U.S. Environmental Protection Agency, 1982), continues to be true today. Only a few studies on the direct effects of acid aerosols have been completed (U.S. Environmental Protection Agency, 1982). The major effects are indirect and occur through the soil (see Section 4.2.3.2).

Nitrogen. Nitrate is observed in both fine and coarse particles. Nitrates from atmospheric deposition represent a substantial fraction of total nitrogen inputs to southeastern forests (Lovett and Lindberg, 1986). However, much of this is contributed by gaseous nitric acid vapor; and a considerable amount of the particulate nitrate is taken up indirectly through the soil. Garner et al. (1989) estimated deposition of nitrogen to forested landscapes in eastern North America at 10 to 55 kg/ha/year for nitrate and 2 to 10 kg/ha/year for ammonium. About half of these values were ascribed to dry deposition.

Driscoll et al. (2003) estimated anthropogenic nitrogen inputs to eight large watersheds in the northeastern United States for the year 1997. Inputs of total nitrogen deposition ranged from 14 kg/ha/year in the Casco Bay watershed, ME to 68 kg/ha/year in the Massachusetts Bay watershed. Atmospheric deposition of nitrogen was the second largest source ranging in amounts from 5 to 10 kg/ha/year (11 to 36% of the total) (Driscoll et al., 2003). Nitrogen deposition in the western United States ranges from 1 to 4 kg/ha/year over much of the region to as high as 30 kg/year downwind of major urban and agricultural areas. An unknown amount of nitrogen deposited to the West Coast originates in Asia (Fenn et al., 2003a).

Atmospheric additions of particulate nitrogen in excess of vegetation needs are lost from the system, mostly in leachate from the soil as nitrate. Managed agricultural ecosystems may be able to utilize deposited particulate nitrogen more efficiently than native ecosystems, although many cultivated systems also lose considerable nitrogen as nitrate in runoff, deep drainage, or soil water. It has proven difficult to quantify direct foliar fertilization by uptake of nitrogen from ambient particles.

There is no doubt that foliar uptake of nitrate can occur, as clearly shown by the efficacy of foliar fertilization in horticultural systems. Potassium nitrate was taken up by the leaves of deciduous fruit trees (Weinbaum and Neumann, 1977) and resulted in increased foliar nitrogen concentrations. Not all forms of nitrogen are absorbed equally, nor are all equally benign. Following foliar application of 2600 ppm of nitrogen as calcium nitrate $\text{Ca}(\text{NO}_3)_2$, $(\text{NH}_4)_2\text{SO}_4$, or urea $[(\text{NH}_2)_2\text{CO}]$ to apple canopies (Rodney, 1952; Norton and Childers, 1954), leaf nitrogen levels were observed to increase to similar levels; however, calcium nitrate and ammonium sulfate caused visible foliar injury, whereas urea did not. Urea is generally the recommended horticultural foliar fertilizer.

The mechanism of uptake of foliarly deposited nitrate is not well established. Nitrate reductase is generally a root-localized enzyme: it is generally not present in leaves, but is inducible there. Induction typically occurs when the soil is heavily enriched in NO_3^- . As the root complement of nitrate reductase becomes overloaded, unreduced nitrate reaches the leaves through the transpiration stream. Nitrate metabolism has been demonstrated in leaf tissue (Weinbaum and Neumann, 1977) following foliar fertilization. Residual nitrate reductase activity in leaves may be adequate to assimilate typical rates of particulate nitrate deposition.

Uptake of nitrate may be facilitated by codeposited sulfur (Karmoker et al., 1991; Turner and Lambert, 1980).

Nitrate reductase is feedback-inhibited by its reaction product NH_4^+ . The common atmospheric aerosol, ammonium nitrate [NH_4NO_3], therefore may be metabolized in two distinct biochemical steps: first the ammonium (probably leaving nitric acid) and then the nitrate. Losses of nitric acid by volatilization during this process, if they occur, have not been quantified.

Direct foliar effects of particulate nitrogen have not been documented. Application of a variety of fine nitrogenous aerosol particles ($0.25\ \mu\text{m}$) ranging from 109 to $244\ \mu\text{g}/\text{m}^3$ nitrogen with or without $637\ \mu\text{g}/\text{m}^3$ sulfur caused no consistent short-term (2- to 5-h) effect on gas exchange in oak, maize, or soybean leaves (Martin et al., 1992).

Although no evidence exists for the direct transfer of nutrient particulate aerosols into foliage, a few studies give insights into the potential for ammonium and nitrate transfer into leaves. Fluxes of both NO_3^- and NH_4^+ , measured in wet deposition and in throughfall plus stemflow in forests, commonly indicate higher fluxes of nitrogen above the canopy (Parker, 1983; Lindberg et al., 1987; Sievering et al., 1996), and imply net foliar uptake. Lovett and Lindberg (1993) reported a linear relationship between inorganic nitrogen fluxes in deposition and throughfall, suggesting that uptake may be considered passive to some extent.

Garten and Hanson (1990) studied the movement of ^{15}N -labeled nitrate and ammonium across the cuticles of red maple (*Acer rubrum*) and white oak (*Quercus alba*) leaves when applied as an artificial rain mixture. Brumme et al. (1992), Bowden et al. (1989), and Vose and Swank (1990) have published similar data for conifers. These studies show the potential for nitrate and ammonium to move into leaves, where it may contribute to normal physiological processes (e.g., amino acid production) (Wellburn, 1990). Garten (1988) showed that internally translocated ^{35}S was not leached readily from tree leaves of yellow poplar (*Liriodendron tulipifera*) and red maple (*Acer rubrum*), suggesting that SO_4^{2-} would not be as mobile as the nitrogen-containing ions discussed by Garten and Hanson (1990). Further, when the foliar extraction method is used, it is not possible to distinguish sources of chemicals deposited as gases or particles (e.g., HNO_3 , NO_2 , NO_3^-), or sources of ammonium (deposited as NH_3 or NH_4^+) (Garten and Hanson, 1990).

Particle deposition contributes only a portion of the total atmospheric nitrogen deposition reaching vegetation; but, when combined with gaseous and precipitation-derived sources, total

nitrogen deposition to ecosystems has been identified as a possible causal factor leading to changes in natural ecosystems (see Section 4.2.2).

Sulfur. Anthropogenic sulfur emissions are > 90% SO₂. Most of the remaining emissions of sulfur is directly as sulfate (U.S. Environmental Protection Agency, 1996a). Sulfur dioxide is hydrophilic and is rapidly hydrated and oxidized to sulfite and bisulfite and then to sulfate, which is ~30-fold less phytotoxic. The ratio of SO₄²⁻:SO₂ increases with the aging of the air mass and, therefore, with distance from the source. Sulfate is sufficiently hygroscopic in humid air that it may exist significantly in the coarse particulate fraction. Because dilution of both SO₂ and particulate SO₄²⁻ occurs with distance from the source, damaging levels of particulate SO₄²⁻ are rarely deposited. In this case, gas to particle conversion is of benefit to local vegetation.

Sulfur is an essential plant nutrient. Low dosages of sulfur serve as a fertilizer, particularly for plants growing in sulfur-deficient soil (Hogan et al., 1998). However, current levels of sulfate deposition reportedly exceed the capacity of most vegetative canopies to immobilize the sulfur (Lindberg, 1992; Johnson, 1984). Nitrogen uptake in forests may be regulated loosely by sulfur availability, but sulfate additions in excess of needs do not typically lead to injury unless deposited in acidic precipitation (Turner and Lambert, 1980).

There are few field demonstrations of foliar sulfate uptake (Krupa and Legge, 1986, 1998). Sulfate in throughfall is often enriched above levels in precipitation. The relative importance of foliar leachate and prior dry-deposited sulfate particles remains difficult to quantify (Cape et al., 1992). Leaching rates are not constant and may respond to levels of other pollutants, including acids. Uptake and foliar retention of gaseous and particulate sulfur are confounded by variable rates of translocation and accessibility of deposited materials to removal and quantification by leaf washing. Following soil enrichment with ³⁵SO₄²⁻ in a Scots pine forest, the apparent contribution of leachate to throughfall was only a few percent following an initial burst of over 90%, because of extreme disequilibrium in the labeling of tissue sulfate pools (Cape et al., 1992).

Olszyk et al. (1989) provide information on the effects of multiple pollutant exposures including particles (NO₃⁻, 142 µg/m³; NH₄⁺, 101 µg/m³; SO₄²⁻, 107 µg/m³). They found that only gaseous pollutants produced direct harmful effects on vegetation for the concentrations documented, but the authors hypothesized that the long-term accumulation of the nitrogen and

sulfur compounds contributed from particle deposition might have adverse effects on plant nutrition over long periods of time. Martin et al. (1992) exposed oak (*Quercus macrocarpa*), soybean (*Glycine max*), and maize (*Zea mays*) plants to acute exposures (2 to 5 h) of aerosols (0.25 μm) containing only nitrate (109 $\mu\text{g}/\text{m}^3$), ammonium and nitrate (244 and 199 $\mu\text{g}/\text{m}^3$), or ammonium and sulfate (179 and 637 $\mu\text{g}/\text{m}^3$). They found that these exposures, which exceeded the range of naturally occurring aerosol concentrations, had little effect on foliar photosynthesis and conductance. Martin et al. (1992) concluded that future investigations should focus on the effects of particles on physiological characteristics of plants following chronic exposures.

Acidic Deposition. The effects of acidic deposition have been accorded wide attention in the media and elsewhere (Altshuller and Linthurst, 1984; Hogan et al., 1998). Probably the most extensive assessment of acidic deposition processes and effects is the NAPAP Biennial Report to Congress: An Integrated Assessment (National Science and Technology Council, 1998). Concern regarding the effects of acidic deposition on crops and forest trees has resulted in extensive monitoring and research. Exposures to acidic rain or clouds can be divided into “acute” exposures to higher ionic concentrations (several $\mu\text{mol}/\text{L}$) and “chronic” long-term repeated exposures to lower concentrations (Cape, 1993). Pollutant concentrations in rainfall have been shown to have little capacity for producing direct effects on vegetation (Altshuller and Linthurst, 1984; Hogan et al., 1998); however, fog and clouds, which may contain solute concentrations up to 10 times those found in rain, have the potential to cause direct effects. More than 80% of the ionic composition of most cloudwater is made up of four major pollutant ions: H^+ , NH_4^+ , NO_3^- , and SO_4^{2-} . Ratios of hydrogen to ammonium and sulfate to nitrate vary from site to site with all four ions usually present in approximately equal concentrations. Available data from plant effect studies suggest that hydrogen and sulfate ions are more likely to cause injury than ions containing nitrogen (Cape, 1993).

The possible direct effects of acidic precipitation on forest trees have been evaluated in experiments on seedlings and young trees. The size of mature trees makes experimental exposure difficult, therefore necessitating extrapolations from experiments on seedlings and saplings; however, such extrapolations must be used with caution (Cape, 1993). Both conifers and deciduous species have shown significant effects on leaf surface structures after exposure to simulated acid rain or acid mist at pH 3.5. Some species have shown subtle effects at pH 4 and

above. Visible lesions have been observed on many species at pH 3 and on sensitive species at pH 3.5 (Cape, 1993). The relative sensitivities of forest vegetation to acidic precipitation based on macroscopic injury have been ranked as follows: herbaceous dicots > woody dicots > monocots > conifers (Percy, 1991).

Huttunen (1994) described the direct effects of acid rain or acidic mist on epicuticular waxes whose ultrastructure is affected by plant genotype and phenotype. The effects of air pollutants on epicuticular waxes of conifers have received greater study than the waxes of other species. Leaf age and the shorter life span of broad-leaved trees make them less indicative of the effects of acid precipitation. Many experimental studies indicate that epicuticular waxes that function to prevent water loss from plant leaves can be destroyed by acid rain over a few weeks (Huttunen, 1994). Epicuticular waxes are crucial to conifers because of their longevity and evergreen foliage. Microscopic observations of epicuticular wax structures have, for a long time, suggested links between acidic deposition and aging. In Norway spruce (*Picea abies*), acid rain causes not only the aging of needles (which in northern conditions normally last from 11 to 14 years) to be shortened, but also accelerates the erosion rate of the waxes as the needles age.

The effects of acidic precipitation and fog on red spruce (*Picea rubens*) have been studied extensively (Schier and Jensen, 1992). Visible foliar injury of the needles in the form of a reddish-brown discoloration has been observed on red spruce seedlings experimentally exposed to acidic mist, but this visible symptom has not been observed in the field. Ultrastructural changes in the epicuticular wax were observed both experimentally and on spruce growing at high elevations. Laboratory studies indicate that visible injury usually does not occur unless the pH is 3 or less (Schier and Jensen, 1992). Cape (1993) reported that, when compared with other species, red spruce seedlings appeared to be more sensitive to acid mist. From studies of conifers and a review of the literature, Huttunen (1994) concluded that acidic precipitation causes direct injury to tree foliage and indirect effects through the soil. The indirect effects of acidic precipitation are discussed in Section 4.2.3.2.

Based on a review of the many studies in the literature involving field and controlled laboratory experiments on crops, Cape (1993) drew a number of conclusions concerning the direct effects of acidic precipitation on crops:

- foliar injury and growth reduction occur below pH 3;
- allocation of photosynthate is altered, with increased shoot to root ratios;
- expanded and recently expanded leaves are most susceptible, and injury occurs first to epidermal cells;
- leaf surface characteristics such as wettability, buffering capacity, and transport of material across the leaf surface contribute to susceptibility and differ among species;
- data obtained from experiments in greenhouses or controlled environmental chambers cannot be used to predict effects on plants grown in the field;
- quantitative data from experimental exposures cannot be extrapolated to field exposures because of differences and fluctuations in concentrations, durations, and frequency of exposure;
- there are large differences in response within species;
- timing of exposure in relation to phenology is of utmost importance;
- plants may be able to recover from or adapt to injurious exposures; and
- sequential exposure to acidic precipitation and gaseous pollutants is unlikely to be more injurious than exposure to individual pollutants.

Studies by Chevone et al. (1986), Krupa and Legge (1986), and Blaschke (1990) differ with the last conclusion of Cape listed above. Their studies indicate that interactions between acidic deposition and gaseous pollutants do occur. Acidity affects plant responses to both O₃ and SO₂. Chevone et al. (1986) observed increased visible injury on soybean and pinto bean when acid aerosol exposure preceded O₃ exposure; whereas linear decreases in dry root weight of yellow poplar occurred as acidity increased with simultaneous exposures to O₃ and simulated acid rain. Krupa and Legge (1986) also noted increased visible injury to pinto bean plants when aerosol exposure preceded O₃ exposure. In none of the studies cited above did acid rain per se produce significant growth changes. In contrast, Blaschke (1990) observed a decrease in ectomycorrhizal frequency and short root distribution caused by acid rain exposure in combination with either SO₂ or O₃.

Trace Elements. All but 10 of the 90 elements that comprise the inorganic fraction of the soil occur at concentrations of < 0.1% (1000 µg/g) and are termed “trace” elements or trace metals. Trace metals with a density greater than 6 g/cm³, referred to as “heavy metals,” are of

particular interest because of their potential toxicity for plant and animals. Although some trace metals are essential for vegetative growth or animal health, they are all toxic in large quantities. Combustion processes produce metal chlorides that tend to be volatile and metal oxides that tend to be nonvolatile in the vapor phase (McGowan et al., 1993). Most trace elements exist in the atmosphere in particulate form as metal oxides (Ormrod, 1984). Aerosols containing trace elements derive predominantly from industrial activities (Ormrod, 1984). Generally, only the heavy metals cadmium, chromium, nickel, and mercury are released from stacks in the vapor phase (McGowan et al., 1993). Concentrations of heavy metals in incinerator fly ash increase with decreasing particle size.

Vegetational surfaces, especially the foliage, present a major reaction and filtration surface to the atmosphere and act to accumulate particles deposited via wet and dry processes as described in Section 4.2.2 (Tong, 1991; Youngs et al., 1993). The chemical constituents of particles deposited on foliar surfaces may be taken up through the leaf surface. The greatest particle loading is usually on the upper leaf surface where particles accumulate in the mid-vein, center portion of the leaves. Additionally, fungal mycelia become particularly abundant on leaf surfaces as the growing season progresses and can be found in intimate association with deposited particles (Smith, 1990c).

Investigations of trace elements present along roadsides and in industrial and urban environments indicate that impressive burdens of particulate heavy metals can accumulate on vegetative surfaces. Foliar uptake of available metals could result in metabolic effects in above-ground tissues. Only a few metals, however, have been documented to cause direct phytotoxicity in field conditions. Copper, Zn, and Ni toxicities have been observed most frequently. Low solubility, however, limits foliar uptake and direct heavy metal toxicity because trace metals must be brought into solution before they can enter into the leaves or bark of vascular plants. In those instances when trace metals are absorbed, they are frequently bound in leaf tissue and are lost when the leaf drops off (Hughes, 1981). Trace metals in mixtures may interact to cause a different plant response when compared with a single element; however, there has been little research on this aspect (Ormrod, 1984). In experiments using chambers, Marchwińska and Kucharski (1987) studied the effects of SO₂ alone and in combination with the PM components Pb, Cd, Zn, Fe, Cu, and Mn obtained from a zinc smelter bag filter. The combined effects of SO₂ and PM further increased the reduction in yield of beans caused by SO₂;

whereas the combination, though severely injuring the foliage, produced little effect on carrots and parsley roots except after long-term exposures (when there was a decrease in root weight).

Trace metal toxicity of lichens has been demonstrated in relatively few cases. Nash (1975) documented zinc toxicity in the vicinity of a zinc smelter near Palmerton, PA. The richness and abundance of lichen species were reduced by ~90% in lichen communities at Lehigh Water Gap near the zinc smelter when compared with those at Delaware Water Gap. Zinc, Cd, and SO₂ were present in concentrations toxic to some species near the smelter; however, toxic zinc concentrations were detected farther away than the detectable limits of SO₂ (Nash, 1975). Experimental data suggest that lichen tolerance to Zn and Cd falls between 200 and 600 ppm (Nash, 1975).

Though there has been no direct evidence of a physiological association between tree injury and exposure to metals, heavy metals have been implicated because their deposition pattern is correlated with forest decline. The role of heavy metals has been assessed using phytochelatin measurements. Phytochelatin are intracellular metal-binding peptides that act as specific indicators of metal stress. Because they are produced by plants as a response to sublethal concentrations of heavy metals, they can indicate that heavy metals play a role in forest decline (Gawel et al., 1996). Concentrations of heavy metals increased with altitude, as did forest decline; and increased concentrations of heavy metals were found across the study region that showed increased levels of forest injury.

Phytochelatin concentrations were measured in red spruce and balsam fir (*Abies balsamea*) needles throughout the 1993 growing season at 1000 m on Whiteface Mountain in New York. Mean foliar concentrations in red spruce were consistently higher than in balsam fir from June until August, with the greatest and most significant difference occurring at the peak of the growing season in mid-July. In July, the phytochelatin concentrations were significantly higher than at any other time measured. Balsam fir also exhibited this peak, but maintained a consistently low level throughout the season. Both the number of dead red spruce trees and phytochelatin concentrations increased sharply with elevation (Gawel et al., 1996). The relationship between heavy metals and the decline of forests in northeastern United States was further tested by sampling red spruce stands showing varying degrees of decline at 1000 m on nine mountains spanning New Hampshire, Vermont, and New York. The collected samples indicated a systematic and significant increase in phytochelatin concentrations associated with

the extent of tree injury. The highest phytochelatin concentrations were measured during 1994 from sites most severely affected by forest decline in the Green Mountains in Vermont and the Adirondack Mountains in New York. These data strongly imply that metal stress causes tree injury and contributes to forest decline in the northeastern United States (Gawel et al., 1996).

One potential direct effect of heavy metals is on the activity of microorganisms and arthropods resident on and in the leaf-surface ecosystem. The fungi and bacteria living on and in the surfaces of leaves play an important role in the microbial succession that prepares leaves for decay and litter decomposition after their fall (U.S. Environmental Protection Agency, 1996b). Numerous fungi were consistently isolated from foliar surfaces at various crown positions from London plane trees (*Platanus hispanica*) growing in roadside environments in New Haven, CT. Those existing primarily as saprophytes included *Aureobasidium pullulans*, *Chaetomium* sp., *Cladosporium* sp., *Epicoccum* sp., and *Philaphora verrucosa*. Those existing primarily as parasites included *Gnomonia platani*, *Pestalotiopsis* sp., and *Pleurophomella* sp. The following cations were tested in vitro for their ability to influence the growth of these fungi: Cd, Cu, Mg, Al, Cr, Ni, Fe, Pb, Na, and Zn. Results indicated variable fungal response with no correlation between saprophytic or parasitic activity and sensitivity to heavy metals. Both linear extension and dry weight data indicated that the saprophytic *Chaetomium* sp. was very sensitive to numerous metals. *Aureobasidium pullulans*, *Epicoccum* sp., and especially *P. verrucosa*, on the other hand, appeared to be much more tolerant. Of the parasites, *G. platani* appeared to be more tolerant than *Pestalotiopsis* sp. and *Pleurophomella* sp. Metals producing the broadest spectrum growth suppression were Fe, Al, Ni, Zn, Mg, and Pb (Smith and Staskawicz, 1977; Smith, 1990c). These in vitro studies employed soluble compounds containing heavy metals. Trace metals probably occur naturally on leaf surfaces as low-solubility oxides, halides, sulfates, sulfides, or phosphates (Clevenger et al., 1991; Koslow et al., 1977). In the event of sufficient solubility and dose, however, changes in microbial community structure on leaf surfaces because of heavy metal accumulation are possible.

Organic Compounds. Volatile organic compounds in the atmosphere are partitioned between the gas and particle phases, depending on the liquid-phase vapor pressure at the ambient atmospheric temperature, the surface area of the particles per unit volume of air, the nature of the particles and of the chemical being adsorbed; and they can be removed from the atmosphere by

wet and dry deposition (McLachlan, 1996a). Materials as diverse as DDT, polychlorinated biphenyls (PCBs), and polynuclear aromatic hydrocarbons (PAHs) are being deposited from the atmosphere on rural, as well as urban, landscapes (Kylin et al., 1994). Motor vehicles emit particles to the atmosphere from several sources in addition to the tailpipe. Rogge et al. (1993b) inventoried the organic contaminants associated with fine particles (diameter $\leq 2.0 \mu\text{m}$) in road dust, brake-lining-wear particles, and tire-tread debris. More than 100 organic compounds were identified in these samples, including N-alkanols, benzoic acids, benzaldehydes, polyalkylene glycol ethers, PAHs, oxy-PAH, steranes, hopanes, natural resins, and other compound classes. A large number of PAHs, ranging from naphthalene (C_{10}H_8) to 5- and 6-ring and higher PAHs, their alkyl-substituted analogues, and their oxygen- and nitrogen-containing derivatives are emitted from motor vehicle sources (Seinfeld, 1989).

Plants may be used as environmental monitors to compare the deposition of PAH, persistent organic pollutants (POPs), or semivolatile organic components (SOCs) between sites (e.g., urban versus rural) (Wagrowski and Hites, 1997; Ockenden et al., 1998; McLachlan, 1999). Vegetation can be used qualitatively to indicate organic pollutant levels as long as the mechanism of accumulation is considered. The substance may enter the plant via the roots or, as noted above, be deposited as a particle onto the waxy cuticle of leaves or be taken up through the stomata. The pathways are a function of the chemical and physical properties of the pollutant, e.g., its lipophilicity, water solubility, vapor pressure (which controls the vapor-particle partitioning) and Henry's law constant; environmental conditions such as ambient temperature and the organic content of the soil; and the plant species, which controls the surface area and lipids available for accumulation (Simonich and Hites, 1995). Ockenden et al. (1998) have observed that, for lipophilic POPs, atmospheric transfer to plant has been the main avenue of accumulation. Plants can differentially accumulate POPs. Results have shown differences between species with higher concentrations in the lichen (*Hypogymnia physiodes*) than in Scots pine (*Pinus sylvestris*) needles. Even plants of the same species, because they have different growth rates and different lipid contents (depending on the habitat in which they are growing), have different rates of sequestering pollutants. These facts confound data interpretations and must be taken into account when considering their use as passive samplers.


Vegetation itself is an important source of hydrocarbon aerosols. Terpenes, particularly α -pinene, β -pinene, and limonene, released from tree foliage may react in the atmosphere to

form submicron particles. These naturally generated organic particles contribute significantly to the blue haze aerosols formed naturally over forested areas (Geron et al., 2000).

The low water solubility with high lipoaffinity of many of these organic xenobiotics strongly control their interaction with the vegetative components of natural ecosystems. The cuticles of foliar surfaces are covered with a wax layer that helps protect plants from moisture and short-wave radiation stress. This epicuticular wax, consisting mainly of long-chain esters, polyesters, and paraffins, has been demonstrated to accumulate lipophilic compounds. Organic air contaminants in the particulate or vapor phase are absorbed to, and accumulate in, the epicuticular wax of vegetative surfaces (Gaggi et al., 1985; Kylin et al., 1994). Direct uptake of organic contaminants through the cuticle or the vapor-phase uptake through the stomata are characterized poorly for most trace organics. The phytotoxicity and toxicity of organic contaminants to soil microorganisms is not well studied (Foster, 1991).

4.2.3.2 Ecosystem Response to Stresses

The ecosystem response to stress begins within the biotic condition attribute at the population level with changes in patterns resulting from the response of sensitive individual plants or animals. The ecosystem response to pollutant deposition is a direct function of the ecosystem's ability to ameliorate resulting change (Strickland et al., 1993). Plant responses, changes in both structural and compositional patterns, and functional ecological processes must be scaled in both time and space and be propagated from the individual to the more complex levels of community interaction to produce observable changes in an ecosystem (see Figure 4-14). Among ecosystem biota, at least three levels of biological interaction are involved: (1) the individual plant and its environment, (2) the population and its environment, and (3) the biological community composed of many species and its environment (Billings, 1978). Individual organisms within a population vary in their ability to withstand the stress of environmental change. The response of individual organisms within a population is based on their genetic constitution (genotype), stage of growth at time of exposure to stress, and the microhabitat in which they are growing (Levin, 1998). The range within which these organisms can exist and function determines the ability of the population to survive. Those able to cope with the stresses survive and reproduce. Competition among the different species results in succession (community change over time) and, ultimately, produces ecosystems composed of



Evaluating Impacts Within a Level of Organization			
Leaf Level	Carbon exchange-1 Carbon pools-2 Needle number and size-3 Needle retention/abscission-4	Tree Level	Height and diameter growth-10 Crown shape and size-11 Tree vigor-12 Mortality-13
Branch Level	Carbon allocation-5 Branch growth-6 Branch morphology-7 Branch vigor-8 Branch retention-9	Stand Level	Productivity-14 Mortality-15 Species composition-16


Evaluating Interactions Between Different Levels of Organization	
	<p>The diagonal arrow indicates the interaction between any two levels of organization. The types of interaction are due to the properties of variability and compensation.</p> <p>A - Refers to the interaction between the leaf and branch levels, where, for example, variability at the branch level determines leaf quantity, and compensation at the leaf level in photosynthesis may compensate for the reduction in foliage amount.</p> <p>B - Refers to the interaction between the branch and the tree, where variability in branches determines initial interception, branch vigor, and branch location in the crown; compensation may be related to increased radiation reaching lower branches.</p> <p>C - Refers to the interaction between the tree and the stand. Both genetic and environmental variability, inter- and intraspecific compensations, and tree historical and competitive synergisms are involved.</p>

Figure 4-14. Effects of environmental stress on forest trees are presented on a hierarchical scale for the leaf, branch, tree, and stand levels of organization. The evaluation of effects within a level of organization are indicated by horizontal arrows. The evaluation of interactions between different levels of organization are indicated by diagonal arrows.

Source: Hinckley et al. (1992).

populations of plant species that have the capability to tolerate the stresses (Rappport and Whitford, 1999; Guderian, 1985).

The number of species in a community usually increases during succession in unpolluted atmospheres. Productivity, biomass, community height, and structural complexity increase. Severe stresses, on the other hand, divert energy from growth and reproduction to maintenance and return succession to an earlier stage (Waring and Schlesinger, 1985). Ecosystems are subject to natural periodic stresses, such as drought, flooding, fire, and attacks by biotic pathogens (e.g., fungi, insects). Extremely severe natural perturbations return succession to an earlier stage; reduce ecosystem structure and functions (i.e., produce a scarcity of life forms and extinguish symbiotic interactions); disrupt the plant processes of photosynthesis and nutrient uptake, carbon allocation, and transformation that are directly related to energy flow and nutrient cycling; shorten food chains; and reduce the total nutrient inventory (Odum, 1993). This transformation, however, sets the stage for recovery that permits the perturbed ecosystem to adapt to changing environments (Holling, 1986). Therefore, these perturbations are seldom more than a temporary setback, and recovery can be rapid (Odum, 1969).

In contrast, anthropogenic stresses usually produce severe, debilitating effects. Severely stressed ecosystems do not recover readily, but may be further degraded (Odum, 1969; Rappport and Whitford, 1999). Anthropogenic stresses can be classified into four main groups:

(1) physical restructuring (e.g., changes resulting from land use), (2) introduction of exotic species, (3) over harvesting, and (4) discharge of toxic substances into the atmosphere, onto land, and into water. Ecosystems usually lack the capacity to adapt to the above stresses and maintain their normal structure and functions unless the stressor is removed (Rappport and Whitford, 1999). These stresses result in a process of ecosystem degradation marked by a decrease in biodiversity, reduced primary and secondary production, and a lower capacity to recover and return to its original state. In addition, there is an increased prevalence of disease, reduced nutrient cycling, increased dominance of exotic species, and increased dominance by smaller, short-lived opportunistic species (Odum, 1985; Rappport and Whitford, 1999).

Discharge of toxic substances into the atmosphere, onto land, and into water can cause acute and chronic stresses; and, once the stress is removed, a process of succession begins that can ultimately return the ecosystem to a semblance of its former structure. Air pollution stress, if acute, is usually short-term and the effects soon visible. Chronic stresses, on the other hand, are

long-term stresses whose effects occur at different levels of ecosystem organization and appear only after long-term exposures, as in the case of acidic deposition in the northeast or ozone in California (Shortle and Bondietti, 1992; U.S. Environmental Protection Agency, 1996b).

The possible effects of air pollutants on ecosystems have been categorized by Guderian (1977) as follows:

- (1) accumulation of pollutants in the plant and other ecosystem components (such as soil and surface- and groundwater),
- (2) damage to consumers as a result of pollutant accumulation,
- (3) changes in species diversity because of shifts in competition,
- (4) disruption of biogeochemical cycles,
- (5) disruption of stability and reduction in the ability of self-regulation,
- (6) breakdown of stands and associations, and
- (7) expansion of denuded zones.

How changes in these functions can result from PM deposition and influence ecosystems is discussed in the following text. It should be remembered that, although the effects of PM are being emphasized, the vegetational components of ecosystems also are responding to multiple stressors from multiple sources.

Response to Direct Effects of Particulate Matter

In the previous section, it was noted that PM affects patterns in the EEA biotic condition and processes in the chemical/physical and ecological processes categories. The presence of PM in the atmosphere may affect vegetation directly, following physical contact with the foliar surface (Section 4.2), but in most cases, the more significant effects are indirect. These effects may be mediated by suspended PM (i.e., through effects on radiation and climate) and by particles that pass through the vegetative canopies to the soil.

The majority of studies dealing with the direct effects of particulate dust and trace metals on vegetation were focused on responses of individual plant species and were conducted in the laboratory or in controlled environments (Saunders and Godzik, 1986). A few have considered the effects of particles on populations, communities, and ecosystems. Most of these focused on ecosystems in industrialized areas that are heavily polluted by deposits of both chemically inert

and active dusts. Effects can result from direct deposition or indirectly from deposition onto the soil. Reductions in growth, yield, flowering, and reproductive processes of plants from PM deposition have been reported (Saunders and Godzik, 1986). Sensitivities of individual species have been associated with changes in composition and structural patterns of natural ecosystems.

Evidence from studies of effects of PM deposition, specifically chemically inert and active dusts indicates that, within a population, plants exhibit a wide range of sensitivity which is the basis for the natural selection of tolerant individuals. Rapid evolution of certain populations of tolerant species at sites with heavy trace element and nitrate deposition has been observed (Saunders and Godzik, 1986). Tolerant individuals present in low frequencies in populations when growing in unpolluted areas have been selected for tolerance at both the seedling and adult stages when exposed to trace metal or nitrate deposition (Ormrod, 1984; U.S. Environmental Protection Agency, 1993). Chronic pollutant injury to a forest community may result in the loss of sensitive species, loss of tree canopy, and maintenance of a residual cover of pollutant-tolerant herbs or shrubs that are recognized as successional species (Smith, 1974) (see Table 4-13). These changes in forest patterns result from altered ecological processes.

Responses of ecosystems to stresses (unless severe or catastrophic) are difficult to determine because the changes are subtle (Garner, 1991). This is particularly true of responses to particles. Changes in the soil may not be observed until accumulation of the pollutant has occurred for 10 or more years, except in the severely polluted areas around heavily industrialized point sources (Saunders and Godzik, 1986). In addition, the presence of other cooccurring pollutants makes it difficult to attribute the effects to PM alone. In other words, the potential for alteration of ecosystem function and structure exists but is difficult to quantify, especially when there are other pollutants present in the ambient air that may produce additive or synergistic responses even though PM concentrations may not be elevated.

Physical Effects

The direct effects of limestone dust on plants and ecosystems has been known for many years. Changes have been observed in both ecosystem patterns and processes. Long-term changes in the structure and composition of the seedling-shrub and sapling strata of an experimental site near limestone quarries and processing plants in Giles County in southwestern Virginia were reported by Brandt and Rhoades (1972, 1973). Dominant trees in the control area,

**TABLE 4-13. ECOSYSTEM FUNCTIONS IMPACTED BY AIR POLLUTION
EFFECTS ON TEMPERATE FOREST ECOSYSTEMS**

Forest Soil and Vegetation: Activity and Response	Ecosystem Consequence and Impact
(1) Forest tree reproduction, alteration, or inhibition	(1) Altered species composition
(2) Forest nutrient cycling, alteration <ul style="list-style-type: none"> a. Reduced litter decomposition b. Increased plant and soil leaching and soil weathering c. Disturbance of microbial symbioses 	(2) Reduced growth, less biomass
(3) Forest metabolism <ul style="list-style-type: none"> a. Decreased photosynthesis b. Increased respiration c. Altered carbon allocation 	(3) Reduced growth, less biomass
(4) Forest stress, alteration <ul style="list-style-type: none"> a. Phytophagous insects, increased or decreased activity b. Microbial pathogens, increased or decreased activity c. Foliar damage increased by direct air pollution influence 	(4) Altered ecosystem stress: increased or decreased insect infestations; increased or decreased disease epidemics; and reduced growth, less biomass, and altered species composition

Source: Smith (1974).

a part of the oak-chestnut association of the eastern deciduous forests of eastern North America, were chestnut oak (*Quercus prinus*), red oak (*Q. rubra*), and red maple (*Acer rubrum*). An abundance of uniformly distributed saplings and seedlings were visible under the tree canopy, and herbs appeared in localized areas in canopy openings. Chestnut oak dominated the area, and the larger trees were 60 to 80 years old. The dusty site was dominated by white oak (*Q. alba*); whereas red oak and tulip poplars (*Liriodendron tulipifera*) were subcodominants. The largest trees were 100 years old and had necrotic leaves, peeling bark, and appeared to be in generally poor condition except for tulip poplars (which thrived in localized areas). The site contained a tangled growth of seedlings and shrubs, a few saplings, and a prevalence of green briar (*Smilax* spp.) and grape (*Vitis* spp.). The sapling strata in the area was represented by red maple, hickory (*Carya* spp.), dogwood (*Cornus florida*), and hop-hornbeam (*Ostrya virginiana*). Saplings of none of the leading dominant trees were of importance in this stratum. The most

obvious form of vegetation in the seedling-shrub stratum, because of their tangled appearance, were dogwood, hop-hornbeam, redbud (*Cercis canadensis*), and sugar maple (*Acer saccharum*).

Crust formation reduced photosynthesis, induced premature leaf fall and destruction of leaf tissues, inhibited growth of new tissue, and reduced the formation of carbohydrate needed for normal growth and storage (Brandt and Rhoades, 1973). The authors (Brandt and Rhoades, 1972), citing Odum (1969), also stated that one result of the accumulation of toxic pollutants in the biosphere as the result of human activities is the simplification of both plant and animal communities. In plant communities, structure is determined by sampling various strata within the community. Each stratum comprises a particular life form (e.g., herbs, seedlings, saplings, trees). Dust accumulation favored growth of some species and limited others. For example, sugar maple was more abundant in all strata of the dusty site when compared with the control site where it was present only as a seedling. The growth of tulip poplar, dogwood, hop-hornbeam, black haw (*Viburnum prunifolium*), and redbud (*C. canadensis*) appeared to be favored by the dust. Growth of conifers and acidophiles such as rhododendron (*Rhododendron maximum*), however, was limited. Although dust accumulation began in 1945, the heaviest accumulation occurred between 1967 and 1972 during the time of the study.

Changes in community composition were associated closely with changes in the growth of the dominant trees. A decrease in density of seedlings and saplings and in mean basal area, as well as lateral growth of red maple, chestnut oak, and red oak, occurred in all strata. On the other hand, all of these characteristics increased in the tulip poplar, which was a subordinate species before dust accumulation began but had assumed dominance by the time of the study. Reduction in growth of the dominant trees had apparently given the tulip poplar a competitive advantage because of its ability to tolerate dust. Changes in soil alkalinity occurred because of the heavy deposition of limestone dust; however, the facilities necessary for critical analysis of the soils were not available. From the foregoing, it is obvious that PM physical effects in the vicinity of limestone quarries and processing plants can affect ecosystems.

Changes in ecosystem structure resulting from exposures to sea salt were cited previously (Section 4.3.1.1). The dominance of live oak (*Quercus virginiana*) as a practically pure stand on Smith Island (Bald Head), NC and along the eastern and southern coast of North Carolina has been explained as due to its tolerance to salt spray. The absence of more inland species is

attributed to their intolerance to salt spray. Wells (1939) termed the long-term stabilization of live oak as “salt spray climax,” a new type of climax.

Acidic Deposition

The major effects of acidic deposition occur through the soil (National Science Technology Council, 1998) and are discussed below under indirect effects. However, included among the direct responses of forest trees to acidic deposition are increased leaching of nutrients from foliage; accelerated weathering of leaf cuticular surfaces; increased permeability of leaf surfaces to toxic materials, water, and disease agents; and altered reproductive processes (Cape, 1993; Altshuller and Linthurst, 1984).

Trace Elements

Possible direct responses of trace elements on vegetation result from their deposition and residence on the phyllosphere (i.e., foliar surfaces). Fungi and other microorganisms living on the leaves of trees and other vegetation play an important role in leaf decomposition after litterfall (Miller and McBride, 1999; Jensen, 1974; Millar, 1974). Possible effects of heavy metals on nutrient cycling and their effects on leaf microflora appear not to have been studied.

A trace metal must be brought into solution before it can enter into the leaves or bark of vascular plants. Low solubility limits entry. In those instances when trace metals are absorbed, they frequently are bound in the leaf tissue and then are lost from the plant when the leaf drops off (Hughes, 1981) and transferred to the litter layer where they can affect litter decomposition, an important source of soil nutrients. Changes in litter decomposition processes influence nutrient cycling in the soil and limit the supply of essential nutrients. Both Cotrufo et al. (1995) and Niklińska et al. (1998) point out that heavy metals affect forest litter decomposition. Cotrufo et al. (1995) observed that decomposition of oak leaves containing Fe, Zn, Cu, Cr, Ni, and Pb was influenced strongly during the early stages by metal contamination. Fungal mycelia were significantly less abundant in litter and soil in contaminated sites when compared with control sites. Niklińska et al. (1998) stated that toxic effects of heavy metals on soil respiration rate have been reported by many scientists, and that, in polluted environments, this results in accumulation of undecomposed organic matter. However, they state that results of experiments should identify the most important “natural” factors affecting soil/litter sensitivity because the

effects of heavy metals on respiration rates are dose-dependent on the dose of heavy metals, the type of litter, types of metals deposited, and the storage time before respiration tests are made.

Trace metals, particularly heavy metals (e.g., Cd, Cu, Pb, Cr, Hg, Ni, Zn) have the greatest potential for influencing forest growth (Smith, 1991). Experimental data indicate that the broadest spectrum of growth suppression of foliar microflora resulted from Fe, Al, and Zn. These three metals also inhibited fungal spore formation, as did Cd, Cr, Mg, and Ni (see Smith, 1990c). In the field, the greatest injury occurs from pollution near mining, smelting, and other industrial sources (Ormrod, 1984). Direct metal phytotoxicity can occur only if the metal can move from the surface into the leaf or directly from the soil into the root.

Organic Compounds

Primary and secondary organic compounds formed in the atmosphere, the effects of some of which are discussed below, have been variably referred to under the following terms: toxic substances, hazardous air pollutants (HAPs), air toxics, semivolatile organic compounds (SVOCs), pesticides, and persistent organic pollutants (POPs). Again, it should be remembered that chemical substances denoted by such headings are not criteria air pollutants controlled by the National Ambient Air Quality Standards under Section 109 of the Clean Air Act (CAA), but rather are controlled under Sect. 112, Hazardous Air Pollutants (U.S. Code, 1991). Their possible effects on humans and ecosystems are discussed in a number of government documents and in many other publications. They are mentioned here because many of the chemical compounds are partitioned between gas and particle phases in the atmosphere. As particles, they can become airborne, be distributed over a wide area, and affect remote ecosystems. Some of the chemical compounds are of concern because they may reach toxic levels in food chains of both animals and humans; whereas others tend to decrease or maintain the same toxicity as they move through the food chain. Some examples of movement through food chains are provided below.

Many chemical compounds from a variety of anthropogenic sources are released into the ambient air (see Section 4.2.1). In the atmosphere, the emitted compounds initially go through a mixing process, and the airborne particles then are distributed over a wide area and ultimately deposited on ecosystem components. Atmospheric deposition of polychlorinated dibenzo-*p*-dioxins and dibenzofurans (PCDD/DFs), as an example, can be divided into three different

forms: (1) dry gaseous, (2) dry particle-bound, and (3) wet deposition. Dry particle-bound deposition occurs when the PM containing the pollutant is deposited on the plant surface; whereas wet deposition ranges from hail to rain to fog and dewfall (McLachlan, 1996b).

Human exposure to PCDD/DFs has been demonstrated to be caused almost exclusively by the ingestion of animal fat from fish, meat, and dairy products. Almost half of all human exposure to PCDD/DFs is caused by consumption of beef and dairy products (McLachlan, 1996b). Cattle obtain most of their PCDD/DFs through grass. Therefore, the grass → cattle → milk/beef pathway is critical for human exposure. It has been shown that root uptake/translocation is an insignificant pathway of PCDD/DFs compared to absorption via aerial plant parts. Wet and dry particle deposition are the most important for the accumulation of the higher chlorinated congeners in vegetation. The persistence of PCDD/DFs in plants has not been investigated extensively; however, biodegradation probably does not occur in that these compounds are found primarily in the lipophilic cuticle and are very resistant to microbial degradation (McLachlan, 1996b). Feed contaminated with soil containing the pollutant can be another source of exposure of beef and dairy cattle, as well as chickens. The PCDD/DFs levels are near a steady state in milk cows and laying hens; however, animals raised for meat production (such as beef cattle and pigs) may accumulate large amounts. Beef cattle and pigs cannot excrete the contaminants in a lipid-rich matrix such as milk or eggs. Thus, all of the PCDD/DFs ingested are stored in the body. In agricultural food chains, there is a biodilution of PCDD/DFs, with the fugacity decreasing by up to three orders of magnitude between the air and cows milk (McLachlan, 1996b). Fürst et al. (1993), based on surveys to determine the factors that influence the presence of PCDD/DF in cows milk, earlier concluded that regardless of which pathway, soil → grass → cow or air → grass → cow, it was the congener of the chemical that was most important.

Chlorinated persistent organic pollutants (POPs), such as PCBs, PCDFs, and PCDDs, can be transported as particles through the atmosphere from industrial and agricultural sources; be brought down via wet and dry deposition in remote regions, such as the Arctic; and have been detected in all levels of the Arctic food chain (Oehme et al., 1995). High concentrations of PCB (1 to 10 ppm) were found in seals, but the concentrations increased to 10 to 100 ppm in polar bears. The polar bear is the top predator in the Arctic and feeds preferentially on ringed seals and, to a lesser extent, on other seal species. Bioconcentration factors of organochlorines in the

Arctic food web, reaching 10^7 for fish and seals, are biomagnified in polar bears (Oehme et al., 1995). Polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans have also been found in seals (Oehme et al., 1995). Milk taken from anesthetized polar bears was also found to contain PCDD/DF. Very little is known regarding the intake of milk by polar bear cubs. However, estimates of the intake of milk containing detectable levels of PCDD/DF and PCB and the additional consumption of seal blubber confirm that these pollutants are passed on to the next generation (Oehme et al., 1995).

Section 112 of the CAA provides the legislative basis for U.S. hazardous air pollutant (HAP) programs. In response to mounting evidence that air pollution contributes to water pollution, Congress included Section 112m (*Atmospheric Deposition to Great Lakes and Coastal Waters*) in the 1990 CAA Amendments that direct the EPA to establish a research program on atmospheric deposition of HAPS to the “Great Waters.”

Actions taken by EPA and others to evaluate and control sources of Great Waters pollutants of concern appear to have positively affected trends in pollutant concentrations measured in air, sediment, and biota. Details concerning these effects may be found in *Deposition of Air Pollutants to the Great Waters*, Third Report to Congress (U.S. Environmental Protection Agency, 2000a). The Third Report (EPA-453/R-00-005, June 2000), like the First and Second Reports to Congress, focuses on 15 pollutants of concern, including pesticides, metal compounds, chlorinated organic compounds, and nitrogen compounds. The new scientific information in the Third Report supports and builds on three broad conclusions presented in the previous two EPA Reports to Congress:

- (1) Atmospheric deposition from human activities can be a significant contributor of toxic chemicals and nitrogen compounds to the Great Waters. The relative importance of atmospheric loading for a particular chemical in a water body depends on many factors (e.g., characteristics of the water body, properties of the chemical, and the kind and amount of atmospheric deposition versus water discharges).
- (2) A plausible link exists between emissions of toxic pollutants of concern into the air above the Great Waters; the deposition of these pollutants (and their transformation products); and the concentrations of these pollutants found in the water, sediments, and biota, especially fish and shellfish. For mercury, fate and transport modeling and exposure assessments predict that the anthropogenic contribution to the total amount of methylmercury in fish is, in part, the result of anthropogenic mercury releases from industrial and combustion sources increasing mercury body burdens (i.e., concentrations) in fish. Also, the consumption of fish is the dominant pathway of

exposure to methylmercury for fish-consuming humans and wildlife. However, what is known about each stage of this process varies with each pollutant (for instance, the chemical species of the emissions and its transformation in the atmosphere).

- (3) Airborne emissions from local as well as distant sources, from both within and outside the United States, contribute pollutant loadings to waters through atmospheric deposition. Determining the relative roles of particular sources — local, regional, national, and possibly global, as well as anthropogenic, natural, and reemission of pollutants — contributing to specific water bodies is complex, requiring careful monitoring, atmospheric modeling, and other analytical techniques.

Response to Indirect Effects of Particulate Matter

The presence of PM in the atmosphere directly affects vegetation following physical contact with foliar surfaces (as discussed above), but in many cases the more significant effects are indirect. These effects may be mediated by suspended PM (i.e., through effects on radiation and climate) and by particles that pass through vegetative canopies to reach the soil. Effects mediated through the atmosphere are considered briefly below and in greater detail later, under Section 4.5.

The major indirect plant responses are chiefly soil-mediated and depend primarily on the chemical composition of the individual stressors deposited in PM. The chemical stressors must be bioavailable in order to produce an effect. The effects of exposures may result in changes in biota patterns and in chemical/physical soil conditions that affect ecological processes, such as nutrient cycling and uptake by plants.

The soil environment (composed of mineral and organic matter, water, air, and a vast array of bacteria, fungi, algae, actinomycetes, protozoa, nematodes, and arthropods) is one of the most dynamic sites of biological interactions in nature (Wall and Moore, 1999; Alexander, 1977). The quantity of organisms in soils varies by locality. Bacteria and fungi are usually most abundant in the rhizosphere, the soil around plant roots that all mineral nutrients must pass through. Bacteria and fungi benefit from the nutrients in the root exudates (chiefly sugars) in the soil and, in turn, they play an essential role by making mineral nutrients available for plant uptake (Wall and Moore, 1999; Rovira and Davey, 1974). Their activities create chemical and biological changes in the rhizosphere by decomposing organic matter and making inorganic minerals available for plant uptake. Bacteria are essential in the nitrogen and sulfur cycles and make these elements available for plant uptake and growth (see Section 4.3.3). Fungi are

directly essential to plant growth. Attracted to their roots by the exudates, they develop mycorrhizae in a symbiotic fungus-plant relationship, that is integral in the uptake of the mineral nutrients (Allen, 1991). The impact in ecosystems of PM, particularly nitrates, sulfates, and metals, is determined by their effects on the growth of the bacteria involved in nutrient cycling and the mycorrhizal fungi involved in plant nutrient uptake.

Atmospheric Turbidity: Effects on Direct Beam and Photosynthetically Active

Radiation. Photosynthetic processes underlie the contribution of vegetative surfaces to nutrient and energy cycling. The characteristics and net receipts of environmental radiation determine the rates of both photosynthesis and the heat-driven process of water cycling. Atmospheric turbidity due to particulate loading can substantially alter the characteristics and net receipts of solar radiation. One measure of atmospheric turbidity, Linke's turbidity factor, T, can be derived as a direct function of light extinction by solid particles. It is defined as the ratio of the total extinction coefficient and the extinction due exclusively to gases:

$$T = \sigma / \sigma_g = 1 + w\sigma_w / \sigma_g + s\sigma_s / \sigma_g \quad (4-6)$$

where s and w are the relative concentrations of dust and water vapor in the atmosphere, and σ_s and σ_w are the wavelength-dependent scattering coefficients for solid, dry particles and water vapor, respectively. The scattering coefficients are in units of inverse distance, such as km^{-1} (Rosenberg et al., 1983). According to this expression, a clean atmosphere would have a turbidity value of 1. Given that turbidity and visibility are both functions of light scattering, the trends in, and physical processes underlying, reduced visibility discussed in Section 4.3 are directly relevant to the discussion of radiative effects on vegetation due to PM.

Turbidity, as defined above, describes the degree of scattering occurring in the atmosphere due to particles and gases. Total, particle-based extinction, however, is the sum of both scattering and absorption. Absorption of short-wavelength solar radiation reduces the amount of radiation reaching the Earth's surface and leads to atmospheric heating. If the absorbing particles re-radiate in the infrared range, some of this energy is lost as long-wave re-radiation to space. The balance of this energy is captured at the surface as down-welling infrared radiation. Canopy temperature and transpirational water use by vegetation are particularly sensitive to

long-wave, infrared radiation. Atmospheric heating by particles reduces vertical temperature gradients, potentially reducing the intensity of atmospheric turbulent mixing. The magnitude of such potential effects on turbulent transport within canopies remains unknown although the damping of eddy transport may inhibit canopy gas exchange. Suppressed tropospheric mixing could also intensify local temperature inversions and increase the severity of pollution episodes (Pueschel, 1993) with direct inhibitory effects on photosynthetic processes.

Atmospheric turbidity increases the intensity of diffuse (sky) radiation (Hoyt, 1978). In a clear atmosphere, diffuse radiation may be on the order of 10% of total solar radiation (Choudhury, 1987). However, in highly turbid, humid conditions, this fraction may increase, to as much as 100% of the incident solar intensity in extreme cases. The direct-radiation diffuse ratio is highest at solar noon and lowest near dawn or dusk when the path length through the atmosphere is longest. The wavelength dependence of particle scattering induces an enrichment of PAR with respect to total or direct beam radiation. The influence of this enrichment on plant life is discussed in the next section.

Aerosols produced by incomplete combustion, from forest fires to specifically anthropogenic processes such as diesel fuel combustion, contain significant fractions of black carbon which absorbs across the solar and terrestrial radiation spectra. The presence of absorbing aerosols reduces the ratio of photosynthetically active radiation to total radiation received at the surface, potentially reducing photosynthetic water uptake efficiency. The net effect of aerosol absorption on the surface depends on the relative magnitudes of the particulate absorption coefficients in the visible and infrared area and on the albedo of the Earth's surface.

The greater effect of particulate loading on visibility and turbidity and, therefore, radiation receipts by the biosphere, is due to scattering. Non-absorbing, scattering aerosols raise the overall albedo of the atmosphere and reduce the amount of radiation reaching the surface by the amount reflected or scattered back into space. Analysis of data collected by a global network of thermopile pyranometers operated by the World Meteorological Organization (WMO) show a 50-year global trend of a 2.7% per decade reduction in the amount of solar radiation reaching the Earth's surface. This has been associated with an increasing global albedo caused by an increasing abundance of atmospheric particles. By evaluating the WMO data set with four different approaches to the statistical analyses, Stanhill and Cohen (2001) have estimated that average global solar radiation receipts have declined by 20 W m^{-2} since 1958. Examples of

individual measurement sites include Barrow, Alaska (71 °N), where the average solar irradiance from 1963 to 1993 was 100.9 W m^{-2} and the estimated linear trend was -0.23% per year; and Jerusalem, Israel (32 °N), where the average solar irradiance from 1954 to 1994 was 244.2 W m^{-2} and the estimated linear trend was -0.37% per year. This phenomenon has been labeled “global dimming.”

Increased Turbidity and Altered Radiative Flux: Effects on Vegetative Processes.

In a detailed canopy-response model (e.g., Choudhury, 1987), radiation is considered in its direct and diffuse components. Foliar interception by canopy elements is considered for both up- and down-welling radiation (a two-stream approximation). In this case, the effect of atmospheric PM on turbidity influences canopy processes both by radiation attenuation and by influencing the efficiency of radiation interception throughout the canopy through conversion of direct to diffuse radiation (Hoyt, 1978). Diffuse radiation is more uniformly distributed throughout the canopy and increases canopy photosynthetic productivity by distributing radiation to lower leaves. The treatment of down-welling direct-beam radiation in the two-stream approach remains an elaboration of the simplified Beer’s Law analogy with solar angle, leaf area distribution, and orientation individually parameterized (Choudhury, 1987). Diffuse down-welling radiation is a function of diffuse and direct radiation at the top of the canopy and penetration within the canopy according to cumulative leaf area density and foliage orientation. Diffuse up-welling radiation results from scattering and reflectance of both direct and diffuse down-welling radiation within the canopy and by the soil.

Rochette et al. (1996) conducted simultaneous measurements of radiation and water use efficiencies by maize and found that, in the absence of water stress and with adequate fertilization, 90% of all variation in crop net photosynthesis (P-n) could be explained by variations in PAR. Alternatively, an evaluation of the available experimental literature and statistics on crop yields by Stanhill and Cohen (2001) indicate that plant productivity is more affected by changes in evapotranspiration induced by changes in the amount of solar radiation plants receive than by changes in the amount of PAR plants receive.

The enrichment in PAR present in diffuse radiation appears, however, to offset a portion of the effect of an increased atmospheric albedo due to atmospheric particles. An observational and theoretical study by Bange et al. (1997) of the level of radiation use efficiency (RUE) of

sunflowers indicated a degree of compensation for reduced incident radiation by a proportional increase in diffuse radiation. Variables measured by Bange et al. (1997) included biomass, phenology, leaf area, canopy light extinction, grain size, and harvest index. Crops subject to reduced direct beam/increased diffuse radiation produced biomass, phenology, leaf area and canopy light extinction at leaves similar to unshaded crops but yielded smaller grains and a lower harvest index. RUE was also seen to improve for soybeans and maize with a proportional increase in diffuse radiation with respect to direct beam (Sinclair et al., 1992; Healey et al., 1998) although the effect on harvest index was not indicated. Gu et al. (2002) compared the relative efficiencies of canopy photosynthesis to diffuse and direct PAR for a Scots pine forest, an aspen forest, a mixed deciduous forest, a tall grass prairie, and a winter wheat crop. They concluded (1) diffuse radiation over direct radiation results in higher light use efficiencies by plant canopies, (2) diffuse radiation has much less tendency to cause canopy photosynthetic saturation, (3) the advantages of diffuse radiation over direct radiation increase with radiation level, (4) temperature as well as vapor pressure deficit can cause different responses in diffuse and direct canopy photosynthesis, indicating that their effects on terrestrial ecosystem carbon assimilation may depend upon radiation regimes, i.e., sky conditions.

The potentially significant effect of regional haze on the yield of crops because of reduction in solar radiation has been examined by Chameides et al. (1999). Using a case study approach, Chameides et al. (1999) studied the effects of regional haze on crop production in China where regional haze is especially severe. A simplified assessment of the direct effect of atmospheric aerosols on agriculture suggests that yields of approximately 70% of crops are being depressed by at least 3 to 5% by regional scale air pollution and its associated haze (Chameides et al., 1999).

Effects of Nitrogen Deposition. Nitrogen is required by all organisms. It is a major constituent of the nucleic acids that determine the genetic character of all living things and the enzyme proteins that drive the metabolic machinery of every living cell (Galloway, 1998; Galloway and Cowling, 2002; U.S. Environmental Protection Agency, 1993). Though nitrogen composes 80% of the total mass of the Earth's atmosphere, it is not biologically available. Nitrogen fixation is accomplished in nature by certain unique organisms that have developed the capability of converting N_2 to biologically active reduced forms of nitrogen such as ammonia,

amines, and amino acids, which are the structural constituents of proteins and nucleic acids (Galloway and Cowling, 2002; Hornung and Langan, 1999; U.S. Environmental Protection Agency, 1993).

Nitrogen has long been recognized as the nutrient most important for plant growth. It is of overriding importance in plant metabolism and, to a large extent, governs the utilization of phosphorus, potassium, and other nutrients. Most of the nitrogen in soils is associated with organic matter. Typically, the availability of nitrogen via the nitrogen cycle controls net primary productivity, and possibly, the decomposition rate of plant litter. Photosynthesis is influenced by nitrogen uptake in that ~75% of the nitrogen in a plant leaf is used during the process of photosynthesis. The nitrogen-photosynthesis relationship is, therefore, critical to the growth of trees and other plants (Hornung and Langan, 1999; Chapin et al., 1987). Plants usually obtain nitrogen directly from the soil through their roots by absorbing NH_4^+ or NO_3^- , or it is formed by symbiotic organisms (e.g., bacteria, blue-green algae) in the roots.

Because nitrogen is not readily available and is usually in short supply, it is the chief element in agricultural fertilizers (Hornung and Langan, 1999). The realization of the importance of nitrogen in crop production resulted in a search for natural nitrogen sources such as guano and nitrate deposits. The invention of the Haber-Bosch process in 1913 made reactive nitrogen (Nr) available for use in food production, and more than half of the food eaten by the peoples of the world today is produced using fertilizer produced by this process (Galloway, 1998; Galloway and Cowling, 2002).

In nature, nitrogen may be divided into two groups: nonreactive (N_2) and reactive (Nr) nitrogen. Reactive N includes all biologically, photochemically, and radioactively active nitrogen compounds in the Earth's atmosphere and biosphere (Galloway et al., 2003). Among those included are the inorganic reduced forms of nitrogen (e.g., NH_3 and NH_4^+), inorganic oxidized forms (e.g., NO_x , HNO_3 , N_2O , and NO_3^-), and organic compounds (e.g., urea, amine, proteins, and nucleic acids) (Galloway et al., 2003).

Food production continues to account for most of the newly created Nr. However, since around 1965, the magnitude of Nr created by humans began to exceed natural terrestrial creation of Nr and its conversion back to N_2 by denitrification. The overall increase in global Nr is the result of three main causes: (1) widespread cultivation of legumes, rice, and other crops that promote conversion of N_2 to organic nitrogen through biological nitrogen fixation;

(2) combustion of fossil fuels, which converts both atmospheric N_2 and fossil nitrogen to reactive NO_x ; and (3) the Haber-Bosch process, which converts nonreactive NH_3 to sustain food production and some industrial activities (Galloway and Cowling, 2002; Galloway et al., 2003).

Reactive nitrogen is now accumulating in the environment on all spatial scales $\frac{1}{m}$ local, regional and global (Galloway, 1998; Galloway and Cowling, 2002; Galloway et al. 2003). As a result, Nr is accumulating in various environmental reservoirs, e.g., the atmosphere, soils and waters (Galloway and Cowling, 2002). The accumulation of Nr in the environment has effects on humans and ecosystems (Rabalais, 2002; van Egmond et al., 2002; Galloway, 1998).

Large uncertainties regarding the rates of Nr accumulation in the various reservoirs limits our ability to determine the temporal and spatial distribution of environmental effects. These uncertainties are of great significance because of the sequential nature of Nr on environmental processes (Galloway and Cowling, 2002). The sequence of transfers, transformations, and environmental effects is referred to as the nitrogen cascade (Figure 4-15) (Galloway and Cowling, 2002; Galloway et al, 2003). A single atom of new NH_x or NO_x can alter a wide array of biogeochemical processes and exchanges among environmental reservoirs.

The results of the Nr cascade in the global system and the wide variety of changes in the nitrogen cycle are both beneficial and detrimental to humans and to ecosystems (Galloway and Cowling, 2002; Galloway et al., 2003). Though the synthetic fertilizers used in cultivation and cultivation-induced bacterial nitrogen fertilization (BNF) sustain a large portion of the world's population, there are consequences: (1) the wide dispersal of Nr by hydrological and atmospheric transport; (2) the accumulation of Nr in the environment because the rates of its creation are greater than the rates of its removal through denitrification to nonreactive N_2 ; (3) Nr creation and accumulation is projected to continue to increase in the future as per capita use of resources by human populations increases; and (4) Nr accumulation contributes to many contemporary environmental problems (Galloway et al., 2003).

Among the contemporary environmental problems as listed in Galloway et al. (2003) are the following:

- increases in Nr lead to production of tropospheric ozone and aerosols and the associated human health problems (Wolfe and Patz, 2002).
- productivity increases in forests and grasslands and then decreases wherever atmospheric Nr deposition increases significantly and critical thresholds are exceeded; Nr additions probably also decrease biodiversity in many natural habitats (Aber et al., 1995);




Figure 4-15. Illustration of the nitrogen cascade showing the movement of the human-produced reactive nitrogen (Nr) as it cycles through the various environmental reservoirs in the atmosphere, terrestrial ecosystems, and aquatic ecosystems.

Source: Galloway et al. (2003).

- reactive nitrogen in association with sulfur is responsible for acidification and loss of biodiversity in lakes and streams in many regions of the world (Vitousek et al., 1997);
- reactive nitrogen is responsible for eutrophication, hypoxia, loss of biodiversity, and habitat degradation in coastal ecosystems. It is now considered the biggest pollution problem in coastal waters (Rabalais, 2002);
- reactive nitrogen contributes to global climate change and stratospheric ozone depletion, both of which have an effect on the health of humans and ecosystems (Cowling et al., 1998).

The effect of increasing nitrogen inputs (e.g., NO_x , NO_3^- , HNO_3) on the nitrogen cycle in forests, wetlands, and aquatic ecosystems is discussed in detail elsewhere (U.S. Environmental Protection Agency, 1993, 1997a; Garner, 1994; World Health Organization, 1997).

The deposition of nitrogen in the United States from human activity has doubled between 1961 and 1997 due mainly to the use of inorganic nitrogen fertilizers and NO_x emissions from fossil fuel combustion, with the largest increase occurring in the 1960s and 1970s (Howarth et al., 2002). Among the most important effects of chronic nitrogen deposition are changes in the composition of plant communities, disruptions in nutrient cycling, increased soil emissions of nitrogenous greenhouse gases, accumulation of nitrogen compounds in the soil with enhanced availability of nitrate or ammonium, soil-mediated effects of acidification, and increased susceptibility to stress factors in plants (Fenn et al., 1998; Bobbink et al., 1998). A major concern is “nitrogen saturation,” the result of the atmospheric deposition of large amounts of particulate nitrates. Nitrogen saturation results when additions to soil background nitrogen (nitrogen loading) exceed the capacity of plants and soil microorganisms to utilize and retain nitrogen (Aber et al., 1989, 1998; Garner, 1994; U.S. Environmental Protection Agency, 1993). Under these circumstances, disruptions of ecosystem functions may result (Hornung and Langan, 1999).

Possible ecosystem responses to nitrate saturation, as postulated by Aber and coworkers (Aber et al., 1989), include (1) a permanent increase in foliar nitrogen and reduced foliar phosphorus and lignin caused by the lower availability of carbon, phosphorus, and water; (2) reduced productivity in conifer stands because of disruptions of physiological function; (3) decreased root biomass and increased nitrification and nitrate leaching; and (4) reduced soil fertility, resulting from increased cation leaching, increased nitrate and aluminum concentrations in streams, and decreased water quality. Nitrate saturation implies that some resource other than nitrogen is limiting biotic function. Water and phosphorus and carbon are the resources most likely to be the secondary limiting factors for plants and microorganisms, respectively (Aber et al., 1989). The appearance of nitrogen in soil solutions is an early symptom of excess nitrogen. In the final stage, disruption of forest structure becomes visible (Garner, 1994).

Changes in nitrogen supply can have a considerable effect on an ecosystem's nutrient balance (Waring, 1987). Large chronic additions of nitrogen disrupt normal nutrient cycling and alter many plant and soil processes involved in nitrogen cycling (Aber et al., 1989). Among the

processes affected are (1) nutrient uptake and allocation in plants, (2) litter production, (3) immobilization (includes ammonification [the release of ammonia] and nitrification [the conversion of ammonia to nitrate during decay of litter and soil organic matter]), and (4) nitrate leaching and trace gas emissions (Figure 4-16) (Aber et al., 1989; Garner 1994).




Figure 4-16. Nitrogen cycle (dotted lines indicate processes altered by nitrogen saturation).

Source: Garner (1994).

Subsequent studies have shown that, although an increase in nitrogen mineralization occurs initially (i.e., the conversion of soil organic matter to nitrogen in available form [see item 3 above]), not nitrogen mineralization rates are reduced under nitrogen-enriched conditions. Aber et al. (1998) hypothesized that mycorrhizal assimilation and exudation, using photosynthate from the host plant as the carbon source, is the dominant process involved in immobilization of nitrogen. In addition, studies suggested that soil microbial communities change from predominantly fungal (mycorrhizal) communities predominately bacterial during nitrate saturation (Aber et al., 1998).

The growth of most forests in North America is limited by the nitrogen supply. Severe symptoms of nitrogen saturation, however, have been observed in high-elevation, nonaggrading spruce-fir ecosystems in the Appalachian Mountains, as well as in the eastern hardwood watersheds at Fernow Experimental Forest near Parsons, WV. Mixed conifer forests and chaparral watersheds with high smog exposure in the Los Angeles Air Basin also are nitrogen saturated and exhibit the highest stream water NO_3^- concentrations for wildlands in North America (Bytnerowicz and Fenn, 1996; Fenn et al., 1998). Forests in southern California, the southwestern Sierra Nevada in central California, and the Front Range in northern Colorado have all been exposed to highly elevated nitrogen deposition, and nitrogen-saturated watersheds have been reported in the above mentioned areas. Annual nitrogen additions through deposition (6 to $11 \text{ kg ha}^{-1} \text{ year}^{-1}$ as through fall) in the southwestern Sierra Nevada are similar to the nitrogen storage ($4 \text{ kg ha}^{-1} \text{ year}^{-1}$) in vegetation growth increment of western forests suggesting that current nitrogen deposition rates may be near the assimilation capacity of the overstory vegetation. Ongoing urban expansion will increase the potential for nitrogen saturation of forests from urban sources (e.g., Salt Lake City, Seattle, Tucson, Denver, central and southern California) unless there are improved emission controls (Fenn et al., 1998).

Atmospherically deposited nitrogen also can act as a fertilizer in soil low in nitrogen. Not all plants, however, are capable of utilizing extra nitrogen, as plants vary in their ability to absorb NH_4^+ and NO_3^- (Chapin, et al., 1987). Inputs of nitrogen to natural ecosystems that alleviate deficiencies and increase growth of some plants can alter competitive relationships and alter species composition and diversity (Ellenberg, 1987; Kenk and Fischer, 1988; U.S. Environmental Protection Agency, 1993).

Not all forest ecosystems react in the same manner to nitrogen deposition. High-elevation alpine watersheds in the Colorado Front Range (Bowman, 2000) and a deciduous forest in Ontario, Canada also are naturally saturated even though nitrogen deposition has been moderate ($\approx 8 \text{ kg ha}^{-1} \text{ year}^{-1}$). The nitrogen-saturated forests in North America, including estimated inputs and outputs, are shown in Table 4-14 (Fenn et al., 1998). The Harvard Forest hardwood stand in Massachusetts, however, has absorbed $> 900 \text{ kg N/ha}$ without significant NO_3^- leaching during an 8-year nitrogen amendment study (Table 4-14) (Fenn et al., 1998). Nitrate leaching losses were high, on the other hand, in the Harvard Forest pine sites suggesting that deciduous forests may have a greater capacity for nitrogen retention. In the 8-year experimental study (1988 to 1996), nitrate leaching was observed in the pine stand after the first year (1989) in the high-nitrogen plots. Further increases were observed in 1995 and 1996, while the hardwood stand did not show significant increases in NO_3^- leaching until 1996. The sharp contrast in the response of the pine and hardwood stands indicates that the mosaic of community types across the landscape must be considered when determining the regional scale response to nitrogen deposition (Magill et al., 2000). Johnson et al. (1991a) reported that measurements showing the leaching of NO_3^- and Al^{+3} from high elevation forests in the Great Smoky Mountains indicate that these forests have reached saturation.

Because the competitive equilibrium of plants in any community is finely balanced, the alteration of one of any number of environmental parameters, (e.g., continued nitrogen additions) can change the vegetation structure of an ecosystem (Bobbink, 1998; Skeffington and Wilson, 1988). Increases in soil nitrogen play a selective role. When nitrogen becomes more readily available, plants adapted to living in an environment of low nitrogen availability will be replaced by plants capable of using increased nitrogen because they have a competitive advantage.

Plant succession patterns and biodiversity are affected significantly by chronic nitrogen additions in some North American ecosystems (Figure 4-17). The location of nitrogen saturated ecosystems in North America, and the steps leading to nitrogen saturation, are indicated on the map in Figure 4-16. Conceptual models of regional nitrogen saturation indicate saturation in New England, in the Colorado alpine ecosystems and in California forests. Fenn et al. (1998) reported that long-term nitrogen fertilization studies in both New England and Europe, as well, suggest that some forests receiving chronic inputs of nitrogen may decline in productivity and

**TABLE 4-14. NITROGEN-SATURATED FORESTS IN NORTH AMERICA,
INCLUDING ESTIMATED N INPUTS AND OUTPUTS**

Location	Forest Type	Elevation (m)	N Input (kg/ha/year)	N Output (kg/ha/year)	Reference
Adirondack Mts. northeastern New York	Northern hardwoods or hardwood/ conifer mix	396-661	9.3 ^a	Stage 1 N loss ^b	Driscoll and Van Dreason (1993)
Catskill Mts., southeastern New York	Mainly hardwood; some eastern hemlock	335-675	10.2 ^a	Stage 1 and 2 N loss ^b	Stoddard (1994)
Turkey Lakes Watershed, Ontario, Canada	Sugar maple and yellow birch	350-400	7.0-7.7 (as throughfall)	17.9-23.6	Foster et al. (1989); Johnson and Lindberg (1992a)
Whitetop Mt., southwestern Virginia	Red spruce	1650	32 ^c	47 ^c	Joslin and Wolfe (1992); Joslin et al. (1992)
Fernow, West Virginia	Mixed hardwood	735-870	15-20	6.1	Gilliam et al. (1996); Peterjohn et al. (1996)
Great Smoky Mts. National Park, Tennessee	American beech	1600	3.1 ^d	2.9	Johnson and Lindberg (1992b)
Great Smoky Mts. National Park, Becking Site, North Carolina	Red spruce	1800	10.3 ^d	19.2	Johnson et al. (1991a)
Great Smokey Mts. National Park, Tower Site, North Carolina	Red spruce	1740	26.6	20.3	Johnson et al. (1991a)
Front Range, Colorado	Alpine tundra, subalpine conifer	3000-4000	7.5-8.0	7.5	Williams et al. (1996)
San Dimas, San Gabriel Mts., southern California	Chapparral and grasslands	580-1080	23.3	0.04-19.4	Riggan et al. (1985)
Camp Paivika, San Bernadino Mts., southern California	Mixed conifer	1600	30	7-26 ^f	Fenn et al. (1996)
Klamath Mts, northern California	Western coniferous	NA	Mainly geologic ^g	NA ^g	Dahlgren (1994)
Thompson Forest, Cascade Mts., Washington	Red alder	220	4.7 plus > 100 as N ₂ fixation	38.9	Johnson and Lindberg (1992b)

^a Estimated total N deposition from wet deposition data is from Driscoll et al. (1991) for the Adirondacks, and from Stoddard and Murdoch (1991) for the Catskills. Total deposition was estimated based on the wet deposition: total N deposition ratio (0.56) at Huntington Forest in the Adirondacks (Johnson and Lindberg, 1992b). Nitrogen deposition can be higher in some areas, especially at high-elevation sites such as Whiteface Mountain (15.9 kg/ha/year; Johnson and Lindberg, 1992b).

^b Stage 1 and 2 of N loss according to the watershed conceptual model of Stoddard (1994). Nitrogen discharge (kg/ha/year) data are not available; only stream water NO₃⁻ concentration trend data were collected.

^c Values appear high compared to other sites, especially N leaching losses. Joslin and Wolfe (1992) concede that "there is considerable uncertainty associated with the estimates of atmospheric deposition and leaching fluxes." However, elevated NO₃⁻ concentrations in soil solution, and lack of a growth response to N fertilization (Joslin and Wolfe, 1994) support the hypothesis that the forest at Whitetop Mountain is N saturated.


^d Estimated total N deposition from throughfall data. Total deposition was estimated based on the throughfall/total N deposition ratio from the nearby Smokies Tower site (Johnson and Lindberg, 1992b).

^e Annual throughfall deposition to the chaparral ecosystem.

^f Nitrogen output is from unpublished streamwater data (Fenn and Poth, 1999). The low value represents a year of average precipitation, and the high value is for 1995, when precipitation was nearly double the long-term average. Nitrogen output includes N export in stream water and to groundwater.

^g Annual input and output data are not known, although N deposition in this forest is probably typical for much of the rural western United States (2-3 kg N/ha/year (Young et al., 1988). Excess N is from weathering of ammonium in mica schist bedrock. The ammonium was rapidly nitrified, leading to high NO₃⁻ concentrations in soil solution (Dahlgren, 1994).

◆ N-Saturated Ecosystems in North America



Review of Ecosystem Effects and Responses to Excess N

<p>1. Nitrogen Inputs:</p> <ul style="list-style-type: none"> ▶ Atmospheric deposition, N₂ fixation, fertilization <p>Nitrogen Retention:</p> <ul style="list-style-type: none"> ▶ Plant biomass and soil organic matter ▶ Soil microbes and woody residues ▶ Abiotic retention <p>Nitrogen Outputs:</p> <ul style="list-style-type: none"> ▶ Hydrologic transport, gaseous emissions from soil ▶ Removal in harvest, fire emissions, and soil erosion 	<p>3. Ecosystems Responses to Excess Nitrogen:</p> <ul style="list-style-type: none"> ▶ Nitrate leaching and export ▶ Eutrophication of estuaries ▶ Toxicity of surface waters ▶ Foliar nutrient responses ▶ Nitrogen mineralization and nitrification ▶ Effects on soil organic matter ▶ Soil acidification, cation depletion, Al toxicity ▶ Greenhouse gas fluxes
<p>2. Characteristics Predisposing Forests to N Saturation:</p> <ul style="list-style-type: none"> ▶ Stand vigor and succession, forest type ▶ Previous land use – stand history ▶ Soil N accumulation ▶ Topography and climate ▶ Nitrogen deposition 	<p>4. Regional N Saturation Conceptual Models:</p> <ul style="list-style-type: none"> ▶ New England forests ▶ California forests ▶ Colorado alpine ecosystems

Figure 4-17. Diagrammatic overview of excess nitrogen (N) in North America.

Adapted from: Fenn et al. (1998).

experience greater mortality. Long-term fertilization experiments at Mount Ascutney, VT suggest that declining coniferous forest stands with slow nitrogen cycling may be replaced by fast-growing deciduous forests that cycle nitrogen rapidly (Fenn et al., 1998).

Atmospheric nitrogen deposition in the northeastern United States is largely a regional problem (Driscoll et al., 2001). In contrast, in the western United States, vast acres of land receive low levels of atmospheric nitrogen deposition but are interspersed with hot spots of elevated nitrogen deposition downwind of large expanding metropolitan centers or large agricultural operations (Fenn et al., 2003b).

Fenn et al. (1998) have documented the major effects of Nr deposition in terrestrial and aquatic ecosystems in the western United States. Primarily these effects are in response to nitrogen enrichment of systems that are naturally nitrogen limited. Included in these effects are increased greenhouse gas emissions, higher nitrogen concentrations in plant tissues, and increased nitrification rates and nitrate (NO_3^-) levels in soils, streams, and lakes (Fenn et al., 2003b). A result of chronic Nr enrichment has resulted in important community-level changes in vegetation, lichens, mycorrhizae, and phytoplankton, occasionally occurring at relatively low levels of nitrogen deposition (3 to 8 kgN/ha/year; Baron et al., 2000).

Developments in recent decades in the Colorado Front range have resulted in increased nitrogen deposition at high-elevation sites since the 1980s. Total deposition values currently range from 4 to 8 kgN/ha/year (Baron et al., 2000). Competition among species resulting in changes in community composition is one of the most notable responses to environmental change (Bowman, 2000). Nitrogen saturation, the result of increased deposition in the alpine tundra of Niwot Ridge in the Front Range of the southern Rockies in Colorado, has changed nitrogen cycling and provided the potential for replacement in plant species by more competitive, faster growing species (Bowman and Steltzer, 1998; Bowman, 2000; Baron et al., 2000). Plants growing in an alpine tundra, as is true of other plants growing in low resource environments (e.g., infertile soil, shaded understory, deserts), have been observed to have certain similar characteristics: a slow grow rate, low photosynthetic rate, low capacity for nutrient uptake, and low soil microbial activity (Bowman and Steltzer, 1998; Bowman, 2000). An important feature of such plants is that they continue to grow slowly and tend to respond even less when provided with an optimal supply and balance of resources (Pearcy et al., 1987; Chapin, 1991). Plants adapted to cold, moist environments grow more leaves than roots as the relative availability of nitrogen increases; however, other nutrients may soon become limiting. These patterns of vegetative development affect the plants capacities to respond to variation in available resources and to environmental stresses such as frost, high winds, and drought.

Preformation of buds 3 to 4 years in advance of emergence, reduced cell numbers, and high biomass allocation to belowground organs also limits the ability of many alpine plants to respond to variations in their environment (Bowman, 2000). However, significant interspecific genetic variation influences the capacity of the alpine species to respond to changes in resource availability. The capacity of subalpine and boreal species in particular, and gymnosperms in general, to reduce nitrates in either roots or leaves appears to be limited. In addition, the ability of trees to use nitrogen varies with the age of the tree and the density of the stand (Waring, 1987).

In experimental studies of nitrogen deposition conducted by Wedin and Tilman (1996) over a 12-year period on Minnesota grasslands, plots dominated by native warm-season grasses shifted to low-diversity mixtures dominated by cool-season grasses at all but the lowest rates of nitrogen addition. Grasslands with high nitrogen retention and carbon storage rates were the most vulnerable to loss of species and major shifts in nitrogen cycling. The shift to low-diversity mixtures was associated with the decrease in biomass carbon to nitrogen (C:N) ratios, increased nitrogen mineralization, increased soil nitrate, high nitrogen losses, and low carbon storage (Wedin and Tilman, 1996). Naeem et al. (1994) experimentally demonstrated under controlled environmental conditions that the loss of biodiversity, genetic resources, productivity, ecosystem buffering against ecological perturbation, and loss of aesthetic and commercially valuable resources also may alter or impair ecosystems services.

The long-term effects of increased nitrogen deposition have been studied in several western and central European plant communities including lowland heaths, species-rich grasslands, mesotrophic fens, ombrotrophic bogs, upland moors, forest-floor vegetation, and freshwater lakes (Bobbink, 1998). Large changes in species composition have been observed in regions with high nitrogen loadings or in field experiments after years of nitrogen addition (Bobbink et al., 1998). The increased input of nitrogen gradually increased the availability of nitrogen in the soil, and its retention because of low rates of leaching and denitrification, resulting in faster litter decomposition and mineralization rates. Faster growth and greater height of nitrophilic species enables these plants to shade out the slower growing species, particularly those in oligotrophic or mesotrophic conditions (Bobbink, 1998; Bobbink et al., 1998). Excess nitrogen inputs to unmanaged heathlands in the Netherlands has resulted in nitrophilous grass species replacing the slower growing heath species (Roelofs et al., 1987; Garner, 1994).

Van Breemen and Van Dijk (1988) noted that over the past several decades the composition of plants in the forest herb layers has been shifting toward species commonly found in nitrogen-rich areas. They also observed a decrease in number of the fruiting bodies of mycorrhizal fungi.

Other studies in Europe point out the effects of excessive nitrogen deposition on mixed-oak forest vegetation along a deposition gradient largely controlled by soil acidity, nitrogen supply, canopy composition, and location of sample plots (Brunet et al., 1998; Falkengren-Grerup, 1998). Results of the study, using multivariate methods, suggest that nitrogen deposition has affected the field-layer vegetation directly by increased nitrogen availability and, indirectly, by accelerating soil acidity. Time series studies indicate that 20 of the 30 field-layer species (nonwoody plants) that were associated most closely with high nitrogen deposition increased in frequency in areas with high nitrogen deposition during the past decades. Included in the field-layer species were many species generally considered nitrophilous; however, there were also several acid-tolerant species (Brunet et al, 1998). In an experimental study involving 15 herbs and 13 grasses, Falkengren-Grerup (1998), observed that species with a high nitrogen demand and a lesser demand for nutrients other than nitrogen were particularly competitive in areas with acidic soils and high nitrogen deposition. The grasses grew better than herbs with the addition of nitrogen. It was concluded that, at the highest nitrogen deposition, growth was limited for most species by the supply of nutrients other than nitrogen; and, at the intermediate nitrogen concentration, the grasses were more efficient than the herbs in utilizing nitrogen. Nihlgård (1985) suggested that excessive nitrogen deposition may contribute to forest decline in other specific regions of Europe. Additionally, Schulze (1989), Heinsdorf (1993), and Lamersdorf and Meyer (1993) attributed magnesium deficiencies in German forests, in part, to excessive nitrogen deposition.

The carbon to nitrogen (C:N) ratio of the forest floor can also be changed by nitrogen deposition over time. This change appears to occur when the ecosystem becomes nitrogen saturated (Gundersen et al., 1998a). Long-term changes in C:N status have been documented in Central Europe and indicate that nitrogen deposition has changed the forest floor. In Europe, low C:N ratios coincide with high deposition regions (Gundersen et al., 1998a). A strong decrease in forest floor root biomass has been observed with increased nitrogen availability. Roots and the associated mycorrhizae appear to be an important factor in the accumulation of organic matter in the forest floor at nitrogen-limited sites. If root growth and mycorrhizal

formation are impaired by excessive nitrogen deposition, the stability of the forest floor vegetation may be affected by stimulating turnover and decreasing the root litter input to the forest floor, thus decreasing the nitrogen that can be stored in the forest floor pool (Gundersen et al., 1998b). Nitrogen-limited forests have a high capacity for deposited nitrogen to be retained by plants and microorganisms competing for available nitrogen (Gundersen et al., 1998b). Nitrate leaching has been correlated significantly with nitrate status but not with nitrate depositions. The forest floor C:N ratio has been used as a rough indicator of ecosystem nitrogen status in mature coniferous forests and the risk of nitrate leaching; analyses of European databases indicate an empirical relationship between forest floor C:N ratio and nitrate leaching (Gundersen et al., 1998a). Nitrate leaching was observed when the deposition received was more than 10 kg N/ha. All of the data sets supported the threshold at which nitrate leaching seems to increase at a C:N ratio of 25. Therefore, to predict the rate of changes in nitrate leaching, it is necessary to be able to predict the rate of changes in the forest floor C:N ratio. Decreased foliar and soil nitrogen and soil C:N ratios, as well as changes in nitrogen mineralization rates, have been observed when comparing responses to nitrogen deposition in forest stands east and west of the Continental Divide in the Colorado Front Range (Baron et al., 2000; Rueth and Baron, 2002). Understanding the variability in forest ecosystem response to nitrogen input is essential to assessing pollution risks (Gundersen et al., 1998a).

The plant root is an important region of nutrient dynamics. The rhizosphere includes the soil that surrounds and is influenced by plant roots (Wall and Moore, 1999). The mutualistic relationship between plant roots, fungi, and microbes is critical for the growth of the organisms involved. The plant provides shelter and carbon; whereas the symbiont provides access to limiting nutrients such as nitrogen and phosphorus. As indicated above, changes in soil nitrogen influence the mycorrhizal-plant relationship. Mycorrhizal fungal diversity is associated with above-ground plant biodiversity, ecosystem variability, and productivity (Wall and Moore, 1999). Aber et al. (1998) showed a close relationship between mycorrhizal fungi and the conversion of dissolved inorganic nitrogen to soil nitrogen. During nitrogen saturation, soil microbial communities change from being fungal, and probably being dominated by mycorrhizae, to being dominated by bacteria. The loss of mycorrhizal function has been hypothesized as the key process leading to increased nitrification and nitrate mobility. Increased nitrate mobility leads to increased cation leaching and soil acidification (Aber et al., 1998).

The interrelationship of above- and below-ground flora is illustrated by the natural invasion of heathlands by oaks (*Quercus robur*). The soil-forming factors under the heath have been vegetation typed during the last 2000 years; whereas the invasion by oaks has been taking place for only a few decades. Clearly, changes in the ground floor and soil morphology takes place when trees colonize heath (Nielsen et al., 1999). The distribution of roots also changed under the three different vegetation types. Under both heather and the Sitka spruce plantation, the majority of roots are confined to the uppermost horizons; whereas under oak, the roots are distributed more homogeneously. There was also a change in the C:N ratio when heather was replaced by oaks. Also, the spontaneous succession of the heath by oaks changed the biological nutrient cycle into a deeper vertical cycle when compared to the heath where the cycle is confined to the upper soil horizons. Soils similar to those described in this Jutland, Denmark study with a mainly organic buffer system seem to respond quickly to changes in vegetation (Nielsen et al., 1999).

The effects of changes in root to shoot relationships in plants were observed in studies of the coastal sage scrub (CSS) community in southern California, which is composed of the drought-deciduous shrubs *Artemisia californica*, *Encelia farinosa*, and *Eriogonum fasciculatum*. The CSS in California has been declining in land area and in shrub density over the past 60 years and is being replaced in many areas by Mediterranean annual grasses (Allen et al., 1998; Padgett et al., 1999; Padgett and Allen, 1999). Nitrogen deposition was considered as a possible cause. Up to 45 kg/ha/year are deposited in the Los Angeles Air Basin (Bytnerowicz and Fenn, 1996). Tracts of land set aside as reserves, which in many cases in southern California are surrounded by urbanization, receive large amounts of nitrogenous compounds from polluted air. The CSS is of particular interest, because some 200 sensitive plant species and several federally listed animal species are found in the area (Allen et al., 1998). Because changes in plant community structure often can be related to increases in the availability of a limiting soil nutrient or other resource, experiments were conducted to determine whether increased nitrogen availability was associated with the significant loss in native shrub cover. Studies indicated that the three native perennial shrubs (*Artemisia californica*, *Eriogonum fasciculatum*, and *Encelia farinosa*) tended to be more nitrophilous than the two exotic annual grasses (*Bromus rubens*, *Avena fatua*) and the weedy pod mustard (*Brassica geniculata*). These results contrast with most models dealing with the adaptation of perennial species to stressful environments (Padgett and Allen, 1999).

If nitrogen were the only variable between the invasive annuals and native shrubs, neither shrubs nor grasses would have a particular advantage. Although CSS shrubs are able to take up nitrogen at high rates, native grasses have a denser seedbank and earlier germination than native species. The native seedlings are unable to compete with dense stands of exotic grasses, and thus are gradually replaced by the grasses following disturbances such as frequent fire (Eliason and Allen, 1997; Cione et al., 2002; Yoshida and Allen, 2001). In addition, nitrogen-induced changes in arbuscular mycorrhizal fungi may also affect the growth of native seedlings. Nitrogen enrichment of the soils has induced a shift in the arbuscular mycorrhizal community composition. Larger-spored fungal species (*Scutellospora* and *Gigaspora*), due to a failure to sporulate, decreased in number with a concomitant proliferation of small-spored species of *Glomus aggregatum*, *G. leptotichum*, and *G. geosporum*, indicating a strong selective pressure for the smaller spored species of fungi (Edgerton-Warburton and Allen, 2000). These results demonstrate that nitrogen enrichment of the soil significantly alters the arbuscular mycorrhizal species composition and richness and markedly decreases the overall diversity of the arbuscular mycorrhizal community. The decline in coastal sage scrub species can, therefore, be directly linked to the decline of the arbuscular mycorrhizal community (Edgerton-Warburton and Allen, 2000).

In addition to excess nitrogen deposition effects on terrestrial ecosystems of the types noted above (e.g., dominant species shifts and other biodiversity impacts), direct atmospheric nitrogen deposition and increased nitrogen inputs via runoff into streams, rivers, lakes, and oceans can noticeably affect aquatic ecosystems as well (Figure 4-15). Estuaries are among the most intensely fertilized ecosystems on Earth, receiving far greater nutrient inputs than other systems. Chesapeake Bay is a prime example (Fenn et al., 1998). Another illustrative example is found in recently reported research (Paerl et al., 2001) characterizing the effects of nitrogen deposition on the Pamlico Sound, NC estuarine complex, which serves as a key fisheries nursery supporting an estimated 80% of commercial and recreational finfish and shellfish catches in the southeastern U.S. Atlantic coastal region. Such direct atmospheric nitrogen deposition onto waterways feeding into the Pamlico Sound or onto the sound itself, combined with indirect nitrogen inputs via runoff from upstream watersheds, contribute to conditions of severe water oxygen depletion; formation of algae blooms in portions of the Pamlico Sound estuarine complex; altered fish distributions, catches, and physiological states; and increases in the

incidence of disease in fish. Under extreme conditions of especially high rainfall rate events (e.g., hurricanes) affecting watershed areas feeding into the sound, the effects of nitrogen runoff (in combination with excess loadings of metals or other nutrients) can be massive — e.g., creation of the widespread “dead-zone” affecting large areas of the Pamlico Sound for many months after hurricane Fran in 1996 and hurricanes Dennis, Floyd, and Irene in 1999 impacted eastern North Carolina.

The primary pathways of nitrogen loss from forest ecosystems are hydrological transport beyond the rooting zone into groundwater or stream water, or surface flows of organic nitrogen as nitrate and nitrogen loss associated with soil erosion (Fenn et al., 1998). Stream water nitrate concentrations have been related to forest successional stage in the eastern United States. Logging and fire histories of an area are major variables determining the capability of a forest stand to retain nitrogen. Nitrogen concentrations were high in mature ecosystems after disturbances such as clearcutting, but lower in mid-successional forests.

Nitrogen saturation of a high elevation watershed in the southern Appalachian Mountains was observed to affect stream water chemistry. High nitrate concentrations have been observed in streams draining undisturbed watersheds in the Great Smoky Mountains National Park in Tennessee and North Carolina. Nitrate concentrations were highest at higher elevations and in areas around old-growth forests that had never been logged (Fenn et al., 1998).

In the Northeast, nitrogen is the element most responsible for eutrophication in coastal waters of the region (Jaworski et al., 1997). There has been a 3- to 8-fold increase in nitrogen flux from 10 watersheds in the northeastern United States since the early 1900s. These increases are associated with the deposition of nitrogen oxide emissions from combustion, which have increased 5-fold. Riverine nitrogen fluxes have been correlated with atmospheric deposition onto their landscapes and with nitrogen oxides emissions into their airsheds. Data from 10 benchmark watersheds with good historical records, indicate that ~36 to 80% of the riverine total nitrogen export, with an average of 64%, was derived directly or indirectly from nitrogen oxide emissions (Jaworski et al., 1997).

Nitrogen saturation of a high-elevation watershed in the southern Appalachian Mountains was observed to affect stream water chemistry. The Great Smoky Mountains in the southeastern United States receive high total atmospheric deposition of sulfur and nitrogen (2,200 Eq/ha/year of total sulfur and approximately 1,990 Eq/ha/year of total nitrogen). A major portion of the

atmospheric loading is from dry and cloud deposition. Extensive surveys conducted in October 1993 and March 1994 indicated that stream pH values were near or below pH 5.5 and that the acid neutralizing capacity (ANC) was below 50 $\mu\text{eq/L}$ at high elevations. Analysis of stream water indicated that nitrate was the dominant anion (Flum and Nodvin, 1995; Nodvin et al., 1995). The study was expanded to the watershed scale with the monitoring of precipitation, throughfall, stream hydrology, and stream chemistry. Nitrogen saturation of the watershed resulted in extremely high exports of nitrate and promoted both chronic and episodic stream acidification in which the nitrate was the dominant ion. Significant exports of base cation was also observed. Nitrification of the watershed soils resulted in elevations of soil solution aluminum concentrations to levels known to inhibit calcium uptake in red spruce (Nodvin et al., 1995).

Excessive nitrogen loss is a symptom of terrestrial ecosystem dysfunction and results in the degradation of water quality with potentially deleterious effects on terrestrial and aquatic ecosystems (Fenn and Poth, 1999). Data from a number of hydrologic, edaphic, and plant indicators indicate that the mixed conifer forests and chaparral systems directly exposed to air pollution from greater Los Angeles are nitrogen saturated. Preliminary data suggests that symptoms of nitrogen saturation are evident in mixed conifer or chaparral sites receiving atmospheric deposition of 20 to 25 kg/N/ha/year (Fenn et al, 1996). Available data clearly indicate that ecosystems with a Mediterranean climate have a limited capacity to retain nitrogen within the terrestrial system (Fenn and Poth, 1999). A 3-year study of stream water NO_3^- concentrations along nitrogen deposition gradients in the San Bernardino Mountains in southern California evaluated stream water quality and whether the stream water concentrations covaried with nitrogen deposition across pollution gradients in the San Bernardino Mountains. Stream water NO_3^- concentrations at Devil Canyon in the San Gabriel Mountains northeast of Los Angeles are the highest reported in North America for forested watersheds (Fenn and Poth, 1999). Five of the six streams monitored maintained elevated NO_3^- throughout the year. Peak nitrate concentrations ranged from 40 to 350 $\mu\text{mol/L}$. In the San Gorgonio Wilderness, an area of low to moderate deposition where 12 streams were sampled, only the five that had the greatest air pollution exposure had high NO_3^- concentrations. These results suggest a strong association between NO_3^- levels exported in stream water and the severity of chronic nitrogen deposition to the terrestrial watersheds. However, nitrogen processing within terrestrial and aquatic systems,

even in areas with high nitrogen deposition, determine stream water NO_3^- concentrations (Fenn and Poth, 1999). The Fernow Experimental Forest in West Virginia, the Great Smoky Mountains National park in Tennessee, and watersheds in southwestern Pennsylvania are the only undisturbed forested sites in North America known to have stream water NO_3^- concentrations within the range of values found at Devil Canyon (Fenn and Poth, 1999).

Effects of Sulfur Deposition. Sulfur is a major component of plant proteins and, as such, is an essential plant nutrient. The most important source of sulfur is sulfate taken up from the soil by plant roots, even though plants can use atmospheric SO_2 (Marschner, 1995). The availability of organically bound sulfur in soils depends largely on microbial decomposition, a relatively slow process. The major factor controlling the movement of sulfur from the soil into vegetation is the rate of release from the organic to the inorganic compartment (May et al., 1972; U.S. Environmental Protection Agency, 1982; Marschner, 1995). Sulfur plays a critical role in agriculture as an essential component of the balanced fertilizers needed to grow and increase worldwide food production (Ceccotti and Messick, 1997). Atmospheric deposition is an important component of the sulfur cycle. This is true not only in polluted areas where atmospheric deposition is very high, but also in areas of low sulfur input. Additions of sulfur into the soil in the form of SO_4^{2-} could alter the important organic-sulfur/organic-nitrogen relationship involved in protein formation in plants. The biochemical relationship between sulfur and nitrogen in plant proteins and the regulatory coupling of sulfur and nitrogen metabolism indicate that neither element can be assessed adequately without reference to the other. A sulfur deficiency reduces nitrate reductase and, to a similar extent, glutamine synthetase activity. Nitrogen uptake in forests, therefore, may be loosely regulated by sulfur availability, but sulfate additions in excess of needs do not necessarily lead to injury (Turner and Lambert, 1980; Hogan et al., 1998).

Only two decades ago, there was little information comparing sulfur cycling in forests with other nutrients, especially nitrogen. With the discovery of deficiencies in some unpolluted regions (Kelly and Lambert, 1972; Humphreys et al., 1975; Turner et al., 1977; Schnug, 1997) and excesses associated with acidic deposition in other regions of the world (Meiwes and Khanna, 1981; Shriner and Henderson, 1978; Johnson et al., 1982a,b), interest in sulfur nutrition and cycling in forests has heightened. General reviews of sulfur cycling in forests have been

written by Turner and Lambert (1980), Johnson (1984), Mitchell et al. (1992a,b), and Hogan et al. (1998). The salient elements of the sulfur cycle as it may be affected by changing atmospheric deposition are summarized by Johnson and Mitchell (1998). Sulfur has become the most important limiting factor in European agriculture because of the desulfurization of industrial emissions (Schnug, 1997).

Most studies dealing with the effects of sulfur deposition on plant communities have been conducted in the vicinity of point sources and have investigated the above-ground effects of SO₂ or the acidifying effects of sulfate on soils (Krupa and Legge, 1998; Dreisinger and McGovern, 1970; Legge, 1980; Winner and Bewley, 1978a,b; Laurenroth and Michunas, 1985; U.S. Environmental Protection Agency, 1982). Krupa and Legge (1986), however, observed a pronounced increase in foliar sulfur concentrations in all age classes of needles of the hybrid pine lodgepole × jack pine (*Pinus contorta* × *P. banksiana*). This vegetation had been exposed to chronic low concentrations of sulfur dioxide (SO₂) and hydrogen sulfide (H₂S) for more than 20 years and, then, to fugitive sulfur aerosol. Observations under the microscope showed no sulfur deposits on the needle surfaces and led to the conclusion that the sulfur in the needles was derived from the soil. The oxidation of elemental sulfur and the generation of protons is well known for the soils of Alberta, Canada. This process is mediated by bacteria of the *Thiobacillus* sp. As elemental sulfur is gradually converted to protonated SO₄, it can be leached downward and readily taken up by plant roots. The activity of *Thiobacillus* sp. is stimulated by elemental sulfur additions (Krupa and Legge, 1986).

Effects of Acidic Deposition on Forest Soils. Acidic deposition over the past quarter of a century has emerged as a critical environmental stress that affects forested landscapes and aquatic ecosystems in North America, Europe, and Asia (Driscoll et al., 2001). Acidic deposition can originate from transboundary air pollution and affect large geographic areas. It is composed of ions, gases, particles derived from gaseous emissions of SO₂, NO_x, NH₃, and particulate emissions of acidifying and neutralizing compounds and is highly variable across space and time. It links air pollution to diverse terrestrial and aquatic ecosystems and alters the interactions of the H⁺ with many elements (e.g., S, N, Ca, Mg, Al, and Hg). Acidic deposition contributes directly and indirectly to biological stress and the degradation of ecosystems and has

played a major role in the recent acidification of soil in some areas of Europe and, to a more limited extent, eastern North America (Driscoll et al., 2001).

Substantial and previously unsuspected changes in soils have been observed in polluted areas of eastern North America, the United Kingdom, Sweden, and central Europe and in less polluted regions of Australia and western North American (reviewed by Johnson et al., 1999 and by Huntington, 2000). In some cases, trends were toward more acidic soils (e.g., Markewitz et al., 1998); in others cases, there were no consistent trends, with some soils showing increases and some showing decreases at different sampling times and some showing no change (e.g., Johnson and Todd, 1998; Trettin et al., 1999; Yanai et al., 1999).

Significant changes in soil chemistry have occurred at many sites in the eastern United States during recent decades. Patterns of change in tree ring chemistry, principally at high elevations sites in the eastern United States, reflect the changing inputs of regional pollutants to forests. A temporal sequence of changes in uptake patterns, and possibly in tree growth, would be expected if significant base cation mobilization and depletion of base cations from eastern forest soils has occurred. Temporal changes in the chemistry of the tree rings of red spruce were examined as indicators of historical changes in the red spruce's chemical environment.

Analysis of changes in wood chemistry from samples across several sites indicate that there have been substantial departures from the expected linear decreases in calcium accumulation patterns in wood. A region-wide calcium increase above expected levels followed by decreasing changes in wood calcium suggest that calcium mobilization began possibly 30 to 40 years ago and has been followed by reduced accumulation rates in wood, presumably associated with decreasing calcium availability in soil (Bondietti and McLaughlin, 1992). The period of calcium mobilization coincides with a region-wide increase in the growth rate of red spruce; whereas the period of decreasing levels of calcium in wood corresponds temporally with patterns of decreasing radial growth at high elevation sites throughout the region during the past 20 to 30 years. The decline in wood calcium suggests that calcium loss may have increased to the point at which base saturation of soils has been reduced. Increases in aluminum and iron typically occur as base cations are removed from the soils by tree uptake (Bondietti and McLaughlin, 1992). The changes are spatially and temporally consistent with changes in the emissions of SO₂ and NO₂ across the region and suggest that increased acidification of soils has occurred.

Studies by Shortle and Bondietti (1992) support the view that changes in soil chemistry in eastern North America forest sites occurred many decades ago, “before anybody was looking.” Sulfur and nitrogen emissions began increasing in eastern North America in the 1920s and continued to increase into the 1980s when sulfur began to decrease but nitrogen emissions did not (Garner et al., 1989). Shortle and Bondietti (1992) present evidence that, from the late 1940s into the 1960s, the mor humus (organic) layer of acid-sensitive forest sites in eastern North America underwent a significant change that resulted in the loss of exchangeable essential base cations and interrupted the critical base nutrient cycles between mature trees and the root-humus complex. The timing of the effect appears to have coincided with the period when the SO_x and NO_x emissions in eastern North America subject to long-range transport were increasing the most rapidly (see above; Shortle and Bondietti, 1992). Although forest ecosystems other than the high-elevation spruce-fir forests are not currently manifesting symptoms of injury directly attributable to acid deposition, less sensitive forests throughout the United States are experiencing gradual losses of base cation nutrients, which, in many cases, will reduce the quality of forest nutrition over the long term (National Science and Technology Council, 1998). In some cases it may not even take decades, because these forests already have been receiving sulfur and nitrogen deposition for many years. The current status of forest ecosystems in different U.S. geographic regions varies, as does their sensitivity to nitrogen and sulfur deposition. Variation in potential future forest responses or sensitivity are caused, in part, by differences in deposition of sulfur and nitrogen, ecosystem sensitivities to sulfur and nitrogen additions, and the responses of soils to sulfur and nitrogen inputs (National Science and Technology Council, 1998).

Acidic deposition has played a major role in recent soil acidification in some areas of Europe and, to a more limited extent, eastern North America. Examples include the study by Hauhs (1989) at Lange Bramke, Germany, which indicated that leaching was of major importance in causing substantial reduction in soil-exchangeable base cations over a 10-year period (1974 to 1984). Soil acidification and its effects result from the deposition of NO_3^- and SO_4^{2-} and associated H^+ . The effects of excessive nitrogen deposition on soil acidification and nutrient imbalances have been well established in Dutch forests (Van Breemen et al., 1982; Roelofs et al., 1985; Van Dijk and Roelofs, 1988). For example, Roelofs et al. (1987) proposed that $\text{NH}_3/\text{NH}_4^+$ deposition leads to heathland changes via two modes: acidification of the soil

with the loss of cations K^+ , Ca^{2+} , and Mg^{2+} ; and nitrogen enrichment that results in “abnormal” plant growth rates and altered competitive relationships. Nihlgård (1985) suggested that excessive nitrogen deposition may contribute to forest decline in other regions of Europe. Falkengren-Grerup (1987) noted that, over about 50 years, unexpectedly large increases in the growth of beech trees (*Fagus sylvatica* L.) were associated with decreases in pH and exchangeable cations in some sites in southernmost Sweden.

Likens et al. (1996, 1998) suggested that soils are changing at the Hubbard Brook Watershed, NH, because of a combination of acidic deposition and reduced base cation deposition. They surmised, based on long-term trends in stream water data, that large amounts of Ca and Mg have been lost from the soil-exchange complex over a 30-year period, from approximately 1960 to 1990. The authors speculate that the declines in base cations in soils may be the cause of recent slowdowns in forest growth at Hubbard Brook. In a follow-up study, however, Yanai et al. (1999) found no significant decline in Ca and Mg in the forest floors at Hubbard Brook over the period 1976 to 1997. They also found both gains and losses in forest floor Ca and Mg between 1980 and 1990 in a regional survey. Thus, they concluded that “forest floors in the region are not currently experiencing rapid losses of base cations, although losses may have preceded the onset of these three studies.” The biogeochemistry of calcium at Hubbard Brook is discussed in detail by Likens et al. (1998).

Hydrogen ions entering a forest ecosystem first encounter the forest canopy, where they are often exchanged for base cations that then appear in throughfall (Figure 4-18 depicts a model of H^+ sources and sinks). Base cations leached from the foliage must be replaced through uptake from the soil, or foliage cations will be reduced by the amounts leached. In the former case, the acidification effect is transferred to the soil where H^+ is exchanged for a base cation at the root-soil interface. The uptake of base cations or NH_4^+ by vegetation or soil microorganisms causes the release of H^+ in order to maintain charge balance; conversely, the uptake of nutrients in anionic form (NO_3^- , SO_4^{2-} , PO_4^{3-}) causes the release of OH^- in order to maintain charge balance. Thus, the net acidifying effect of uptake is the difference between cation and anion uptake. The form of ions taken up is known for all nutrients but nitrogen, because either NH_4^+ or NO_3^- can be used by the vegetation. In that nitrogen is the nutrient taken up in the greatest quantities, the uncertainty in the ionic form of nitrogen utilized creates great uncertainty in the overall H^+ budget for soils (Johnson, 1992).




Figure 4-18. Schematic of sources and sinks of hydrogen ions in a forest.

Source: Taylor et al. (1994).

The cycles of base cations differ from those of N, P, and S in several respects. The fact that calcium, potassium, and magnesium exist primarily as cations in solution, whereas N, P, and S exist primarily as anions, has major implications for the cycling of the nutrients and the effects of acid deposition on these cycles. The most commonly accepted model of base cation cycling in soils is one in which base cations are released by the weathering of primary minerals to cation exchange sites, making the cation available for either plant uptake or leaching (Figure 4-18). The introduction of H⁺ by atmospheric deposition or by internal processes will affect the fluxes of Ca, K, and Mg via cation exchange or weathering processes. Therefore, soil leaching is often

of major importance in cation cycles, and many forest ecosystems show a net loss of base cations (Johnson, 1992).

Two basic types of soil change are involved: (1) a short-term intensity type change resulting from the concentrations of chemicals in soil water and (2) a long-term capacity change based on the total content of bases, aluminum, and iron stored in the soil (Reuss and Johnson, 1986; Van Breemen et al., 1983). Changes in intensity factors can have a rapid effect on the chemistry of soil solutions. Rapid changes in intensity resulting from the addition of sulfur and nitrogen in acidic deposition can cause immediate increases in acidity and the mobilization of aluminum in soil solutions. Increased aluminum concentrations and an increase in the Ca:Al ratio in soil solution have been linked to a significant reduction in the availability of essential base cations to plants, an increase in plant respiration, and increased biochemical stress (National Science and Technology Council, 1998).

Capacity changes are the result of many factors acting over long time periods. The content of base cations (calcium, magnesium, sodium, and potassium) in soils results from additions via atmospheric deposition, decomposition of vegetation, and geologic weathering; losses may occur through plant uptake and leaching. Increased leaching of base cations may result in nutrient deficiencies in soils, as has been happening in some sensitive forest ecosystems (National Science and Technology Council, 1998).

Aluminum toxicity is a possibility in acidified soils. Atmospheric deposition (or any other source of mineral anions) can increase the concentration of Al, especially Al^{3+} , in soil solution without causing significant soil acidification (Johnson and Taylor, 1989). Aluminum can be brought into soil solution in two ways: (1) by acidification of the soil and (2) by increasing the total anion and cation concentration of the soil solution. The introduction of mobile, mineral acid anions to an acid soil will cause increases in the concentration of aluminum in the soil solution, but extremely acid soils in the absence of mineral acid anions will not produce a solution high in aluminum. An excellent review of the relationships among the most widely used cation-exchange equations and their implications for the mobilization of aluminum into soil solution is provided by Reuss (1983).

A major concern has been that soil acidity would lead to nutrient deficiency. Calcium is essential for root development and the formation of wood, and it plays a major role in cell membrane integrity and cell wall structure. Aluminum concentrations in the soil can influence

forest tree growth in regions where acidic deposition and natural acidifying processes increase soil acidity. Acidic deposition mobilizes calcium and magnesium, which are essential for root development and stem growth. Mobilized aluminum can also bind to fine root tips of red spruce, further limiting calcium and magnesium uptake (Shortle and Smith, 1988; Shortle et al., 1997).

There is abundant evidence that aluminum is toxic to plants. Upon entering tree roots, aluminum accumulates in root tissues (Thornton et al., 1987; Vogt et al., 1987a, b). Reductions in calcium uptake have been associated with increased aluminum uptake (Clarkson and Sanderson, 1971), and a number of studies suggest that the toxic effect of aluminum on forest trees could be caused by a Ca^{2+} deficiency (Shortle and Smith, 1988; Smith, 1990a). Mature trees have a high calcium requirement relative to agriculture crops (Rennie, 1955). Shortle and Smith (1988) attributed the decline of red spruce in eight stands across northern New England from Vermont to Maine to an imbalance of Al^{3+} and Ca^{2+} in fine root development.

To be taken up from the soil by roots, Ca must be dissolved in soil water (Lawrence and Huntington, 1999). Aluminum in soil solution reduces Ca uptake by competing for binding sites in the cortex of fine roots. Tree species may be adversely affected if high aluminum to nutrient ratios create a nutrient deficiency by limiting the uptake of calcium and magnesium (Shortle and Smith, 1988; Garner, 1994). Acid deposition, by lowering the pH of Al-rich soil, can increase Al concentrations in soil water through dissolution and ion exchange processes. Aluminum is more readily taken up than is Ca because of its greater affinity for negatively charged surfaces and, when present in the forest floor, Al tends to displace adsorbed Ca, causing it to be more readily leached. The continued buildup of Al in the forest floor layer, where nutrient uptake is greatest, can lower the efficiency of Ca uptake when the Ca:Al ratio in soil water is less than one (Lawrence and Huntington, 1999). A reduction in Ca uptake suppresses cambial growth and reduces the rate of wood (annual ring) formation, decreases the amount of functional sapwood and live crown, and predisposes trees to disease and injury from stress agents when the functional sapwood becomes less than 25% of cross-sectional stem area (Smith, 1990a). A 1968 Swedish report to the United Nations postulated a decrease in forest growth of ~1.5%/year when the Ca:Al ratio in soil water is less than one (Lawrence and Huntington, 1999). The concern that acidification and nutrient deficiency may result in forest decline remains today.

Acidic deposition has been firmly implicated as a causal factor in the northeastern high-elevation decline of red spruce (DeHayes et al., 1999). The frequency of freezing injury of red

spruce has increased over the past 40 years, a period that coincides with increase emissions of sulfur and nitrogen oxides and acidic deposition (DeHayes et al., 1999). Studies indicate that there is a significant positive association between cold tolerance and foliar Ca in trees exhibiting deficiency in foliar Ca. Most of the calcium in conifer needles is insoluble calcium oxalate and pectate crystals, which are of little physiological importance. It is the labile calcium ions in equilibrium within the plasma membrane that are of major physiological importance (DeHayes et al., 1999). The membrane-associated pool of calcium (mCa), although a relatively small fraction of total foliar ion pools, strongly influences the response of cells to changing environmental conditions. The plant plasma membrane plays a critical role in mediating cold acclimation and low-temperature injury. The leaching of Ca associated with acidic deposition is considered to be the result of cation exchange due to exposure to the H⁺ ion. The studies of DeHayes et al. (1999) demonstrated that the direct deposition of acidic deposition on needles represents a unique environmental stress, in that it preferentially removes mCa, which is not readily replaced in the autumn. They propose that direct acidic deposition on red spruce foliage preferentially displaces those Ca ions specifically associated with the plasma membranes of mesophyll cells resulting in the reduction of mCa and the destabilizing of plasma membranes and depletion of messenger Ca ions. Further, DeHayes et al. (1999) state that their studies raise the strong possibility that acid rain alteration of the mCa and membrane integrity is not unique to red spruce but has been demonstrated in many other northern temperate forest tree species, including: yellow birch (*Betula alleghaniensis*), white spruce (*Picea glaucus*), red maple (*Acer rubrum*), eastern white pine (*Pinus strobus*), and sugar maple (*Acer saccharum*). Assessments of mCa, membrane integrity, and the effects of other secondary stressors have not yet been made for these species.

Seasonal and episodic acidification of surface waters have been observed in the eastern United States, Canada, and Europe (Hyer et al., 1995). In the Northeast, the Shenandoah National Park in Virginia, and the Great Smoky Mountains, episodic acidification has been associated with the nitrate ion (Driscoll et al., 2001; Hyer et al., 1995; Eshleman et al., 1995). The short-term acid episodes occur during spring snowmelts and large precipitation events (Driscoll et al., 2001). Episodic acidification of surface waters has usually been considered to be a transient loss of acid neutralizing capacity associated with snowmelt/rainfall runoff and, as such, represents a short-term (hours to weeks) effect considered to be distinguishable from the

chronic long-term (years to centuries) changes in acidity. Studies of both episodic and chronic acidification of surface waters indicate that acidification can have long-term adverse effects on fish populations and result in the decline of species richness and the abundance of zooplankton and macroinvertebrates (Driscoll et al., 2001; Eshleman et al., 1995). Nitrogen saturation of soils and the slow release of nitrates has been shown to inhibit the recovery of acid sensitive systems (Driscoll et al., 2001). The acidification of aquatic ecosystems and effects on aquatic biota are discussed in more detail in the EPA document Air Quality Criteria for Nitrogen Oxides (U.S. Environmental Protection Agency, 1993).

Air pollution is not the sole cause of soil change. High rates of acidification are occurring in less polluted regions of the western United States and Australia because of internal soil processes, such as tree uptake of nitrate and nitrification associated with excessive nitrogen fixation (Johnson et al., 1991b). Many studies have shown that acidic deposition is not a necessary condition for the presence of extremely acidic soils, as evidenced by their presence in unpolluted, even pristine, forests of the northwestern United States and Alaska (Johnson et al., 1991b). Soil can become acidic when H^+ ions attached to NH_4^+ or HNO_3 remain in the soil after nitrogen is taken up by plants. For example, Johnson et al. (1982b) found significant reductions in exchangeable K^+ over a period of only 14 years in a relatively unpolluted Douglas fir Integrated Forest Study (IFS) site in the Washington Cascades. The effects of acid deposition at this site were negligible relative to the effects of natural leaching (primarily of carbonic acid) and nitrogen tree uptake (Cole and Johnson, 1977). Even in polluted regions, numerous studies have shown the importance of tree uptake of NH_4^+ and NO_3^- in soil acidification. Binkley et al. (1989) attributed the marked acidification (pH decline of 0.3 to 0.8 units and base saturation declines of 30 to 80%) of abandoned agricultural soil in South Carolina over a 20-year period to NH_4^+ and NO_3^- uptake by a loblolly pine plantation.

An interesting example of uptake effects on soil acidification is that of Al uptake and cycling (Johnson et al., 1991b). Aluminum accumulation in the leaves of coachwood (*Ceratopetalum apetalum*) in Australia has been found to have a major effect on the distribution and cycling of base cations (Turner and Kelly, 1981). The presence of *C. apetalum* as a secondary tree layer beneath brush box (*Lophostemon confertus*) was found to lead to increased soil exchangeable Al^{3+} and decreased soil exchangeable Ca^{2+} (Turner and Kelly, 1981). The

constant addition of aluminum-rich litter fall obviously has had a substantial effect on soil acidification, even if base cation uptake is not involved directly.

Given the potential importance of particulate deposition to the base cation status of forest ecosystems, the findings of Driscoll et al. (1989, 2001) and Hedin et al. (1994) are especially relevant. Driscoll et al. (1989, 2001) noted a decline in both SO_4^{2-} and base cations in both atmospheric deposition and stream water over the past two decades at Hubbard Brook Watershed, NH. The decline in SO_4^{2-} deposition was attributed to a decline in emissions, and the decline in stream water SO_4^{2-} was attributed to the decline in sulfur deposition. Hedin et al. (1994) reported a steep decline in atmospheric base cation concentrations in both Europe and North America over the past 10 to 20 years. The reductions in SO_2 emissions in Europe and North America in recent years have not been accompanied by equivalent declines in net acidity related to sulfate in precipitation. These current declines in sulfur deposition have, to varying degrees, been offset by declines in base cations and may be contributing “to the increased sensitivity of poorly buffered systems.” Analysis of the data from the IFS supports the authors’ contention that atmospheric base cation inputs may seriously affect ecosystem processes. Johnson et al. (1994) analyzed base cation cycles at the Whiteface Mountain IFS site in detail and concluded that Ca losses from the forest floor were much greater than occurred historically, based on historical changes in forest floor Ca observed in an earlier study. Further, the authors suggested that the difference between historical and current net loss rates of forest floor Ca may be caused by sharply reduced atmospheric inputs of Ca after about 1970 and may be exacerbated by sulfate leaching (Johnson et al., 1994).

The Ca:Al molar ratio has been suggested to be a valuable ecological indicator of an approximate threshold beyond which the risk of forest injury from Al stress and nutrient imbalances increases (Cronan and Grigal, 1995). The Ca:Al ratio also can be used as an indicator to assess forest ecosystem changes over time in response to acidic deposition, forest harvesting, or other process that contribute to acid soil infertility. This ratio, however, may not be a reliable indicator of stress in areas with both high atmospheric deposition of ammonium and magnesium deficiency via antagonism involving ammonium rather than aluminum and in areas with soil solutions with calcium concentrations greater than $500 \mu\text{mol/L}$ (National Science and Technology Council, 1998). Cronan and Grigal (1995), based on a review of the literature, have made the following estimates for determining the effects of acidic deposition on tree growth or

nutrition: (a) forests have a 50% risk of adverse effects if the Ca:Al ration is 1.0; (b) the risk is 75% if the ratio is 0.5:1.0; and (c) the risk approaches 100% if the ratio is 0.2:1.0. The Ca:Al ratio of soil solution provides only an index of the potential for Al stress. Cronan and Grigal (1995) stated that the overall uncertainty of the Ca:Al ratio associated with a given probability ratio is considered to be approximately $\pm 50\%$. Determination of thresholds for potential forest effects requires the use of four successive measurement endpoints in the soil, soil solution, and plant tissue: (a) soil base saturation less than 15% of effective cation exchange capacity; (b) soil solution Ca:Al molar ratio less than 1.0:1.0 for 50% risk; (c) fine roots tissue Ca:Al molar ratio less than 0.2:1.0 for 50% risk; and (d) foliar tissue Ca:Al molar ratio less than 12.5:1.0 for 50% risk. The application of the Ca:Al ratio indicator for assessment and monitoring of forest health risks has been recommended for sites or in geographic regions where the soil base saturation is $< 15\%$.

Critical Loads. The critical loads framework for assessing the effects of atmospheric deposition originated in Europe where the concept has generally been accepted as the basis for abatement strategies to reduce or prevent injury to the functioning and vitality of forest ecosystems caused by long-range transboundary chronic acidic deposition (Løkke et al., 1996). The critical load has been defined as a “quantitative estimate of an exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge” (Løkke et al., 1996). This concept is useful for estimating the amounts of pollutants that sensitive ecosystems can absorb on a sustained basis without experiencing measurable degradation.

The response to pollutant deposition of an ecosystem is a direct function of the level of sensitivity of the ecosystem to the pollutant and its capability to ameliorate change. The estimation of ecosystem critical loads requires an understanding of how an ecosystem will respond to different loading rates in the long term. The approach can be of special value for ecosystems receiving chronic deposition of pollutants such as nitrogen and sulfur. A program was designed to develop and evaluate a framework for setting critical loads of nitrogen and sulfur in the United States in 1989 (Strickland et al., 1993). A flexible six-step approach has been outlined for use with the critical load framework (Figure 4-19). These steps are (1) selection of ecosystem components, indicators, and characterization of the resource;




Figure 4-19. Key elements of proposed framework for determining critical loads for nitrogen and sulfur in the United States.

Source: Strickland et al. (1993).

(2) definition of the functional subregions; (3) characterization of deposition within each of the subregions; (4) definition of an assessment endpoint; (5) selection and application of models; and (6) mapping of projected ecosystem responses. The approach permits comparability in ecosystem characteristic and data availability (Strickland et al., 1993).

Ecological endpoints or indicators are measurable characteristics related to the structure, composition, or functioning of ecological systems (i.e., indicators of condition). One or more measurable endpoints are associated with each of the EEA elements listed in Table 4-1. These assessment endpoints represent a formal expression of the environmental value that is to be protected. If the assessment endpoint is to be used as a regulatory limit, it should be socially relevant. Selection of a specific ecosystem for study will depend on the severity of the problem of concern for a region. Time scales of response must be considered in selecting and evaluating ecosystem response(s) to changes in atmospheric deposition. Responses of aquatic ecosystems

to depositions can occur quickly. Surface water acidification associated with nitrate leaching should respond to decreases in nitrogen loading in a short period of time. However, changes in growth responses of vegetation resulting from soil nutrient imbalances may require years or decades to detect. The focus of concern should be on the populations within an ecosystem that are sensitive to nitrogen and sulfur deposition (Hunsaker et al., 1993).

Biogeochemical indicators for monitoring forest nitrogen status have been proposed by Fenn and Poth (1998). Because nitrogen is a major constituent of all forms of life and is cycled through a complex web of processes involving many biotic and abiotic mechanisms, evaluating forest nitrogen status is a challenge. Indicators of ecosystems at risk of nitrogen saturation should include those that can be identified when nitrogen availability exceeds biotic demand. Such indicators typically should monitor parameters that are normally at background or low levels in nitrogen-limited systems and should be those that commonly respond to excess nitrogen in a wide range of ecosystems (Fenn and Poth, 1998). Such indicators include foliar nitrogen, nutrient ratios (N:P, N:cation), foliar nitrate, foliar $\delta^{15}\text{N}$, arginine concentration, soil C:N ratio, NO_3^- in soil extracts or in soil solution, and flux rates of nitrogenous trace gases from soil (Fenn et al., 1998). The cardinal indicator or manifestation of nitrogen saturation in all ecosystem types, including California forests and chaparral, is increased and prolonged NO_3^- loss below the main rooting zone and in stream water. Seasonal patterns of stream water nitrate concentrations are especially good indicators of watershed nitrogen status (Fenn and Poth, 1998).

In Europe, the elements used in the critical load concept include: a biological indicator, a chemical criterion, and a critical value. The biological indicator is the organism used to indicate the status of the receptor ecosystem; the chemical criterion is the parameter that results in harm to the biological indicator; and the critical value is the value of the chemical criterion below which no significant harmful response occurs to the biological indicator (Løkke et al., 1996). Trees, and sometimes other plants, are used as biological indicators in the case of critical loads for forests. The critical load calculation using the current methodology is essentially an acidity/alkalinity mass balance calculation. The chemical criterion must be expressible in terms of alkalinity. Initially, the Ca:Al ratio was used; but, recently, the (Ca+Mg+K):Al ratio has been used (Løkke et al., 1996).

Ideally, changes in acidic deposition should result in changes in the status of the biological indicator used in the critical load calculation. However, the biological indicator is the integrated response to a number of different stresses. Furthermore, some other organisms are more sensitive to acid deposition than trees. At high concentrations, Al^{3+} is known to be toxic to plants, inhibiting root growth and, ultimately, plant growth and performance (Løkke et al., 1996; National Science and Technology Council, 1998). Sensitivity to Al varies considerably between species and within species because of changes in nutritional demands and physiological status that are related to age and climate. Experiments have shown that there are large variations in Al sensitivity, even among ecotypes.

Mycorrhizal fungi have been suggested as possible biological indicators by Løkke et al. (1996) because they are intimately associated with tree roots, depend on plant assimilates, and play an essential role in plant nutrient uptake, thereby influencing the ability of their host plants to tolerate different anthropogenically generated stresses. Mycorrhizas and fine roots are an extremely dynamic component of below-ground ecosystems and can respond rapidly to stress. They have a relatively short lifespan, and their turnover appears to be strongly controlled by environmental factors. Changes in mycorrhizal species composition, or the loss of dominant mycorrhizal species in areas where diversity is already low, may lead to increased susceptibility of plants to stress (Løkke et al., 1996). Stress affects the total amount of carbon fixed by plants and modifies carbon allocation to biomass, symbionts, and secondary metabolites. The physiology of carbon allocation has also been suggested as an indicator of anthropogenic stress (Andersen and Rygiewicz, 1991). Because mycorrhizal fungi are dependent for their growth on the supply of assimilates from the host plants, stresses that shift the allocation of carbon reserves to the production of new leaves at the expense of supporting tissues will be reflected rapidly in decreased fine root and mycorrhizal biomass (Winner and Atkinson, 1986). Decreased carbon allocation to roots also affects soil carbon and rhizosphere organisms. Soil dwelling animals are important for decomposition, soil aeration, and nutrient redistribution in the soil. They contribute to decomposition and nutrient availability mainly by increasing the accessibility of dead plant material to microorganisms. Earthworms decrease in abundance, and in species number, in acidified soils (Løkke et al., 1996).

Effects of Wet and Dry Deposition on Biogeochemical Cycling — the Integrated Forest Study. The Integrated Forest Study (IFS; Johnson and Lindberg, 1992a) has provided the most extensive data set available on wet and dry deposition, as well as on deposition effects on the cycling of elements in forest ecosystems. The overall patterns of deposition and cycling have been summarized by Johnson and Lindberg (1992a), and the reader is referred to that reference for details. The following is a summary of particulate deposition, total deposition, and leaching in the IFS sites.

Particulate deposition in the IFS was separated at the 2- μm level; however, the decision was made to include total particulate deposition in this analysis and may include the deposition of particles larger than 10 μm . Particulate deposition contributes considerably to the total impact of base cations to most of the IFS sites. On average, particulate deposition contributes 47% to total Ca deposition (range: 4 to 88%), 49% of total K deposition (range: 7 to 77%), 41% to total Mg deposition (range: 20 to 88%), 36% to total Na deposition (range: 11 to 63%), and 43% to total base cation deposition (range: 16 to 62%). Of total particulate deposition, the vast majority (> 90%) is > 2 μm in size.

Figures 4-20 through 4-23 summarize the deposition and leaching of Ca, Mg, K, and total base cations for the IFS sites. As noted in the original synthesis (Johnson and Lindberg, 1992a), measurements indicated annual gains of base cations for some sites (i.e., total deposition > leaching), some losses (total deposition < leaching) at others, and some approximately in balance. Not all cations follow the same pattern at each site. For example, a net accumulation of Ca occurs at the Coweeta, TN, Durham (Duke), NC and Bradford Forest, FL sites (Figure 4-20), whereas accumulation of K was noted at the Duke, Florida, Thompson, WA, Huntington Forest, NY, and White Face Mountain, NY sites (Figure 4-22). Magnesium accumulated only at the Florida site (Figure 4-21), and only at the Florida site is there a clear net accumulation of total base cations (Figure 4-23).

As noted previously, the factors affecting net Ca accumulation or loss include the soil-exchangeable cation composition; base cation deposition rate; the total leaching pressure because of atmospheric sulfur and nitrogen inputs, as well as natural (carbonic and organic) acids; and biological demand (especially for potassium). At the Florida site, which has a very cation-poor, sandy soil (derived from marine sand), the combination of all these factors leads to net base cation accumulation from atmospheric deposition (Johnson and Lindberg, 1992a).




Figure 4-20. Calcium deposition in > 2- μm particles, < 2- μm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. CP = *Pinus strobus*, Coweeta, TN; DL = *Pinus taeda*, Durham (Duke), NC; GS = *Pinus taeda*, B. F. Grant Forest, GA; LP = *Pinus taeda*, Oak Ridge, TN; FS = *Pinus eliottii*, Bradford Forest, FL; DF = *Psuedotsuga menziesii*, Thompson, WA; RA = *Alnus rubra*, Thompson WA; NS = *Picea abies*, Nordmoen, Norway; HF = northern hardwood, Huntington Forest, NY; MS = *Picea rubens*, Howland, ME; WF = *Picea rubens*, Whiteface Mountain, NY; and ST = *Picea rubens*, Clingman's Dome, NC.

The site showing the greatest net base cation losses, the red alder stand in Washington state, is under extreme leaching pressure by nitrate produced by excessive fixation by that species (Van Miegroet and Cole, 1984). In the red spruce site in the Smokies, the combined effects of SO_4^{2-} and NO_3^- leaching are even greater than in the red alder site (Figure 4-24), but a considerable proportion of the cations leached from this extremely acid soil consist of H^+ and Al^{3+} rather than of base cations (Johnson and Lindberg, 1992a). Thus, the red spruce site in the Smokies is approximately in balance with respect to calcium and total base cations, despite the very high leaching pressure at this site (Figures 4-20 and 4-23).




Figure 4-21. Magnesium deposition in > 2-µm particles, < 2-µm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.




Figure 4-22. Potassium deposition in > 2-µm particles, < 2-µm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.




Figure 4-23. Base cation deposition in > 2-µm particles, < 2-µm particles, and wet forms (upper bars) and leaching (lower bars) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviation.

The relative importance of particulate-base-cation deposition varies widely with site and cation and is not always related to the total deposition rate. The proportion of Ca deposition in particulate form ranges from a low of 4% at the Whiteface Mountain site to a high of 88% at the Maine site (Figure 4-20). The proportion of K deposition as particles ranges from 7% at the Smokies site to 77% at the Coweeta site (Figure 4-22). The proportion of total base cation deposition ranges from 16% at the Whiteface site to 62% at the Maine site (Figure 4-23). Overall, particulate deposition at the site in Maine accounted for the greatest proportion of Ca, Mg, K, and base cation deposition (88, 88, 57, and 62%, respectively) even though total deposition was relatively low. At some sites, the relative importance of particulate deposition varies considerably by cation. At the Whiteface Mountain site, particulate deposition accounts for 4, 20, and 40% of Ca, Mg, and K deposition, respectively. At the red spruce site in the Smokies, particulate deposition accounts for 46, 26%, 7% of Ca, Mg, and K deposition, respectively.




Figure 4-24. Total cation leaching (total height of bar) balanced by SO₄²⁻ and NO₃⁻ estimated from particulate deposition (assuming no ecosystem retention, particulate sulfur and nitrogen) and by other sources (both deposition and internal) of SO₄²⁻ and NO₃⁻ (other sulfur and nitrogen sources) and by other anions in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.

As indicated in the IFS synthesis, SO₄²⁻ and NO₃⁻ leaching often are dominated by atmospheric sulfur and nitrogen (Johnson and Lindberg, 1992a). The exceptions to this are in cases where natural nitrogen inputs are high (i.e., the nitrogen-fixing red alder stand), as are NO₃⁻-leaching rates even though nitrogen deposition is low, and where soils adsorb much of the atmospherically deposited SO₄²⁻ thus reducing SO₄²⁻ leaching compared to atmospheric sulfur input.

Sulfate and NO₃⁻ leaching have a major effect on cation leaching in many of the IFS sites (Johnson and Lindberg, 1992a). Figure 4-24 shows the total cation leaching rates of the IFS sites and the degree to which cation leaching is balanced by SO₄²⁻ + NO₃⁻ deposition. The SO₄²⁻ and NO₃⁻ fluxes are subdivided further into that proportion potentially derived from particulate sulfur and nitrogen deposition (assuming no ecosystem retention, a maximum effect) and other sulfur and nitrogen sources (i.e., wet and gaseous deposition, internal production).

As noted in the IFS synthesis, total SO_4^{2-} and NO_3^- inputs account for a large proportion (28 to 88%) of total cation leaching in most sites. The exception is the Georgia loblolly pine site where there were high rates of HCO_3^- and Cl^- leaching (Johnson and Lindberg, 1992a). The role of particulate sulfur and nitrogen deposition in this leaching is generally very small (< 10%), however, even if it is assumed that there is no ecosystem sulfur or nitrogen retention.

It was previously noted in this chapter that the contribution of particles to total deposition of nitrogen and sulfur at the IFS sites is lower than that for base cations. On average, particulate deposition contributes 18% to total nitrogen deposition (range: 1 to 33%) and 17% to total sulfur deposition (range: 1 to 30%). Particulate deposition contributes only a small amount to total H^+ deposition (average = 1%; range: 0 to 2%). (It should be noted, however, that particulate H^+ deposition in the > 2 μm fraction was not measured.)

Based on the IFS data, it appears that particulate deposition has a greater effect on base cation inputs to soils than on base cation losses associated with the inputs of sulfur, nitrogen, and H^+ . It cannot be determined what fraction of the mass of these particles are < 10 μm , but only a very small fraction is < 2 μm . These inputs of base cations have considerable significance, not only to the base cation status of these ecosystems, but also to the potential of incoming precipitation to acidify or alkalize the soils in these ecosystems. As noted above, the potential of precipitation to acidify or alkalize soils depends on the ratio of base cations to H^+ in deposition, rather than simply on the input of H^+ alone. In the case of calcium, the term “lime potential” has been applied to describe this ratio; the principle is the same with respect to Mg and K. Sodium is a rather special case, in that it is a poorly absorbed cation and leaching tends to balance input over a relatively short term.

Net balances of base cations tell only part of the story as to potential effects on soils; these net losses or gains must be placed in the perspective of the soil pool size. One way to express this perspective is to compare soil pool sizes with the net balances. This comparison is made for exchangeable pools and net balances for a 25-year period as shown in Figures 4-25 to 4-27. It is readily seen that the net leaching losses of cations pose no threat in terms of depleting soil-exchangeable Ca^{2+} , K^+ , or Mg^{+2} within 25 years at the Coweeta, Duke, Georgia, Oak Ridge, or Douglas-fir sites. However, there is a potential for significant depletion at the red alder, Whiteface Mountain, and Smokies red spruce sites.




Figure 4-25. Soil exchangeable Ca²⁺ pools and net annual export of Ca²⁺ (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.




Figure 4-26. Soil exchangeable Mg²⁺ pools and net annual export of Mg²⁺ (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.




Figure 4-27. Soil exchangeable K^+ pools and net annual export of K^+ (deposition minus leaching times 25 years) in the Integrated Forest Study sites. See Figure 4-20 for site abbreviations.

The range of values for soil-exchangeable turnover is very large, reflecting variations in both the size of the exchangeable pool and the net balance of the system. Soils with the highest turnover rates are those most likely to experience changes in the shortest time interval, other things being equal. Thus, the Whiteface Mountains, Smokies, and Maine red spruce sites, the Thompson red alder site, and the Huntington Forest northern hardwood site appear to be most sensitive to change. The actual rates, directions, and magnitudes of change that may occur in these soils (if any) will depend on weathering inputs and vegetation outputs, in addition to deposition and leaching. It is noteworthy that each of the sites listed above as sensitive has a large store of weatherable minerals, whereas many of the other soils (with larger exchangeable cation reserves) have a small store of weatherable minerals (e.g., Coweeta white pine, Duke loblolly pine, Georgia loblolly pine, and Oak Ridge loblolly pine) (Johnson and Lindberg, 1992a; April and Newton, 1992).

Base cation inputs are especially important to the Smokies red spruce site because of potential aluminum toxicity and calcium and magnesium deficiencies. Johnson et al. (1991a)

found that soil-solution Al concentrations occasionally reached levels found to inhibit Ca uptake and cause changes in root morphology in solution culture studies of red spruce (Raynal et al., 1990). In a follow-up study, Van Miegroet et al. (1993) found a slight, but significant, growth response to Ca and Mg fertilizer in red spruce saplings near the Smokies red spruce site. Joslin et al. (1992) reviewed soil and solution characteristics of red spruce in the southern Appalachians, and it appears that the IFS site is rather typical of observations for that area.

Wesselink et al. (1995) reported on the complicated interactions among changing deposition and soils at a beech and spruce forest in Solbing, Germany (including repeated sampling of soil exchangeable base cation pools) from 1969 to 1991 and compared these results with those of a simulation model. They identified three basic stages of change in this ecosystem. During Stage I, there was increased deposition of sulfur and constant deposition of base cations, causing increased base cation leaching and reduced base saturation in the soils. During Stage II, sulfur deposition is reduced; soil solution SO_4^{2-} and base cation leaching decline accordingly, but base saturation continues to decrease. During Stage III, two alternative scenarios are introduced: (a) sulfur deposition continues to decline, whereas base cation deposition stays constant; or (b) both sulfur and base cation deposition decline. Under Stage III-a, sulfate and base cation leaching continue to decline, and base saturation begins to increase as base cations displace exchangeable Al and cause it to transfer to the gibbsite pool. Under Stage III-b, this recovery in base saturation is overridden by the reduction in base cation deposition.

The IFS project, for the first time, accurately quantified atmospheric deposition inputs to nutrient cycles using state-of-the-art techniques to measure wet and dry deposition. The main aim of the project was to determine effects of atmospheric deposition on nutrient status of a variety of forest ecosystems and to determine if these effects are in any way related to current or potential forest decline. Acidic deposition appears to be having a significant effect on nutrient cycling in most of the forest ecosystems studied in the IFS project. The exceptions were the relatively unpolluted Douglas fir, red alder, and Findley Lakes sites in Washington state. The nature of the effects, however, varies from one location to another (Johnson, 1992). In all but the relatively unpolluted Washington sites, atmospheric deposition was having a significant (often overwhelming) effect on cation leaching from the soils. In general, nutrient budget data from IFS and the literature suggest that the susceptibility of southeastern sites to base cation depletion

from soils and the development of cation deficiencies by that mechanism appears to be greater than in northern sites (Johnson, 1992).

Atmospheric deposition may have significantly affected the nutrient status of some IFS sites through the mobilization of Al. Soil-solution Al levels in the Smokies sites approach and sometimes exceed levels noted to impede cation uptake in solution culture studies. It is therefore possible that the rates of base cation uptake and cycling in these sites have been reduced by soil solution Al levels. To the extent that atmospheric deposition has contributed to these elevated soil-solution Al levels, it has likely caused a reduction in base cation uptake and cycling rates at these sites. Nitrate and sulfate are the dominant anions in the Smokies sites, and nitrate pulses are the major cause of Al pulses in soil solution (Johnson, 1992). The connection between Al mobilization and forest decline is not clear. The decline in red spruce has been more severe in the Northeast than in the Southeast; yet all evidence indicates that Al mobilization is most pronounced in the southern Appalachians. However, the Whiteface Mountain site which was selected for study because it was in a state of decline had soil solution levels lower than in the Smokies, which are not in a visibly obvious state of decline (there was no dieback other than the fir killed by the balsam wooly adelgid and no needle yellowing). Thus, Al mobilization continues to be worthy of further study (Johnson, 1992).


The simple calculations shown above give some idea of the importance of particulate deposition in these forest ecosystems, but they cannot account for the numerous potential feedbacks between vegetation and soils nor for the dynamics through time that can influence the ultimate response. One way to examine some of these interactions and dynamics is to use simulation modeling. The nutrient cycling model (NuCM) has been developed specifically for this purpose and has been used to explore the effects of atmospheric deposition, fertilization, and harvesting on some of the IFS sites (Johnson et al., 1993). The NuCM model is a stand-level model that incorporates all major nutrient cycling processes (uptake, translocation, leaching, weathering, organic matter decay, and accumulation).

Johnson et al. (1999) used the NuCM model to simulate the effects of reduced S, N, and base cation deposition on nutrient pools, fluxes, soil, and soil solution chemistry in two contrasting southern Appalachian forest ecosystems: the red spruce and Coweeta hardwood sites from the IFS project. The scenarios chosen for these simulations included no change; 50%

N and S deposition; 50% base cation deposition; and 50% N, S, and base cation deposition (50% N, S, base cation). The NuCM simulations suggested that, for the extremely acid red spruce site, S and N deposition is the major factor affecting soil-solution Al concentrations and base cation deposition is the major factor affecting soil solution base cation concentrations. The effects of S and N deposition were largely through changes in soil solution SO_4^{2-} and NO_3^- and, consequently, mineral acid anion (MAA) concentrations rather than through changes in soils. This is illustrated in Figures 4-28a,b and 4-29a,b, which show simulated soil-solution mineral-acid anions, base cations, Al, and soil base saturation in the B horizon in the red spruce site. The 50% S and N scenario caused reductions in soil solution SO_4^{2-} , NO_3^- and, therefore, MAA concentrations, as expected. This, in turn, caused short-term reductions in base cation concentrations. However, by the end of the 24-year simulation, base cations in the 50% S, N scenario were nearly as high as in the no-change scenario because base saturation had increased and the proportion of cations as Al decreased. The 50% base cation scenario had virtually no effect on soil solution SO_4^{2-} , NO_3^- and, therefore, MAA concentrations, as expected, but did cause a long-term reduction in base cation concentrations. This was caused by a long-term reduction in base saturation (Figure 4-29a,b). Thus, the effects of base cation deposition were solely through changes in soils rather than through changes in soil solution MAA, as postulated by Driscoll et al. (1989). In the less acidic Coweeta soil, base saturation was high and little affected by the scenario cited above; Al was unimportant; and S and N deposition had a much greater effect than base cation deposition in all respects (Figure 4-30a,b).

In summary, Johnson et al. (1999) found that the results of the red spruce simulations, in part, supported the hypothesis of Driscoll et al. (1989) in that: base cation deposition can have a major effect on base cation leaching through time in an extremely acid system. This effect occurred through changes in the soil exchanger and not through changes in soil solution MAA concentration. On the other hand, S and N deposition had a major effect on Al leaching at the Noland Divide site. This occurred primarily because of changes in soil solution MAA concentration. At the less acidic Coweeta site, base cation deposition had a minor effect on soils and soil solutions; whereas S and N deposition had delayed but major effects on base cation leaching because of changes in SO_4^{2-} and MAA concentrations.

Red Spruce



4-138

Figure 4-28a. Simulated soil solution mineral acid anions in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).




Figure 4-28b. Simulated soil solution base cations in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).





Figure 4-29a. Simulated soil solution Al in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).

Red Spruce



4-141

Figure 4-29b. Simulated soil solution soil base saturation in the red spruce site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).

Coweeta




Figure 4-30a. Simulated soil solution mineral acid anions in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).




Figure 4-30b. Simulated soil solution base cations in the Coweeta site with no change, 50% N and S deposition, and 50% base cation deposition.

Source: Redrawn from Johnson et al. (1999).

Effects of Trace Elements

Trace metals are natural elements that are ubiquitous in small (trace) amounts in soils, ground water, and vegetation. Many are essential elements required for growth by plants and animals as micronutrients. Naturally occurring surface mineralizations can produce metal concentrations in soils and vegetation that are as high, or higher, than those in the air and deposited near man-made sources (Freedman and Hutchinson, 1981). The occurrence and concentration of trace metals in any ecosystem component depend on the sources of the metal (i.e., via the soil or as a particulate). Even when air pollution is the primary source, continued deposition can result in the accumulation of trace metals in the soil (Martin and Coughtrey, 1981). Many metals are deposited and bound into soils by chemical processes and are not available to plants (Saunders and Godzik, 1986).

When aerial deposition is the primary source of metal particles, both the chemical form and particle size deposited determine the heavy metal concentration in the various ecosystem components (Martin and Coughtrey, 1981). Human activities introduce heavy metals into the atmosphere and have resulted in the deposition of antimony, cadmium, chromium, copper, lead, molybdenum, nickel, silver, tin, vanadium, and zinc (Smith, 1990c). Extensive evidence indicates that heavy metals deposited from the atmosphere to forests accumulate either in the richly organic forest floor, or in the soil layers immediately below the floor, areas where the interaction of roots and soil is greatest. The greater the depth of soil, the lower the metal concentration. The accumulation of metal in the soil layers where the biological activity is greatest, therefore, has the potential to be toxic to roots and soil organisms and to interfere with nutrient cycling (Smith, 1990c). The shallow-rooted plant species are most likely to take up metals from the soil (Martin and Coughtrey, 1981). Though all metals can be directly toxic at high levels, only copper, nickel, and zinc have been documented as frequently being toxic. Toxicity due to cadmium, cobalt, and lead has been seen only under unusual conditions (Smith, 1990c). Exposures to the above metals at lower concentrations have the potential, over the long term, to interfere with the nutrient-cycling processes when they affect mycorrhizal function.

Biological accumulation of metals through the plant-herbivore and litter-detritivore chains can occur. A study of Cd, Pb, and Zn concentrations in earthworms suggested that Cd and Zn were bioaccumulated in earthworms, but Pb was not. Studies indicate that heavy metal deposition onto the soil, via food chain accumulation, can lead to excessive levels and toxic

effects in certain animals. Cadmium appears to be relatively mobile within terrestrial food chains; however, the subsequent mobility of any metal after it is ingested by a herbivorous animal depends on the site of accumulation within body tissues. Although food chain accumulation of a metal may not in itself cause death, it can reduce the breeding potential in a population (Martin and Coughtrey, 1981).

Case studies have revealed that the deposition of Cu and Zn particles around a brassworks can result in an accumulation of incompletely decomposed litter. In one study, litter accumulation was reported up to 7.4 km from the stack of a primary smelter in southeastern Missouri. Similar results were reported around a metal smelter at Avonmouth, England. In the latter case, litter accumulation was associated closely with concentrations specifically of Cd, as well as with those of Pb, Cu, and Zn (Martin and Coughtrey, 1981). Accumulations of metals emitted in PM were also reported in soil litter close to a metal smelter at Palmerton, PA in 1975 and 1978. The continued presence of Cd, Pb, Zn, and Cu in the upper soil horizons (layers) were observed 6 years after the smelter terminated operation in 1980. Metal levels were highest near the smelter. Experimental data (using mesh bags containing litter) supports the hypothesis that reduced decomposition occurs close to heavy metal sources.

Accumulation of heavy metals in litter presents the greatest potential for interference with nutrient cycling. Microorganisms are essential for the decomposition of organic matter and soil fertility. Toxic effects on the microflora can be caused by Zn, Cd, and Cu. Addition of a few mg/kg of soil of Zn can inhibit the more sensitive microbial processes (van Beelen and Doelman, 1997). Experiments by Kandeler et al. (1996) indicated that microbial biomass and enzyme activities decreased with increasing heavy metal concentrations. The amount of decrease varied among the enzymes, with those involved in carbon cycling being least affected, the activities of the enzymes involved in the cycling of N, P, and S (especially arylsulfatase and phosphatase) were dramatically affected.

Metal accumulation in litter can be found mainly around brass works and Pb and Zn smelters. Invertebrates inhabiting soil litter may also accumulate metals. Earthworms from roadsides were shown to contain elevated concentrations of Cd, Ni, Pb, and Zn; but, interference with earthworm activity was not cited (Martin and Coughtrey, 1981). It has been shown, however, that when soils are acidic, earthworm abundance decreases, and bioaccumulation of metals from soil may increase exponentially with decreasing pH (Løkke et al, 1996). Organisms

that feed on earthworms living in soils with elevated levels of Cd, Ni, Pb, and Zn for extended periods may accumulate Pb and Zn to toxic levels (Martin and Coughtrey, 1981). Increased concentrations of heavy metals have been found in a variety of small mammals living in areas with elevated heavy metal concentrations in the soils. Furthermore, increased amounts of metal in body tissues were seen in both amphibians and mammals. Levels of Cd in the kidneys and liver of white-tailed deer (*Odocoileus virginianus*) were five times higher at Palmerton, PA than in those from 180 km southwest (downwind). The abnormal amounts of metal in the tissues of terrestrial vertebrates and the absence or low abundance of wildlife at Palmerton indicated that ecological processes within 5 km of the Zn smelter continued to be markedly influenced even 6 years after its closing (Storm et al., 1994).

Studies by Babich and Stotsky (1978) support the concept that increased accumulation of litter in metal-contaminated areas is due to the effects on the microorganismal populations. Cadmium toxicity to microbial populations was observed to decrease and prolong logarithmic rates of microbial population increase, to reduce microbial respiration and fungal spore formation and germination, to inhibit bacterial transformation, and to induce abnormal morphologies. Smith (1991) reported the effects of Cd, Cu, Ni, and Zn on the symbiotic activity of fungi, bacteria, and actinomycetes. The formation of mycorrhizae by *Glomus mosseae* with onions was reduced when Zn, Cu, Ni, and Cd was added to the soil. The relationship of the fungus with white clover, however, was not changed. It was suggested that the effect of heavy metals on vesicular-arbuscular mycorrhizal fungi will vary from host to host (Gildon and Tinker, 1983). Studies with ericoid plants indicated that, in addition to *Calluna vulgaris*, mycorrhizae also protect *Vaccinium macrocarpa* and *Rhodendron ponticum* from heavy metals (Bradley et al., 1981). Heavy metals tend to accumulate in the roots, thereby lessening shoot toxicity.

The effects of sulfur deposition on litter decomposition in the vicinity of smelters must also be considered. Metal smelters emit SO₂ as well as heavy metals. Altered litter decomposition rates have been well documented near SO₂ sources (Prescott and Parkinson, 1985). The presence of sulfur in litter has been associated with reduced microbial activity (Bewley and Parkinson, 1984). In addition, the effects of sulfur on the symbiotic activity of fungi, bacteria, and actinomycetes were reported by Smith (1990d).

The potential pathways of accumulation of trace metals in terrestrial ecosystems, as well as the possible consequences of trace metal deposition on ecosystem functions, is summarized in

Figure 4-31. The generalized trophic levels found in an ecosystem and the various physiological and biological processes that could be affected by trace metals are shown in the figure.

Reduction in physiological processes can affect productivity, fecundity, and mortality (Martin and Coughtrey, 1981). Therefore, any effects on the structure and function of an ecosystem are likely to occur through the soil and litter (Tyler, 1972).




Figure 4-31. Relationship of plant nutrients and trace metals with vegetation. Compartments (roman numerals) represent potential storage sites; whereas arrows (arabic numerals) represent potential transfer routes.

Certain species of plants are tolerant of metal-contaminated soils (e.g., soils from mining activities) (Antonovics et al., 1971). Certain species of plants also have been used as bioindicators of metals (e.g., *Astragalus* is an accumulator of selenium). The sources of both

macroelements and trace metals in the soil of the Botanical Garden of the town of Wroclow, Poland, were determined by measuring the concentrations of the metals in *Rhododendron catawbiense*, *Ilex aquifolium*, and *Mahonia aquifolium* growing in a garden and comparing the results with the same plant species growing in two botanical gardens in nonpolluted areas. Air pollution deposition was determined to be the source of metals in plants rather than soils contamination (Samecka-Cymerman and Kempers, 1999).

The effects of Pb in ecosystems are discussed in the EPA document *Air Quality Criteria for Lead* (U.S. Environmental Protection Agency, 1986). Studies have shown that there is cause for concern in three areas in which ecosystems may be extremely sensitive to Pb: (1) delay of decomposition because of inhibition by Pb of the activity of some decomposer microorganisms and invertebrates, (2) subtle shifts toward Pb-tolerant plant populations, and (3) Pb in the soil and on the surfaces of vegetation where it may circumvent the processes of biopurification. The problems cited above arise because Pb is deposited on the surface of vegetation, accumulates in the soil, and is not removed by the surface and ground water of the ecosystem (U.S. Environmental Protection Agency, 1986).

4.2.4 Urban Ecosystems

Humans dominate Earth's ecosystems. Evidence accumulating from anthropological and archeological research indicates that human influence has been pervasive for thousands of years (Grimm et al., 2000). Major human effects on the environment probably began as early as 12,000 to 15,000 years ago and continue to be a major influence on all natural ecosystems. Human activities have been more intense in cities, suburbs, exurbs and in supporting hinterlands (Grimm et al., 2000). Interest has increased recently in the study of urban ecological systems.

Vitousek et al. (1997) pointed out that understanding a human-dominated planet requires that the human dimensions of global change; that is, the social, cultural, and other drivers of human actions need to be included in ecological analyses. Therefore, humans must be integrated into models for more complete understanding of extant ecological systems.

In the past, ecological plant or animal studies conducted in urban settings used traditional ecological approaches and considered humans as agents of disturbance. Although the term "urban ecosystem" has been used to describe human-dominated ecosystems, it does not adequately take into account the developmental history, sphere of influence, and potential

impacts required to understand the true nature of an urban ecosystem (McIntyre, et al., 2000). Because urbanization is both an ecological and a social phenomenon, urban ecology implicitly recognizes the role humans play in developing unique systems. Therefore, if urban ecology is to be a truly interdisciplinary field, both social and natural sciences must be integrated into the study of urban ecosystems (McIntyre, et al., 2000).

Although the study of ecological phenomena in urban environments is not a new area of science, the concept of the city as an ecosystem is relatively new for the field of ecology (Grimm et al., 2000). There is a wealth of information on the terrestrial components of urban ecological systems. However, much of it is organized from the perspective of ecology *in* cities while the more comprehensive perspective identified as ecology *of* cities is needed (Pickett et al., 2001). The basic questions addressed by the literature of ecology *in* cities relate to how ecological patterns and processes differ in cities as compared with other environments. What is the effect of the city (i.e., a concentration of human population and activities) on the ecology of organisms inside and outside its boundary and influence? The concept of ecology *of* cities has to do with how aggregated parts make up the whole, i.e., how cities process energy or matter relative to their surroundings (Grimm et al., 2000). The latter concept includes primary production, species richness, biogeophysical budgets, ecosystem patterns and processes, and an open definition of urban ecosystems that incorporates the exchanges of materials and influence between cities and surrounding landscapes (Pickett et al., 2001). If ecosystems are to be understood, there is a need for a new integrative ecology that explicitly incorporates human decisions, culture, institutions, and economic systems (Grimm et al., 2000). This makes an ecological approach to land use planning essential to maintain long-term sustainability of ecosystem benefits, services, and resources (Zipperer et al., 2000). The ecological and social effects of “edge city” need to be studied as they may be greater than the previous patterns of suburbanization. The classical ecosystem approach and a patch dynamic approach are needed to understand and manage the dynamics of urban and urbanizing ecosystems (Zipperer et al., 2000).

There has been little work on the rates of atmospheric deposition to urban ecosystems despite extensive data on concentrations and chemical reactions of air pollutants in cities. A search of the literature produced no references that dealt with the effects of PM deposition. Lovett et al. (2000), however, reported that urban ecosystems are likely to be subjected to large rates of deposition of anthropogenic pollutants. Decades of research on urban air quality

indicate that cities are often sources of NO_x , SO_x , and dust, among many other pollutants. Some of these air pollutants are major plant nutrients (e.g., nitrogen) and may be affecting nutrient cycles in plant-dominated areas in and around cities, but have not been studied. The gases and particles in urban air can increase atmospheric deposition within and downwind of the city. Studying deposition rates of atmospheric pollutants in urban areas can provide quantitative estimates of amounts of gaseous and particulate air pollutants removed by urban vegetation.

To determine the patterns of atmospheric deposition and throughfall in the vicinity of a large city, Lovett et al. (2000) measured bulk deposition, oak forest throughfall, and particulate dust at sites along a transect within and to the north of New York City. They observed that concentrations and fluxes of NO_3^- , NH_4^+ , Ca^{2+} , Mg^{2+} , SO_4^{2-} , and Cl^- in throughfall all declined significantly with increasing distance from the city, while H^+ concentration and flux increased significantly with increasing distance from the city. Most of the change in concentrations and fluxes occurred within 45 km of the city. Additionally, it was observed that throughfall nitrogen was twice as high in the urban areas compared with suburban and rural areas. Most of the dry deposition of nitrate was from gaseous nitrogen oxides. As mentioned earlier, the effects of the atmospheric deposition of the particulate pollutants was not mentioned.

McDonnell et al. (1997) in a 10-year study of ecosystem processes along an urban-rural gradient included plant litter dynamics and nitrogen cycling of two key components of a forest ecosystem: litter decomposition and heavy metal levels in soil and foliar litter. Foliar litter decomposition integrates many features of the abiotic and biotic environment. It is an important site of heavy metal incorporation into ecosystems and provides both a habitat and a resource for fungi, bacteria, and invertebrates. Litter decomposition integrates the effects of resource quality, environmental factors, and activities of decomposer organisms on nutrient cycling and serves as an easily measured indicator of the effect of urbanization on an important ecosystem function. McDonnell et al. (1997) noted that levels of heavy metals were higher in the foliar litter of urban forest soils than in rural forest soils. The levels in urban forest stands approached or exceeded the levels reported to affect soil invertebrates, macrofungi, and soil microbial processes. The urban forests exhibited reduced fungal biomass and microarthropod densities when compared to rural stands. These results supported the concept that urban forests have depauperate communities because of anthropogenic stress resulting from poor air quality due to high levels of SO_2 , SO_4^{2-} , O_3 and NO_3^- ; elevated levels of soil- and forest-floor-heavy metals; and low water

availability such as those caused by hydrophobic soils (McDonnell et al., 1997). Thus, forests at the urban end of the gradient exhibited reduced fungal and microarthropod populations and poorer leaf quality than the more rural forests. The potential effect of these conditions on the ecosystem processes of decomposition and nitrogen cycling in urban forests appeared to be ameliorated by two other anthropogenic factors: increased average temperatures caused by the heat island effect and the introduction and successful colonization of earthworms in the urban forests (McDonnell et al., 1997).

McDonnell et al. (1997) observed that the changes in forest nitrogen dynamics were related to increased anthropogenic nitrogen deposition in an urban environment. The studies of Aber et al. (1989) in the northeastern United States on forest nitrogen dynamics demonstrated that elevated nitrogen deposition over many years results in increased nitrification and the mineralization of more nitrogen than can be taken up by plants and microorganisms. Nitrification can lead to decreases in fine root biomass and increases in nitrate leaching below the root zone. These effects of nitrogen deposition were not related to inputs from a specific source such as PM.

While studies of heavy metal deposition in or near cities have been performed, they do not cite the effects of metals in the soil. Pouyat and McDonnell (1991) discussed heavy metal accumulations in forest soils along an urban-rural gradient in southeastern New York. Variations in the amounts of Zn, Cu, Ni, and Cd appeared to be indicative of a pattern of atmospheric deposition near point sources (Section 4.3.2.6). The concentrations of heavy metals in forest floor and soils corresponded closely with the urban-rural land use gradient. Again, as in the study by Lovett et al. (2000), the pollutants were highest near the urban end of the gradient and declined toward rural sites, with Pb, Ni and Cu highest near the urban end.

The air quality of the region around East St. Louis has been of concern due to industries in the area (Kaminski and Landsberger, 2000a), which include ferrous and nonferrous metal smelters (Pb, Zn, Cu, and Al), coal-fired power plants, producers of organic and inorganic chemicals, municipal waste incinerators, and petroleum refineries. The city is also in the path of diverse plumes from refineries to the north, coal-fired power plants to the west, and nonferrous smelters to the south. Concentrations of heavy metals and metalloids (As, Cd, Cu, Hg, Pb, Sb, Zn) in the soil provided a basis for analysis (Kaminski and Landsberger, 2000b). These studies of the extent of long-term metal deposition on the soil surface and depth of soil contamination,

as well as the leaching dynamics of heavy metals were made to determine possible effects on biota uptake or groundwater contamination. The effects on biota are not mentioned; however, the soils in the area acted as a sink and there was little groundwater mobility (Kaminski and Landsberger, 2000b).

4.3 AIRBORNE PARTICLE EFFECTS ON VISIBILITY

4.3.1 Introduction

Visibility may be thought of as the degree to which the atmosphere is transparent to visible light (National Research Council, 1993). The beauty of scenic vistas in many parts of the United States is often diminished by haze that reduces contrast, washes out colors, and renders distant landscape features indistinct or invisible. This degradation of visibility is due primarily to the scattering and absorption of light by fine particles suspended in the atmosphere. One quantitative measure of visibility, used traditionally by meteorologists, is the visual range, defined as the farthest distance at which a large black object can be distinguished against the horizon sky (U.S. Environmental Protection Agency, 1979).

In August 1977, Congress amended the Clean Air Act (CAA) to establish as a national goal “the prevention of any future and remedying of any existing impairment of visibility in mandatory Class I Federal areas (many national parks and wilderness areas), which impairment results from manmade air pollution” (Title I Part C Section 169A, U.S. Code [1990]). The 1977 Amendments also included provisions requiring applicants for new major source permits to assess the potential for their projects to cause adverse effects on air quality-related values (e.g., visibility) in nearby Class I areas. In 1980, the EPA established regulatory requirements under Section 169A to address Class I protection from “reasonably attributable” visibility impairment, i.e., visibility impairment attributable to a single source or small group of sources.

The CAA, as amended in 1990 (Section 169B), required EPA both to conduct research on regional visibility impairment and to establish the Grand Canyon Visibility Transport Commission (GCVTC). The GCVTC was charged with assessing and providing advice to help preserve clear days and to improve visibility in the 16 national parks and wilderness areas located on the Colorado Plateau. The GCVTC also was mandated to provide recommendations to the U.S. EPA for reduction of visibility impairment due to regional haze, described as any

perceivable change in visibility (light extinction, visual range, contrast, or coloration) from that which would have existed under natural conditions and is caused predominantly by a combination of many anthropogenic sources over a wide geographical area (U.S. EPA, 1999a). In July 1999, EPA published the Regional Haze Rule (Federal Register, 1999). The regulation established a program for the improvement and protection of visibility in the 156 protected Class I parks and wilderness areas and included the establishment of baseline and current visibility conditions and the tracking of changes in visibility conditions over time. Implementation of the regional haze regulations is supported by EPA's PM_{2.5} monitoring network and an expanded Interagency Monitoring of Protected Visual Environments (IMPROVE) network. The PM_{2.5} monitoring network and the IMPROVE network are described briefly later in this section and in more detail elsewhere (National Park Service, 1998; Evans and Pitchford, 1991; U.S. Environmental Protection Agency, 2000b; U.S. Environmental Protection Agency, 2001a).

The objective of the visibility discussion in this section is to provide a brief description of the fundamentals of atmospheric visibility and to summarize the linkage between PM and visibility. Visibility is affected by air quality and, unlike the PM concentration, is not a property of an element of volume in the atmosphere. Visibility can be quantified only for a sight path and depends on the illumination of the atmosphere and the direction of view. However, the concentration of particles in the atmosphere plays a key role in determining visibility. Therefore, visibility impairment may be controlled by control of particle concentrations. The relationships between particles, other factors, and visibility impairment are described in this section. For a more detailed discussion on visibility, the reader is referred to the 1996 *Air Quality Criteria for Particulate Matter* (U.S. Environmental Protection Agency, 1996a); the Recommendations of the Grand Canyon Visibility Transport Commission (Grand Canyon Visibility Transport Commission, 1996); the National Research Council (National Research Council, 1993); the National Acid Precipitation Assessment Program (Trijonis et al., 1991); Interim Findings on the Status of Visibility Research (U.S. Environmental Protection Agency, 1995a); Visibility: Science and Regulation (Watson, 2002), and reports summarizing visibility science and data from the IMPROVE visibility monitoring network (Malm, 2000; Sisler, 1996; Sisler et al., 1993).

4.3.2 Factors Affecting Atmospheric Visibility

The visual perception of a distant object is influenced by a large number of factors including human vision (the eye), the brain's response to signals received from the eye, the interaction of light with the atmosphere (e.g., atmospheric illumination, path and transmitted radiance, contrast, and optical properties), and atmospheric pollution from natural and anthropogenic sources. Detailed discussion of this full range of topics can be found in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) and other general references (Malm, 1999; Watson, 2002). This section focuses only on those topics that have been addressed by more recent research, including atmospheric illumination, the optical properties of gases and particles in the atmosphere, and the effects of relative humidity on the optical properties of particles.

4.3.2.1 Optical Properties of the Atmosphere and Atmospheric Particles

Atmospheric particles and gases attenuate image-forming light as it travels from a viewed object to an observer. The fractional attenuation of light per unit distance is known as the light extinction coefficient. The light extinction coefficient, b_{ext} , is expressed in units of one over length, for example inverse kilometers (km^{-1}) or inverse megameters (Mm^{-1}). The light extinction coefficient can be expressed as the sum of the light scattering and light absorption coefficients of particles and gases:

$$b_{ext} = b_{ap} + b_{ag} + b_{sg} + b_{sp} \quad (4-7)$$

where the subscripts p and g signify particles and gases, and s and a signify scattering and absorption.

The light extinction coefficient can be measured with a reasonable degree of accuracy or can be calculated with the size, composition, shape, and the orientation of the particles. The light extinction coefficient is influenced by meteorological conditions and optical properties along the sight path.

Relationship Between Light, Targets, and Objects in a Sight Path

The appearance of a distant object is determined by light from two sources: the light reflected from the object itself (initial radiance) and the light reflected by the intervening atmosphere (path radiance). Human vision and the brain's response to signals received from the eye can distinguish between objects by contrast or differences in the radiance of adjacent objects. Light reflected by objects is attenuated by scattering and absorption as it travels through the atmosphere toward the observer. The portion that reaches the observer is the transmitted radiance.

During the daytime, the sight path is illuminated by the direct rays of the sun, diffuse skylight, light that has been reflected from the surface of the Earth, etc. Some of this illumination is scattered toward the observer by the air molecules and PM in the sight path. The accumulation of the light scattered into the sight path is the path radiance or air light. The path radiance significantly influences the light transmitted by the object being viewed. As the path radiance increases, the light transmitted by an object decreases.

The transmitted radiance carries the information about the object; the path radiance only carries information about the intervening atmosphere and is often quite featureless. When the transmitted radiance is dominant, visibility is good. Conversely, when the path radiance is dominant, visibility is poor. In a dense fog, the transmitted radiance from nearby objects can be seen, but the transmitted radiance from more distant objects is completely overwhelmed by the path radiance (i.e., the light scattered by the fog). Distant objects are lost in the white (or gray) of the fog (Gazzi et al., 2001).

Figure 4-32 illustrates the radiance seen by an observer looking at a hillside or through the aperture of a measurement instrument. The radiance that enters the eye of the observer (or the aperture of a measurement instrument) is known as the apparent radiance (i.e., the sum of the transmitted and path radiance). The competition between the transmitted radiance and the path radiance determines visibility.

Light Absorption and Scattering by Gases

In the ambient atmosphere, the only visible-light-absorbing gas of any consequence is nitrogen dioxide (NO₂), which primarily absorbs blue light and, if present in sufficient concentration across a sight path, contributes to the yellow or brown color seen in urban hazes.




Figure 4-32. Light reflected from a target toward an observer. The intervening atmosphere scatters a portion of this light out of the sight path and scatters light from the sun into the sight path. Some particles and gases also absorb a portion of the light from the target. The light scattered into the sight path increases with distance from the target, whereas the light transmitted from the target decreases with distance from the target. The visual range is the closest distance between the target and the observer at which the transmitted light can no longer be distinguished from the light scattered into the sight path.

Source: Watson and Chow (1994).

Usually the absorption by NO_2 is much smaller than the scattering by particles that are typically present in polluted environments, such as urban areas. The most common exception to this situation of relatively small NO_2 absorption is in effluent plumes from combustion facilities where the particles are effectively removed but the nitrogen oxide (NO), which can convert rapidly to NO_2 , is not removed. Except for such particle-depleted NO plumes, the light absorption coefficient for gases is usually ignored in determinations of the light extinction.

Light scattering by gases in the atmosphere is described by the Rayleigh scattering theory (van de Hulst, 1981) and is referred to as Rayleigh scattering. The magnitude of Rayleigh scattering depends on the gas density of the atmosphere and varies from about 9 Mm^{-1}

to 11 Mm^{-1} for most locations of interest, depending primarily on site elevation. To simplify comparisons of light extinction coefficient values among sites at a variety of elevations, a standard value of 10 Mm^{-1} is often used for the Rayleigh scattering component (Malm, 2000).

Light Absorption and Scattering by Particles

Light scattering by particles tends to dominate light extinction except under pristine atmospheric conditions when Rayleigh scattering by gas molecules is the largest contributor. If the particle size, refractive index, and shape are known, the extinction coefficient can be calculated. For particles of sizes similar to the wavelength of visible light, Mie equations for homogeneous spheres can be used to calculate the scattering and absorption of individual particles.

Absorption by particles is primarily caused by elemental carbon (also referred to as soot or light-absorbing carbon) generated by the incomplete combustion of fossil fuels. Some minerals in crustal particles also absorb light and can be a significant factor during fugitive dust episodes.

Most particle absorption data are determined by measuring light transmission or reflection of particles captured on filter media. Absorption estimates made in this way are sensitive to the filter substrate used, the optical configuration of the transmission measurement, particle loading on the filter, and particle scattering albedo with the result that there are significant uncertainties for measurements of filtered particles (Horvath, 1993). Another approach to estimating aerosol light absorption is by subtracting concurrent light scattering measurements made with a nephelometer from light extinction measurements made with a transmissometer. Substantial uncertainty in this difference approach results from the assumption that the point measurement of scattering is representative of the scattering over a long path (1 to 10 km) that is typically required for transmissometer measurements. A recently field-tested prototype photoacoustic spectrometer designed to determine absorption of suspended aerosol and an enclosed-folded path transmissometer offer hope for resolving the problems of the filter-based and difference approaches to the measurement of light absorption by particles (Arnott et al., 1999).

The relationship between elemental carbon concentration and particle absorption can be calculated using Mie equations for particles with known size distribution, particle density, index of refraction, shapes, and for various internal mixtures with non-absorbing aerosol materials (Fuller et al., 1999). Mie equations are used to determine the efficiency factors for extinction

Q_{ext} , Q_{scat} , and Q_{abs} . The efficiency factors represent the fraction of light falling on a circle with the same diameter and index of refraction as the particle. While such application of this theory can provide a range of absorption efficiencies for various model aerosol distributions, it is rare that sufficiently detailed particle characterization data for ambient aerosols are available. Also, although elemental carbon is the strongest and most common of the absorbing particles, light absorption by elemental carbon particles can be reduced when the particle is covered by other chemical species (Dobbins et al., 1994) or may be enhanced when coated with a non-absorbing refractive material such as ammonium sulfate (Fuller et al., 1999).

More commonly, estimates of elemental carbon absorption efficiency are empirically determined from the ratios of, or the slopes of regression analysis fits to, absorption coefficient and corresponding elemental carbon concentration measurements. Use of the regression approach permits the inclusion of crustal component concentrations as a second dependent parameter, so that crustal absorption can also be estimated. Uncertainties in the absorption efficiency determined empirically are a combination of the measurement uncertainties for the absorption coefficients, elemental carbon concentrations, and, where used, the crustal concentrations. In reviews of estimates of elemental carbon light absorption mass efficiency (i.e., the absorption coefficient per carbon mass concentration), Horvath (1993) and Liousse et al. (1993) found values ranging from 2 to 17 m²/g. Moosmüller et al. (1998) showed that by limiting the absorption coefficient estimates to those using photoacoustic methods, the absorption efficiency shows a wavelength dependence with highest values (17 m²/g) at the shortest wavelength used ($\lambda = 0.42 \mu\text{m}$) and lowest values (3 m²/g) at the longest wavelengths used ($\lambda = 0.8 \mu\text{m}$). The center of the visible light wavelength ($\lambda = 0.53 \mu\text{m}$) yielded elemental carbon absorption efficiencies values of ~ 10 m²/g, a commonly used value for elemental carbon absorption efficiency. Fuller et al. (1999) suggested that isolated spheres of light absorbing carbon have a specific absorption of less than 10 m²/g. Light absorption by carbon particles will be greater than 10 m²/g only if the particles are internally mixed and the occluding particles are sufficiently large. Absorption values for graphitic and amorphous carbon for primary sizes typical of diesel soot are ~ 5 m²/g.

Particle scattering tends to dominate light extinction except under pristine atmospheric conditions when Rayleigh scattering by gas molecules is the largest contributor. Light-scattering by particles has been reported to account for 68 to 86% of the total extinction coefficient in

several cities in California (Eldering et al., 1994). When light-scattering increases, visibility is impaired because of a decrease in the transmitted radiance and an increase in the path radiance. The single most important factor that determines the amount of light scattered by a particle is its size, as shown in Figure 4-33 (based on Mie calculations). The maximum single-particle scattering efficiency (i.e., scattering per cross-sectional area of a particle) is associated with particles with diameters of about the wavelength of visible light (centered at 0.53 μm). For particles that are small compared to the wavelength of light, the single-particle scattering efficiency is low. For particles larger than the wavelength, the single-particle scattering efficiency initially decreases with diameter and then fluctuates around a value of two as size increases. However, a larger particle always scatters more light than a smaller particle because particle cross-sectional area increases faster with diameter than does the decrease in single-particle scattering efficiency at any point on the scattering efficiency curve. The mass scattering efficiency (i.e., the scattering per mass concentration) peaks for particles that are about 0.5 μm to 0.8 μm in diameter. Smaller particles are much less efficient at scattering light, but the mass of particles increases with particle size faster than the increase in the amount of light they scatter.




Figure 4-33. Light-scattering efficiency factor (per cross-sectional area), Q , for a homogeneous sphere with an index of refraction of 1.50 as a function of the size parameter, $\alpha = \pi D/\lambda$.

Source: Penndorf (1958).

Use of the Mie equation to calculate light scattering or the light scattering efficiency of particles in the atmosphere is severely limited by the general lack of sufficiently detailed particle characterization data. At a minimum, size-resolved particle composition data (e.g., aerosol collected on an 8-stage impactor) are needed to permit meaningful Mie scattering calculations. The chemical composition provides clues to the appropriate particle density and index of refraction, while the size distribution is inferred by fitting a distribution function to the concentration for each stage. Assumptions are still necessary to address the particle component mixture characteristics of the aerosol. Resulting scattering calculations can be compared to directly measured particle extinction to assess the reasonableness of the Mie calculations.

Reported calculated dry scattering efficiencies for sulfates range from 1.2 to 5.6 m²/g. Sulfate scattering efficiencies have been reported to increase by a factor of two when the size distribution increased from 0.15 to 0.5 μm (McMurry et al., 1996). Calculated scattering efficiencies for carbon particles ranged from 0.9 to 8.1 m²/g. A scattering efficiency of 1.0 and 0.6 m²/g was reported for soil and coarse mass, respectively (U.S. Environmental Protection Agency, 1996a; Sisler and Malm, 2000).

Figure 4-34 gives the volume-specific light scattering efficiency in units of μm⁻¹ as a function of particle diameter. The light scattering coefficient is derived by multiplying the volume-specific light scattering efficiency factor by the volume concentration. The mass-specific light scattering efficiency can be obtained by dividing the values for the curves by the density of the PM.

Similar results have been produced in field nephelometer measurements of ambient particle scattering. A variety of nephelometer configurations, unrestricted or size-selective inlets, and the control of sample air temperature and relative humidity, permit the composite scattering properties of ambient aerosol to be directly observed (Day et al., 1997). When sample-controlled nephelometer data are combined with collocated particle speciation data, composite particle scattering efficiency values for ambient aerosol can be empirically derived (Malm et al., 2000).

The scattering efficiency for particles has been reported by White et al. (1994) for dry particles < 2.5 μm (2.4 and 2.5 m²/g) and coarse particles (0.34 to 0.45 m²/g). Other reported values for coarse particles include 0.4 and 0.6 m²/g (White and Macias, 1990; Trijonis et al., 1987). Nephelometer measurements for light scattering by coarse particles is underestimated




Figure 4-34. Volume-specific light-scattering efficiency as a function of particle diameter D_p . The calculations were performed for the indicated indices of refraction and a wavelength of 550 nm. For large particle diameters, the scattering efficiencies tend toward a value of $3/D_p$. Mass-specific light-scattering efficiencies (in units of m^2/g) can be obtained by dividing the values of the curves by the particle density (in units of g/cm^3).

Source: U.S. Environmental Protection Agency (1996a).

(White et al., 1994). Chow et al. (2002a) reported scattering efficiencies of 2 to 3 m^2/g but in some cases $> 5 m^2/g$ for dry particles $< 2.5 \mu m$.

4.3.2.2 Relative Humidity Effects on Particle Size and Light-Scattering Properties

The ability of some commonly occurring chemical components of atmospheric aerosol to absorb water from the vapor phase has a significant effect on particle light scattering. Hygroscopic particles, which typically include sulfuric acid, the various ammonium sulfate salts, ammonium nitrate, and sodium chloride, change size by the accumulation and loss of water as

they maintain equilibrium with the vapor phase as a function of changes in relative humidity. For some materials (e.g., sulfuric acid), the growth is continuous and reversible over the entire range of relative humidity. For other materials, water absorption begins abruptly for a dry particle at a specific relative humidity known as the deliquescent point (e.g., ~80% for ammonium sulfate) and continues as relative humidity increases. There is a hysteresis effect with these materials in that, once wet, the relative humidity can be reduced below the deliquescent point until crystallization occurs at a substantially lower relative humidity (e.g., ~30% for ammonium sulfate). Figure 4-35 shows the water vapor growth curve for ammonium sulfate.




Figure 4-35. Particle growth curve as a function of relative humidity (RH) showing deliquescent growth of ammonium sulfate [(NH₄)₂ SO₄] particles at the deliquescent point (A, about 80% RH), reversible hygroscopic growth of ammonium sulfate solution droplets at RH > 80%, and hysteresis (the droplet remains supersaturated as the RH decreases below 80%) until the crystallization point (B, about 38% RH) is reached.

Source: Adapted from National Research Council (1993) and Tang (1980).

The water growth behavior for hygroscopic materials commonly found in atmospheric aerosol in pure form or in some mixtures is generally well known as a result of laboratory measurements (Tang and Munkelwitz, 1994; Tang, 1997). Models that calculate water growth of mixtures from known solubility properties of many common water-soluble chemicals have long been available (Zdanovskii, 1948) and have been successfully applied to determine growth for particles with known composition (Saxena and Peterson, 1981; Pilinis et al., 1995; Saxena et al., 1993).

The water growth of individual ambient particles can be directly measured using a humidity-controlled tandem differential mobility analyzer (TDMA) (McMurry and Stolzenburg, 1989; Zhang et al., 1993). Inferences can be made about the mixtures of soluble and insoluble particle components by comparing TDMA-measured growth and size-resolved aerosol composition data with water growth model predictions (Pitchford and McMurry, 1994; Zhang et al., 1993; Saxena et al., 1995). A practical limitation of TDMA measurements in studying aerosol optical properties is that particles $> 0.5 \mu\text{m}$ are not well measured by this approach.

Accounting for water growth of atmospheric aerosols is important in determining visibility, because particles containing hygroscopic or deliquescent materials change size, index of refraction, and, hence, scattering efficiency, with changing relative humidity. The nonlinear nature of particle growth curves for hygroscopic aerosols means that substantial light scattering changes result from modest relative humidity changes under humid conditions (relative humidity $> 90\%$). The magnitude of the water growth effect on light scattering for ambient aerosols can be directly measured with humidity-controlled nephelometer measurements (Day et al., 1997). Measurements of water growth effects on light scattering are compared to the results of water growth and Mie scattering models applied to size-resolved composition data using various mixture assumptions to infer average mixture and other aerosol characteristics (Malm et al., 2000a,b).

While the importance of inorganic hygroscopic particles is well understood, the role of organic compounds in particle water growth has been the subject of recent investigations. In their interpretation of TDMA and particle composition data from two locations, Saxena et al. (1995) made the case that organic components of the aerosol enhanced water absorption by particles at a remote desert location and retarded water absorption at an urban location. They

speculated that the latter might be due to hydrophobic organic material coatings on inorganic hygroscopic particles.

Whereas some of the thousands of organic compounds that are in atmospheric aerosols are known to be hygroscopic and a significant fraction of the organic aerosol material is known to be water soluble, there is a lack of water absorption data for most organic compounds. The incomplete water solubility data, combined with incomplete data on the abundance of the numerous organic compounds in ambient aerosols, means that organic water growth model calculations are not a reasonable approach for assessing the importance of water growth by organic aerosol components in the atmosphere. To overcome this constraint, Saxena et al. (1995) compared organic concentration to the difference between total aerosol water measured by TDMA and model-estimated water for the inorganic hygroscopic aerosol components. In contrast, Pitchford and McMurry (1994) used the same remote location data set and showed that, on 6 of the 8 sampling days, water uptake by the sulfates and nitrates could account for all of the measured water absorption.

Swietlicki et al. (1999) made TDMA measurements in northern England and found that growth takes place in two modes, one mode being less hygroscopic than the other. They concluded that growth could be attributed to the inorganic content of the aerosol. Cocker et al. (2001) measured hygroscopic properties of Pasadena, CA aerosol and concluded that growth factors increased when forest fires were present. McDow et al. (1995) measured water uptake by diesel soot, automobile exhaust, and wood smoke particles. They found all three emission types absorbed water: the wood smoke sample weight increased by about 10% as sample relative humidity was increased; whereas diesel soot sample weight increased by only 2 to 3%. Chughtai et al. (1999) examined the hydration characteristics of a number of anthropogenic and natural organic materials. They found surface water adsorption increased with age and surface oxidation. Hemming and Seinfeld (2001) evaluated the relative hygroscopicity of different organics, the differences in the amount of water taken up by mixtures, and the individual components of the mixtures in their pure state using the UNIFAC. They found that mixtures take up less water than the individual components in the pure state. The relative hygroscopicity of atmospheric organics was diacids > monoacids > alcohols > carbonyls. Analysis of humidity-controlled and size-resolved chemistry data from the Great Smoky Mountains and Grand Canyon National Parks (Malm et al., 1997; Malm and Kreidenweis, 1997; Malm et al., 2000a) showed

that ambient organic aerosol are, at most, weakly hygroscopic to within the measurement uncertainty and modeling assumptions.

A more detailed discussion of the effects of relative humidity on the size distribution of ambient particles appears in Chapter 2 of this document.

4.3.3 Relationships Between Particles and Visibility

Visibility, referring to the appearance of scenic elements in an observer's line of sight, depends on more than the optical characteristics of the atmosphere. Numerous scene and lighting characteristics are important to this broad definition of visibility. However, under a variety of viewing conditions, visibility reduction or haziness is directly related to the extinction coefficient.

Light extinction, the sum of the light scattered and absorbed by particles and gases, is frequently used to estimate the effect of air pollution on visibility. Light extinction is usually quantified using the light extinction coefficient, i.e., the sum of the light scattering and absorption coefficients for gases and particles (see Section 4.3.2.1).

The influence of particles on visibility degradation is dependent on the particle size, composition, and solubility (Pryor and Steyn, 1994). Fine particles (particles with mass mean diameters $\leq 2.5 \mu\text{m}$) scatter more light than coarse particles. Fine particle species include sulfates (assumed to be ammonium sulfate), nitrates (assumed to be ammonium nitrate), organics, light-absorbing carbon, and soil (Malm et al., 1994). Of the fine particle species, sulfates and nitrates are the most hygroscopic and require the use of a relative humidity adjustment factor. The effect of particle light extinction can be determined by totaling the scattering and absorption of light by multiplying the mass-specific efficiency values and the mass concentration for each of the particle species. The effect of relative humidity and the relative humidity adjustment factors are discussed in Section 4.3.2.2.

Visibility is measured by human observation, the light extinction coefficient (light scattering and absorption by particles and gases), and parameters related to the light extinction coefficient (visual range, deciview) and fine particle mass concentrations. The equation for the light extinction coefficient, light extinction by particles is stated in the discussion of the IMPROVE Program.

The visual range method of visibility measurement, commonly taken to be the greatest distance that a large dark object (e.g., a mountain in shadow) can be seen against the background sky (Middleton, 1952), was developed for and continues to function well as an aid in military operations and transportation safety. Visual range is inversely proportional to the light extinction. Visual range can be calculated from a point measurement of light, assuming that the atmosphere and the illumination over the sight path is uniform and the threshold contrast is 2% of the extinction coefficient

$$\text{Visual Range} = K/b_{\text{ext}} \quad (4-8)$$

where visual range is in kilometers, b_{ext} is in km^{-1} , and a threshold contrast of 2% is assumed. If b_{ext} is in Mm^{-1} , the Koschmieder constant becomes 3,912.

An index of haziness, expressed in deciview (dv) units, is also very simply related to the light extinction coefficient (Pitchford and Malm, 1994).

$$\text{Haziness (dv)} = 10 \ln(b_{\text{ext}} / 10 \text{ Mm}^{-1}) \quad (4-9)$$

An important characteristic of this visibility index is that it is more nearly linearly related to perceived changes in haze level than either visual range or light extinction. A change of 1 or 2 dv in uniform haze under many viewing conditions will be seen as a small but noticeable change in the appearance of a scene, regardless of the initial haze condition.

Figure 4-36 illustrates the relationship of light extinction in Mm^{-1} , deciview index, and visual range in kilometers. Although the deciview is related to extinction, it is scaled in such a way that is perceptually correct (Fox et al., 1999).

The amount of light that an aerosol of a given mass concentration scatters depends on the particle size distribution. Several studies and reports (Chow et al., 2002a; Samuels et al., 1973; Waggoner and Weiss, 1980; Waggoner et al., 1981) indicate that the mass concentration of particles of $< 3 \mu\text{m}$ in diameter correlate well with light scattering and visibility when the measurements are made under dry conditions. The question of whether the relationship between




Figure 4-36. Comparison of extinction (Mm^{-1}) and visual range (km).

Source: Fox et al. (1999).

light scattering and fine particle mass varies too greatly between geographical locations to allow the use of fine particle mass as a metric for visibility effects has been considered. McMurry (2000), citing a study by Charlson et al. (1968), stated that the ratio of dry scattering coefficient to the dry fine particle mass concentration from different geographical locations did not vary a great deal. However, the use of either fine particle mass or the individual major components to determine particle-related light scattering requires measurements, estimates, or assumptions about concurrent ambient relative humidity in order to accurately estimate the relationship between ambient fine particles and light extinction.

Figure 4-37 shows the relationship between fine particle mass and calculated light extinction. The figure, as reported in Chow et al (2002b), was generated using data reported by Samuels et al. (1973). According to Samuels et al. (1973), there was a direct correlation between particle mass concentration, light scattering, and visibility. However, there were large standard errors in the scattering coefficient.

Most routine aerosol monitoring programs and many special study visibility characterization programs were designed to measure the major aerosol components (Malm et al., 1994; Tombach and Thurston, 1994; Watson et al., 1990); they were not designed to determine the microphysical and chemical characteristics of these species. However, the inherent limitations of estimating aerosol optical properties from bulk aerosol measurements have been addressed, at least in part, by a number of authors. For instance, Ouimette and Flagan (1982) have shown using basic theoretical considerations that if an aerosol is mixed externally (i.e., separate particles contain the major aerosol components) or, if internally mixed, the index of




Figure 4-37. Proportionality of observed daytime haziness to fine particle mass concentration in Los Angeles. Visual ranges are 8-h averages of hourly human observations, plotted as extinction according to Koschmieder formula. Mass concentrations are from 8-h samples collected behind a cyclone with 3- μm cut point. Relative humidities were $\leq 70\%$.

Source: Chow et al. (2002b).

refraction is not a function of composition or size and the aerosol density is independent of volume, then

$$b_{sp} = \sum \alpha_i m_i \quad (4-10)$$

where α_i is the specific mass scattering efficiency and m_i is the mass of the individual aerosol species.

Sloane (1983, 1984, 1986), Sloane and Wolff (1985), and more recently, Lowenthal et al. (1995) and Malm and Kreidenweiss (1997) have shown that differences in estimated specific scattering between external and internal model assumptions are usually less than about 10%. In the absence of detailed microphysical and chemical information of ambient particles, the

above studies demonstrate that a reasonable estimate of aerosol extinction can be achieved by assuming each species is externally mixed.

The latest IMPROVE network report (Malm, 2000) included calculated aerosol light extinction for each of the five major fine fraction particle ($PM_{2.5}$) components, coarse fraction mass ($PM_{10-2.5}$), and Rayleigh scattering by gases and summed them for an estimate of total light extinction in Mm^{-1} using the following algorithm:

$$\begin{aligned}
 b_{ext} = & (3)f(RH) [SULFATE] \\
 & +(3)f(RH) [NITRATE] \\
 & +(1.4) [ORGANIC CARBON] \\
 & +(10) [LIGHT ABSORBING CARBON] \\
 & +(1) [SOIL] \\
 & +(0.6) [COARSE PM] \\
 & +10 \text{ (for Rayleigh scattering by gases)}
 \end{aligned}
 \tag{4-11}$$

where each PM term is the product of a constant dry extinction efficiency for that species, the mass concentration of the species, and, for sulfate and nitrate, an adjustment factor that is a function of relative humidity to account for their hygroscopic behavior. The relative humidity adjustment term for sulfate and nitrate, shown in Figure 4-38, is based upon the ammonium sulfate growth curve, shown in Figure 4-28, smoothed between the upper and lower curves of the hysteresis loop for the relative humidity range of 30 to 80%. The extinction efficiencies for soil and coarse mass used in this algorithm are taken from a literature review by Trijonis et al. (1987). The extinction efficiency for light absorbing (elemental) carbon of $10 \text{ m}^2/\text{g}$ is consistent with the value reported by Moosmüller et al. (1998) corresponding to $\lambda = 0.53$ in the middle of the visible light spectrum. The dry extinction efficiencies of $3 \text{ m}^2/\text{g}$ for sulfate and nitrate species and $4 \text{ m}^2/\text{g}$ for organic species are based on literature reviews by Trijonis et al. (1991) and by White (1991). Trijonis' best estimate for sulfates is $2.5 \text{ m}^2/\text{g}$ with an uncertainty of a factor of 2, while White's average low and high estimates for the rural West are 3.0 and $3.7 \text{ m}^2/\text{g}$. For organics, Trijonis estimated a dry extinction efficiency of $3.75 \text{ m}^2/\text{g}$ with an uncertainty of a factor of 2, and White's range for the rural West is 1.8 to $4.1 \text{ m}^2/\text{g}$. Malm et al.




Figure 4-38. Relative humidity adjustment factor, $f(RH)$, for ammonium sulfate as a function of relative humidity.

Source: Malm (2000).

(1996) and Malm (2000) used this algorithm to successfully reconstruct light scattering at a total of eleven IMPROVE monitoring sites.

Malm (2000) used additional sophisticated aerosol size, composition, and microphysical data from a special study at the Great Smoky Mountains National Park to compare the performance of a number of models for calculating light extinction. He found that the simplest approach adequately predicted for periods of low light scattering but under-predicted by about 30% during periods of high sulfate concentration. The greatest improvement over the simple model was obtained by including the degree of sulfate ammoniation in the model. This produced better estimates of the extinction coefficient over the entire range. Table 4-15 lists various visibility metrics and methods for visibility measurement.

TABLE 4-15. VISIBILITY MEASUREMENT TECHNIQUES

Visibility Metric and Method	Measurement Principle
Visual Range	Human observation of prevailing visibility. Targets are selected at known distances from an observer. Nighttime targets require lights and may differ from daytime targets. Each hour, the observer records the distance (i.e., visual range) corresponding to the furthest target that is visible. This method provides the longest history of visibility measurements in the United States, as it was used at most U.S. airports from 1948 to 1995.
Light extinction (b_{ext})	Directly measures the radiance of a constant light source (transmitter) after the light travels over a finite atmospheric path.
Long-path transmissometer	The transmittance of the path is calculated by dividing the measured radiance by the calibrated initial intensity of the light source. The average extinction of the path is calculated from the transmittance and length of the path.
Short-path transmissometer	Starting in early 1990s, many airports replaced human observations with automated sensors (i.e., the Automated Surface Observing System [ASOS], Automated Weather Observing System [AWOS]) to measure 1-min-average light extinction. The visibility sensor measures forward scattering using a xenon flash lamp source. Instruments can provide measurements up to 48 km, but they are not recorded beyond 16 km.
<u>Contrast transmittance:</u> Teleradiometer	A telescope is focused on a distant target and the background. Changes in radiance are measured by photodiode detectors. Measurements can be made at different wavelengths (e.g., 450, 550, and 630 nm) using narrow band filters. Measures effects of particles of all sizes. Sensitive to variations in inherent contrast (e.g., bright or dark cloud behind the target) and nonuniform illumination conditions. Because path radiance depends on how the atmosphere is illuminated, contrast transmittance represents human perception better than air quality.
Photographs and time-lapse film	A densitometer measures light transmitted through different portions of a color slide. The film's light-response function (the gamma curve) determines target-sky contrast. Computerized photographic simulations can change contrast transmittance for different meteorological and atmospheric conditions, and can be used to judge how people react to these changes.
Particle scattering ($b_{scat,p}$): Integrating nephelometer	Air is drawn into a nephelometer chamber that is illuminated with white or filtered (typically 500-550 nm) light. Light is detected at 90° to the direction of illumination to measure the amount scattered out of the light path. Chamber dimensions limit the integrated arc to ~10-170° instead of a full 0-180°, which results in the underestimation of some forward scattering from coarse particles. Nephelometers are calibrated with gases of known indices of refraction. Particles (especially hygroscopic and volatile species) may be modified as they pass through the chamber, which is inadvertently heated by the illumination source.

TABLE 4-15 (cont'd). VISIBILITY MEASUREMENT TECHNIQUES

Visibility Metric and Method	Measurement Principle
<p><u>Particle absorption ($b_{abs,p}$):</u> Aethalometer or particle soot absorption photometer</p>	<p>Particles are collected on a quartz-fiber filter tape. The change in transmittance at selected wavelengths (e.g., 880 nm) across the filter before and after sampling is measured or compared with the reference area. When the filter spot darkens, a new portion of the tape is moved into the sampling position. Assumes a constant relationship between black carbon (BC) mass and quartz filter transmittance of $\sim 19\text{m}^2/\text{g}$. Assumes a relationship of $10\text{ m}^2/\text{g}$ between BC absorption and concentration.</p>
<p>Photoacoustic spectroscopy</p>	<p>Particles absorb energy from a modulated laser ($\sim 514.5\text{ nm}$) and transfer heat to the surrounding air. Expansion of the heated gas produces sound waves (acoustic signals) that are proportional to the amount of absorbed energy. These are detected by a high-sensitivity microphone. Absorption appears to vary with illumination wavelength.</p>
<p>Filter transmittance, reflectance</p>	<p>Uses densitometer, integrating plate, or integrating sphere spectrophotometer to measure how much light is transmitted or reflected. The difference in the logarithms of light transmission through the filter before and after sampling is proportional to the particle deposit. Light transmission of particles collected on a filter may overestimate light absorption because some of the incident light is scattered within the filter and by other particles in the deposit. There are no absolute calibration standards for densitometry, integrating plate, or sphere methods.</p>
<p>Suspension of insoluble elemental carbon</p>	<p>Particles collected on a Nuclepore polycarbonate-membrane filter are extracted in 30% isopropanol/70% distilled deionized water to form a suspension of insoluble EC particles. Using a spectrophotometer, light transmission (e.g., 400-650 nm, peaking at 575 nm) is measured through the liquid extract.</p>
<p>Component extinction (b_{ext} sum of $b_{scat,p}$, $b_{scat,g}$, $b_{abs,p}$, and $b_{abs,g}$)</p>	<p>The sum of clean air scattering estimated from temperature and pressure, NO_2 absorption estimated from NO_2 concentrations, particle scattering measured by nephelometer, and particle absorption ($b_{abs,p}$) measured by one of the $b_{abs,p}$ methods. Measurements are at a single location rather than along a sight path.</p>
<p><u>Chemical extinction (b_{ext}):</u> Filter measurements for SO_4^{2-}, NO_3^-, organics, EC, fine soil, and coarse mass, plus clear air scattering</p>	<p>Six aerosol chemical components are used to calculate chemical extinction.</p>

4-172

Adapted from: Watson (2002).

4.3.4 Photographic Modeling of Visibility Impairment

None of the visibility indices communicate visibility associated with various aerosol conditions as well as directly seeing their effects on a scene. Photographic modeling for the representation of haze can be useful in portraying changes in visibility specifically due to changes in air pollutant concentrations. Photographic modeling holds constant the effects of sun angle, cloud cover, and relative humidity and is a cost-effective method of evaluating various air quality scenarios. Photographic modeling is difficult to do with actual photographs because of the range of possible conditions in the same scene over multiple days; and, over time, photographs can be expensive to produce. Another limitation in using photographic models for representation of haze is that haze is assumed to be uniformly distributed throughout the scene and selected conditions are idealized, so the full range of conditions that occur in a scene are not represented.

Eldering et al. (1996) proposed the use of a model that uses simulated photographs from satellite and topographic images to evaluate the effect of atmospheric aerosols and gases on visibility. Use of this model requires ground-based photography and data concerning the size distribution and chemical composition of atmospheric aerosols, NO₂ concentration, temperature, and relative humidity for a clear day. Light extinction and sky color are then calculated based on differences in aerosol size distribution, NO₂ concentration, temperature, and relative humidity. The images created represent natural landscape elements.

Molenaar et al. (1994) provides a discussion of existing visual air quality simulation methods based on techniques under development for the past 20 years. A photograph taken on a very clean, cloud-free day serves as the base image. The photograph is taken during the season and at the same time of day as the scene to be modeled. The light extinction represented by the scene is derived from aerosol and optical data associated with the day the image was taken, or it is estimated from contrast measurements of features in the image. The image is then digitized to assign an optical density to each picture element (pixel) for the wavelength bands of interest. A detailed topographic map and an interactive image-processing display system is used to determine the specific distance, elevation angle, and azimuth angle for each element in the picture with respect to the observer's position.

Various models are employed to allow the presentation of different air quality scenarios. The output from atmospheric aerosol models (e.g., extinction, scattering coefficients, single-

scattering albedo, and scattering phase matrix) is incorporated into radiative transfer models to calculate the changes in radiant energy (path radiance, image radiance, sky radiance, terrain radiance) caused by scattering and absorption by gases and particles as it passes through the atmosphere. Atmospheric aerosol models are also used to model the effect of relative humidity on the visual air quality (Molenar et al., 1994).

Molenar et al. (1994) has developed a system call WinHaze that permits the viewing of computer-generated uniform hazes superimposed on digitized scenic photographs of both remote and urban scenes. The program simulates changes in visual air quality imagery from user-specified changes in optical parameters (e.g., σ_{ext} , visual range, or deciview values) or aerosol species concentrations. WinHaze includes imaging for various Class I national parks and wilderness areas and for Boston, MA; Dallas, TX; Denver, CO; Fort Collins, CO; Phoenix, AZ; and Tucson, AZ. The computer software is available through the IMPROVE website (<http://vista.cira.colostate.edu/improve/>).

4.3.5 Visibility Monitoring Methods and Networks

Visibility monitoring studies measure the properties of the atmosphere either at the sampler inlets (point measurements), as is the case with air quality measurements, or by determining the optical properties of a sight path through the atmosphere (path measurements). Instrumental methods for measuring visibility are generally of three types: (1) direct measurement of light extinction of a sight path using a transmissometer, (2) measurement of light scattering at one location using an integrating nephelometer, and (3) measurement of ambient aerosol mass concentration and composition (Mathai, 1995).

The largest instrumental visibility monitoring network in the United States is the Automated Surface Observing System (ASOS), commissioned by the National Weather Service, Federal Aviation Administration, and Department of Defense for use at more than 900 airports. The system is designed to objectively measure the clarity of the air versus the more subjective evaluations of human observations. The system provides real-time data for airport visibility.

The visibility sensor, instead of measuring how far one can see, measures the clarity of the air using a forward-scatter visibility meter. The clarity is then converted to what would be perceived by the human eye using a value called Sensor Equivalent Visibility (SEV). Values derived from the sensor are not affected by terrain, location, buildings, trees, lights, or cloud

layers near the surface. The amount of moisture, dust, snow, rain, and particles in the light beam will affect the amount of light scattered. The sensor transmits 1-minute values based on rolling 10-minute periods. Hourly visibility range data are available only at a quantized resolution of 18 binned ranges with a visual range of up to 10 miles. The value provides a generally accurate and representative visibility measurement within a 2 to 3 mile radius of the site. The forward scatter meter was found to correlate fairly well with extinction coefficient measurements from the Optec Transmissometer (National Weather Service, 1998).

Visibility data from the ASOS network is reported in terms of visual range in increments of 1/4 to 1 statute mile. Visual range conditions exceeding 10 miles are truncated to 10 miles for real-time reporting purposes. Data is not extensively archived at ASOS locations; however, researchers are able to download the raw data directly from certain sites. In addition, since 1998, the raw visibility data (including light extinction measurements corresponding to visual ranges exceeding 10 miles) have been archived for a number of sites. Visual range measurements beyond 10 miles may be used to derive PM concentrations except in clean environments. These data are available from the National Climatic Data Center.

The largest monitoring network that includes both visibility and aerosol conditions is the IMPROVE network. This network was formed in 1987 as a collaborative effort between Federal, regional, and state organizations responsible for protection of visibility in the 156 mandatory Class I Federal areas (national parks and wilderness areas) and other areas of interest to land management agencies, states, tribes, and other organizations (National Park Service, 1998; U.S. Environmental Protection Agency, 1995b, 1996a, 1999b; Eldred et al., 1997; Perry et al., 1997; Sisler and Malm, 2000). It is predominantly a rural-based network with more than 140 sites across the country. The primary monitoring objectives of the IMPROVE program are to document current visibility conditions in the mandatory Class I areas, to identify anthropogenic chemical species and emission sources of visibility impairment through the collection of speciated PM_{2.5} data, and to document long-term trends for assessing progress towards elimination of anthropogenic visibility impairment. The IMPROVE network has also been involved in visibility related research, including the advancement of visibility monitoring instrumentation and analysis techniques and visibility monitoring and source attribution field studies (National Park Service, 1998; Evans and Pitchford, 1991).

Visibility monitoring under the IMPROVE network can be divided into three categories: aerosol, optical, and scene. Twenty-four hour PM_{2.5} and PM₁₀ aerosol samples are collected by filters at least every third day. The PM_{2.5} samples are analyzed to determine the mass concentration of the major particulate constituents (sulfates, nitrates, organic carbon compounds, elemental carbon, chlorides, and crustal elements) and for elements that indicate sources of visibility-impairing particles (trace elements and ions). Optical monitoring provides a direct measurement of light scattering and absorption. Color photographic imaging documents the appearance of the scene under a variety of air quality and illumination conditions (U.S. Environmental Protection Agency, 1999b). It is anticipated that all data generated by the IMPROVE network will be added to the AIRS database.

The EPA has deployed a new national monitoring network (Federal Reference Method Monitoring network) designed to assess PM_{2.5} concentrations and composition. There are over 1,000 monitoring sites in operation and many sites report data to the AIRS. Analyses of these data are expected to provide a more complete understanding of visibility conditions, in particular urban visibility, across the country. The PM_{2.5} monitoring effort has been coordinated with visibility monitoring efforts currently in place to maximize the benefits of all of the monitoring programs (U.S. Environmental Protection Agency, 1997b, 2000b, 2001a).

The Northeast States for Coordinated Air Use Management (NESCAUM) has established a real-time visibility monitoring network (CAMNET) using digital photographic imaging. There is currently digital photographic imaging for five urban locations (Boston, MA; Burlington, VT; Hartford, CT; Newark, NJ; and New York City, NY), and two rural locations (Acadia National Park, ME and Mt. Washington, NH). The visibility images are updated every 15 minutes. Near real-time air pollution and meteorological data are updated every hour. Archived images will be available for studies of the visual effects of PM air pollution in the Northeast. CAMNET may be accessed at www.hazecam.net (Northeast States for Coordinated Air Use Management, 2002; Leslie, 2001).

The Midwest Regional Planning Organization, in cooperation with a number of other groups, has also developed a real-time visibility camera network (hazecam). The camera network includes several urban (Chicago, IL; Indianapolis, IN; and Cincinnati, OH) and rural locations (Seney NWR, MI; Mayville, WI; and Isle Royale National Park, MI/Grand Portage, MN). The Midwest hazecam can be found at <http://www.mwhazecam.net>.

4.3.6 Visibility Impairment: Trends and Current Conditions

In the United States, visibility impairment is caused by particles primarily composed of sulfates, nitrates, organic compounds, carbon soot, and crustal dust. Visibility is best in Alaska and the western Great Basin. Moderate levels of light extinction are common on the Pacific Coast, including the western slopes of the Sierra Nevadas in California and the Cascade Range in Oregon and Washington. Visibility is most impaired in the areas encompassing and adjacent to the Ohio and Tennessee River Valleys. Visibility gradually improves along the Atlantic seaboard northeast of New York City (Watson, 2002). Natural visibility in the East and West is about 75 to 150 km (45 to 90 miles) and 200 to 300 km (120 to 180 miles) (U.S. Environmental Protection Agency, 2001b).

4.3.6.1 Trends in Visibility Impairment

Trends in visibility impairment or haziness often are used as indicators of trends in fine particles mass. Observations of visual range, obtained by the National Weather Service and available through the National Climatic Data Center of the National Oceanic and Atmospheric Administration provide one of the few truly long-term, daily records of impairment related to air pollution. After some manipulation including correction for relative humidity effects, the visual range data can be used as an indicator of fine mode particle pollution. The data reduction process and analyses of resulting trends have been reported by Schichtel et al. (2001), Husar et al. (1994), Husar and Wilson (1993), and Husar et al. (1981).

There are many statistical approaches to estimating trends. These approaches include simple correlation and regression analyses, time-series analyses, and methods based on nonparametric statistics. A discussion and comparison of the methods for the detection of linear trends is provided in Hess et al. (2001). Schimek (1981) provides a discussion of nonlinear trends. In its annual air quality trends report, the EPA characterized trends using a nonparametric regression analysis approach commonly referred to as the Theil test (U.S. Environmental Protection Agency, 1998; Hollander and Wolfe, 1973).

Regional Trends and Class I Areas

The two largest contributors to visibility impairment are sulfates and carbon-based particles. In the East, sulfates are responsible for 60 to 86% of the visibility impairment. The

sulfate contribution decreases further west but is still responsible for between 25 to 50% of the visibility impairment. Carbon-based particles are responsible for 10 to 18% of the visibility impairment in the East and 25 to 40% in the West. Nitrates account for only 7 to 16% of the light extinction in the East but are responsible for between 5 to 45% of the light extinction in the West. Crustal material can be a major contributor in the West, accounting for 5 to 25% of the light extinction. Elemental carbon also is contributor to light extinction, but to a lesser degree (U.S. Environmental Protection Agency, 2001a,b).

The EPA designated five regional groupings as part of the regional haze program. The regions are Northeast (Mid-Atlantic/Northeast Visibility Union), Southeast (Visibility Improvement State and Tribal Association of the Southeast), Central (Central States Regional and Air Partnership), Midwest (Midwest Regional Planning Organization), and West (Western Regional Air Partnership). The regional groupings serve as consensus organizations comprised of states, tribes, and federal agencies coordinating the implementation of the regional haze rule.

Using hourly prevailing daytime visibility data from human observations at weather stations, Schichtel et al. (2001) observed that haziness declined approximately 10% across the United States between 1980 and 1995. The decrease in haziness was highest in the southeastern United States with a 20% decrease in the 90th percentile light extinction and a 12% decrease in the 75th percentile over the 15-year period. There was a 17% decrease in the 90th percentile and a 9% decrease in the 75th percentile over the eastern United States. Over the eastern United States, haziness was greatest during the summer months. The greatest visibility impairment was adjacent to the Appalachian Mountains in Tennessee and the Carolinas (extinction coefficient of $> 0.2 \text{ km}^{-1}$ equivalent to 6 miles). During the cold season, elevated haze (extinction coefficient of $> 0.2 \text{ km}^{-1}$) was seen between the Great Lakes and the Ohio River Valley, over the gulf states between Texas and Florida, along the coast from North Carolina to New Jersey, and along the Pacific coast, particularly central and south California.

Visibility impairment or haziness in the southeastern United States from sulfate emissions is greatest in the humid summer months because of the ability of sulfate to absorb atmospheric water vapor. Summer haziness increased in the southeastern United States from the 1950s to 1980 along with increasing SO_2 emissions. A statistically significant increase in summer sulfate concentrations was noted in two Class I areas in the eastern United States (Shenandoah and the Great Smoky Mountains) from 1982 to 1992 (Eldred et al., 1993; Cahill et al., 1996). During

that time, the majority of the Southwest showed decreasing sulfur concentrations (Eldred et al., 1993; Eldred and Cahill, 1994). Increasing summer sulfate concentrations were later shown at those two locations by Iyer et al. (2000).

Limited visibility measurements are available for the upper Midwest region (Illinois, Indiana, Michigan, Ohio, and Wisconsin and the tribal lands located in those states). The Midwest Regional Planning Organization conducted an initial assessment of the regional haze problem in this region using existing reports and available air quality data for four major urban areas (St. Louis, Chicago, Detroit, Cincinnati) and Class I areas. The “worst” and “best” visibility days occur throughout the year. Particulate sulfates were the major contributors to light extinction during the summer months, and nitrates dominated on the worst visibility days during the winter and fall in both urban and Class I areas. Organics also were significant contributors to light extinction in urban areas. Higher PM_{2.5} concentrations were correlated with poorer visibility in the southern portion of the region (Midwest Regional Planning Organization, 2001).

The EPA’s National Air Quality and Emission Trends Report summarized an estimate of the regional trends and current conditions in 35 Class I areas and one urban area (Washington, DC), using chemical concentrations data from the IMPROVE network (U.S. Environmental Protection Agency, 2001a). The visibility trends analysis is an aggregate of 10 eastern Class I areas and 26 western Class I areas. Trends were presented for annual average values for the clearest (“best”) 20% , middle (“typical”) 20%, and haziest (“worst”) 20% of the days monitored each year. The visibility trends, given in changes in deciview values, for the eastern and western sites are illustrated in Figures 4-39a and 4-39b. From the figures, it can be seen that the haziest days in the West are equivalent to the best days in the East. In the East, there was a 16% (1.5 deciview) improvement in haziness on the clearest days since 1992. Improvements in visibility were noted in the East for the haziest days. However, based on monitoring data for 1999, visibility remains significantly impaired, with a visual range of 23 km for the haziest days compared to a mean visual range of 84 km for the clearest days. A 25% and a 14% improvement in visibility impairment were seen for the clearest and middle days in the West, respectively; whereas conditions for the haziest days degraded by 18.5% (1.7 deciviews) between 1997 and 1999, but were relatively unchanged compared to 1990 conditions (U.S. Environmental Protection Agency, 2001a).




Figure 4-39a. Aggregate visibility trends (in deciviews) for 10 eastern Class 1 areas.




Figure 4-39b. Aggregate visibility trends (in deciviews) for 26 western Class 1 areas.

Source: U.S. Environmental Protection Agency (2001a).

Figures 4-40a and 4-40b illustrate aggregate trends in aerosol light extinction, including trends by major aerosol component for the haziest 20% of days monitored for the 10 eastern Class I areas from 1992 to 1999 and the haziest 20% of days monitored for the 26 western Class I areas from 1990 to 1999. The National Ambient Air Trends Report also includes a number of maps characterizing aerosol light extinction and key components at 36 IMPROVE sites (all rural except Washington, DC) for 1997 through 1999 (U.S. Environmental Protection Agency, 2001a).

Urban Trends.

Most of the available visibility measurements, with the exception of the airport visual range measurements, focus on the impact of pollution on visibility in scenic vistas and regional haze (Class I areas). Many urban metropolitan areas monitor daily visibility conditions. These findings are generally not available in a published form and may not distinguish between pollution- and weather-related effects. Although the EPA Regional Haze Rule addresses visibility impairment in Class I areas and calls for states to establish goals for improving visibility in these areas and to develop long-term strategies for reducing emissions of air pollutants that cause visibility impairment, the steps states take to implement the regulation will also improve visibility and health in broad areas across the country.

Kleeman et al. (2001), citing previously published studies, provided an historical description of visibility conditions in Southern California from the early 1930s. Based on airport observation data for 1932 to 1949, visibility conditions began to decrease in Los Angeles with the advent of industrialization and population growth. Visibility conditions were worse during the 1940s than the 1930s, with the lowest visibility conditions occurring between 1944 and 1947. During this period, there was nearly a complete loss of extremely good visibility days. Between 1943 and 1947, the number of extremely good visibility days during the summer season dropped from 21% to 0.2%. Between 1950 and 1961, deteriorating visibility conditions extended eastward from Los Angeles along the corridor adjacent to the foothills of the San Gabriel and San Bernardino Mountains. The visual range in the areas nearest Los Angeles was ≤ 3 miles for more than 140 days per year when the relative humidity was $< 70\%$. Further east of Los Angeles, past Ontario and San Bernardino, the visibility was ≤ 3 miles for 110 days per year during the same time period. Improvements in visibility conditions have been made since the




Figure 4-40. (a) Eastern class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component; (b) Western class I area aggregate trends in aerosol light extinction on the 20% haziest days, including trends by major aerosol component.

Source: U.S. Environmental Protection Agency (2001a).

1970s. The greatest improvements occurred in the western Los Angeles Basin. In Ontario, the average number of days per year when the visual range was > 10 miles was 99 between 1976 and 1978 and increased to 113 between 1988 and 1990 (Kleeman et al., 2001).

In contrast to the Los Angeles Basin and Ontario area, Denver and surrounding areas have experienced high pollution episodes since the 1970s. Climate changes contributed to the high pollution episodes by trapping cooler air under a cover of warm air, causing the pollution to remain stagnant over the area and producing a brown cloud comprising a variety of pollutants including nitrogen and sulfur oxides, as well as grit and dust. Debate over the cause of the increasing pollution in the Denver area and the controversy over converting coal-fired power plants to a cleaner natural gas system led to the initiation of the 1973 Denver Air Pollution Study, 1978 Denver Winter Haze Study, 1987-1988 Metro Denver Brown Cloud Study, and the 1993 Denver Brown Cloud Modeling Study. In 1990, in an effort to improve air quality, Denver adopted a visibility standard of 0.076 km^{-1} (units of atmospheric extinction per kilometer; 20.1 deciviews) averaged over 4 hours. While this is a step towards reducing air pollution, the Denver region still exceeds the visibility standard 50 to 80 times per year (Lloyd, 2002).

During the fall and winter of 1988 to 1990, a major air quality study was conducted in Phoenix, AZ to address degrading visibility conditions in Phoenix and other urban areas. The objectives were to (1) develop a data base of visibility, air quality, and meteorological measurements; (2) establish quantitative relationships between light extinction and emission sources; and (3) evaluate measurement systems for short-term and long-term monitoring in Phoenix. The major contributors to light extinction in Phoenix were residual wood burning, primary mobile source emissions, and secondary ammonium nitrate (Chow et al., 1990).

The Arizona Department of Environmental Quality has conducted optical measurements of visibility in Tucson since 1993 and in Phoenix since 1994. The measurements are divided into the mean of the “dirtiest” 20% of all hours, the mean of all hours, and the mean of the “cleanest” 20% of all hours for the entire day and for the 5:00 to 11:00 a.m. period. Figures 4-41a,b represent the trends in visibility conditions from 1993 to 2001 for Tucson and from 1994 to 2001 for Phoenix. Visibility on the dirtiest days in the Phoenix metropolitan area has not changed since visibility monitoring started; but, visibility on the best days has significantly degraded. There is a seasonally related effect on visibility in Phoenix: the dirtiest 20% of all hourly light extinction and the mean of all hours are more pronounced during the winter and fall months




Figure 4-41a. Light extinction trends in Tucson, Arizona from 1993 to 2002.

Source: Arizona Department of Environmental Quality (2002).




Figure 4-41b. Light extinction trends in Phoenix, Arizona from 1994 to 2001.

Source: Arizona Department of Environmental Quality (2002).

(Arizona Department of Environmental Quality, 2002). Daytime visibility is worst during the morning. Samples taken from 1994 to 1996 showed that organic and elemental carbon dominated visibility impairment during the dirtiest and cleanest days, whereas ammonium nitrate was an important contributor during the 20% dirtiest days (Neuroth and Heisler, 2000).

The State of Virginia compared visibility trends at five locations both with and without pre-1948 airport visual range data. Average annual airport visual-range-data from Roanoke (1936 to 1998), DC Reagan National (1930 to 1998), Richmond (1942 to 1998), Lynchburg (1935 to 1998), and Elkins, WV (1936 to 1994) were used in the analysis. Trends in visibility conditions were dependent on the baseline year used for visibility measurements. When airport visual range data from 1948 are used as the baseline data for visibility conditions, the trend in visual range shows declining conditions for all sites. When pre-1948 data are included in the analysis, visibility conditions basically are unchanged or improved (except for Elkins). When 1948 data are used as the baseline for visibility trends measurements, visual range decreases at all locations. Average annual visual ranged varied between 7+ miles and 8+ miles for all sites (Virginia Climate Advisory, 2000).

4.3.6.2 Current Conditions

Current visibility conditions have been well-characterized for Class I areas using updated data from the IMPROVE network (U.S. Environmental Protection Agency, 2001a; Malm, 2000; IMPROVE, 1998). During recent decades, daytime visibility conditions at all major airports throughout the United States were recorded hourly by human observation. These data were used to determine current visibility conditions and visibility trends in the United States, as well as the spatial distribution of visibility conditions (Trijonis et al., 1991). The use of human observation is being replaced by an automated observation system, the Automated Surface Observing System (ASOS). More than 900 airports are currently commissioned. In addition, the EPA has deployed a new national monitoring network to assess $PM_{2.5}$ concentrations and composition.

More detailed information on visibility conditions for urban and suburban areas will become more widely available as data from the national $PM_{2.5}$ speciation monitoring network and the ASOS airport visibility network are further analyzed. Efforts are currently underway to develop a web-based system to allow the use of the high resolution ASOS data in air quality

monitoring and assessment programs. The objectives are to collect and quality control an archive of the ASOS visibility data, to deliver processed hourly visibility data to public and air quality communities, and to use the web-based system to support the acquisition and dissemination of visibility data (Falke, 2001).

4.3.7 Societal Impacts of Particulate Matter Visibility Effects

Society recognizes the need to impose remedies for repairing and preventing further anthropogenic pollutant-related effects on visibility conditions (Ely et al., 1991; Davidson et al., 2000a). Information about how the public perceives and values improvements in visibility comes from both economic studies and from local or state initiatives to adopt local visibility goals and strategies.

4.3.7.1 Economic Studies

Various methods have been utilized to help determine the economic valuation of changes in visibility. Where possible, direct economic valuation can be determined using marketplace cost estimates. Avoided-cost methods estimate the costs of pollution by using the expenditures that are made necessary by pollution damage. As an example, if ambient levels of PM result in increased frequency of building cleaning or repainting, the appropriately calculated increase in these costs is a reasonable estimate of true economic damage. Benefits associated with reductions in the pollution levels are then represented by the avoided costs of these damages.

Estimating the benefits of clear skies is a more difficult and less precise exercise, because, although the public values aesthetic views, they are not directly bought and sold in the marketplace. However, there are several methods available to economists to estimate the economic impact of these kinds of changes in environmental conditions (Freeman, 1993). These methods include hedonic valuation or pricing, contingent valuation and contingent choice, and travel cost (Johnson and Desvousges, 1997; Hanley and Spash, 1993). The primary methods used to date for the valuation of visibility have been the hedonic price and contingent valuation methods (Hanley and Spash, 1993). However, this is not an exact science, and there are still issues and limitations associated with each of these methods.

Hedonic pricing can be used to estimate economic valuations for environmental effects that have a direct effect on market values. It relies on the measurement of differentials in property values under various environmental quality conditions including air pollution and environmental amenities, such as aesthetic views. The hedonic method works by analyzing the way that market prices change with changes in environmental quality. Part of the economic costs imposed by PM-related reductions in visibility can be estimated by looking at the differences in sales price between otherwise identical houses that have different degrees of visibility impairment.

The contingent valuation method (CVM) is the most widely used method for estimating value changes in both visibility and ecosystem functions (Hanley and Spash, 1993; Chestnut, 1997; Watson and Chow, 1993). The CVM creates hypothetical markets for goods and services that have no market-determined price. As part of the evaluation, individuals are shown photographs with perceivable differences in visibility levels. Carefully structured surveys are administered to estimate the amount of compensation equivalent to a given change in environmental quality or, equivalently, how much an individual would be willing to pay (WTP) for improvements in environmental quality or willing to accept (WTA) existing conditions without further deterioration. There is an extensive scientific literature and body of practice on both this theory and technique; however, there are still concerns about the use of this technique for quantitative purposes.

The travel-cost method estimates can be used to estimate the value of recreational benefits of an ecosystem based on the environmental quality at the site. The travel-cost method uses information on actual behavior rather than responses to hypothetical scenarios. The time and travel expenses incurred to visit a site represents the price of access to the site. The willingness to pay to maintain the site is determined by the number of times the individual visits the site at different travel costs.

The effects of PM on visibility may differ widely between urban residential and recreational areas. Therefore, separate estimates are needed to account for impacts associated with changes in visibility in residential and recreational (Class I) areas. Chestnut and Dennis (1997) compared the findings of the more recent studies on the economic impact of changes in regional haze using the contingent valuation method in residential areas in several eastern cities and in Los Angeles and San Francisco and using the hedonic value method in Los Angeles and

San Francisco. The findings of the contingent valuation studies are shown in Table 4-16. Findings using the contingent and hedonic methods of valuation for Los Angeles and San Francisco are compared in Table 4-17.

Using the contingent valuation method, Chestnut and Rowe (1990) found that 83% of those individuals responding to a survey on visibility were willing to pay to improve visibility in the national parks. Survey participants were selected from California, Arizona, Missouri, New York, and Virginia. The national parks from three regions (California, southwestern United States, and southeastern United States) were considered in different versions of the survey. The survey included questions on past and future visitations to national parks, potential pollution effects from human activities outside of the park, three hypothetical visibility scenarios, socioeconomic characteristics, and various photographic presentations of visibility conditions within the parks. Higher responses were noted for residents residing in the state or region where the national park was located; responses for males and the elderly were generally lower and there was a direct correlation between household income and the response.

Using the results from the Chestnut and Rowe (1990) study, Chestnut and Dennis (1997) calculated an extinction coefficient of 85 for in-state residents and 50 for out-of-state residents. These extinction coefficients were suggested to represent an annual willingness to pay per household of \$15 and \$9, in 1994 dollars, for a 20% improvement in visual range.

4.3.7.2 Public Perception and Attitude Studies

An initiative in Denver to address public concerns about visibility impairment began with a series of visibility-related studies in the 1970s through the 1980s, leading to the adoption of a visibility standard for the city of Denver in 1990. This standard is based on a light extinction level of 0.076 km^{-1} , averaged over four daylight hours, reflecting the short-term nature of the perception of changes in visibility conditions. This standard is equivalent to a visual range of approximately 50 km. The approach used to develop this standard relied upon citizen judgments about acceptable and unacceptable levels of visual air quality (Ely et al., 1991). In Phoenix, a study was conducted between 1988 and 1990 to address degrading visibility conditions (Chow et al., 1990). This work led the Arizona Department of Environmental Quality to establish a Blue Sky Index, which focuses on days in which the visual range, averaged over six daylight hours, is 40 km or more. This target is based on a method very similar to that used in Denver for

**TABLE 4-16. RESIDENTIAL VISIBILITY CONTINGENT VALUATION
STUDY RESULTS**

City	Mean WTP (\$1994)	Starting-Ending Visual Range (VR) (miles)	Extinction Coefficient	WTP for 20% Change in VR (\$)	Reference
Atlanta	Unadj. \$44	17.6-20	346	\$63	McClelland et al. (1993)
Chicago	Partial \$28		222		
	Full \$20		159		
Chicago	-\$361	38233	416	\$76	Tolley et al. (1986)
	\$346	38247			
	\$430	38259			
Atlanta	-\$301	38327	469	\$86	
	\$289	38342			
	\$432	12-32			
Boston	-\$222	18-13	422	\$77	
	\$212	18-28			
	\$262	18-38			
Mobile	-\$240	38264	312	\$57	
	\$257	38279			
	\$302	38289			
Washington, DC	-\$356	15-10	635	\$116	
	\$366	15-25			
	\$465	15-35			
Cincinnati	-\$88	38233	120	\$22	
	\$87	38248			
	\$98	38258			
Miami	-\$152	38359	256	\$47	
	\$136	38370			
	\$160	38380			
Cincinnati	\$198	11.4-16.4	602	\$110	Rae (1983)

Source: Chestnut and Dennis (1997).

TABLE 4-17. RESIDENTIAL VISIBILITY VALUATION STUDY RESULTS FOR LOS ANGELES AND SAN FRANCISCO

City	Method	Mean WTP	Starting-Ending Visual Range (VR) (miles)	Extinction Coefficient	WTP for 20% Change in VR	Reference
Los Angeles	CVM	\$130	2-12	1191328	\$22	Brookshire et al. (1979)
		\$333	2-28		\$242	
Los Angeles	Property Value	\$183	12-28	1191328	\$242	Trijonis et al. (1985)
			18.6-16.3		\$245-647	
San Francisco	CVM	\$211	16.3-18.3	1191328	\$242	Loehman et al. (1985)
		\$124			\$496-552	
San Francisco	Property Value			1191328	\$496-552	Trijonis et al. (1985)

Adapted from: Chestnut and Dennis (1997).

obtaining citizens' judgments as to acceptable levels of visual air quality (BBC Research and Consulting, 2003). While in practice these standard target values are exceeded many times per year in these areas, they reflect a reasonable degree of consistency in the outcome of the approach used to characterize the value that citizens in these two urban areas place on visual air quality.

Studies conducted in other locations have resulted in similar visibility threshold determinations, convergent on a minimal visual range of 40 to 60 km. For example, using a methodology similar to that used by Ely et al. (1991) in Denver, Pryor (1996) conducted a study of the perception of acceptable visibility conditions in two suburban locations in the lower Fraser Valley in British Columbia, CN. This study reflected public judgments about the acceptability of visual ranges of approximately 40 km to 60 km. Other notable visibility protection initiatives have resulted in state visibility standards that are within this range, including standards for the Lake Tahoe area (Molenaar, 2000) and the state of Vermont.

4.4 PARTICULATE MATTER EFFECTS ON MATERIALS

Effects of air pollution on materials are related to both aesthetic appeal and physical damage. Studies have demonstrated that particles, primarily carbonaceous compounds, cause soiling of commonly used building materials and culturally important items, such as statues and works of art. Physical damage from the dry deposition of air pollutants, such as PM (especially sulfates and nitrates) and SO₂, and the absorption or adsorption of corrosive agents on deposited particles also can result in the acceleration of naturally occurring weathering processes of man-made building and cultural materials.

In the atmosphere, PM may be “primary,” existing in the same form in which it was emitted, or “secondary,” formed by the chemical reactions of free, absorbed, or dissolved gases. The major constituents of atmospheric PM are sulfate, nitrate, ammonium, and hydrogen ions; particle-bound water; elemental carbon; a great variety of organic compounds; and crustal material. A substantial fraction of the fine particle mass, particularly during the warmer months, is secondary sulfate and nitrate. Sulfates may be formed by the gas-phase conversion of SO₂ to H₂SO₄ by OH radicals and aqueous-phase reactions of SO₂ with H₂O₂, O₃, or O₂. During the day, NO₂ may be converted to nitric acid (HNO₃) by reacting with OH radicals. Nitrogen dioxide also can be oxidized to HNO₃ by a sequence of reactions initiated by O₃. A more detailed discussion of the atmospheric chemistry of PM appears in Chapter 2 of this document.

4.4.1 Corrosive Effects of Particles and Sulfur Dioxide on Man-Made Surfaces

This section (a) summarizes information on exposure-related effects on materials associated with sulfur-containing pollutants (formed by chemical reactions of SO₂ with other atmospheric pollutants) as addressed in the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996a) and (b) presents relevant information derived from very limited research conducted and published since completion of that document. The effects of nitrates on man-made building materials and naturally occurring cultural materials were discussed in an earlier EPA Nitrogen Oxides Criteria Document (U.S. Environmental Protection Agency, 1993).

4.4.1.1 Metals

Metals undergo natural weathering processes in the absence of environmental pollutants. The additive effects of pollutants on the natural weathering processes depend on the nature of the pollutant, the deposition rate (the uptake of a pollutant by the material's surface), and the presence of moisture. The influence of the metal-protective corrosion film, the presence of other surface electrolytes, the orientation of the metal surface, the presence of surface moisture, and the variability in the electrochemical reactions also contributes to pollutant exposure effects on metal surfaces.

Several studies demonstrate the importance of the duration of surface wetness (caused by dew and fog condensation and rain) on metals. Surface moisture facilitates the deposition of pollutants, especially SO₂, and promotes corrosive electrochemical reactions on metals (Haynie and Upham, 1974; Sydberger and Ericsson, 1977). Of critical importance is the formation of hygroscopic salts on the metal, increasing the duration of surface wetness and, thereby, enhancing the corrosion process.

The effect of temperature on the rate of corrosion is complex. Under normal temperature conditions, temperature does not have an effect on the rate of corrosion; but when the temperature decreases, the relative humidity increases and the diffusivity decreases. Pitchford and McMurry (1994) and Zhang et al. (1993) demonstrated particle-size-related effects of relative humidity. The corrosion rate decreases as the temperature approaches freezing, because ice inhibits the diffusion of SO₂ to the metal surface and minimizes electrochemical processes (Haynie, 1980; Bieffer, 1981; Sereda, 1974).

The metal-protective corrosion film (i.e., the rust layer on metal surfaces) provides some protection against further corrosion. The effectiveness of the corrosion film in slowing down the corrosion process is affected by the solubility of the corrosion layer and the concentration and deposition rate of pollutants. If the metal-protective corrosion film is insoluble, it may add some protection against acidic pollutants. An atmospheric corrosion model that considers the formation and dissolution of the corrosion film on galvanized steel was proposed by Spence et al. (1992). The model considers the effects of SO₂, rain acidity, and duration of wetness on the rate of corrosion. Although the model does not specifically characterize particle effects, the contribution of particulate sulfate was considered in model development.

Whether suspended particles actually increase metal corrosion is not clear. Several studies have suggested that suspended particles promote the corrosion of metals (Goodwin et al., 1969; Barton, 1958; Sanyal and Singhania, 1956; Baedecker et al., 1991); however, other studies have not demonstrated a correlation between particle exposure and metal corrosion (Mansfeld, 1980; Edney et al., 1989). Walton et al. (1982) suggested that catalytic species within several species in fly ash promote the oxidation of SO_x to a corrosive state. Still other researchers indicate that the catalytic effect of particles is not significant and that the corrosion rate is dependent on the conductance of the thin-film surface electrolytes during periods of wetness. Soluble particles likely increase the solution conductance (Skerry et al., 1988; Askey et al., 1993).

The corrosion of most ferrous metals (iron, steel, and steel alloys) increases with increasing SO_2 exposure. Steels are susceptible to corrosion when exposed to SO_2 in the absence of protective organic or metallic coatings. Studies on the corrosive effects of SO_2 on steel indicate that the rate of corrosion increases with increasing SO_2 and is dependent on the deposition rate of the SO_2 (Baedecker et al., 1991; Butlin et al., 1992a). The corrosive effects of SO_2 on aluminum is exposure-dependent, but appears to be insignificant (Haynie, 1976; Fink et al., 1971; Butlin et al., 1992a). The rate of formation of the patina (protective covering) on copper can take as long as five years and is dependent on the SO_2 concentration, deposition rate, temperature, and relative humidity (Simpson and Horrobin, 1970). Further corrosion is controlled by the availability of copper to react with deposited pollutants (Graedel et al., 1987). Butlin et al. (1992a), Baedecker et al. (1991), and Cramer et al. (1989) reported an average corrosion rate of 1 $\mu\text{m}/\text{year}$ for copper; however, less than a third of the corrosion was attributed to SO_2 exposure, suggesting that the rate of patina formation was more dependent on factors other than SO_2 . A report by Strandberg and Johansson (1997) showed relative humidity to be the primary factor in copper corrosion and patina formation. The results of the studies on particles and SO_2 corrosion of metals are summarized in Table 4-18.

4.4.1.2 Painted Finishes

Exposure to air pollutants affects the durability of paint finishes by promoting discoloration, chalking, loss of gloss, erosion, blistering, and peeling. Evidence exists that particles can damage painted finishes by serving as carriers for corrosive pollutants (Cowling

TABLE 4-18. CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON METALS

Metal	Exposure Conditions	Comments	Source	
Mild Steel Galvanized Steel	Specimens exposed to SO ₂ and O ₃ under natural and artificial conditions, and to NO ₂ under natural conditions. SO ₂ concentrations ranged from 2.1-60 µg/m ³ . Annual average concentrations were about 20 µg/m ³ . Meteorological conditions were unaltered. Specimens exposed at 29 sites for 2 years for mild steel and 1 year for galvanized steel.	Steel corrosion was dependent on long-term SO ₂ exposure. The corrosion rate was about 50 µm/year for mild steel specimens for most industrial sites, but ranged from 21 to 71 µm/year. The corrosion rate ranged from 1.45 to 4.25 µm/year for galvanized steel. The authors concluded that rainfall also may have a significant effect on galvanized steel based on a corrosion rate of 3.4 µm/year seen at a very wet site.	Butlin et al. (1992a)	
Zinc	Rolled zinc specimens exposed at various sites around the country (rural, industrialized, marine) for up to 20 years. Actual pollutant exposures not reported.	The highest corrosion rates were associated with industrialized environments and marine environments in direct contact with salt spray.	Showak and Dunbar (1982)	
4-194	Zinc	Specimens exposed at 5 sites for 1-5 years. Average SO ₂ concentrations ranged from 2 ± 4 to 15 ± 17 ppb (5.2 ± 10.4 to 39.3 ± 44.5 µg/m ³). PM concentrations ranged from 14 to 60 µg/m ³ . Highest pollutant concentrations recorded at 1-year exposure site.	Average corrosion rate ranged from 0.63-1.33 µm per year. The highest corrosion was noted in the most industrialized area. However, the corrosion rates did not differ significant regardless of the SO ₂ concentration, suggesting that SO ₂ exposure may not be the dominant factor in zinc corrosion.	Baedecker et al. (1991) Cramer et al. (1989)
Carbon Steel Weathering Steel	See Baedecker et al. (1991) above for exposure conditions.	Average corrosion rate for samples exposed for 5 years ranged from 6.6-12.8 µm/year for carbon steel and 3.7-5.0 µm/year for weathering steel. Highest corrosion rate noted for samples exposed for 1 year.	Baedecker et al. (1991) Cramer et al. (1989)	
Aluminum	See Baedecker et al. (1991) above for exposure conditions.	Corrosion rate was very low at all sites and ranged from 0.036-0.106 µm/year.	Baedecker et al. (1991)	
Aluminum	See Butlin et al. (1992a) above for exposure conditions.	Corrosion greater on the underside of specimens, possibly because of lack of washoff and increased PM in area. Maximum corrosion rate was 0.85 µm/year. Pit depths of up to 72 µm were noted after 2 years of exposure.	Butlin et al. (1992a)	

TABLE 4-18 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON METALS

Metal	Exposure Conditions	Comments	Source
Copper	See Baedecker et al. (1991) above for exposure conditions.	Average corrosion rate for 3- and 5-year exposures was about 1 $\mu\text{m}/\text{year}$ but the soluble portion was less than a third of that which could be contributed to SO_2 exposure. Dry deposition of SO_2 was not as important in patina formation as wet deposition of H^+ .	Baedecker et al. (1991)
Copper	See Butlin et al. (1992a) above for exposure conditions.	Majority of test sites showed a corrosion rate of $1 \pm 0.2 \mu\text{m}/\text{year}$. The corrosion rate was $1.48 \mu\text{m}/\text{year}$ at the site receiving the most rainfall. The lowest corrosion rate, $0.66 \mu\text{m}/\text{year}$, was associated with low rainfall, low SO_2 .	Butlin et al. (1992a)
Copper	Specimens exposed to 4-69 ppb (10.4-180.7 $\mu\text{g}/\text{m}^3$) and 1.0 ppm (2,618.7 $\mu\text{g}/\text{m}^3$) SO_2 for 20 h at various relative humidities.	SO_2 had no effect on copper when relative humidity was < 75%. Increasing relative humidity increases patina formation in presence of trace SO_2 . No SO_2 -related effects were noted on copper specimens exposed to high SO_2 regardless of the percent relative humidity.	Strandberg and Johansson (1997)
Copper	Specimens exposed artificially to 0.49 ± 0.01 ppm ($187 \pm 3.8 \mu\text{g}/\text{m}^3$) SO_2 for 4 weeks at 70 and 90% relative humidity.	Corrosive effect of SO_2 on copper increased with increasing relative humidity.	Eriksson et al. (1993)
Iron	Specimens from restorations of Acropolis monuments over many years. The oldest specimens were 142 years old. Other specimens used for monument restoration after 1950.	Specimens used after 1950s had an oxidation rate 25% greater than those specimens before 1950. The accelerated oxidation had a negative impact on the structure, producing a quicker formation of cracks inside the marble.	Zuburtikudis and Triantafyllou (2001)

and Roberts, 1954) or by staining and pitting of the painted surfaces (Fochtman and Langer, 1957; Wolff et al., 1990).

The erosion rate of oil-based house paint has been reported to be enhanced by exposure to SO₂ and high humidity. An erosion rate of 36.71 ± 8.03 μm/year was noted for oil-based house paint samples exposed to SO₂ (78.6 μg/m³), O₃ (156.8 μg/m³), and NO₂ (94 μg/m³), and low humidity (50%) (Spence et al., 1975). The erosion rate increased with increased SO₂ and humidity. The authors concluded that SO₂ and humidity accounted for 61% of the erosion. Acrylic and vinyl coil coatings show less pollutant-related erosion. Erosion rates range from 0.7 to 1.3 μm/year and 1.4 to 5.3 μm/year, respectively. Similar findings on SO₂-related erosion of oil-based house paints and coil coatings have been reported by other researchers (Davis et al., 1990; Yocom and Grappone, 1976; Yocom and Upham, 1977; Campbell et al., 1974). Several studies have suggested that the effect of SO₂ is caused by its reaction with extender pigments such as calcium carbonate and zinc oxide (Campbell et al., 1974; Xu and Balik, 1989; Edney, 1989; Edney et al., 1988, 1989). However, Miller et al. (1992) suggested that calcium carbonate acts to protect paint substrates. Another study indicated that exposure to SO₂ can increase the drying time of some paints by reacting with certain drying oils and will compete with the auto-oxidative curing mechanism responsible for crosslinking the binder (Holbrow, 1962).

4.4.1.3 Stone and Concrete

Numerous studies suggest that air pollutants can enhance the natural weathering processes on building stone. The development of crusts on stone monuments has been attributed to the interaction of the stone's surface with sulfur-containing pollutants, wet or dry deposition of atmospheric particles, and dry deposition of gypsum particles from the atmosphere. Because of a greater porosity and specific surface, mortars have a greater potential for reacting with environmental pollutants (Zappia et al., 1998). Details on these studies are discussed in Table 4-19. The stones most susceptible to the deteriorating effects of sulfur-containing pollutants are the calcareous stones (limestone, marble, and carbonated cement). Exposure-related damage to building stones result from the formation of salts in the stone that are subsequently washed away during rain events, leaving the stone surface more susceptible to the effects of pollutants. Dry deposition of sulfur-containing pollutants promotes the formation of gypsum on the stone's surface. Gypsum is a gray to black crusty material comprised mainly

TABLE 4-19. CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Vermont marble	Runoff water was analyzed from seven summer storms. SO ₂ concentration stated to be low.	Between 10 and 50% of calcium in runoff water estimated from gypsum formation from dry deposition of SO ₂ .	Schuster et al. (1994)
Marble sandstone	Analysis of runoff water for five slabs test-exposed to ambient conditions at a angle of 30° to horizontal.	Pollutant exposure related erosion was primarily caused by dry deposition of SO ₂ and nitric acid between rain events and wet deposition of hydrogen ion. Recession estimates ranged from 15-30 μm/year for marble and 25-45 μm/year for limestone. A large portion of the erosion results from the reaction of CO ₂ with the calcium in the stone.	Baedecker et al. (1992)
Limestone	Ambient air conditions. Exposure ranged from 70-1,065 days. Averaged pollutant exposure ranged from 1.4-20.4 ppb (3.7-53.4 μg/m ³) SO ₂ ; 4.1-41.1 ppb NO _x ; 2.4-17.4 ppb (4.5-32.7 μg/m ³) NO ₂ ; 10.1-25.6 ppb (19.8-50.2 μg/m ³) O ₃ .	Increased stone weight loss with increased SO ₂ . Rainfall did not significantly affect stone degradation. Stone loss associated with SO ₂ exposure estimated to be 24 μm/year. Slight trend in decreasing stone loss with increasing length of exposure.	Webb et al. (1992)
Portland limestone White Mansfield dolomitic sandstone Monk's Park limestone	Experimental tablets exposed under sheltered and unsheltered ambient air conditions. Exposure for 1 and 2 years.	Significant correlations existed between the mean annual SO ₂ concentration, rainfall volume, and hydrogen ion loading and the weight changes.	Butlin et al. (1992b)
Sandstones (calcite and noncalcite stones)	Ambient air; low concentrations of sulfates, SO ₂ , and nitrates; RH sufficient to produce condensation on stones rarely occurred.	Insignificant differences in erosion rate found between calcite and noncalcite sandstone. Moisture affected the rate of pollutant deposition and enhanced susceptibility to pollutant related erosion. Rain events given as primary factor affecting stone erosion. Pollutant related erosion judged to be insignificant.	Petuskey et al. (1995)
Limestones Sandstones Marble Granite Basalt	Ambient air; urban and rural locations in Mediterranean.	Crusts on stones were found to contain two layers. Top layer, usually black in color, composed of gypsum between 40 and 400 μm thick. Innermost layer, ranging from brown to orange in color, primarily consisted of calcite, between 10 and 600 μm thick. Gypsum-rich layer thought to be the result of sulfation of the calcitic layer by atmospheric pollutants or dry or wet deposition of atmospheric dust.	Garcia-Vallès et al. (1998)
Portland limestone Massangis Jaune Roche limestone White Mansfield dolomitic	Samples exposed to SO ₂ , NO ₂ , and NO at 10 ppmv, both with and without O ₃ and under dry (coming to equilibrium with the 84% RH) or wetted with CO ₂ -equilibrated deionized water conditions. Exposure was for 30 days.	In the absence of moisture, little reaction is seen. Sulfur dioxide is oxidized to sulfates in the presence of moisture. The effect is enhanced in the presence of O ₃ . Massangis Jaune Roche limestone was the least affected by the pollutant exposure. Crust lined pores of specimens exposed to SO ₂ .	Haneef et al. (1993)

TABLE 4-19 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Monk's Park limestone Portland limestone	Samples exposed for 2 months under both sheltered and unsheltered conditions. Mean daily atmospheric SO ₂ concentration was 68.7 µg/m ³ and several heavy rainfalls.	Significant amounts of gypsum were noted on the Portland stone. Sheltered stones also showed soiling by carbonaceous particles and other combustion products. Etch holes and deep etching was noted in some of the exposed unsheltered samples.	Viles (1990)
Carrara marble Travertine Tranistone	Sample exposed in laboratory to 3 ppm SO ₂ and 95% RH at 25 °C for 150 days. Samples were coated with three carbonaceous particle samples from combustion sources, and with activated carbon and graphite.	Exposure to particles from combustion processes enhanced sulfation of calcareous materials by SO ₂ because of metal content of particles.	Sabbioni et al. (1996)
Carrara marble Georgia marble	Samples exposed in sheltered ambient environment for 6, 12, or 20 months.	Carrara marble found to be more reactive with SO ₂ than Georgia marble, possibly because of the compactness of the Georgia marble. Greater effects noted when samples were also exposed to NO ₂ .	Yerrapragada et al. (1994)
Carrara marble	Samples exposed for 6 months (cold and hot conditions) in ambient environment. PM concentrations ranged from 57.3 to 116.7 µg/m ³ (site 1) and 88 to 189.8 µg/m ³ (site 2). Some exposures also were associated with high SO ₂ , NO, and NO ₂ .	Pollutant exposed samples showed increased weight gain over that expected from natural weathering processes. There was a blackening of stone samples exposed to carbonaceous rich PM.	Realini et al. (1995)
Monk's Park limestone Portland limestone	Samples artificially exposed to fly-ash containing 1,309.3 µg/m ³ SO ₂ (0.5 ppm) at 95% RH and 25 °C for 81 or 140 days. Fly-ash samples from five different sources were used in the study.	Exposure to fly-ash did not enhance oxidation of SO ₂ to sulfates. Mineral oxides in fly ash contributed to sulphation of CaCO ₃ .	Hutchinson et al. (1992)
Lime mortar Pozzolan mortar Cement mortar	Samples exposed to 7,856 µg/m ³ (3 ppm) SO ₂ at 100% RH and 25 °C for 30, 60, or 90 days; samples sprayed with bidistilled water every 7 days to simulate rainfall.	Exposure to SO ₂ produced significant quantities of calcium sulfite and calcium sulfate on specimens; however, the amount produced was dependent of the porosity, specific surface, and alkalinity of the sample.	Zappia et al. (1994)
Limestone Travertine marble	Samples exposed under actual ambient air conditions at two locations in Rome. Monitoring data obtained for SO ₂ , NO, NO ₂ , and total suspended particulates (TSP) but not reported. Exposure was for four seasons.	TSP exposure increased the cleaning frequency for stone monuments. Monuments are soiled proportionately overtime, based on brightness values. Horizontal surfaces showed higher graying values because of particle sediment.	Lorusso et al. (1997)

TABLE 4-19 (cont'd). CORROSIVE EFFECTS OF PARTICULATE MATTER AND SULFUR DIOXIDE ON STONE

Stone	Exposure Conditions	Comments	Source
Limestone Quartz-cemented sandstone Calcite-cemented sandstone Granite Brick	Samples from structures exposed for varying periods of time under ambient air conditions. Samples selected because of black layer on surface.	Black layers were found to be primarily comprised of iron compounds, quartz, silicate, soot, and dirt.	Nord and Ericsson (1993)
Carrara marble	Samples from structures taken from monuments in Venice, Italy. Samples included those shielded from running water (unwashed), areas at the interface of the washed and unwashed areas, and washed areas. Samples of fog and rain near the monument were also taken for analysis.	Sulfate was higher in fog compared to other ions. The average concentration of chlorides and sulfates were 7-16 times higher in fog than rain. The degree of sulphonation on stone samples from areas shielded from running water was < 40%. At the interface between washed and unwashed surfaces and at washed areas, sulphonation was > 40%. Dendrite shaped crust from the transformation of calcium carbonate into gypsum was found on samples shielded from rain. Samples from the washed areas displayed superficial granular disaggregation, a natural process of deterioration ascribed to natural agents; however, the process was accelerated due to the ambient air sulfates.	Fassina et al. (2001)
Limestone Sandstone	Samples of ancient grey crust formed between 1180 and 1636 on the Church of Saint Trophime in Arks and formed between 1530 and 1187 on the Palazzo d'Accursio in Bologna.	Crust samples contained calcite, soil dust, carbonaceous particles, and gypsum crystals.	Ausset et al. (1998)
Carrara marble Travertine marble Trani limestone Portland limestone Lime mortar Pozzolan mortar Cement mortar	Samples of the stones and mortars were representative of those used in the past and currently for new construction and restorations. Samples were exposed for 6, 12, and 24 months under ambient conditions in Milan.	Mortars were more reactive than the stones. Of the mortars, cement and pozzolan mortar were more reactive than the lime mortar. Carrara marble was the least reactive of the stones. The maximum amount of degradation was found in areas sheltered from rain.	Zappia et al. (1998)
Lime mortar	Sample of black crust taken from Zamboni Tower Gate.	Exposure to environmental pollutants caused the formation of two separate layers on the mortar: an outer thin surface black crust composed of gypsum and carbonaceous particles and the inner composed of products from the dissolution and sulphation of the carbonate matrix in the mortar.	Sabbioni et al. (1998)
Carrara marble	Samples of crust removed from Milan General Hospital, built around 1937.	Gypsum main component of crust followed by carbonaceous particles and iron oxides. Estimated rate of crust formation was 2-5 µm/year. Total amount of gypsum formed over the lifetime of exposure was 5-13 mg/cm ² , an estimated 0.2 mg/cm ² /year.	Bugini et al. (2000)

of calcium sulfate dihydrate from the reaction of calcium carbonate (calcite) in the stone with atmospheric SO₂ and moisture (relative humidities exceeding 65%) according to the following reaction.



The sulfate anions formed in the moist air reacts with the Ca²⁺ through diffusion processes forming the gypsum (Zuburtikudis and Triantafyllou, 2001). Approximately 99% of the sulfur in gypsum is sulfate because of the sulfonation process caused by the deposition of SO₂ aerosol. Sulfites also are present in the gypsum layer as an intermediate product (Sabbioni et al., 1996; Ghedini et al., 2000; Gobbi et al., 1998; Zappia et al., 1998). Gypsum is more soluble than calcite and is known to form on limestone, sandstones, and marble when exposed to SO₂. Gypsum also has been reported to form on granite stone by replacing silicate minerals with calcite (Schiavon et al., 1995). Gypsum occupies a larger volume than the original stone, causing the stone's surface to become cracked and pitted. The rough surface serves as a site for deposition of airborne particles. As the gypsum grows, it loosens and falls apart (Zuburtikudis and Triantafyllou, 2001).

The dark colored gypsum is created by surface deposition of carbonaceous particles (noncarbonate carbon) from combustion processes occurring in the area (Sabbioni, 1995; Saiz-Jimenez, 1993; Ausset et al., 1998; Hermosin and Saiz-Jimenez, 2000), trace metals contained in the stone, dust, and numerous other anthropogenic pollutants. After analyzing damaged layers of several stone monuments, Zappia et al. (1993) found that the dark-colored damaged surfaces contained 70% gypsum and 20% noncarbonate carbon. The lighter colored, damaged layers were exposed to rain and contained 1% gypsum and 4% noncarbonate carbon. It is assumed that rain removes reaction products, permitting further pollutant attack on the stone monument and likely redeposits some of the reaction products at rain runoff sites on the stone. After sulfur compounds, carbon was reported to be the next highest element in the dark crust on historical monuments in Rome. Elemental carbon and organic carbon accounted for 8 and 39% of the total carbon in the black crust samples. The highest percentage of carbon, carbonate carbon, was derived from the carbonate matrix in the stones. The high ratio of organic carbon to elemental carbon indicates the presence of a carbon source other than combustion processes

(Ghedini et al., 2000). Cooke and Gibbs (1994) suggested that stones damaged during times of higher ambient pollution exposure likely would continue to exhibit a higher rate of decay, termed the “memory effect,” than newer stones exposed under lower pollution conditions. Increased stone damage also has been associated with the presence of sulfur-oxidizing bacteria and fungi on stone surfaces (Garcia-Vallès et al., 1998; Young, 1996; Saiz-Jimenez, 1993; Diakumaku et al., 1995).

Dissolution of gypsum on the stone’s surface initiates structural changes in the crust layer. Garcia-Vallès et al. (1998) proposed a double mechanism: the dissolution of the gypsum, in the presence of sufficient moisture, followed by recrystallization inside fissures or pores. In the event of limited moisture, the gypsum is dissolved and recrystallizes at its original location. According to the authors, this would explain the gypsum-rich crustal materials on stone surfaces sheltered from precipitation.

Moisture was found to be the dominant factor in stone deterioration for several sandstones (Petuskey et al., 1995). Dolske (1995) reported that the deteriorative effects of sulfur-containing rain events, sulfates, and SO₂ on marble were largely dependent on the shape of the monument or structure rather than the type of marble. The author attributed the increased fluid turbulence over a non-flat vertical surface versus a flat surface to the increased erosion. Sulfur-containing particles also have been reported to enhance the reactivity of Carrara marble and Travertine and Trani stone to SO₂ (Sabbioni et al., 1992). Particles with the highest carbon content had the lowest reactivity.

The rate of stone deterioration is determined by the pollutant and the pollutant concentration, the stone’s permeability and moisture content, and the pollutant deposition velocity. Dry deposition of SO₂ between rain events has also been reported to be a major causative factor in pollutant-related erosion of calcareous stones (Baedecker et al., 1991; Dolske, 1995; Cooke and Gibbs, 1994; Schuster et al., 1994; Hamilton et al., 1995; Webb et al., 1992). Sulfur dioxide deposition increases with increasing relative humidity (Spiker et al., 1992), but the pollutant deposition velocity is dependent on the stone type (Wittenburg and Dannecker, 1992), the porosity of the stone, and the presence of hygroscopic contaminants.

Although it is clear from the available information that gaseous pollutants (dry deposition of SO₂ in particular) will promote the decay of some types of stones under specific conditions, carbonaceous particles (noncarbonate carbon) may help to promote the decay process by aiding

in SO₂ transformation to a more acidic species (Del Monte and Vittori, 1985). Several authors have reported enhanced sulfation of calcareous material by SO₂ in the presence of particles containing metal oxides (Sabbioni et al., 1996; Hutchinson et al., 1992).

4.4.2 Soiling and Discoloration of Man-Made Surfaces

Ambient particles can cause soiling of man-made surfaces. Soiling has been defined as the deposition of particles of less than 10 µm on surfaces by impingement. Soiling generally is considered an optical effect, that is, soiling changes the reflectance from opaque materials and reduces the transmission of light through transparent materials. Soiling can represent a significant detrimental effect, requiring increased frequency of cleaning of glass windows and concrete structures, washing and repainting of structures, and, in some cases, reducing the useful life of the object. Particles, especially carbon, may also help catalyze chemical reactions that result in the deterioration of materials during exposure.

It is difficult to determine the accumulated particle levels that cause an increase in soiling. Soiling is dependent on the particle concentration in the ambient environment, particle size distribution, the deposition rate, and the horizontal or vertical orientation and texture of the surface being exposed (Haynie, 1986). The chemical composition and morphology of the particles and the optical properties of the surface being soiled will determine the time at which soiling is perceived (Nazaroff and Cass, 1991). Carey (1959) reported that the average observer could observe a 0.2% surface coverage of black particles on a white background. Work by Bellan et al. (2000) indicated that it would take a 12% surface coverage by black particles before there is 100% accuracy in identifying soiling. Sharples et al. (2001) studied the effect of air pollution, moisture, and the function of the room/building on glazing daylight transmittance for a number of building windows. They found that the direct soiling load to a window was dependent on the immediate external and internal environment. For instance, there was only a 10% reduction in daylight transmittance for windows from an office building that had not been cleaned for 5 years compared to clean windows. The reduction in transmittance for windows in a swimming pool complex was in excess of 20% due to soiling of the interior surface. For most office buildings, there was a reduction of glazing transmittance ranging from 3 to 10%, with most windows showing about a 3% reduction. The rate at which an object is soiled increases linearly with time; however, as the soiling level increases, the rate of soiling decreases. The

buildup of particles on a horizontal surface is counterbalanced by an equal and opposite depletion process, which is based on the scouring and washing effects of wind and rain (Schwar, 1998).

4.4.2.1 Stones and Concrete

Most of the research evaluating the effects of air pollutants on stone structures has concentrated on gaseous pollutants. The deposition of the sulfur-containing pollutants is associated with the formation of gypsum on the stone (see Section 4.5.1.3). The dark color of gypsum is attributed to soiling by carbonaceous particles from nearby combustion processes. A lighter gray colored crust is attributed to soil dust and metal deposits (Ausset et al., 1998; Camuffo, 1995; Moropoulou et al., 1998). Realini et al. (1995) recorded the formation of a dark gypsum layer and a loss of luminous reflection in Carrara marble structures exposed for 1 year under ambient air conditions. Dark areas of gypsum were found by McGee and Mossitti (1992) on limestone and marble specimens exposed under ambient air conditions for several years. The black layers of gypsum were located in areas shielded from rainfall; whereas particles of dirt were concentrated around the edges of the gypsum formations. Lorusso et al. (1997) attributed the need for frequent cleaning and restoration of historic monuments in Rome to exposure to total suspended particulates. They also concluded that, based on a decrease in brightness (graying), surfaces are soiled proportionately over time; however, graying is higher on horizontal surfaces because of sedimented particles. Davidson et al. (2000b) evaluated the effects of air pollution exposure on a limestone structure on the University of Pittsburgh campus, using estimated average TSP levels in the 1930s and 1940s and actual values for the years 1957 to 1997. Monitored levels of SO₂ were available for the years 1980 to 1998. Based on the available data concerning pollutant levels and photographs, it was thought that soiling began while the structure was under construction. With decreasing levels of pollution, the soiled areas have been slowly washed away, the process taking several decades, leaving a white, eroded surface.

4.4.2.2 Household and Industrial Paints

Few studies are available that evaluate the soiling effects of particles on painted surfaces. Particles composed of elemental carbon, tarry acids, and various other constituents are

responsible for the soiling of structural painted surfaces. Coarse-mode particles ($> 2.5 \mu\text{m}$) initially contribute more soiling of horizontal and vertical painted surfaces than do fine-mode particles ($< 2.5 \mu\text{m}$), but are more easily removed by rain (Haynie and Lemmons, 1990). The accumulation of fine particles likely promotes remedial action (i.e., cleaning of the painted surfaces); whereas coarse-mode particles are primarily responsible for soiling of horizontal surfaces. Rain interacts with coarse particles, dissolving the particle and leaving stains on the painted surface (Creighton et al., 1990; Haynie and Lemmons, 1990). Haynie and Lemmons (1990) proposed empirical predictive equations for changes in surface reflectance of gloss-painted surfaces that were exposed protected and unprotected from rain while oriented horizontally or vertically.

Early studies by Parker (1955) and Spence and Haynie (1972) demonstrated an association between particle exposure and the increased frequency of cleaning of painted surfaces. Particle exposures also caused physical damage to the painted surface (Parker, 1955). Unsheltered painted surfaces are initially more soiled by particles than sheltered surfaces, but the effect is reduced by rain washing. Reflectivity is decreased more rapidly on glossy paint than on flat paint (Haynie and Lemmons, 1990). However, surface chalking of the flat paint was reported during the exposure. The chalking interfered with the reflectance measurements for particle soiling. Particle composition measurements that were taken during exposure of the painted surfaces indicated sulfates to be a large fraction of the fine mode and only a small fraction of the coarse mode. Although no direct measurements were taken, fine-mode particles likely also contained large amounts of carbon and possibly nitrogen or hydrogen (Haynie and Lemmons, 1990).

4.5 ATMOSPHERIC PARTICULATE MATTER, CLIMATE CHANGE, AND EFFECTS ON SOLAR UV-B RADIATION TRANSMISSION

Atmospheric particles alter the amount of solar radiation transmitted through the Earth's atmosphere. The absorption of solar radiation by atmospheric particles, together with the trapping of infrared radiation emitted by the Earth's surface by certain gases, enhances the heating of the Earth's surface and lower atmosphere i.e., the widely-known "greenhouse effect." Increases in the atmospheric concentration of these gases due to human activities may lead to

impacts, due to climate change, on human health and the environment. Lesser consequences of airborne particles include alterations in the amount of ultraviolet solar radiation (especially UV-B, 290 to 315 nm) penetrating through the Earth's atmosphere and reaching its surface where UV radiation can exert various effects on human health, plant and animal biota, and other environmental components.

The effects of atmospheric PM on the transmission of electromagnetic radiation emitted by the sun at ultraviolet and visible wavelengths, and by the Earth at infrared wavelengths, depend on the radiative properties (extinction efficiency, single-scattering albedo, and asymmetry parameter) of the particles, which depend, in turn, on the size and shape of the particles, the composition of the particles, and the distribution of components within individual particles. In general, the radiative properties of particles are size- and wavelength-dependent, with the extinction cross section tending toward its maximum when the particle radius is similar to the wavelength of the incident radiation. This means that fine particles present mainly in the accumulation mode would be expected to exert a greater influence on the transmission of electromagnetic radiation than would coarse particles. The chemical components of particles can be crudely summarized in terms of the broad classes identified in Chapter 2 of this document. These classes include nitrate, sulfate, mineral dust, elemental carbon, organic carbon compounds (e.g., PAHs), and metals derived from high-temperature combustion or smelting processes. The major sources of these components are shown in Table 3-9 of Chapter 3 in this document.

Knowledge of the effects of PM on the transfer of radiation in the visible and infrared spectral regions is needed for assessing relationships between particles and climate change processes, as well as environmental and biological effects. Knowledge of the factors controlling the transfer of solar radiation in the ultraviolet spectral range is needed to assess potential the biological and environmental effects associated with exposure to UV-B radiation. Climate change processes, their potential to affect human and environmental health, and their potential relationships to atmospheric PM are discussed below. Solar ultraviolet radiation processes and the related effects of atmospheric PM are then summarized and discussed in the next section.

4.5.1 Atmospheric Particle Interactions with Solar and Terrestrial Radiation Related to Climate Change

4.5.1.1 The Projected Impacts of Global Climate Change

The study of atmospheric processes involved in mediating global climate change and its potential consequences for human health and global ecosystems is an area of active research. The most thorough evaluation of currently available science regarding climate change is the Third Assessment Report (TAR) of the Intergovernmental Panel on Climate Change (IPCC, 2001a). Earlier assessments include those conducted by the United Nations Environment Program (UNEP, 1986), the World Meteorological Organization (WMO, 1988), the EPA (U.S. Environmental Protection Agency, 1987), and others (e.g., Patz et al., 2000 a,b). The reader is referred to these documents for more complete discussion of climate change science. An abbreviated list of the IPCC conclusions, to date, and a short discussion of the potential impacts of climate change on human health and welfare is provided here to serve as the context for the discussion of the role of PM in climate.

The IPCC TAR (2001a) notes that the increasing body of observations indicates that the Earth is warming and that other climate changes are underway. These observations include the global surface temperature record for the time period beginning in the year 1860, the satellite temperature record begun in 1979, changes in snow and ice cover recorded since the 1950s, sea level measurements taken throughout the 20th century, and sea surface temperature observations recorded since the 1950s. Other evidence includes a marked increase over the past 100 years in the frequency, intensity and persistence of the zonal atmospheric circulation shifts known as the El Nino-Southern Oscillation (ENSO). ENSO events occur when the tropical ocean has accumulated a large, localized mass of warm water that interrupts cold surface currents along South America, altering precipitation and temperature patterns in the tropics, subtropics and the midlatitudes.

Atmospheric concentrations of greenhouse gases (GHGs), which trap solar energy within the climate system, are continuing to increase due to human activities. These activities will continue to change the composition of the atmosphere throughout the 21st century. The IPCC TAR describes the scientific evidence that ties the observed increase in GHGs over the past 50 years to human activities (IPCC, 2001b).

The IPCC (1998, 2001b) reports also describe the results of general circulation model (GCM) studies that predict that human activities will alter the climate system in a manner that is likely to lead to marked global and regional changes in temperature, precipitation, and other climate properties. Global mean sea level is expected to increase, as will the number of extreme weather events including floods and droughts, thus inducing changes in soil moisture. These changes will directly impact human health, ecosystems, and global economic sectors, e.g., hydrology and water resources, food and fiber production, etc., (IPCC 1998, 2001b). Table 4-20 summarizes these projected impacts. Wide variations in the course and net impacts of climate change in different geographic areas can be expected. In general, projected climate change impacts can be expected to represent additional stressors on those natural ecosystems and human societal systems already impacted by increasing resource demands, unsustainable resource management practices, and pollution — with wide variation likely across regions and nations in their ability to cope with the consequent alterations in ecological balances, in the availability of adequate food, water, and clean air, and in human health and safety. However, although many regions are likely to experience severe adverse impacts of climate change (some possibly irreversible), certain climate change impacts may be beneficial in some regions. For example, ecosystems that may benefit from warmer temperatures or increased CO₂ fertilization include west coast coniferous forests and some western rangelands. In addition, reduced energy requirements for heating, reduced road salting and snow-clearance costs, and longer open-water seasons in northern channels and ports may benefit communities in the northern latitudes. The IPCC report, “The Regional Impacts of Climate Change” (IPCC, 1998), describes the projected effects of human-induced climate change on the different regions of the globe, including Africa, the Arctic and Antarctic, the Middle East and arid Asia, Australasia, Europe, Latin America, North America, the small island nations, temperate Asia, and tropical Asia.

It must be borne in mind that while current climate models are successful in simulating present annual mean climate and the seasonal cycle on continental scales, they are less successful at regional scales. Clouds and humidity, essential factors in defining local and regional (“sub-grid”) climate, are significantly uncertain (IPCC, 2001a). Due to modeling uncertainties, both in reproducing regional climate and in predicting future economic activity, the projected impacts discussed above are also uncertain.

TABLE 4-20. EXAMPLES OF IMPACTS RESULTING FROM PROJECTED CHANGES IN EXTREME CLIMATE EVENTS

Projected changes during the 21st Century in Extreme Climate Phenomena and their Likelihood ^a	Representative Examples of Projected Impacts ^b (all high confidence of occurrence in some areas ^c)
<i>Simple Extremes</i>	
Higher maximum temperatures; more hot days and heat waves ^d over nearly all land areas (<i>very likely</i> ^a)	<ul style="list-style-type: none"> • Increased incidence of death and serious illness in older age groups and urban poor • Increased heat stress in livestock and wildlife • Shift in tourist destinations • Increased risk of damage to a number of crops • Increased electric cooling demand and reduced energy supply reliability
Higher (increasing) minimum temperatures; fewer cold days, frost days, and cold waves ^d over nearly all land areas (<i>very likely</i> ^a)	<ul style="list-style-type: none"> • Decreased cold-related human morbidity and mortality • Decreased risk of damage to a number of crops, and increased risk to others • Extended range and activity of some pest and disease vectors • Reduced heating energy demand
More intense precipitation events (<i>very likely</i> ^a over many years)	<ul style="list-style-type: none"> • Increased flood, landslide, avalanche, and mudslide damage • Increased soil erosion • Increased flood runoff could increase recharge of some floodplain aquifers • Increased pressure on government and private flood insurance systems and disaster relief
<i>Complex Extremes</i>	
Increased summer drying over most mid-latitude continental interiors and associated risk of drought (<i>likely</i> ^a)	<ul style="list-style-type: none"> • Decreased crop yields • Increased damage to building foundations caused by ground shrinkage • Decreased water resource quantity and quality • Increased risk of forest fire
Increased tropical cyclone peak wind intensities, mean and peak precipitation intensities (<i>likely</i> ^a over some areas) ^e	<ul style="list-style-type: none"> • Increased risk to human life, risk of infections, disease epidemics, and many other risks • Increased coastal erosion and damage to coastal buildings and infrastructure • Increased damage to coastal ecosystems such as coral reefs and mangroves
Intensified droughts and floods associated with El Niño events in many different regions (<i>likely</i> ^a) (see also under droughts and intense precipitation events)	<ul style="list-style-type: none"> • Decreased agricultural and rangeland productivity in drought- and flood-prone regions • Decreased hydropower potential in drought-prone regions
Increased Asian summer monsoon precipitation variability (<i>likely</i> ^a)	<ul style="list-style-type: none"> • Increased flood and drought magnitude and damages in temperate and tropical Asia
Increased intensity of mid-latitude storms (little agreement between current models) ^d	<ul style="list-style-type: none"> • Increased risks to human life and health • Increased property and infrastructure losses • Increased damage to coastal ecosystems

^aLikelihood refers to judgmental estimates of confidence used by TAR WGI: *very likely* (90-99% chance); *likely* (66-90% chance). Unless otherwise stated, information on climate phenomena is taken from the Summary for Policymakers, TAR WGI. TAR WGI = Third Assessment Report of Working Group 1 (IPCC, 2001a).

^bThese impacts can be lessened by appropriate response measures.

^cHigh confidence refers to probabilities between 67 and 95%.

^dInformation from TAR WGI, Technical Summary.

^eChanges in regional distribution of tropical cyclones are possible but have not been established.

Source: IPCC (2001b).

Findings from the IPCC TAR (2001a), Regional Impacts Assessment (1998), and other regional assessments illustrate well the considerable uncertainties and difficulties in projecting likely climate change impacts on regional or local scales. The findings predict a mixture of positive and negative projected potential climate change impacts for U.S. regions differing widely among regions. Projections of region-specific climate change impacts are complicated by the need to evaluate potential effects of local- or regional-scale changes in key air pollutants.

4.5.1.2 Airborne Particle Relationships to Global Warming and Climate Change

Atmospheric particles both scatter and absorb incoming solar radiation. Visibility reduction is caused by particle scattering in all directions, whereas climate effects are mainly caused by particle scattering in the upward direction. Upward scattering of solar radiation reduces the total amount of energy received by the Earth system, leading to surface cooling. The effect on climate due to upward scattering and to absorption of radiation by aerosol can be roughly quantified as a “radiative forcing” (Houghton et al., 1990). Global and regional climate (at equilibrium) is defined by the balance between a large number of “positive” and “negative” forcings induced by different components of the Earth system. The Earth system responds to these forcings in a potentially complex way due to feedback mechanisms that are theorized but difficult to model. In the absence of information about climate feedbacks, radiative forcing values for many components of the climate system are estimated as a tool for approximating their relative importance in climate change. Forcing estimates for various classes of atmospheric particles are derived on the basis of climate modeling studies and reported by the IPCC (IPCC, 2001b).

Particulate matter appears to play a significant role in defining climate on both global and regional scales. Significant reductions over the past 50 years in solar radiation received at the Earth’s surface on a globally averaged basis correlate with increases in atmospheric aerosol (Stanhill and Cohen, 2001). While this correlation seems clear, quantifying the cooling and warming effects of aerosol in relation to GHG-related warming is difficult. Aerosols complicate the prediction of climate change due to their spatial and temporal inhomogeneities and uncertain radiative properties. However, inclusion of modeled atmospheric sulfate concentrations substantially improved the agreement between modeled and observed surface temperatures (Kiehl and Briegleb, 1993). Haywood et al. (1999) also found that the inclusion of

anthropogenic aerosols results in a significant improvement between calculations of reflected sunlight at the top of the atmosphere and satellite observations in oceanic regions close to sources of anthropogenic PM. On a regional scale, the suspected influence of aerosols upon climate relates to regional hydrological cycles. Evidence is accumulating that pollution aerosols reduce precipitation frequency by clouds, potentially leading to drought in some parts of the world (Ramanathan et al., 2001).

Greenhouse Gases, Particulate Matter, and the Earth's Radiative Equilibrium

According to simple radiative transfer theory, at thermal equilibrium, the Earth's surface should be near $-15\text{ }^{\circ}\text{C}$. This is the temperature of a theoretical "black body" that is receiving and then re-emitting 342.5 W m^{-2} , the globally averaged amount of solar radiation absorbed and then re-emitted by the Earth as infrared terrestrial radiation. In fact, satellite observations indicate that the Earth's average planetary temperature is remarkably close to the theoretical black body value at $-18\text{ }^{\circ}\text{C}$, a temperature at which liquid water ordinarily does not exist.

At its surface, however, the Earth's average temperature is $+15\text{ }^{\circ}\text{C}$. The $33\text{ }^{\circ}\text{C}$ temperature differential between Earth's planetary temperature and its surface temperature is due to the existence of infrared radiation-absorbing components in the atmosphere, i.e., GHGs including carbon dioxide, methane, several other trace gases, and some types of particles and clouds. The phenomenon of planetary surface warming due to the atmospheric absorption and re-emission of infrared radiation is popularly known as the "greenhouse effect" (Arrhenius, 1896; Schneider, 1992). Radiation trapped by the Earth's atmosphere is reflected back to its surface, with some small fraction absorbed by dark atmospheric particles. The fraction of this radiation that is not directly re-emitted as long-wave terrestrial radiation transforms into heat energy that drives the atmospheric processes that form the basis of weather and climate. Eventually this energy is transformed a second time, to terrestrial radiation, and is re-emitted as part of the process that maintains Earth's radiative equilibrium.

The evidence of a physical relationship between radiatively active air pollutants (GHGs and particles) and solar energy retained by the Earth system suggests that continuously changing atmospheric concentrations of air pollutants, along with other alterations to the climate system due to human activities, may be shifting the Earth's radiative equilibrium.

Radiatively active gases in the atmosphere are largely responsible for the greenhouse effect, although some light-absorbing particles and clouds contribute to atmospheric and surface warming (IPCC, 2001a). The majority of clouds and particles play a role in counteracting the greenhouse effect by increasing the degree to which the Earth is able to reflect solar radiation, i.e. its albedo. Successful modeling of the Earth's climate and, therefore, assessment of the degree of human-induced climate change and development of appropriate policy depends on the high quality information on the relative efficiencies, amounts, and spatial and temporal distributions of the various radiatively active components of the atmosphere involved in absorbing and/or reflecting solar and terrestrial radiation.

“Forcing” and the Earth’s Radiative Balance

A measure of the relative influence of a given component of the climate system on the Earth's radiative balance is its “radiative forcing”. Radiative forcing, in W m^{-2} , is a quantity that was developed by the climate modeling community as a first order-only means of estimating relative effects of anthropogenic and natural processes on the surface-troposphere system. No more precise metric has yet been found to replace radiative forcing as a measure of impact of upon global climate (IPCC, 2001a). The convention for this quantity assigns a negative forcing to climate system components that reflect solar radiation back into space and assigns a positive forcing to those that enhance the greenhouse effect or otherwise act to enhance the heat-absorbing capacity of the Earth system. Purely reflective atmospheric aerosol, snow-covered land surfaces, and dense sea ice provide a negative forcing, while black carbon-containing atmospheric aerosol, GHGs, and dark ocean surfaces provide a positive forcing of the climate system.

The IPCC reports included estimated values for forcing by the individual, radiatively active gas and particle-phase components of the atmosphere. These estimates are derived primarily through expert judgment, incorporating the results of peer-reviewed modeling studies. Uncertainty ranges are assigned that reflect the range of modeled values reported in these studies. According to the available research on climate forcing by aerosol, the panel has provided estimates for sulfate, organic, black carbon, biomass burning, and mineral dust aerosol. The current estimate of forcing due to long-lived, well-mixed GHGs accumulated in the atmosphere from the preindustrial era (~1750) through the year 2000 is $+2.4 \text{ W m}^{-2}$ (IPCC,

2001a). In contrast, forcing due to a sulfate aerosol-related increase in planetary albedo has been assigned a value of -0.4 W m^{-2} . Biomass burning and fossil-fuel-related organic aerosol also increase the Earth's reflectivity and are estimated to contribute a -0.2 W m^{-2} and -0.1 W m^{-2} forcing, respectively. Fossil-fuel black carbon is expected to warm the atmosphere, resulting in an estimated $+0.2 \text{ W m}^{-2}$ forcing. No estimate for forcing by nitrate aerosol has been proposed due to the wide discrepancies in current global modeling results and the difficulties associated with obtaining accurate ambient samples of nitrate concentrations and size distributions. Likewise, no specific estimate has been offered for forcing by mineral dust aerosol introduced into the atmosphere due to human activities beyond the assignment of a tentative range of -0.6 to $+0.4 \text{ W m}^{-2}$. The estimated forcing and associated uncertainty for each aerosol type is shown in relation to forcing estimates for the known GHGs along with an indication of the level of confidence in each of these estimates in Figure 4-42.

The relationship between perturbations to the Earth's radiative balance and climate is complicated by various feedbacks within the climate system. An example would be the positive feedback associated with melting sea ice. As sea ice melts with increasing surface temperatures, the dark ocean surface is revealed which absorbs, rather than reflects, solar radiation. Such a feedback increases the rate of surface warming. The role of feedbacks in determining the sensitivity of climate to changes in radiative forcing is described in detail by the IPCC in its third assessment report (IPCC, 2001a).

One possible feedback worth consideration in the discussion of the role of aerosols in climate may result from the possible sensitivity of aerosol number and mass to atmospheric temperature (Hemming and Seinfeld, 2001). Increasing atmospheric temperatures may result in a reduction of aerosol as semivolatile organic and inorganic aerosol constituents evaporate, leading to a change in aerosol forcing. As described in Chapter 2 of this document, and further discussed below, ambient aerosols are known to contain complex chemical mixtures of both scattering and absorbing materials. The feedback that may result from this phenomenon will depend upon whether the aerosols become more absorbing or more reflective with the loss of semivolatile material. Research is currently underway to evaluate both the role of temperature in determining aerosol mass and in defining the link between air quality and climate, but no literature yet exists to support an assessment of these effects. The following discussion,




Figure 4-42. Estimated global mean radiative forcing exerted by gas and various particle phase species for the year 2000, relative to 1750.

Source: IPCC (2001a).

therefore, will focus solely on the relative forcing properties of aerosols, for which a body of scientific research is available for consideration.

The physical and chemical properties of atmospheric aerosols, and their regional distribution and temporal nature, all play a role in determining the degree to which they force climate. These details are described, below, followed by a description of the “indirect” effect of aerosols due to changes in cloud properties, and then by brief discussions of the sources of uncertainty in determining aerosol-related climate forcing and of the link to human health and the environment.

The Physics of Particulate Matter and Its Climate Effects

Four wavelength-dependent physical properties of ambient aerosols are needed to calculate their optical depth and, thus, their radiative properties: the mass light-scattering efficiency (a_{sp}), the functional dependence of light-scattering on relative humidity $f(\text{RH})$, the single-scattering albedo (ω_0), and the scattering asymmetry parameter (g); (Charlson et al., 1992; Penner et al., 1994). Direct forcing by aerosols is especially sensitive to single-scattering albedo. Depending on the color of the underlying surface (i.e. the surface albedo) small changes in ω_0 can change the sign of the calculated aerosol forcing (Hansen et al., 1997).

The wavelength-dependent phase function and scattering and absorption coefficients are calculated using Mie scattering theory. These values are then used to calculate mass light-scattering efficiency, single-scattering albedo, and the asymmetry factor. Mie calculations require the ratio of particle size versus wavelength and the complex refractive index of the particle — a composition-dependent property (Salby, 1996). The relationship between light-scattering and relative humidity likewise depends upon composition, as water absorption depends upon the presence of hygroscopic material within the particle. Therefore, according to the current understanding of aerosol optics, good information about composition and size distribution is needed to successfully predict aerosol-related forcing.

The Chemistry of Particulate Matter and Its Climate Effects

Although forcing estimates are reported for specific aerosol classes (i.e., sulfate, black carbon, dust, etc.) it is understood that shortly after emission, primary aerosols undergo chemical transformation in the atmosphere. These transformations occur through the partitioning of gas phase compounds, by coagulation with another aerosol, or through a combination of both processes (Seinfeld and Pandis, 1998). While the radiative properties of nonabsorbing aerosol, such as sulfates or nitrates, are primarily sensitive to particle size, the radiative properties of aerosols containing absorbing constituents (i.e., black carbon and mineral aerosols) are also sensitive to chemical composition and mixing state. Studies of refractive index changes as a function of composition have shown that the type of mixing present both in the population of aerosols (i.e., “internal” versus “external” mixtures) and the extent of mixing within individual aerosols influence aerosol optical properties when absorbing material is present (Fuller et al., 1999). Modeled estimates for radiative forcing by black carbon-containing aerosols do, in fact,

range widely depending on whether the aerosol population is assumed to be internally or externally mixed and whether the absorbing carbon is uniformly distributed in the particle or whether it exists as a core surrounded by nonabsorbing material. The IPCC Third Assessment Report (2001a) provides a tabulation of studies and their forcing estimates for black carbon as it exists in different mixing states. For example, Haywood and Shine (1995) calculated that externally mixed fossil-fuel black carbon forces climate by $+0.2 \text{ W m}^{-2}$. Jacobson (2000) assumed that if black carbon exists as a solid core contained within an otherwise nonabsorbing droplet, the global forcing will be $+0.54 \text{ W m}^{-2}$.

Chemically and physically detailed models at high spatial resolution are required to accurately represent the chemical transformation and size evolution of aerosol within the ambient atmosphere. Such models, however, are impractical for global scale climate modeling. Parameterizations are formulated to represent processes occurring at spatial and temporal scales that are too fine for climate models (i.e., “sub-grid” processes). Several important fundamental chemical and physical processes, however, are not yet well-enough defined to assure reasonable parameterization in large scale climate models. Prediction of the organic and black carbon content of ambient aerosols and their associated radiative properties, as described above, and the aerosol-induced changes in cloud properties, remain especially weak.

Modeling of the effect of black carbon aerosols on climate, to date, has been done on the basis of limited and poor quality data regarding total global emissions, aerosol composition, and the mixing state of ambient aerosols. Appendix 2B.2 of Chapter 2 in this document describes the many problems associated with the most commonly used organic carbon/elemental carbon measurement method (thermal-optical reflectance and transmittance). For example, elemental carbon comprises a relatively small fraction of total carbon in biomass burning-related aerosol which is, nevertheless, an important source of black carbon in the atmosphere due to the extensive use of biofuels in developing countries and to the prevalence of natural and agriculture-related open biomass burning (Ludwig et al., 2003). The concentrations of elemental carbon in these aerosols are close to the detection limits of thermal-optical measurement systems, adding a high degree of uncertainty to the reported values. Thermal-optical measurements also do not provide information about the specific chemical composition of carbonaceous aerosol that is needed to calculate refractive indices. Furthermore, the technique is

based on filter collection, which only provides an average concentration of organic and elemental carbon and no information about mixing states.

The magnitude of the errors introduced into estimates of black carbon-related radiative forcing due to the lack of detailed information about optical properties of ambient black carbon aerosol cannot yet be estimated. The forcing estimate of $+0.2 \text{ W m}^{-2}$ provided by the IPCC is based upon a summary of the available modeling estimates, some of which are founded on simplistic assumptions about the composition and mixing state of ambient BC aerosol.

New methods, such as photoacoustic spectroscopy, hold promise for detecting quantities of light absorbing materials in ambient aerosol streams (Arnott et al., 1999; Moosmuller et al., 2001). The photoacoustic method is capable of identifying the absolute fraction of a given ambient aerosol sample that absorbs radiation at a given wavelength at high temporal resolution. The method does not alter the composition of the aerosol sample through heating and oxidation, thus introduces no chemical artifacts. Present photoacoustic instruments detect absorption at wavelengths of interest for visibility studies, but the measurement principle can be adapted to climate studies through the selection of climate-forcing relevant wavelengths. Specific chemical speciation at the organic and inorganic compound level coupled with photoacoustic measurements of the light absorbing properties of ambient aerosol would provide the best available test of atmospheric chemistry models attempting to simulate the evolution in aerosol radiative properties between emission and atmospheric removal.

The Regional and Temporal Dependence of Aerosol Forcing

In addition to information about the composition and size distributions of ambient aerosol, details regarding atmospheric concentrations on a spatial and temporal basis are needed to estimate climate forcing. Unlike the long-lived GHGs, aerosols have lifetimes averaging only days to weeks or less, leading to inhomogeneity in regional and global-scale concentrations. Industrial processes and other human activities that produce air pollution aerosols also vary on a seasonal, monthly, weekly, or diurnal basis, introducing other complications into forcing estimates (IPCC, 2001a). Peak aerosol concentrations, along with the greatest temporal variation, exist near emissions sources.

Deviations from global mean forcing estimates can be very large on the regional scale. For instance, Tegen et al. (1996) found that local radiative forcing exerted by dust raised from

disturbed lands ranges from -2.1 W m^{-2} to $+5.5 \text{ W m}^{-2}$ over desert areas and their adjacent seas. The largest regional values of radiative forcing caused by anthropogenic sulfate are about -3 W m^{-2} in the eastern United States, south central Europe, and eastern China (Kiehl and Briegleb, 1993). These regional maxima in aerosol forcing are at least a factor of 10 greater than their global mean values shown in Figure 4-41. By comparison, regional maxima in forcing by the well-mixed GHGs are only about 50% greater than their global mean value (Kiehl and Briegleb, 1993). Thus, the estimates of local radiative forcing by particles also are large enough to completely cancel the effects of GHGs in many regions and to cause a number of changes in the dynamic structure of the atmosphere that still need to be evaluated. A number of anthropogenic pollutants whose distributions are highly variable are also effective greenhouse absorbers. These gases include O_3 and, possibly HNO_3 , C_2H_4 , NH_3 , and SO_2 , all of which are not commonly considered in radiative forcing calculations (Wang et al., 1976). High-ozone values are found downwind of urban areas and areas where there is biomass burning. However, Van Dorland et al. (1997) found that there may not be much cancellation between the radiative effects for ozone and for sulfate because both species have different seasonal cycles and show significant differences in their spatial distribution.

“Indirect” Effects of Particulate Matter on Climate

Aerosols directly affect climate by scattering and absorbing solar and terrestrial radiation. Depending on chemical composition, they can also nucleate new cloud droplets. For a given total liquid water content (LWC), increasing cloud droplet number means smaller droplets that scatter solar radiation more effectively, reduce the amount of precipitation from the cloud, and, consequently, increase cloud lifetime. Observational evidence exists for this impact of hygroscopic aerosol on both warm and ice cloud properties. Increased albedo and increased cloud lifetime effects are treated separately by the climate modeling community. Particle-induced increases in cloud albedo are referred to as the “first” indirect effect, and changes to the cloud lifetime due to reduced precipitation are referred to as the “second” indirect effect. The highly uncertain estimated forcing due to the effects of particles on cloud albedo is given as a range from 0 to $+2.0 \text{ W m}^{-2}$ (IPCC, 2001a). A further effect, referred to as the “semi-indirect effect” is the reduction in cloud reflectivity due to the inclusion of black carbon-containing aerosol within the cloud drops themselves or as interstitial aerosol (Hansen et al., 2000).

Sulfate aerosols, especially those > 50 nm in diameter, are believed to alter clouds to the largest extent due to their efficient nucleation of cloud drops and ice crystals (Twomey, 1974). Organic aerosols that contain highly oxidized carbon compounds may be similarly efficient in nucleating cloud droplets (Novakov and Penner, 1993). Both satellite and in-situ aircraft observations reinforce the hypothesis that pollutant aerosols increase cloud reflectivity and lifetime (Ramanathan et al., 2001).

An important consequence of this property of aerosols on regional climate includes suppression of rain over polluted regions. Satellite observations show that precipitation occurs only outside of pollution tracks, while clouds within pollution tracks show a reduction in effective cloud drop radius to below the precipitation threshold (Rosenfeld, 1999). Desert dust also appears to alter the microphysical properties of clouds, suppressing precipitation from warm clouds while nucleating ice crystals in cold clouds.

While climate models are not yet equipped for modeling the effect of aerosols on regional and global hydrological cycles, it has been proposed that aerosols will reduce precipitation efficiency over land where anthropogenic activities provide a major source of cloud condensation nuclei. Several studies using coupled ocean-atmosphere general circulation models support the possibility of a “spin down” effect upon hydrological cycles due only to a reduction in surface radiation receipts from sulfate aerosol scattering. When indirect effects are included, the reduction in precipitation rates from clouds is large enough to reverse the effect of GHG-related forcing (Ramanathan et al., 2001).

Sources of Uncertainty in Aerosol Forcing Estimates

Uncertainties in calculating the direct effect of airborne particles arise from a lack of knowledge concerning their vertical and horizontal variability, their size distribution, chemical composition, and the distribution of components within individual particles. For instance, gas-phase sulfur species may be oxidized to form a layer of sulfate around existing particles in continental environments or they may be incorporated in sea-salt particles (e.g., Li-Jones and Prospero, 1998). In either case, the radiative effects of a given mass of the sulfate will be much lower than if pure sulfate particles were formed. It must also be stressed that the overall radiative effect of particles at a given location is not simply determined by the sum of effects

caused by individual classes of particles because of radiative interactions between particles with different radiative characteristics and with gases.

That calculations of the indirect effects of particles on climate are subject to much larger uncertainties than are calculations of their direct effects reflects uncertainties in a large number of chemical and microphysical processes that determine aerosol chemistry, size distribution, and the number of droplets within a cloud. A complete assessment of the radiative effects of PM will require computationally intensive calculations that incorporate the spatial and temporal behavior of particles of varying composition that have been emitted from, or formed from precursors emitted from, different sources. Refining the values of model-input parameters (such as improving emissions estimates) may be as important as improving the models per se in calculations of direct radiative forcing (Pan et al., 1997) and indirect radiative forcing (Pan et al., 1998). However, the uncertainties associated with the calculation of radiative effects of particles will likely remain much larger than those associated with well-mixed GHGs.

Aerosol-Related Climate Effects, Human Health, and the Environment

The specific impacts on human health and the environment due to aerosol effects on the climate system can not be calculated with confidence given the present difficulty in accurately modeling an aerosol's physical, chemical, and temporal properties and its regionally dependent atmospheric concentration levels, combined with the difficulties of projecting location-specific increases or decreases in anthropogenic emissions of atmospheric particles (or their precursors). However, substantial qualitative information, from observation and modeling, indicates that aerosol forces climate both positively and negatively, globally and regionally, and may be negatively impacting hydrological cycles on a regional scale. Global and other regional scale impacts are suspected on the basis of current, though uncertain, modeling studies suggesting that climate change in general can have positive and negative effects on human health, human welfare, and the environment.

4.5.2 Atmospheric Particulate Matter Effects on the Transmission of Solar Ultraviolet Radiation Transmission: Impacts on Human Health and the Environment

4.5.2.1 Potential Effects of Increased Ultraviolet Radiation Transmission

The transmission of solar UV-B radiation through the Earth's atmosphere is controlled by ozone, clouds, and particles. The depletion of stratospheric ozone caused by the release of anthropogenically produced chlorine (Cl)- and bromine (Br)-containing compounds has resulted in heightened concern about potentially serious increases in the amount of solar UV-B radiation (SUVB) reaching the Earth's surface. SUVB is also responsible for initiating the production of hydroxyl radicals ($\cdot\text{OH}$) that oxidize a wide variety of volatile organic compounds, some of which can deplete stratospheric ozone (e.g., CH_3Cl , CH_3Br), absorb terrestrial infrared radiation (e.g., CH_4), and contribute to photochemical smog formation (e.g., C_2H_4 , C_5H_8).

Increased penetration of SUVB to the Earth's surface as the result of stratospheric ozone depletion continues to be of much concern because of projections of consequent increased surface-level SUVB exposure and associated potential negative impacts on human health, plant and animal biota, and man-made materials. Several summary overviews (Kripke, 1989; Grant, 1989; Kodama and Lee, 1994; Van der Leun et al., 1995, 1998) of salient points related to stratospheric ozone depletion and bases for concern provide a concise introduction to the subject. Only a brief summary will be given here. Stratospheric ozone depletion results from (a) anthropogenic emissions of certain trace gases having long atmospheric residence times, e.g., chlorofluorocarbons (CFCs), carbon tetrachloride (CCl_4), and Halon 1211 ($\text{CF}_2\text{Cl Br}$) and 1301 (CF_3Br) — which have atmospheric residence times of 75 to 100 years, 50 years, 25 years, and 110 years, respectively; (b) their tropospheric accumulation and gradual transport, over decades, up to the stratosphere, where (c) they photolyze to release Cl and Br that catalyze ozone destruction; leading to (d) stratospheric ozone depletion. Such ozone depletion is most marked over Antarctica during spring in the southern hemisphere, to a less marked but still significant extent over the Arctic polar region during late winter and spring in the northern hemisphere, and to a lesser extent, over midlatitude regions during any season. Given the long time frame involved in the transport of such gases to the stratosphere and their long residence times there, any effects already seen on stratospheric ozone are likely to have been caused by the atmospheric loadings of trace gases from anthropogenic emissions over the past few decades. Ozone-depleting gases already in the atmosphere will continue to affect stratospheric ozone

concentrations well into the 21st century. Shorter-lived gases, such as CH₃Br, also exert significant ozone depletion effects.

The main types of deleterious effects hypothesized as likely to result from stratospheric ozone depletion and consequent increased SUVB penetration through the Earth's atmosphere include the following:

- (1) *Direct Human Health Effects*, such as skin damage (sunburn), leading to more rapid aging and increased incidence of skin cancer; ocular effects (retinal damage and increased cataract formation, possibly leading to blindness); and suppression of some immune system components (contributing to skin cancer induction and spread to nonirradiated skin areas, as well as possibly increasing susceptibility to certain infectious diseases).
- (2) *Agricultural/Ecological Effects*, mediated largely through altered biogeochemical cycling resulting in consequent damaging impacts on terrestrial plants (leading to possible reduced yields of rice, other food crops, and commercially important trees, as well as to biodiversity shifts in natural terrestrial ecosystems); and deleterious effects on aquatic life (including reduced ocean zooplankton and phytoplankton as important base components of marine food-chains supporting the existence of commercially important, edible fish and other seafood, as well as to other aquatic ecosystem shifts).
- (3) *Indirect Human Health and Ecological Effects*, mediated through increased tropospheric ozone formation (and consequent exacerbation of surface-level, ozone-related health and ecological impacts) and alterations in the concentrations of other important trace species, most notably the hydroxyl radical and acidic aerosols.
- (4) *Other Types of Effects*, such as faster rates of polymer weathering because of increased UV-B radiation and other effects on man-made commercial materials and cultural artifacts, secondary to climate change or exacerbation of air pollution problems.

Extensive qualitative and quantitative characterizations of stratospheric ozone depletion processes and projections of their likely potential impacts on human health and the environment have been the subjects of periodic (1988, 1989, 1991, 1994, 1998) international assessments carried out under WMO and UNEP auspices since the 1987 signing of the Montreal Protocol on Substances that Deplete the Ozone Layer. For more detailed up-to-date information, the reader is referred to recent international assessments of (a) processes contributing to stratospheric ozone depletion and the status of progress towards ameliorating the problem (WMO, 1999) and (b) revised qualitative and quantitative projections of potential consequent human health and environmental effects (UNEP, 1998, 2000).

Of considerable importance is the growing recognition, as reflected in these newer assessments, of the impacts of enhanced solar radiation on biogeochemical cycles (see, for example, Zepp et al. [1998]). As noted in that paper, the effects of UV-B radiation (both in magnitude and direction) on trace gas (e.g., CO) emissions and mineral nutrient cycling are species specific and can affect a variety of processes. These include, for example, changes in the chemical composition of living plant tissue, photodegradation of dead plant matter (e.g., ground litter), release of CO from vegetation previously charred by fire, changes in microbial decomposer communities, and effects on nitrogen-fixing microorganisms and plants. In addition, changes in the amount and composition of organic matter, caused by enhanced UV-B penetration, affect the transmission of solar ultraviolet and visible radiation through the water column. These changes in light quality broadly impact the effects of UV-B on aquatic biogeochemical cycles. Enhanced UV-B levels cause both positive and negative effects on microbial activities in aquatic ecosystems that can affect nutrient cycling and the uptake or release of GHGs. Thus, there are emerging complex issues regarding interactions and feedbacks between climate change and changes in terrestrial and marine biogeochemical cycles because of increased UV-B penetration to the Earth's surface.

In contrast to the above types of negative impacts projected as likely to be associated with increased UV-B penetration to Earth's surface, some research results suggest possible beneficial effects of increased UV-B radiation. For example, a number of U.S. and international studies have focused on the protective effects of UV-B radiation with regard to the incidence of cancers other than skin cancer. One of the first of these studies investigated potential relationships between sunlight, vitamin D, and colon cancer (Garland and Garland, 1980). More recent studies continue to provide suggestive evidence that UV-B radiation may be protective against several types of cancer and some other diseases. For example, Grant (2002) conducted a number of ecologic-type epidemiologic studies, which suggest that UV-B radiation, acting through the production of vitamin D, is a risk-reduction factor for mortality due to several types of cancer, including cancer of the breast, colon, ovary, and prostate, as well as non-Hodgkin lymphoma. Other related studies that provide evidence for protective effects of UV-B radiation include Gorham et al. (1989, 1990); Garland et al. (1990); Hanchette and Schwartz (1992); Ainsleigh (1993); Lefkowitz and Garland (1994); Hartge et al. (1996); and Freedman et al. (1997).

As noted in the above detailed international assessments, since the signing of the Montreal Protocol, much progress has been made in reducing emissions of ozone depleting gases, leading to estimates that the maximum extent of stratospheric ozone depletion may have leveled off during recent years. This is expected to be followed by a gradual lessening of the problem and its impacts during the next half-century. However, the assessments also note that the modeled projections are subject to considerable uncertainty (see, for example, UNEP [2000]). Varying potential roles of atmospheric particles, discussed below, are among salient factors complicating predictive modeling efforts.

4.5.2.2 Airborne Particle Effects on Atmospheric Transmission of Solar Ultraviolet Radiation

A given amount of ozone in the lower troposphere has been shown to absorb more solar radiation than an equal amount of ozone in the stratosphere, because of the increase in its effective optical path produced by Rayleigh scattering in the lower atmosphere (Brühl and Crutzen, 1988). The effects of particles are more complex. The impact of particles on the SUVB flux throughout the boundary layer are highly sensitive to the altitude of the particles and to their single-scattering albedo. Even the sign of the effect can reverse as the composition of the particle mix changes from scattering to absorbing types (e.g., from sulfate to elemental carbon or PAHs) (Dickerson et al., 1997). In addition, scattering by particles also may increase the effective optical path of absorbing molecules, such as ozone, in the lower atmosphere.

The effects of particles present in the lower troposphere on the transmission of SUVB have been examined both by field measurements and by radiative transfer model calculations. The presence of particles in urban areas modifies the spectral distribution of solar irradiance at the surface. Shorter wavelength (i.e., ultraviolet) radiation is attenuated more than visible radiation (e.g., Peterson et al., 1978; Jacobson, 1999). Wenny et al. (1998) also found greater attenuation of SUVB than SUVA (315 to 400 nm). However, this effect depends on the nature of the specific particles involved and, therefore, is expected to depend strongly on location. Lorente et al. (1994) observed an attenuation of SUVB ranging from 14 to 37%, for solar zenith angles ranging from about 30° to about 60°, in the total (direct and diffuse) SUVB reaching the surface in Barcelona during cloudless conditions on very polluted days (aerosol scattering optical depth at 500 nm, $0.46 \lesssim \tau_{500 \text{ nm}} \lesssim 1.15$) compared to days on which the turbidity of urban air was similar to that for rural air ($\tau_{500 \text{ nm}} \lesssim 0.23$).

Particle concentrations that can account for these observations can be estimated roughly by combining Koschmieder's relation for expressing visual range in terms of extinction coefficient with one for expressing the mass of PM_{2.5} particles in terms of visual range (Stevens et al., 1984). By assuming a scale height (i.e., the height at which the concentration of a substance falls off to 1/e of its value at the surface) of 1 km for PM_{2.5}, an upper limit of 30 µg/m³ can be derived for the clear case and between 60 and 150 µg/m³ for the polluted case. Estupiñán et al. (1996) found that summertime haze under clear sky conditions attenuates SUVB between 5 and 23% for a solar zenith angle of 34° compared to a clear-sky day in autumn. Mims (1996) measured a decrease in SUVB by about 80% downwind of major biomass burning areas in Amazonia in 1995. This decrease in transmission corresponded to optical depths at 340 nm ranging from 3 to 4. Justus and Murphey (1994) found that SUVB reaching the surface decreased by about 10% because of changes in aerosol loading in Atlanta, GA from 1980 to 1984. In addition, higher particle levels in Germany (48 °N) may be responsible for the greater attenuation of SUVB than seen in New Zealand (Seckmeyer and McKenzie, 1992).

In a study of the effects of nonurban haze on SUVB transmission, Wenny et al. (1998) derived a very simple regression relation between the measured aerosol optical depth at 312 nm,

$$\ln(\text{SUVB solar noon}) = -0.1422 \tau_{312\text{nm}} - 0.138, R^2 = 0.90, \quad (4-13)$$

and the transmission of SUVB to the surface at solar noon. In principle, values of $\tau_{312\text{nm}}$ could be found from knowledge of the aerosol optical properties and visual range values. Wenny et al. (1998) also found that absorption by particles accounted for 7 to 25% of the total (scattering + absorption) extinction. Relations such as the one above are strongly dependent on local conditions and should not be used in other areas without knowledge of the differences in aerosol properties. Although all of the above studies reinforce the idea that particles play a major role in modulating the attenuation of SUVB, none included measurements of ambient PM concentrations, so direct relations between PM levels and SUVB transmission could not be determined.

Vuilleumier et al. (2001) concluded that variations in aerosol scattering and absorption were responsible for 97% of the variability in the optical depth measured at seven wavelengths from 300 to 360 nm at Riverside, CA from 1 July to 1 November 1997. Similar measurements

made at nearby Mt. Wilson, located above the main surface haze layer, showed that 80% of the variations in optical depth were still driven by variations in aerosol scattering and absorption. The remainder of the variability in optical depth was attributed mainly to variability in ozone under clear-sky conditions. However, these results cannot be extrapolated to other locations, because these effects are coupled and nonlinear and, thus, are not straightforward. They depend on the concentrations of these species and on the physical and chemical characteristics of the particles. Hence, any quantitative statements regarding the relative importance of particles and ozone will be location-specific.

Liu et al. (1991) roughly estimated the overall effects on atmospheric transmission of SUVB of increases of anthropogenic airborne particles that have occurred since the beginning of the industrial revolution. Based on (a) estimates of the reduction in visibility from about 95 km to about 20 km over nonurban areas in the eastern United States and in Europe, (b) calculations of optical properties of airborne particles found in rural areas to extrapolate the increase in extinction at 550 to 310 nm, and (c) radiative transfer model calculations, Liu et al. (1991) concluded that the amount of SUVB reaching Earth's surface likely has decreased from 5 to 18% since the beginning of the industrial revolution. This was attributed mainly to scattering of SUVB back to space by sulfate-containing particles. Radiative transfer model calculations have not been done for urban particles.

Although aerosols are expected to decrease the flux of SUVB reaching the surface, scattering by particles is expected to result in an increase in the actinic flux within and above the aerosol layer. However, when the particles significantly absorb SUVB, a decrease in the actinic flux is expected. Actinic flux is the radiant energy integrated over all directions at a given wavelength incident on a point in the atmosphere and is the quantity needed to calculate rates of photolytic reactions in the atmosphere. Blackburn et al. (1992) measured attenuation of the photolysis rate of ozone and found that aerosol optical depths near unity at 500 nm reduced ozone photolysis rate by as much as a factor of two. Dickerson et al. (1997) showed that the photolysis rate for NO_2 , a key parameter for calculating the overall intensity of photochemical activity, could be increased within and above a scattering aerosol layer extending from the surface, although it would be decreased at the surface. This effect is qualitatively similar to what is seen in clouds, where photolysis rates are increased in the upper layers of a cloud and above the cloud (Madronich, 1987). For a simulation of an ozone episode that occurred during July

1995 in the Mid-Atlantic region, Dickerson et al. (1997) calculated ozone increases of up to 20 ppb compared to cases that did not include the radiative effects of particles in urban airshed model (UAM-IV) simulations. In contrast, Jacobson (1998) found that particles may have caused a 5 to 8% decrease in O₃ levels during the Southern California Air Quality Study in 1987. Absorption by organic compounds and nitrated inorganic compounds was hypothesized to account for the reductions in UV radiation intensity.

The photolysis of ozone in the Hartley bands also leads to production of electronically excited oxygen atoms, O(¹D), that then react with water vapor to form hydroxyl radicals. Thus, enhanced photochemical production of ozone is accompanied by the scavenging of species involved in greenhouse warming and stratospheric depletion. However, these effects may be neutralized, or even reversed, by the presence of absorbing material in the particles. Any evaluation of the effects of particles on photochemical activity, therefore, will depend on the composition of the particles and will also be location-specific.

Further complicating any straightforward evaluation of UV-B penetration to specific areas of the Earth's surface are the influences of clouds, as discussed by Erlick et al. (1998), Frederick et al. (1998), and Soulen and Fredrick (1999). The transmission of solar UV and visible radiation is highly sensitive to cloud type, cloud amount, and the extent of their external or internal mixing with cloud droplets. Even in situations of very low atmospheric PM (e.g., over Antarctica), interannual variations in cloudiness over specific areas can be as important as ozone levels in determining ultraviolet surface irradiation, with net impacts varying monthly to seasonally (Soulen and Fredrick, 1999). Evaluations of the effects of changes in the transmission of solar UV-B radiation to the surface have been performed usually for cloud-free or constant cloudiness conditions.

Given the above considerations, the quantification of projected effects of variations in atmospheric PM on human health or on the environment related to the effects of particles on the transmission of solar UV-B radiation would require location-specific evaluations that take into account (a) composition, concentration, and internal structure of the particles; (b) temporal variations in atmospheric mixing height and depths of layers containing the particles; and (c) the abundance of ozone and other absorbers within the planetary boundary layer and the free troposphere. The outcome of such modeling effects would likely vary from location to location in terms of increased or decreased surface level UV-B exposures because of location-specific

changes in atmospheric PM concentrations or composition. For example, to the extent that any location-specific scattering by airborne PM were to affect the directional characteristics of ultraviolet radiation at ground level, and thereby enhance radiation incident from low angles (Dickerson et al., 1997), the biological effectiveness (whether deleterious or beneficial) of resulting ground-level UV-B exposures could be enhanced. Airborne PM also can reduce the ground-level ratio of photorepairing radiation (UV-A and short-wavelength visible) to damaging UV-B radiation. Lastly, PM deposition is a major source of PAHs in certain freshwater lakes and coastal areas; and the adverse effects of solar UV are enhanced by the uptake of PAHs by aquatic organisms. Thus, although airborne PM may, in general, tend to reduce ground-level UV-B, its net effect in some locations may be to increase UV damage to certain aquatic and terrestrial organisms, as discussed by Cullen and Neale (1997).

4.6 SUMMARY AND KEY CONCLUSIONS

4.6.1 Particulate Matter Effects on Vegetation and Ecosystems

The first section of this chapter assessed and characterized the overall ecological integrity and indicated the status of ecosystems within the United States affected by the deposition of the anthropogenic stressors associated with PM. There are six essential environmental attributes (EEAs) — landscape condition, biotic condition, and chemical/physical characteristics, and ecological processes, hydrology/geomorphology, and natural disturbance regimes that can be used to provide a hierarchical framework for characterizing ecosystem status. The first three can be separated into “patterns” and the last three into “processes.” The ecological processes create and maintain the ecosystem elements in patterns. The patterns, in turn, affect how the ecosystem processes are expressed. Patterns at the higher level of biological organization emerge from the interactions and selection processes at localized levels. Changes in patterns or processes result in changes in the status and functioning of an ecosystem. The relationships among the EEAs are complex because all are interrelated (i.e., changes in one EEA may affect, directly or indirectly, every other EEA).

Human existence on Earth depends on the life-support services provided by the interaction of the different EEAs. Ecosystem processes and patterns provide the functions that maintain clean water, clean air, a vegetated earth, and a balance of organisms that enable humans to

survive. The benefits they impart include the absorption and breakdown of pollutants, cycling of nutrients, binding of soil, degradation of organic waste, maintenance of a balance of gases in the air, regulation of radiation balance, climate, and fixation of solar energy. Concern has arisen in recent years regarding biodiversity and the integrity of ecosystems. Because the activities of few ecosystems on Earth today are not influenced by humans, understanding the changes in biodiversity and nutrient cycling resulting from anthropogenic PM deposition are of great importance.

The PM whose effects on vegetation and ecosystems is discussed in this section is not a single pollutant but represents a heterogeneous mixture of particles differing in origin, size, and chemical constituents. The effects of exposure to a given mass concentration of PM of particular size may, depending on the particular mix of deposited particles, lead to widely differing phytotoxic responses.

The deposition of PM onto vegetation and soil, depending on its chemical composition (acid/base, trace metal, or nutrients, e.g., nitrates or sulfates), can produce direct or indirect responses within an ecosystem. The ecosystem response to pollutant deposition is a direct function of the level of sensitivity of the ecosystem and its ability to ameliorate resulting change. Changes in ecosystem structural patterns and the functioning of ecological processes, must be scaled in both time and space and propagated to the more complex levels of community interaction to produce observable ecosystem changes.

The stressors of greatest environmental significance are particulate nitrates and sulfates whose indirect effects occur primarily via their deposition onto the soil. Upon entering the soil environment, they can alter the ecological processes of energy flow and nutrient cycling, inhibit nutrient uptake, change ecosystem structure, and affect ecosystem biodiversity. The soil environment is one of the most dynamic sites of biological interaction in nature. It is inhabited by microbial communities of bacteria, fungi, and actinomycetes. Bacteria are essential participants in the nitrogen and sulfur cycles that make these elements available for plant uptake. Fungi in association with plant roots form mycorrhizae, a symbiotic relationship that is integral in mediating plant uptake of mineral nutrients. Changes in the soil environment that influence the role of the bacteria in nutrient cycling and fungi in making minerals available for plant utilization, determine plant and ultimately ecosystem response.

The effects on the growth of plants resulting from the deposition of nitrates and sulfates and the acidifying effect of the associated H^+ ion in wet and dry deposition are the most important environmentally. Nitrogen is of overriding importance in plant metabolism and, to a large extent, governs the utilization of phosphorus, potassium, and other nutrients. Typically, the availability of nitrogen via the nitrogen cycle controls net primary productivity, and possibly, the decomposition rate of plant litter. Plants usually obtain nitrogen directly from the soil by absorbing NH_4^+ or NO_3^- through their roots, or it is formed by symbiotic organisms in their roots. Plants vary in their ability to absorb ammonium and nitrate from the soil.

Although nitrogen as molecular nitrogen (N_2) is the most abundant element in the atmosphere, it only becomes available to support the growth of plants after it is converted into reactive forms. In nature, nitrogen may be divided into two groups: reactive (Nr) and nonreactive (N_2). Reactive Nr includes the inorganic reduced forms of nitrogen (e.g., ammonia [NH_3] and ammonium [NH_4^+]), inorganic oxidized forms (e.g., nitrogen oxides [NO_x], nitric acid [HNO_3], nitrous oxide [N_2O], and nitrate [NO_3^-]), and organic compounds (e.g., urea, amine, proteins, and nucleic acids).

Reactive nitrogen is now accumulating in the environment on all spatial scales — local, regional, and global. The three main causes of the increase in global Nr is the result of the (1) widespread cultivation of legumes, rice, and other crops that promote conversion of N_2 to organic nitrogen through biological nitrogen fixation; (2) combustion of fossil fuels, which converts both atmospheric N_2 and fossil nitrogen to reactive NO_x ; and (3) the Haber-Bosch process, which converts nonreactive NH_3 to sustain food production and some industrial activities. The major changes in the nitrogen cycle due to the cited causes can be both beneficial and detrimental to the health and welfare of humans and ecosystems.

Reactive nitrogen can be widely dispersed and accumulate in the environment when the rates of its formation exceed the rates of removal via denitrification. Reactive nitrogen creation and accumulation is projected to increase as per capita use of resources by human populations increases. The cascade of environmental effects of increases in Nr include the following: (1) production of tropospheric ozone and aerosols that induce human health problems; (2) increases in the productivity in forests and grasslands followed by decreases wherever deposition increases significantly and exceeds critical loads; Nr additions probably also decrease biodiversity in many natural habitats; (3) acidification and loss of biodiversity in lakes and

streams in many regions of the world in association with sulfur; (4) eutrophication (now considered the biggest pollution problem in coastal waters), hypoxia, loss of biodiversity, and habitat degradation in coastal ecosystems; and (5) contributes to global climate change and stratospheric ozone depletion, which can in turn affect ecosystems and human health (see Table 4-21).

TABLE 4-21. EFFECTS OF REACTIVE NITROGEN

<p><i>Direct effects of Nr on ecosystems include:</i></p> <ul style="list-style-type: none"> – Increased productivity of Nr-limited natural ecosystems. – Ozone-induced injury to crop, forest, and natural ecosystems and predisposition to attack by pathogens and insects. – Acidification and eutrophication effects on forests, soils, and freshwater aquatic ecosystems. – Eutrophication and hypoxia in coastal ecosystems. – N-saturation of soils in forests and other natural ecosystems. – Biodiversity losses in terrestrial and aquatic ecosystems and invasions by N-loving weeds. – Changes in abundance of beneficial soil organisms that alter ecosystem functions.
<p><i>Indirect effects of Nr on other societal values include:</i></p> <ul style="list-style-type: none"> – Increased wealth and well being of human populations in many parts of the world. – Significant changes in patterns of land use. – Regional hazes that decrease visibility at scenic vistas and airports. – Depletion of stratospheric ozone by N₂O emissions. – Global climate change induced by emissions of N₂O and formation of tropospheric ozone. – Damage to useful materials and cultural artifacts by ozone, other oxidants, and acid deposition. – Long-distance transport of Nr, which causes harmful effects in countries distant from emission sources and/or increased background concentrations of ozone and fine PM.
<p><i>In addition to these effects, it is important to recognize that:</i></p> <ul style="list-style-type: none"> – The magnitude of Nr flux often determines whether effects are beneficial or detrimental. – All of these effects are linked by biogeochemical circulation pathways of Nr. – Nr is easily transformed among reduced and oxidized forms in many systems. Nr is easily distributed by hydrologic and atmospheric transport processes.

Changes in nitrogen supply can have a considerable effect on an ecosystem's nutrient balance. Large chronic additions of nitrogen influence normal nutrient cycling and alter many plant and soil processes involved in nitrogen cycling. "Nitrogen saturation" results when Nr concentrations exceed the capacity of a system to use it. Saturation implies that some resource other than nitrogen is limiting biotic function. Water and phosphorus for plants and carbon for microorganisms are most likely to be secondary limiting factors. The appearance of nitrogen in

soil solution is an early symptom of excess nitrogen. In the final stage, disruption of ecosystem structure becomes apparent.

Possible ecosystem responses to nitrogen saturation include (1) permanently increased foliar nitrogen and reduced foliar phosphorus and lignin caused by the lower availability of carbon, phosphorus, and water; (2) reduced productivity in conifer stands because of disruptions of physiological function; (3) decreased root biomass and increased nitrification and nitrate leaching; (4) reduced soil fertility, resulting from increased cation leaching, increased nitrate and aluminum concentrations in streams; (5) decreased water quality; and (6) changes in soil microbial communities from predominantly fungal (mycorrhizal) communities to those dominated by bacteria during saturation.

Although soils of most North American forest ecosystems are nitrogen-limited, some exhibit severe symptoms of nitrogen saturation (See Table 4-14). They include the high-elevation, nonaggrading spruce-fir ecosystems in the Appalachian Mountains; the eastern hardwood watersheds at the Fernow Experimental Forest near Parsons, WV; forests in southern California, the southwestern Sierra Nevada in Central California; and the Front Range in northern Colorado. The mixed conifer forest and chaparral watershed with high smog exposure in the Los Angeles Air Basin exhibit the highest stream water NO_3^- concentrations for wildlands in North America.

Increases in soil nitrogen play a selective role in ecosystems. Plants adapted to living in an environment of low nitrogen availability will be replaced by nitrophilic plants capable of using increased nitrogen because they have a competitive advantage when nitrogen becomes more readily available. Plant succession patterns and biodiversity are affected significantly by chronic nitrogen additions in some North American ecosystems. Long-term nitrogen fertilization studies in both New England and Europe suggest that some forests receiving chronic inputs of nitrogen may decline in productivity and experience greater mortality. Declining coniferous forest stands with slow nitrogen cycling may be replaced by deciduous fast-growing forests that cycle nitrogen.

Linked to the nitrogen cascade is the deposition of Nr and sulfates and the associated hydrogen ion in acidic precipitation, a critical environmental stress that affects forest landscapes and aquatic ecosystems in North America, Europe, and Asia. Composed of ions, gases, and particles derived from gaseous emissions of sulfur dioxide (SO_2), nitrogen oxides (NO_x),

ammonia (NH₃) and particulate emissions of acidifying and neutralizing compounds, acidic precipitation varies highly across time and space. Its deposition and the resulting soil acidity can lead to nutrient deficiencies and to high aluminum-to-nutrient ratios that limit uptake of calcium and magnesium and create a nutrient deficiency. Aluminum accumulation in root tissue can reduce calcium uptake and causes Ca²⁺ deficiencies. Tree species can be adversely affected if altered Ca:Al ratios impair the uptake of calcium or magnesium. Calcium is essential in the formation of wood and the maintenance of the primary plant tissues necessary for tree growth. Studies suggest that the decline of red spruce stands in Vermont may be related to the Ca:Al ratio (Johnson and Lindberg, 1992b).

The evidence of the effects of wet and dry particulate deposition SO₄⁻ and Nr species on nutrient cycling in forest ecosystems is provided by the IFS. The deposition data from the study illustrates several important aspects of the atmospheric exposure characteristics across a wide elevational gradient and over a wide spatial scale. Atmospheric deposition plays a significant role in the biogeochemical cycles at all IFS sites, but is most important in the east at high-elevation sites. The flux of the sulfate ion, Nr compounds, and H⁺ ions in throughfall at all sites is dominated by atmospheric deposition. Atmospheric deposition may have significantly affected the nutrient status of some IFS sites through the mobilization of aluminum by impeding cation uptake. Nitrates and sulfate are the dominant anions in the Smokies. Pulses of nitrates are the major causes of aluminum pulses in soil solutions. However, the connection between aluminum mobilization and forest decline is not clear, hence, aluminum mobilization presents a situation worthy of further study.

Notable impacts of excess nitrogen deposition also have been observed with regard to aquatic systems. For example, atmospheric nitrogen deposition into soils in watershed areas feeding into estuarine sound complexes (e.g., the Pamlico Sound of North Carolina) appear to contribute to excess nitrogen flows in runoff (especially during and after heavy rainfall events such as hurricanes) from agricultural practices or other uses (e.g., fertilization of lawns or gardens). Massive influxes of such nitrogen into watersheds and sounds can lead to dramatic decreases in water oxygen and increases in algae blooms that can cause extensive fish kills and damage to commercial fish and sea food harvesting.

The critical loads concept is useful for estimating the amounts of pollutants (e.g., Nr and acidic precipitation) that sensitive ecosystems can absorb on a sustained basis without

experiencing measurable degradation. The estimation of ecosystem critical loads requires an understanding of how an ecosystem will respond to different loading rates in the long term and can be of special value for ecosystems receiving chronic deposition of Nr and sulfur independently and as acidic deposition when in combination.

Changes in the soil can result from the deposition of heavy metals. Exposures to heavy metals are highly variable, depending whether deposition is by wet or dry processes. Few (e.g., Cu, Ni, Zn) have been documented to have direct phytotoxicity under field conditions. Exposure to coarse particles of natural origin and elements such as Fe and Mg are more likely to occur via dry deposition, while fine particles of atmospheric origin are more likely to contain elements such as Ca, Cr, Pb, N, and V. Ecosystems immediately downwind of major emissions sources such as power generating, industrial, or urban complexes can receive locally heavy inputs. By negatively affecting litter decomposition, heavy-metal accumulation presents the greatest potential for influencing nutrient cycling. Microbial populations decreased and logarithmic rates of microbial increase were prolonged as a result of cadmium toxicity. Additionally, the presence of Ca, Cu, and Ni were observed to impair the symbiotic activity of fungi, bacteria, and actinomycetes.

Phytochelatin produced by plants as a response to sublethal concentrations of heavy metals are indicators of metal stress and can be used to indicate that heavy metals are involved in forest decline. The increasing concentrations of phytochelatin across regions and at greater altitudes associated with greater levels of forest injury implicate them in forest decline.

The ambient concentration of particles, the parameter for which there is most data (Chapter 3), is, at best, an indicator of exposure. The amount of PM entering the immediate plant environment and deposited onto the plant surfaces or soil in the vicinity of the roots, determines the biological effect. Three major routes are involved during the wet and dry deposition processes: (1) precipitation scavenging in which particles are deposited in rain and snow; (2) occult (fog, cloud water, and mist interception); and (3) dry deposition, a much slower, yet more continuous removal to plant surfaces.

Deposition of PM on the surfaces of above-ground plant parts can have either a physical and/or a chemical effect. Particles transferred from the atmosphere to plant surfaces may cause direct effects if they (1) reside on the leaf, twig, or bark surface for an extended period; (2) are taken up through the leaf surface; or (3) are removed from the plant via resuspension to the

atmosphere, washing by rainfall, or litter-fall with subsequent transfer to the soil. Ecosystem effects have been observed in the neighborhoods of limestone quarries.

Secondary organics formed in the atmosphere have been variously subsumed under the following terms: toxic substances, pesticides, HAPS, air toxics, SVOCs, and POPs. The substances alluded to under the above headings are controlled under CAA Sect.112, Hazardous Air Pollutants, not as criteria pollutants controlled by NAAQS under CAA Sections 108 and 109 (U.S. Code, 1991). Their possible effects on humans and ecosystems are discussed in many other government documents and publications. They are noted in this chapter because, in the atmosphere, many of the chemical compounds are partitioned between gas and particle phases and are deposited as PM. As particles, they become airborne and can be distributed over a wide area and affect remote ecosystems. Some of the chemical compounds are of concern to humans because they may reach toxic levels in the food chains of both animals and humans, whereas others tend to decrease or maintain the same toxicity as they move through food chains.

An important characteristic of fine particles is their ability to affect the flux of solar radiation passing through the atmosphere directly, by scattering and absorbing solar radiation, and, indirectly, by acting as cloud condensation nuclei that, in turn, influence the optical properties of clouds. Regional haze has been estimated to diminish surface solar visible radiation by approximately 8%. Crop yields have been reported as being sensitive to the amount of sunlight received, and crop losses have been attributed to increased airborne particle levels in some areas of the world.

Key conclusions and findings about PM-related effects on vegetation and ecosystems include the following:

- A number of ecosystem-level conditions (e.g., nitrogen saturation, terrestrial and aquatic acidification, coastal eutrophication) that can lead to negative impacts on human health and welfare have been associated with chronic, long-term exposure of ecosystems to elevated inputs of compounds containing Nr, sulfur and/or associated hydrogen ions.
- Some percentage of total ecosystem inputs of these chemicals is contributed by deposition of atmospheric particles, although the percentage greatly varies temporally and geographically and has not generally been well quantified.
- Unfortunately, our ability to relate ambient concentrations of PM to ecosystem response is hampered by the following significant data gaps and uncertainties:

- The lack of a long-term, historic database of annual speciated PM deposition rates in sensitive ecosystems precludes establishing relationships between PM deposition (exposure) and ecosystem response at this time. Except for data from the IMPROVE network and from some CASTNet sites, much of the PM monitoring effort has focused on urban or near urban areas, rather than on sensitive ecosystems.
- Modeled deposition rates, used in the absence of monitored data, can be highly uncertain, since there are a multitude of factors that influence the amounts of PM that get deposited from the air onto sensitive receptors. These factors include the mode of deposition (wet, dry, occult), windspeed, surface roughness/stickiness, elevation, particle characteristics (e.g., size, shape, chemical composition, etc.), and relative humidity.
- Each ecosystem is unique from all others, since each has developed within a context framed by the topography, underlying bedrock, soils, climate, meteorology, hydrologic regime, natural and land use history, species associations that co-occur at that location (i.e., soil organisms, plants, etc.), and successional stage. Because of this variety, and insufficient baseline data on each of these features for most ecosystems, it is currently impossible to extrapolate with much confidence any effect from one ecosystem to another, or to predict an appropriate “critical load.” Thus, a given PM deposition rate or load of nitrates in one ecosystem may produce entirely different responses than the same deposition rate at another location.
- There remain large uncertainties associated with the length of residence time of Nr in a particular ecosystem component or reservoir, and thus, its impact on the ecosystem as it moves through the various levels of the N cascade.
- As additional PM speciated air quality and deposition monitoring data become available, there is much room for fruitful research into the areas of uncertainty identified above.

4.6.2 Particulate Matter-Related Effects on Visibility

Visibility is defined as the degree to which the atmosphere is transparent to visible light and the clarity and color fidelity of the atmosphere. Visual range is the farthest distance a black object can be distinguished against the horizontal sky. Visibility impairment is any humanly perceptible change in visibility. For regulatory purposes, visibility impairment, characterized by light extinction, visual range, contrast, and coloration, is classified into two principal forms: (1) “reasonably attributable” impairment, attributable to a single source or small group of sources, and (2) regional haze, any perceivable change in visibility caused by a combination of many sources over a wide geographical area.

Visibility is measured by human observation, light scattering by particles, the light extinction-coefficient, and parameters related to the light-extinction coefficient (visual range and deciview scale), and fine PM mass concentrations.

The air quality within a sight path will affect the illumination of the sight path by scattering or absorbing solar radiation before it reaches the Earth's surface. The rate of energy loss with distance from a beam of light is the light extinction coefficient. The light extinction coefficient is the sum of the coefficients for light absorption by gases (σ_{ag}), light scattering by gases (σ_{sg}), light absorption by particles (σ_{ap}), and light scattering by particles (σ_{sp}). Corresponding coefficients for light scattering and absorption by fine and coarse particles are σ_{sfp} and σ_{afp} , and σ_{scp} and σ_{acp} , respectively. Visibility within a sight path longer than approximately 100 km (60 mi) is affected by the change in the optical properties of the atmosphere over the length of the sight path.

Visual range was developed for, and continues to be used as, an aid in military operations and to a lesser degree in transportation safety. Visual range is commonly taken to be the greatest distance a dark object can be seen against the background sky. The deciview is an index of haziness. A change of 1 or 2 deciviews is seen as a noticeable change in the appearance of a scene.

Visibility impairment is associated with airborne particle properties, including size distributions (i.e., fine particles in the 0.1- to 1.0- μm size range) and aerosol chemical composition, and with relative humidity. With increasing relative humidity, the amount of moisture available for absorption by particles increases, thus causing the particles to increase in both size and volume. As the particles increase in size and volume, the light scattering potential of the particles also generally increases. Visibility impairment is greatest in the eastern United States and Southern California. In the eastern United States, visibility impairment is caused primarily by light scattering by sulfate aerosols and, to a lesser extent, by nitrate particles and organic aerosols, carbon soot, and crustal dust. Up to 86% of the haziness in the eastern United States is caused by atmospheric sulfate. Farther West, scattering contributions to visibility impairment decrease from 25 to 50%. Light scattering by nitrate aerosols is the major cause of visibility impairment in southern California. Nitrates contribute about 45% to the total light extinction in the West and up to 17% of the total extinction in the East. Organic particles are the second largest contributors to light extinction in most U.S. areas. Organic carbon is the greatest

cause of light extinction in the West, accounting for up to 40% of the total extinction and up to 18% of the visibility impairment in the East. Coarse mass and soil, primarily considered “natural extinction,” is responsible for some of the visibility impairment in the West, accounting for up to 25% of the light extinction.

Key conclusions and findings about PM-related effects on visibility include the following:

- Airborne particles degrade visibility by scattering and absorbing light. These optical properties can be well characterized in terms of a light extinction coefficient, which is the fractional attenuation of light per unit distance. The extinction coefficient produced by a given distribution of particle sizes and compositions is strictly proportional to the particle mass concentration.
- The efficiency with which different particles attenuate light depends upon particle size, with fine particles in the accumulation mode being much more important in causing visibility impairment than coarse-mode particles. Thus, it is fine-particle mass concentrations that tend to drive extinction coefficients in polluted air.
- The spatial and temporal variability in the observed extinction coefficient per mass of fine particles is mainly due to the effects of particle-bound water, which varies with relative humidity and is removed by drying when ambient $PM_{2.5}$ mass concentrations are measured using the Federal Reference Method.
- Improved relationships between ambient fine particle levels and visibility impairment have been developed as a result of refinements in algorithms that relate particle size and composition and relative humidity to light extinction, and thus, to visual range, as well as the availability of much expanded databases of $PM_{2.5}$ concentrations and related compositional information and higher resolution visibility data.
- Various local initiatives to address visibility impairment have demonstrated the usefulness of approaches now being used to evaluate public perceptions and attitudes about visibility impairment and public judgments as to the importance of standards to improve visibility relative to current conditions.
- Various such initiatives, conducted in areas with notable scenic vistas (e.g., Denver, CO), have resulted in local standards that reflect what might be referred to as “adverse thresholds” associated with a minimum visual range of approximately 40 to 60 km.
- These various local standards take into account that visibility impairment is an instantaneous effect of ambient $PM_{2.5}$ levels and that the public primarily values enhanced visibility during daylight hours. These considerations are reflected in local standards that are based on sub-daily averaging times (e.g., 4 to 6 hours), typically averaged across mid-day hours.

- This general convergence of visual range values and averaging times that have been determined to be acceptable to the public in a number of such locations suggests that these values and averaging times are relevant for consideration in assessing the need for a national secondary standard to protect visibility in such areas.

4.6.3 Particulate Matter-Related Effects on Materials

Building materials (metals, stones, cements, and paints) undergo natural weathering processes from exposure to environmental elements (wind, moisture, temperature fluctuations, sunlight, etc.). Metals form a protective film of oxidized metal (e.g., rust) that slows environmentally induced corrosion. On the other hand, the natural process of metal corrosion from exposure to natural environmental elements is enhanced by exposure to anthropogenic pollutants, particularly SO₂, that render the protective film less effective.

Dry deposition of SO₂ enhances the effects of environmental elements on calcareous stones (limestone, marble, and cement) by converting calcium carbonate (calcite) to calcium sulfate dihydrate (gypsum). The rate of deterioration is determined by the SO₂ concentration, the stone's permeability and moisture content, and the deposition rate; however, the extent of the damage to stones produced by the pollutant species apart from the natural weathering processes is uncertain. Sulfur dioxide also has been found to limit the life expectancy of paints by causing discoloration, loss of gloss, and thinning of the paint film layer.

A significant detrimental effect of particle pollution is the soiling of painted surfaces and other building materials. Soiling changes the reflectance of opaque materials and reduces the transmission of light through transparent materials. Soiling is a degradation process that requires remediation by cleaning or washing, and, depending on the soiled surface, repainting. Available data on pollution exposure indicates that particles can result in increased cleaning frequency of the exposed surface and may reduce the usefulness of the soiled material. Attempts have been made to quantify the pollutants exposure levels at which materials damage and soiling have been perceived. However, to date, insufficient data are available to advance our knowledge regarding perception thresholds with respect to pollutant concentration, particle size, and chemical composition.

4.6.4 Effects of Atmospheric Particulate Matter on Global Warming Processes and Transmission of Solar Ultraviolet Radiation

The physical processes (i.e., scattering and absorption) responsible for airborne particle effects on transmission of solar visible and ultraviolet radiation are the same as those responsible for visibility degradation. Scattering of solar radiation back to space and absorption of solar radiation determine the effects of an aerosol layer on solar radiation.

Atmospheric particles greatly complicate projections of future trends in global warming processes because of emissions of GHGs; consequent increases in global mean temperature; resulting changes in regional and local weather patterns; and mainly deleterious (but sometimes beneficial) location-specific human health and environmental effects. The body of available evidence, ranging from satellite to in situ measurements of aerosol effects on radiation receipts and cloud properties, is strongly indicative of an important role in climate for aerosols. This role, however, is poorly quantified. No significant advances have been made in reducing the uncertainties assigned to forcing estimates provided by the IPCC for aerosol-related forcing, especially for black carbon-containing aerosol. The IPCC characterizes the scientific understanding of GHG-related forcing as “high” in contrast to that for aerosol, which it describes as “low” to “very low.”

Quantification of the effect of anthropogenic aerosols on hydrological cycles requires more information than is presently available regarding ecosystems responses to reduced solar radiation and other changes occurring in the climate system. However, several global-scale studies indicate that aerosol cooling alone can slow down the hydrological cycle, while cooling plus the nucleation of additional cloud droplets can dramatically reduce precipitation rates.

In addition to direct climate effects through the scattering and absorption of solar radiation, particles also exert indirect effects on climate by serving as cloud condensation nuclei, thus affecting the abundance and vertical distribution of clouds. The direct and indirect effects of particles appear to have significantly offset global warming effects caused by the buildup of GHGs on a globally-averaged basis. However, because the lifetime of particles is much shorter than the time required for complete mixing within the northern hemisphere, the climate effects of particles generally are felt much less homogeneously than are the effects of long-lived GHGs.

Any effort to model the impacts of local alterations in particle concentrations on projected global climate change or consequent local and regional weather patterns would be subject to considerable uncertainty.

Atmospheric particles also complicate estimation of potential future impacts on human health and the environment projected as possible to occur because of increased transmission of SUVB through the Earth's atmosphere, secondary to stratospheric ozone depletion due to anthropogenic emissions of chlorofluorocarbons (CFCs), halons, and certain other gases. The transmission of SUVB radiation is affected strongly by atmospheric particles. Measured attenuations of UV-B under hazy conditions range up to 37% of the incoming solar radiation. Measurements relating variations in PM mass directly to SUVB transmission are lacking. Particles also can affect the rates of photochemical reactions occurring in the atmosphere, e.g., those involved in catalyzing tropospheric ozone formation. Depending on the amount of absorbing substances in the particles, photolysis rates either can be increased or decreased. Thus, the atmospheric particle effects on SUVB radiation vary depending on the size and composition of particles and can differ substantially over different geographic areas, and from season to season over the same area. Any projection of the effects of location-specific airborne PM alterations on increased atmospheric transmission of solar UV radiation (and associated potential human health or environmental effects) due to stratospheric ozone-depletion would, therefore, also be subject to considerable uncertainty.

REFERENCES

- Abdullah, U. M.; Iqbal, M. Z. (1991) Response of automobile, stone and cement particulate matters on stomatal clogging of plants. *Geobios (Jodhpur, India)* 18: 196-202.
- Aber, J. D.; Nadelhoffer, K. J.; Steudler, P.; Melillo, J. M. (1989) Nitrogen saturation in northern forest ecosystems: excess nitrogen from fossil fuel combustion may stress the biosphere. *BioScience* 39: 378-386.
- Aber, J. D.; Magill, A.; McNulty, S. G.; Boone, R. D.; Nadelhoffer, K. J.; Downs, M.; Hallett, R. (1995) Forest biogeochemistry and primary production altered by nitrogen saturation. *Water Air Soil Pollut.* 85: 1665-1670.
- Aber, J.; McDowell, W.; Nadelhoffer, K.; Magill, A.; Berntson, G.; Kamakea, M.; McNulty, S.; Currie, W.; Rustad, L.; Fernandez, I. (1998) Nitrogen saturation in temperate forest ecosystems. *BioScience* 48: 921-934.
- Ainsleigh, H. G. (1993) Beneficial effects of sun exposure on cancer mortality. *Prev. Med.* 22: 132-140.
- Alexander, M. (1977) Introduction to soil microbiology. 2nd ed. New York, NY: John Wiley & Sons; pp. 225-230; 239-246; 272-286; 355-356.
- Allen, M. F. (1991) The ecology of mycorrhizae. Cambridge, MA: Cambridge University Press.
- Allen, A. G.; Harrison, R. M.; Nicholson, K. W. (1991) Dry deposition of fine aerosol to a short grass surface. *Atmos. Environ. Part A* 25: 2671-2676.
- Allen, E. R.; Cabrera, N.; Kim, J.-C. (1994) Atmospheric deposition studies in rural forested areas of the southeastern United States. Presented at: 87th annual meeting & exhibition of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-WP88.02.
- Allen, E. B.; Padgett, P. E.; Bytnerowicz, A.; Minich, R. (1998) Nitrogen deposition effects on coastal sage vegetation of southern California. USDA Forest Service Gen. Tech. Rep. PSW-GTR-166, pp. 131-139.
- Altshuller, A. P.; Linthurst, R. A., eds. (1984) Acidic deposition phenomenon and its effects: critical assessment review papers; volume I, atmospheric sciences; volume II, effects sciences. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development; EPA report nos. EPA/600/8-83/016aF-bF. Available from: NTIS, Springfield, VA; PB85-100030 and PB85-100048.
- Andersen, C. P.; Rygiewicz, P. T. (1991) Stress interactions and mycorrhizal plant response: understanding carbon allocation priorities. *Environ. Pollut.* 73: 217-244.
- Anderson, J. B.; Baumgardner, R. E.; Mohnen, V. A.; Bowser, J. J. (1999) Cloud chemistry in the eastern United States, as sampled from three high-elevation sites along the Appalachian Mountains. *Atmos. Environ.* 33: 5105-5114.
- Aneja, V. P.; Murthy, A. B. (1994) Monitoring deposition of nitrogen-containing compounds in a high-elevation forest canopy. *Air Waste* 44: 1109-1115.
- Antonovics, J.; Bradshaw, A. D.; Turner, R. G. (1971) Heavy metal tolerance in plants. In: Cragg, J. B., ed. *Advances in ecological research*. Volume 7. London, United Kingdom: Academic Press; pp. 1-85.
- April, R.; Newton, R. (1992) Mineralogy and mineral weathering. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag; pp. 378-425. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. *Ecological studies: analysis and synthesis*: v. 91).
- Arizona Department of Environmental Quality. (2002) Air quality report. A.R.S. § 49-424.10. Phoenix, AZ: Arizona Department of Environmental Quality; Air Quality Assessment Section. Available: <http://www.azdeq.gov/function/forms/download/2002/aq.pdf> [29 September, 2004].
- Arnott, W. P.; Moosmüller, H.; Rogers, C. F.; Jin, T.; Bruch, R. (1999) Photoacoustic spectrometer for measuring light absorption by aerosol; instrument description. *Atmos. Environ.* 33: 2845-2852.
- Arrhenius, S. (1896) On the influence of carbonic acid in the air upon the temperature of the ground. *Philos. Mag.* 41: 237-276.
- Askey, A.; Lyon, S. B.; Thompson, G. E.; Johnson, J. B.; Wood, G. C.; Sage, P. W.; Cooke, M. J. (1993) The effect of fly-ash particulates on the atmospheric corrosion of zinc and mild steel. *Corros. Sci.* 34: 1055-1081.
- Ausset, P.; Bannery, F.; Del Monte, M.; Lefevre, R. A. (1998) Recording of pre-industrial atmospheric environment by ancient crusts on stone monuments. *Atmos. Environ.* 32: 2859-2863.
- Ayensu, E.; Van R. Claasen, D.; Collins, M.; Dearing, A.; Fresco, L.; Gadgil, M.; Gitay, H.; Glaser, G.; Juma, C.; Krebs, J.; Lenton, R.; Lubchenco, J.; McNeely, J. A.; Mooney, H. A.; Pinstrip-Andersen, P.; Ramos, M.; Raven, P.; Reid, W. V.; Samper, C.; Sarukhan, J.; Schei, P.; Tundisi, J. G.; Watson, R. T.; Guanhua, X.; Zakri, A. H. (1999) International ecosystem assessment. *Science (Washington, DC)* 286: 685-686.
- Babich, H.; Stotzky, G. (1978) Effects of cadmium on the biota: influence of environmental factors. In: Perlman, D., ed. *Advances in applied microbiology*, volume 23. New York, NY: Academic Press; pp. 55-117.

- Bache, D. H. (1979a) Particle transport within plant canopies - I. A framework for analysis. *Atmos. Environ.* 13: 1257-1262.
- Bache, D. H. (1979b) Particulate transport within plant canopies - II. Prediction of deposition velocities. *Atmos. Environ.* 13: 1681-1687.
- Baedecker, P. A.; Edney, E. O.; Moran, P. J.; Simpson, T. C.; Williams, R. S. (1991) Effects of acidic deposition on materials. In: Irving, P. M., ed. *Acidic deposition: state of science and technology, volume III: terrestrial, materials, health and visibility effects*. Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 19).
- Baedecker, P. A.; Reddy, M. M.; Reimann, K. J.; Sciammarella, C. A. (1992) Effects of acidic deposition on the erosion of carbonate stone—experimental results from the U.S. National Acid Precipitation Assessment Program (NAPAP). *Atmos. Environ. Part B* 26: 147-158.
- Baldocchi, D. (1988) A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. *Atmos. Environ.* 22: 869-884.
- Bange, M. P.; Hammer, G. L.; Rickert, K. G. (1997) Effect of radiation environment on radiation use efficiency and growth of sunflower. *Crop Sci.* 37: 1208-1214.
- Baron, J. S.; Rueth, H. M.; Wolfe, A. M.; Nydick, K. R.; Allstott, E. J.; Minear, J. T.; Moraska, B. (2000) Ecosystem responses to nitrogen deposition in the Colorado front range. *Ecosystems* 3: 352-368.
- Barrie, L. A.; Anlauf, K.; Wiebe, H. A.; Fellin, P. (1984) Acidic pollutants in air and precipitation at selected rural locations in Canada. In: Hicks, B. B., ed. *Deposition both wet and dry: [papers presented at the American Chemical Society acid rain symposium]; March 1982; Las Vegas, NV*. Boston, MA: Butterworth Publishers; pp. 15-35. (Teasley, J. I., ed. *Acid precipitation series: v. 4*).
- Barton, K. (1958) The influence of dust on atmospheric corrosion of metals. *Werkst. Korros.* 8/9: 547-549.
- BBC Research & Consulting. (2003) Phoenix area visibility survey. Phoenix, AZ: Arizona Department of Environmental Quality. Available: http://www.azdeq.gov/enviro/air/download/vis_021903f.pdf [17 September, 2004].
- Beier, C. (1991) Separation of gaseous and particulate dry deposition of sulfur at a forest edge in Denmark. *J. Environ. Qual.* 20: 460-466.
- Beier, C.; Gundersen, P.; Rasmussen, L. (1992) A new method for estimation of dry deposition of particles based on throughfall measurements in a forest edge. *Atmos. Environ. Part A* 26: 1553-1559.
- Bellan, L. M.; Salmon, L. G.; Cass, G. R. (2000) A study on the human ability to detect soot deposition onto works of art. *Environ. Sci. Technol.* 34: 1946-1952.
- Bennett, J. H.; Hill, A. C. (1973) Absorption of gaseous air pollutants by a standardized plant canopy. *J. Air Pollut. Control Assoc.* 23: 203-206.
- Bennett, J. H.; Hill, A. C. (1975) Interactions of air pollutants with canopies of vegetation. In: Mudd, J. B.; Kozlowski, T. T., eds. *Responses of plants to air pollution*. New York, NY: Academic Press; pp. 273-306.
- Bewley, R. J. F.; Parkinson, D. (1984) Effects of sulphur dioxide pollution on forest soil microorganisms. *Can. J. Microbiol.* 30: 179-185.
- Biefer, G. J. (1981) Atmospheric corrosion of steel in the Canadian Arctic. *Mater. Perform.* 20: 16-19.
- Billings, W. D. (1978) *Plants and the ecosystem*. 3rd ed. Belmont, CA: Wadsworth Publishing Company, Inc.; pp. 1-62, 83-108.
- Binkley, D.; Valentine, D.; Wells, C.; Valentine, U. (1989) An empirical analysis of the factors contributing to 20-year decrease in soil pH in an old-field plantation of loblolly pine. *Biogeochemistry* 8: 39-54.
- Blackburn, T. E.; Bairai, S. T.; Stedman, D. H. (1992) Solar photolysis of ozone to singlet *D* oxygen atoms. *J. Geophys. Res. [Atmos.]* 97: 10,109-10,117.
- Blaschke, H. (1990) Mycorrhizal populations and fine root development on Norway spruce exposed to controlled doses of gaseous pollutants and simulated acidic rain treatments. *Environ. Pollut.* 68: 409-418.
- Bobbink, R. (1998) Impacts of tropospheric ozone and airborne nitrogenous pollutants on natural and semi-natural ecosystems: a commentary. *New Phytol.* 139: 161-168.
- Bobbink, R.; Hornung, M.; Roelofs, J. G. M. (1998) The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *J. Ecol.* 86: 717-738.
- Bondietti, E. A.; McLaughlin, S. B. (1992) Evidence of historical influences of acidic deposition on wood and soil chemistry. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag; pp. 358-377. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. *Ecological studies analysis and synthesis: v. 91*).
- Bormann, F. H.; Likens, G. E. (1979) Catastrophic disturbance and the steady state in northern hardwood forests. *Am. Sci.* 67: 660-669.

- Bowden, R. D.; Geballe, G. T.; Bowden, W. B. (1989) Foliar uptake of ^{15}N from simulated cloud water by red spruce (*Picea rubens*) seedlings. *Can. J. For. Res.* 19: 382-386.
- Bowman, W. D. (2000) Biotic controls over ecosystem response to environmental change in alpine tundra of the Rocky Mountains. *Ambio* 29: 396-400.
- Bowman, W. D.; Steltzer, H. (1998) Positive feedbacks to anthropogenic nitrogen deposition in Rocky Mountain alpine tundra. *Ambio* 27: 514-517.
- Boyce, S. G. (1954) The salt spray community. *Ecol. Monogr.* 24: 29-67.
- Bradley, R.; Burt, A. J.; Read, D. J. (1981) Mycorrhizal infection and resistance to heavy metal toxicity in *Calluna vulgaris*. *Nature (London)* 292: 335-337.
- Brandt, C. J.; Rhoades, R. W. (1972) Effects of limestone dust accumulation on composition of a forest community. *Environ. Pollut.* 3: 217-225.
- Brandt, C. J.; Rhoades, R. W. (1973) Effects of limestone dust accumulation on lateral growth of forest trees. *Environ. Pollut.* 4: 207-213.
- Brook, J. R.; Zhang, L.; Di-Giovanni, F.; Padro, J. (1999) Description and evaluation of a model of deposition velocities for routine estimates of air pollutant dry deposition over North America. Part I: model development. *Atmos. Environ.* 33: 5037-5051.
- Brookshire, D. S.; d'Arge, R. C.; Schulze, W. D.; Thayer, M. D. (1979) Methods development for assessing air pollution control benefits: volume II. alternative benefit measures of air pollution control in the south coast air basin of southern California. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development; report no. EPA-600/6-79-001b.
- Brühl, C.; Crutzen, P. J. (1988) On the disproportionate role of tropospheric ozone as a filter against solar UV-B radiation. *Geophys. Res. Lett.* 16: 703-706.
- Brumme, R.; Leimcke, U.; Matzner, E. (1992) Interception and uptake of NH_4 and NO_3 from wet deposition by above-ground parts of young beech (*Fagus sylvatica* L.) trees. *Plant Soil* 142: 273-279.
- Brunet, J.; Diekmann, M.; Falkengren-Grerup, U. (1998) Effects of nitrogen deposition on field layer vegetation in south Swedish oak forests. *Environ. Pollut. (Oxford, U.K.)* 102(suppl. 1): 35-40.
- Bugini, R.; Tabasso, M. L.; Realini, M. (2000) Rate of formation of black crusts on marble. A case study. *J. Cult. Heritage* 1: 111-116.
- Businger, J. A. (1986) Evaluation of the accuracy with which dry deposition can be measured with current micrometeorological techniques. *J. Clim. Appl. Meteorol.* 25: 1100-1124.
- Butlin, R. N.; Coote, A. T.; Devenish, M.; Hughes, I. S. C.; Hutchens, C. M.; Irwin, J. G.; Lloyd, G. O.; Massey, S. W.; Webb, A. H.; Yates, T. J. S. (1992a) Preliminary results from the analysis of metal samples from the National Materials Exposure Programme (NMEP). *Atmos. Environ. Part B* 26: 199-206.
- Butlin, R. N.; Coote, A. T.; Devenish, M.; Hughes, I. S. C.; Hutchens, C. M.; Irwin, J. G.; Lloyd, G. O.; Massey, S. W.; Webb, A. H.; Yates, T. J. S. (1992b) Preliminary results from the analysis of stone tablets from the National Materials Exposure Programme (NMEP). *Atmos. Environ. Part B* 26: 189-198.
- Bytnerowicz, A.; Fenn, M. E. (1996) Nitrogen deposition in California forests: a review. *Environ. Pollut.* 92: 127-146.
- Bytnerowicz, A.; Miller, P. R.; Olszyk, D. M. (1987a) Dry deposition of nitrate, ammonium and sulfate to a *Ceanothus crassifolius* canopy and surrogate surfaces. *Atmos. Environ.* 21: 1749-1757.
- Bytnerowicz, A.; Miller, P. R.; Olszyk, D. M.; Dawson, P. J.; Fox, C. A. (1987b) Gaseous and particulate air pollution in the San Gabriel Mountains of southern California. *Atmos. Environ.* 21: 1805-1814.
- Cahill, T. A.; Eldred, R. A.; Wakabayashi, P. H. (1996) Trends in fine particle concentrations at Great Smoky Mountains National Park. For presentation at: Annual meeting of the Air & Waste Management Association; June; Nashville, TN. Pittsburgh, PA: Air & Waste Management Association; paper no. 96-MP1A.05.
- Campbell, G. G.; Schurr, G. G.; Slawikowski, D. E.; Spence, J. W. (1974) Assessing air pollution damage to coatings. *J. Paint Technol.* 46: 59-71.
- Camuffo, D. (1995) Physical weathering of stones. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. *Sci. Total Environ.* 167: 1-14.
- Cape, J. N. (1993) Direct damage to vegetation caused by acid rain and polluted cloud: definition of critical levels for forest trees. *Environ. Pollut.* 82: 167-180.
- Cape, J. N.; Sheppard, L. J.; Fowler, D.; Harrison, A. F.; Parkinson, J. A.; Dao, P.; Paterson, I. S. (1992) Contribution of canopy leaching to sulphate deposition in a Scots pine forest. *Environ. Pollut.* 75: 229-236.
- Carey, W. F. (1959) Atmospheric deposits in Britain: a study of dinginess. *Int. J. Air Pollut.* 2: 1-26.

- Ceccotti, S. P.; Messick, D. L. (1997) A global review of crop requirements, supply, and environmental impact on nutrient sulphur balance. In: Cram, W. J.; De Kok, L. J.; Stulen, I.; Brunold, C.; Rennenberg, H., eds. Sulphur metabolism in higher plants. Leiden, The Netherlands: Backhuys Publishers; pp. 155-163.
- Chamberlain, A. C. (1975) The movement of particles in plant communities. In: Monteith, J. L., ed. Vegetation and the atmosphere: v. 1, principles. London, United Kingdom: Academic Press; pp. 155-203.
- Chamberlain, A. C.; Little, P. (1981) Transport and capture of particles by vegetation. In: Grace, J.; Ford, E. D.; Jarvis, P. G., eds. Plants and their atmospheric environment: the 21st symposium of the British Ecological Society; 1979; Edinburgh, United Kingdom. Oxford, United Kingdom: Blackwell Scientific Publications; pp. 147-173.
- Chameides, W. L.; Yu, H.; Liu, S. C.; Bergin, M.; Zhou, X.; Mearns, L.; Wang, G.; Kiang, C. S.; Saylor, R. D.; Luo, C.; Huang, Y.; Steiner, A.; Giorgi, F. (1999) Case study of the effects of atmospheric aerosols and regional haze on agriculture: an opportunity to enhance crop yields in China through emission controls? *Proc. Natl. Acad. Sci. U. S. A.* 96: 13626-13633.
- Chapin, F. S., III. (1991) Integrated responses of plants to stress: a centralized system of physiological responses. *BioScience* 41: 29-36.
- Chapin, F. S., III; Ruess, R. W. (2001) The roots of the matter. *Nature (London)* 411: 749-752.
- Chapin, F. S., III; Bloom, A. J.; Field, C. B.; Waring, R. H. (1987) Plant responses to multiple environmental factors. *BioScience* 37: 49-57.
- Chapin, F. S., III; Sala, O. E.; Burke, I. C.; Grime, J. P.; Hooper, D. U.; Lauenroth, W. K.; Lombard, A.; Mooney, H. A.; Mosier, A. R.; Naeem, S.; Pacala, S. W.; Roy, J.; Steffen, W. L.; Tilman, D. (1998) Ecosystem consequences of changing biodiversity. *BioScience* 48: 45-52.
- Charlson, R. J.; Ahlquist, N. C.; Horvath, H. (1968) On the generality of correlation of atmospheric aerosol mass concentration and light scatter. *Atmos. Environ.* 2: 455-464.
- Charlson, R. J.; Schwartz, S. E.; Hales, J. M.; Cess, R. D.; Coakley, J. A., Jr.; Hansen, J. E.; Hofmann, D. J. (1992) Climate forcing by anthropogenic aerosols. *Science (Washington, DC)* 255: 423-430.
- Chestnut, L. (1997) Methodology for estimating values for changes in visibility at national parks [draft memorandum]. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; contract no. 68-D-98-001; EPA air docket A-96-56, document no. VI-B-09-(ooo).
- Chestnut, L. G.; Dennis, R. L. (1997) Economic benefits of improvements in visibility: acid rain. Provisions of the 1990 clean air act amendments. *J. Air Waste Manage. Assoc.* 47: 395-402.
- Chestnut, L. G.; Rowe, R. D. (1990) Preservation values for visibility protection at the national parks (draft final report). Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Economic Analysis Branch.
- Chevone, B. I.; Herzfeld, D. E.; Krupa, S. V.; Chappelka, A. H. (1986) Direct effects of atmospheric sulfate deposition on vegetation. *J. Air Pollut. Control Assoc.* 36: 813-815.
- Choudhury, B. J. (1987) Relationships between vegetation indices, radiation absorption, and net photosynthesis evaluated by a sensitivity analysis. *Remote Sens. Environ.* 22: 209-233.
- Chow, J. C.; Watson, J. G.; Pritchett, L.; Lowenthal, D.; Frazier, C.; Neuroth, G.; Evans, K.; Moon, D. (1990) Wintertime visibility in Phoenix, Arizona. Presented at: the 83rd annual meeting & exhibition; June; Pittsburgh, PA. Pittsburgh, PA: Air & Waste Management Association; paper no. 90-66.6.
- Chow, J. C.; Watson, J. G.; Lowenthal, D. H.; Richards, L. W. (2002a) Comparability between PM_{2.5} and particle light scattering measurements. *Environ. Monit. Assess.* 79: 29-45.
- Chow, J. C.; Bachmann, J. D.; Wierman, S. S. G.; Mathai, C. V.; Malm, W. C.; White, W. H.; Mueller, P. K.; Kumar, N.; Watson, J. G. (2002b) Visibility: science and regulation. *J. Air Waste Manage. Assoc.* 52: 973-999.
- Chughtai, A. R.; Williams, G. R.; Atteya, M. M. O.; Miller, N. J.; Smith, D. M. (1999) Carbonaceous particle hydration. *Atmos. Environ.* 33: 2679-2687.
- Cione, N. K.; Padgett, P. E.; Allen, E. B. (2002) Restoration of a native shrubland impacted by exotic grasses, frequent fire and nitrogen deposition in southern California. *Restor. Ecol.* 10: 376-384.
- Clarke, J. F.; Edgerton, E. S.; Martin, B. E. (1997) Dry deposition calculations for the clean air status and trends network. *Atmos. Environ.* 31: 3667-3678.
- Clarkson, D. T.; Sanderson, J. (1971) Inhibition of the uptake and long-distance transport of calcium by aluminum and other polyvalent cations. *J. Exp. Bot.* 22: 837-851.
- Clevenger, T. E.; Saiwan, C.; Koirtiyohann, S. R. (1991) Lead speciation of particles on air filters collected in the vicinity of a lead smelter. *Environ. Sci. Technol.* 25: 1128-1133.
- Clough, W. S. (1975) The deposition of particles on moss and grass surfaces. *Atmos. Environ.* 9: 1113-1119.

- Cocker, D. R., III; Whitlock, N. E.; Flagen, R. C.; Seinfeld, J. H. (2001) Hygroscopic properties of Pasadena, California aerosol. *Aerosol Sci. Technol.* 35: 637-647.
- Cole, D. W.; Johnson, D. W. (1977) Atmospheric sulfate additions and cation leaching in a Douglas fir ecosystem. *Water Resour. Res.* 13: 313-317.
- Collett, J. L., Jr.; Hoag, K. J.; Sherman, D. E.; Bator, A.; Richards, L. W. (1999) Spatial and temporal variations in San Joaquin Valley fog chemistry. *Atmos. Environ.* 33: 129-140.
- Cooke, R. U.; Gibbs, G. B. (1994) Crumbling heritage? Studies of stone weathering in polluted atmospheres. *Atmos. Environ.* 28: 1355-1356.
- Costanza, R.; d'Arge, R.; De Groot, R.; Farber, S.; Grasso, M.; Hannon, B.; Limburg, K.; Naeem, S.; O'Neill, R. V.; Paruelo, J.; Raskin, R. G.; Sutton, P.; Van Den Belt, M. (1997) The value of the world's ecosystem services and natural capital. *Nature (London)* 387: 253-259.
- Cotrufo, M. F.; De Santo, A. V.; Alfani, A.; Bartoli, G.; De Cristofaro, A. (1995) Effects of urban heavy metal pollution on organic matter decomposition in *Quercus ilex L.* woods. *Environ. Pollut.* 89: 81-87.
- Cowling, J. E.; Roberts, M. E. (1954) Paints, varnishes, enamels, and lacquers. In: Greathouse, G. A.; Wessel, C. J., eds. *Deterioration of materials: causes and preventive techniques*. New York, NY: Reinhold Publishing Corporation; pp. 596-645.
- Cowling, E. B.; Erisman, J. W.; Smeulders, S. M.; Holman, S. C.; Nicholson, B. M. (1998) Optimizing air quality management in Europe and North America: justification for integrated management of both oxidized and reduced forms of nitrogen. *Environ. Pollut.* 102(suppl. 1): 599-608.
- Cramer, S. D.; McDonald, L. G.; Bhagia, G.; Flinn, D. R.; Linstrom, P. J.; Carter, J. P. (1989) Effects of acid deposition on the atmospheric corrosion of structural metals. Research Triangle Park, NC: U.S. Environmental Protection Agency; internal report. Incorporated in: Irving, P. M., ed. *Acidic Deposition: State of Science and Technology. Volume III: Terrestrial, Materials, Health and Visibility Effects; report 19, section 2*. Washington, DC: National Acid Precipitation Assessment Program.
- Creighton, P. J.; Lioy, P. J.; Haynie, F. H.; Lemmons, T. J.; Miller, J. L.; Gerhart, J. (1990) Soiling by atmospheric aerosols in an urban industrial area. *J. Air Waste Manage. Assoc.* 40: 1285-1289.
- Cronan, C. S.; Grigal, D. F. (1995) Use of calcium/aluminum ratios as indicators of stress in forest ecosystems. *J. Environ. Qual.* 24: 209-226.
- Cullen, J. J.; Neale, P. J. (1997) Biological weighting functions for describing the effects of ultraviolet radiation on aquatic ecosystems. In: Häder, D.-P., ed. *The effects of ozone depletion on aquatic ecosystems*. Austin, TX: Academic Press, pp. 97-118.
- Dahlgren, R. A. (1994) Soil acidification and nitrogen saturation from weathering of ammonium-bearing rock. *Nature (London)* 368: 838-841.
- Daily, G. C. (1997) Introduction: what are ecosystem services? In: Daily, G. C., ed. *Nature's services: societal dependence on natural ecosystems*. Washington, DC: Island Press; pp. 1-10.
- Daily, G. C.; Ehrlich, P. R. (1999) Managing Earth's ecosystems: an interdisciplinary challenge. *Ecosystems* 2: 277-280.
- Darley, E. F. (1966) Studies on the effect of cement-kiln dust on vegetation. *J. Air Pollut. Control Assoc.* 16: 145-150.
- Dasch, J. M. (1987) Measurement of dry deposition to surfaces in deciduous and pine canopies. *Environ. Pollut.* 44: 261-277.
- Dasch, J. M.; Cadle, S. H. (1985) Wet and dry deposition monitoring in southeastern Michigan. *Atmos. Environ.* 19: 789-796.
- Dasch, J. M.; Cadle, S. H. (1986) Dry deposition to snow in an urban area. *Water Air Soil Pollut.* 29: 297-308.
- Dässler, H.-G.; Ranft, H.; Rehn, K.-H. (1972) Zur Widerstandsfähigkeit von Gehölzen gegenüber Fluorverbindungen und Schwefeldioxid [The susceptibility of woody plants exposed to fluorine compounds and SO₂]. *Flora (Jena)* 161: 289-302.
- Davidson, C. I.; Friedlander, S. K. (1978) A filtration model for aerosol dry deposition: application to trace metal deposition from the atmosphere. *J. Geophys. Res.* 83: 2343-2352.
- Davidson, C. I.; Wu, Y.-L. (1989) Dry deposition of trace elements. In: Pacyna, J.M.; Ottar, B., eds. *Control and fate of atmospheric trace metals: proceedings of the NATO advanced research workshop on fate and control of toxic metals in the atmosphere; September 1988; Oslo, Norway*. Boston, MA: Kluwer Academic Publishers; pp. 147-202. (NATO ASI series C, mathematical and physical sciences: v. 268).
- Davidson, C. I.; Wu, Y.-L. (1990) Dry deposition of particles and vapors. In: Lindberg, S. E.; Page, A. L.; Norton, S. A., eds. *Acidic precipitation: v. 3, sources, deposition, and canopy interactions*. New York, NY: Springer-Verlag; pp. 103-216.

- Davidson, C. I.; Miller, J. M.; Pleskow, M. A. (1982) The influence of surface structure on predicted particle dry deposition to natural grass canopies. *Water Air Soil Pollut.* 18: 25-43.
- Davidson, K.; Deck, L.; McCubbin, D.; Post, E. (2000a) Out of sight: the science and economics of visibility impairment. Cambridge, MA: Abt Associates, Inc.
- Davidson, C. I.; Tang, W.; Finger, S.; Etyemezian, V.; Striegel, M. F.; Sherwood, S. I. (2000b) Soiling patterns on a tall limestone building: changes over 60 years. *Environ. Sci. Technol.* 34: 560-565.
- Davis, G. D.; Shaw, B. A.; Arah, C. O.; Fritz, T. L.; Moshier, W. C.; Simpson, T. C.; Moran, P. J.; Zankel, K. L. (1990) Effects of SO₂ deposition on painted steel surfaces. *Surf. Interface Anal.* 15: 107-112.
- Day, D. E.; Malm, W. C.; Kreidenweis, S. M.; Tree, R. M. (1997) Aerosol light scattering measurements: a comparison of differently configured optec nephelometers. In: *Visual air quality: aerosols and global radiation balance: proceedings of a specialty conference sponsored by the Air & Waste Management Association and the American Geophysical Union*; September; Bartlett, NH. Pittsburgh, PA: Air & Waste Management Association; pp. 952-962. (A&WMA publication VIP-76).
- DeHayes, D. H.; Schaberg, P. G.; Hawley, G. J.; Strimbeck, G. R. (1999) Acid rain impacts on calcium nutrition and forest health. *BioScience* 49: 789-800.
- Del Monte, M.; Vittori, O. (1985) Air pollution and stone decay: the case of Venice. *Endeavour* 9: 117-122.
- Diakumaku, E.; Gorbushina, A. A.; Krumbein, W. E.; Panina, L.; Soukharjevski, S. (1995) Black fungi in marble and limestones--an aesthetical, chemical and physical problem for the conservation of monuments. In: Saiz-Jimenez, C., ed. *The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation*; January 1994; Sevilla, Spain. *Sci. Total Environ.* 167: 295-304.
- Dickerson, R. R.; Kondragunta, S.; Stenchikov, G.; Civerolo, K. L.; Doddridge, B. G.; Holben, B. N. (1997) The impact of aerosols on solar ultraviolet radiation and photochemical smog. *Science (Washington, DC)* 278: 827-830.
- Dobbins, R. A.; Mulholland, G. W.; Bryner, N. P. (1994) Comparison of a fractal smoke optics model with light extinction measurements. *Atmos. Environ.* 28: 889-897.
- Dolske, D. A. (1988) Dry deposition of airborne sulfate and nitrate to soybeans. *Environ. Pollut.* 53: 1-12.
- Dolske, D. A. (1995) Deposition of atmospheric pollutants to monuments, statues, and buildings. *Sci. Total Environ.* 167: 15-31.
- Dolske, D. A.; Gatz, D. F. (1984) Field intercomparison of sulfate dry deposition monitoring and measurement methods: preliminary results. In: Hicks, B. B., ed. *Deposition both wet and dry: [papers presented at the American Chemical Society acid rain symposium]*; March 1982; Las Vegas, NV. Boston, MA: Butterworth Publishers; pp. 121-131. (Teasley, J. I., ed. *Acid precipitation series: v. 4*).
- Draaijers, G. P. J.; Erisman, J. W. (1993) Atmospheric sulphur deposition to forest stands: throughfall estimates compared to estimates from inference. *Atmos. Environ. Part A* 27: 43-55.
- Draaijers, G. P. J.; Ivens, W. P. M. F.; Bleuten, W. (1988) Atmospheric deposition in forest edges measured by monitoring canopy throughfall. *Water Air Soil Pollut.* 42: 129-136.
- Draaijers, G. P. J.; van Ek, R.; Meijers, R. (1992) Research on the impact of forest stand structure on atmospheric deposition. *Environ. Pollut.* 75: 243-249.
- Dreisinger, B. R.; McGovern, P. C. (1970) Monitoring atmospheric sulphur dioxide and correlating its effects on crops and forests in the Sudbury area. In: Linzon, S. N., ed. *Impact of air pollution on vegetation conference*; April; Toronto, ON, Canada. Pittsburgh, PA: Air Pollution Control Association.
- Driscoll, C. T.; Van Dreason, R. (1993) Seasonal and long-term temporal patterns in the chemistry of Adirondack lakes. *Water Air Soil Pollut.* 67: 319-344.
- Driscoll, C. T.; Wyskowski, B. J.; DeStaffan, P.; Newton, R. M. (1989) Chemistry and transfer of aluminum in a forested watershed in the Adirondack region of New York, USA. In: Lewis, T. E., ed. *Environmental chemistry and toxicology of aluminum*. Chelsea, MI: Lewis Publishers, Inc.; pp. 83-105.
- Driscoll, C. T.; Newton, R. M.; Gubala, C. P.; Baker, J. P.; Christensen, S. W. (1991) Adirondack mountains. In: Charles, D. F., ed. *Acidic deposition and aquatic ecosystems: regional case studies*. New York, NY: Springer-Verlag; pp. 133-202.
- Driscoll, C. T.; Lawrence, G. B.; Bulger, A. J.; Butler, T. J.; Cronan, C. S.; Eagar, C.; Lambert, K. F.; Likens, G. E.; Stoddard, J. L.; Weathers, K. C. (2001) Acidic deposition in the northeastern United States: sources and inputs, ecosystem effects, and management strategies. *BioScience* 51: 180-198.
- Driscoll, C. T.; Whittall, D.; Aber, J.; Boyer, E.; Castro, M.; Cronan, C.; Goodale, C. L.; Groffman, P.; Hopkinson, C.; Lambert, K.; Lawrence, G.; Ollinger, S. (2003) Nitrogen pollution in the northeastern United States: sources, effects, and management options. *BioScience* 53: 357-374.
- Duan, B.; Fairall, C. W.; Thomson, D. W. (1988) Eddy correlation measurements of the dry deposition of particles in wintertime. *J. Appl. Meteorol.* 27: 642-652.

- Duyzer, J. H.; Bouman, A. M. M.; Van Aalst, R. M.; Diederer, H. S. M. A. (1987) Assessment of dry deposition fluxes of NH_3 and NH_4^+ over natural terrains. In: Asman, W. A. H.; Diederer, H. S. M. A., eds. Ammonia and acidification: proceedings of a symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 97-106.
- Eaton, J. S.; Likens, G. E.; Bormann, F. H. (1973) Throughfall and stemflow chemistry in a northern hardwood forest. *J. Ecol.* 61: 495-508.
- Edgerton, E. S.; Lavery, T. F.; Boksleitner, R. P. (1992) Preliminary data from the USEPA dry deposition network: 1989. *Environ. Pollut.* 75: 145-156.
- Edgerton-Warburton, L. M.; Allen, E. B. (2000) Shifts in arbuscular mycorrhizal communities along an anthropogenic gradient nitrogen deposition gradient. *Ecol. Appl.* 10: 484-496.
- Edney, E. O. (1989) Paint coatings: controlled field and chamber experiments. Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/3-89/032. Available from: NTIS, Springfield, VA; PB89-189849.
- Edney, E. O.; Stiles, D. C.; Corse, E. W.; Wheeler, M. L.; Spence, J. W.; Haynie, F. H.; Wilson, W. E., Jr. (1988) Field study to determine the impact of air pollutants on the corrosion of galvanized steel. *Mater. Perform.* 27: 47-50.
- Edney, E. O.; Cheek, S. F.; Corse, E. W.; Spence, J. W.; Haynie, F. H. (1989) Atmospheric weathering caused by dry deposition of acidic compounds. *J. Environ. Sci. Health Part A* 24: 439-457.
- Edwards, R. S.; Claxton, S. M. (1964) The distribution of air-borne salt of marine origin in the Aberystwyth area. *J. Appl. Ecol.* 1: 253-263.
- El-Shobokshy, M. S. (1985) The dependence of airborne particulate deposition on atmospheric stability and surface conditions. *Atmos. Environ.* 19: 1191-1197.
- Eldering, A.; Cass, G. R.; Moon, K. C. (1994) An air monitoring network using continuous particle size distribution monitors: connecting pollutant properties to visibility via Mie scattering calculations. *Atmos. Environ.* 28: 2733-2749.
- Eldering, A.; Hall, J. R.; Hussey, K. J.; Cass, G. R. (1996) Visibility model based on satellite-generated landscape data. *Environ. Sci. Technol.* 30: 361-370.
- Eldred, R. A.; Cahill, T. A. (1994) Trends in elemental concentrations of fine particles at remote sites in the United States of America. *Atmos. Environ.* 28: 1009-1019.
- Eldred, R. A.; Cahill, T. A.; Feeney, P. J. (1993) Comparison of independent measurements of sulfur and sulfate in the IMPROVE network. Presented at: 86th annual meeting and exhibition of the Air & Waste Management Association; June; Denver, CO. Pittsburgh, PA: Air & Waste Management Association; paper no. 93-RA-110.02.
- Eldred, R. A.; Cahill, T. A.; Flocchini, R. G. (1997) Composition of $\text{PM}_{2.5}$ and PM_{10} aerosols in the IMPROVE network. In: Special issue on aerosols and atmospheric optics—part I, [published papers from the international specialty conference; September 1994; Snowbird, UT]. *J. Air Waste Manage. Assoc.* 47: 194-203.
- Eliason, S. A.; Allen, E. B. (1997) Exotic grass competition in suppressing native shrubland re-establishment. *Restor. Ecol.* 5: 245-255.
- Ellenberg, H. (1987) Floristic changes due to eutrophication. In: Asman, W. A. H.; Diederer, S. M. A., eds. Ammonia and acidification: proceedings of a symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 301-308.
- Eller, B. M. (1977) Road dust induced increase of leaf temperature. *Environ. Pollut.* 13: 99-107.
- Ely, D. W.; Leary, J. T.; Stewart, T. R.; Ross, D. M. (1991) The establishment of the Denver Visibility Standard. Presented at: 84th annual meeting & exhibition of the Air & Waste Management Association; June; Vancouver, British Columbia. Pittsburgh, PA: Air & Waste Management Association; paper no. 91-48.4.
- Eriksson, P.; Johansson, L.-G.; Strandberg, H. (1993) Initial stages of copper corrosion in humid air containing SO_2 and NO_2 . *J. Electrochem. Soc.* 140: 53-59.
- Erismann, J. W. (1993) Acid deposition onto nature areas in the Netherlands; part II. Throughfall measurements compared to deposition estimates. *Water Air Soil Pollut.* 71: 81-99.
- Erismann, J.-W.; Vermetten, A. W. M.; Asman, W. A. H.; Waijers-Ijpelaar, A.; Slanina, J. (1988) Vertical distribution of gases and aerosols: the behaviour of ammonia and related components in the lower atmosphere. *Atmos. Environ.* 22: 1153-1160.
- Erismann, J. W.; Van Pul, A.; Wyers, P. (1994) Parametrization of surface resistance for the quantification of atmospheric deposition of acidifying pollutants and ozone. *Atmos. Environ.* 28: 2595-2607.

- Erismann, J. W.; Draaijers, G.; Duyzer, J.; Hofschreuder, P.; Van Leeuwen, N.; Romer, F.; Ruijgrok, W.; Wyers, P.; Gallagher, M. (1997) Particle deposition to forests--summary of results and application. *Atmos. Environ.* 31: 321-332.
- Erlick, C.; Frederick, J. E.; Saxena, V. K.; Wenny, B. N. (1998) Atmospheric transmission in the ultraviolet and visible: aerosols in cloudy atmospheres. *J. Geophys. Res.* 103: 31,541-31,556.
- Eshleman, K. N.; Miller-Marshall, L. M.; Webb, J. R. (1995) Long-term changes in episodic acidification of streams in Shenandoah National Park, Virginia (U.S.A.). *Water Air Soil Pollut.* 85: 517-522.
- Estupiñán, J. B.; Raman, S.; Crescenti, G. H.; Streicher, J. J.; Barnard, W. F. (1996) Effects of clouds and haze on UV-B radiation. *J. Geophys. Res.* 101: 16,807-16,816.
- Evans, E. G.; Pitchford, M. (1991) Federal visibility monitoring in the eastern United States. Presented at: 84th annual meeting and exhibition of the Air & Waste Management Association; June; Vancouver, BC, Canada. Pittsburgh, PA: Air & Waste Management Association; paper no. 91-49.6.
- Fahey, T. J.; Yavitt, J. B.; Joyce, G. (1988) Precipitation and throughfall chemistry in *Pinus contorta* ssp. *latifolia* ecosystems, southeastern Wyoming. *Can. J. For. Res.* 18: 337-345.
- Falke, S. (2001) Web-based visibility information system. Available: <http://capita.wustl.edu/datawarehouse/datasets/ASOS/WebVIS.htm> (28 May 2003).
- Falkengren-Grerup, U. (1987) Long-term changes in pH of forest soils in southern Sweden. *Environ. Pollut.* 43: 79-90.
- Falkengren-Grerup, U. (1998) Nitrogen response of herbs and graminoids in experiments with simulated acid soil solution. *Environ. Pollut.* 102(suppl. 1): 93-99.
- Farber, R. J.; Welsing, P. R.; Rozzi, C. (1994) PM₁₀ and ozone control strategy to improve visibility in the Los Angeles Basin. *Atmos. Environ.* 28: 3277-3283.
- Farmer, A. M. (1993) The effects of dust on vegetation--a review. *Environ. Pollut.* 79: 63-75.
- Fassina, V.; Favaro, M.; Crivellari, F.; Naccari, A. (2001) The stone decay of monuments in relation to atmospheric environment. *Ann. Chim.* 91: 767-774.
- Federal Register. (1999) Regional haze regulations; final rule. *F. R.* (July 1) 64: 35,713-35,774.
- Fenn, M. E.; Keifer, J. W. (1999) Throughfall deposition of nitrogen and sulfur in a Jeffrey pine forest in the San Gabriel Mountains, Southern California. *Environ. Pollut.* 104: 179-187.
- Fenn, M. E.; Poth, M. A. (1998) Indicators of nitrogen status in California forests. U.S.D.A. Forest Service Gen. Tech. Rep. PSW-GTR-166.
- Fenn, M. E.; Poth, M. A. (1999) Temporal and spatial trends in streamwater nitrate concentrations in the San Bernardino Mountains, southern California. *J. Environ. Qual.* 28: 822-836.
- Fenn, M. E.; Poth, M. A.; Johnson, D. W. (1996) Evidence for nitrogen saturation in the San Bernardino Mountains in southern California. *For. Ecol. Manage.* 82: 211-230
- Fenn, M. E.; Poth, M. A.; Aber, J. D.; Baron, J. S.; Bormann, B. T.; Johnson, D. W.; Lemly, A. D.; McNulty, S. G.; Ryan, D. F.; Stottlemeyer, R. (1998) Nitrogen excess in North American ecosystems: predisposing factors, ecosystem responses, and management strategies. *Ecol. Appl.* 8: 706-733.
- Fenn, M. E.; Haeuber, R.; Tonnesen, G. S.; Baron, J. S.; Grossman-Clarke, S.; Hope, D.; Jaffe, D. A.; Copeland, S.; Geiser, L.; Rueth, H. M.; Sickman, J. O. (2003a) Nitrogen emissions, deposition, and monitoring in the western United States. *BioScience* 53: 391-403.
- Fenn, M. E.; Baron, J. S.; Allen, E. B.; Rueth, H. M.; Nydick, K. R.; Geiser, L.; Bowman, W. D.; Sickman, J. O.; Meixner, T.; Johnson, D. W.; Neitlich, P. (2003b) Ecological effects of nitrogen deposition in the western United States. *BioScience* 53: 404-420.
- Fink, F. W.; Buttner, F. H.; Boyd, W. K. (1971) Final report on technical-economic evaluation of air-pollution corrosion costs on metals in the U.S. Columbus, OH: Battelle Memorial Institute, Columbus Laboratories; EPA contract no. CPA 70-86; EPA report no. APTD-0654. Available from: NTIS, Springfield, VA; PB-198-453.
- Flum, T.; Nodvin, S. C. (1995) Factors affecting streamwater chemistry in the Great Smoky Mountains, USA. *Water Air Soil Pollut.* 85: 1707-1712.
- Fochtman, E. G.; Langer, G. (1957) Automobile paint damaged by airborne iron particles. *J. Air Pollut. Control Assoc.* 6: 243-247.
- Foltescu, V. L.; Isakson, J.; Selin, E.; Stikans, M. (1994) Measured fluxes of sulphur, chlorine and some anthropogenic metals to the Swedish west coast. *Atmos. Environ.* 28: 2639-2649.
- Foster, J. R. (1991) Effects of organic chemicals in the atmosphere on terrestrial plants. In: Moser, T. J.; Barker, J. R.; Tingey, D. T., eds. *Ecological exposure and effects of airborne toxic chemicals: an overview*. Corvallis, OR: U.S. Environmental Protection Agency, Environmental Research Laboratory; pp. 60-89; EPA report no. EPA-600/3-91-001. Available from: NTIS, Springfield, VA; PB91-148460.

- Foster, N. W.; Nicolson, J. A.; Hazlett, P. W. (1989) Temporal variation in nitrate and nutrient cations in drainage waters from a deciduous forest. *J. Environ. Qual.* 18: 238-244.
- Fowler, D.; Cape, J. N.; Unsworth, M. H. (1989) Deposition of atmospheric pollutants on forests. *Philos. Trans. R. Soc. London B* 324: 247-265.
- Fowler, D.; Duyzer, J. H.; Baldocchi, D. D. (1991) Inputs of trace gases, particles and cloud droplets to terrestrial surfaces. *Proc. R. Soc. Edinburgh Sect. B: Biol. Sci.* 97: 35-59.
- Fox, D. G.; Malm, W. C.; Mitchell, B.; Fisher, R. W. (1999) Where there's fire, there's smoke: fine particulate and regional haze. *EM* (November): 15-24.
- Frederick, J. E.; Qu, Z.; Booth, C. R. (1998) Ultraviolet radiation at sites on the Antarctic coast. *J. Photochem. Photobiol. B* 68: 183-190.
- Freedman, B.; Hutchinson, T. C. (1981) Sources of metal and elemental contamination of terrestrial environments. In: Lepp, N. W., ed. *Effect of heavy metal pollution on plants. Volume 2: metals in the environment.* London, United Kingdom: Applied Science Publishers; pp. 35-94.
- Freeman, A. M., III. (1993) *The measurement of environmental and resource values: theory and methods.* Washington, DC: Resources for the Future; pp. 1-27.
- Freedman, D. M.; Zahm, S. H.; Dosemeci, M. (1997) Residential and occupational exposure to sunlight and mortality from non-Hodgkin's lymphoma: composite (threefold) case-control study. *Br. Med. J.* 314: 1451-1455.
- Freudenburg, W. R.; Alario, M. (1999) What ecologists can learn from nuclear scientists. *Ecosystems* 2: 286-291.
- Fuller, K. A.; Malm, W. C.; Kreidenweis, S. M. (1999) Effects of mixing on extinction by carbonaceous particles. *J. Geophys. Res. [Atmos.]* 104: 15,941-15,954.
- Fürst, P.; Krause, G. H. M.; Hein, D.; Delschen, T.; Wilmers, K. (1993) PCDD/PCDF in cow's milk in relation to their levels in grass and soil. *Chemosphere* 27: 1349-1357.
- Gaggi, C.; Bacci, E.; Calamari, D.; Fanelli, R. (1985) Chlorinated hydrocarbons in plant foliage: an indication of the tropospheric contamination level. *Chemosphere* 14: 1673-1686.
- Gallagher, M. W.; Choularton, T. W.; Morse, A. P.; Fowler, D. (1988) Measurements of the size dependence of cloud droplet deposition at a hill site. *Q. J. R. Meteorol. Soc.* 114: 1291-1303.
- Galloway, J. N. (1998) The global nitrogen cycle: changes and consequences. *Environ. Pollut.* 102(suppl. 1): 15-24.
- Galloway, J. N.; Cowling, E. B. (2002) Reactive nitrogen and the world: 200 years of change. *Ambio* 31: 64-71.
- Galloway, J. N.; Aber, J. D.; Erisman, J. W.; Seitzinger, S. P.; Howarth, R. W.; Cowling, E. B.; Cosby, B. J. (2003) The nitrogen cascade. *BioScience* 53: 341-356.
- Garcia-Vallès, M.; Vendrell-Saz, M.; Molera, J.; Blazquez, F. (1998) Interaction of rock and atmosphere: patinas on Mediterranean monuments. *Environ. Geol. (Berlin, Ger.)* 36: 137-149.
- Garland, J. A. (1978) Dry and wet removal of sulphur from the atmosphere. *Atmos. Environ.* 12: 349-362.
- Garland, C.; Garland, F. (1980) Do sunlight and vitamin D reduce the likelihood of colon cancer? *Int. J. Epidemiol.* 9: 227-231.
- Garland, F. C.; Garland, C. F.; Gorham, E. D.; Young, J. F. (1990) Geographic variation in breast cancer mortality in the United States: a hypothesis involving exposure to solar radiation. *Prev. Med.* 19: 614-622.
- Garner, J. H. B. (1991) Ozone exposure and nitrogen loading in the forests of eastern North America. In: Berglund, R. L.; Lawson, D. R.; McKee, D. J., eds. *Tropospheric ozone and the environment: papers from an international conference; March 1990; Los Angeles, CA.* Pittsburgh, PA: Air & Waste Management Association; pp. 289-310. (A&WMA transactions series no. TR-19).
- Garner, J. H. B. (1994) Nitrogen oxides, plant metabolism, and forest ecosystem response. In: Alscher, R. G.; Wellburn, A. R., eds. *Plant responses to the gaseous environment: molecular, metabolic and physiological aspects, [3rd international symposium on air pollutants and plant metabolism]; June 1992; Blacksburg, VA.* London, United Kingdom: Chapman & Hall; pp. 301-314.
- Garner, J. H. B.; Pagano, T.; Cowling, E. B. (1989) An evaluation of the role of ozone, acid deposition, and other airborne pollutants in the forests of eastern North America. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station; general technical report SE-59.
- Garten, C. T., Jr. (1988) Fate and distribution of sulfur-35 in yellow poplar and red maple trees. *Oecologia* 76: 43-50.
- Garten, C. T., Jr.; Hanson, P. J. (1990) Foliar retention of ¹⁵N-nitrate and ¹⁵N-ammonium by red maple (*Acer rubrum*) and white oak (*Quercus alba*) leaves from simulated rain. *Environ. Exp. Bot.* 30: 333-342.
- Garten, C. T., Jr.; Bondietti, E. A.; Lomax, R. D. (1988) Contribution of foliar leaching and dry deposition to sulfate in net throughfall below deciduous trees. *Atmos. Environ.* 22: 1425-1432.
- Gawel, J. E.; Ahner, B. A.; Friedland, A. J.; Morel, F. M. M. (1996) Role for heavy metals in forest decline indicated by phytochelatin measurements. *Nature (London)* 381: 64-65.

- Gazzi, M.; Georgiadis, T.; Vicentini, V. (2001) Distant contrast measurements through fog and thick haze. *Atmos. Environ.* 35: 5143-5149.
- Geron, C.; Rasmussen, R.; Arnts, R. R.; Guenther, A. (2000) A review and synthesis of monoterpene speciation from forests in the United States. *Atmos. Environ.* 34: 1761-1781.
- Ghedini, N.; Gobbi, G.; Sabbioni, C.; Zappia, G. (2000) Determination of elemental and organic carbon on damaged stone monuments. *Atmos. Environ.* 34: 4383-4391.
- Gildon, A.; Tinker, P. B. (1983) Interactions of vesicular-arbuscular mycorrhizal infection and heavy metals in plants: I. the effects of heavy metals on the development of vesicular-arbuscular mycorrhizas. *New Phytol.* 95: 247-261.
- Gilliam, F. S.; Adams, M. B.; Yurish, B. M. (1996) Ecosystem nutrient responses to chronic nitrogen inputs at Fernow Experimental Forest, West Virginia. *Can. J. For. Res.* 26: 196-205.
- Gobbi, G.; Zappia, G.; Sabbioni, C. (1998) Sulphite quantification on damaged stones and mortars. *Atmos. Environ.* 32: 783-789.
- Goodwin, J. E.; Sage, W.; Tilly, G. P. (1969) Study of erosion by solid particles. *Proc. Inst. Mech. Eng.* 184: 279-292.
- Gorham, E. D.; Garland, C. F.; Garland, F. C. (1989) Acid haze air pollution and breast and colon cancer mortality in 20 Canadian cities. *Can. J. Public Health* 80: 96-100.
- Gorham, E. D.; Garland, F. C.; Garland, C. F. (1990) Sunlight and breast cancer incidence in the USSR. *Int. J. Epidemiol.* 19: 820-824.
- Goulder, L. H.; Kennedy, D. (1997) Valuing ecosystem services: philosophical bases and empirical methods. In: Daily, G. C. *Nature's Services: Societal Dependence on Natural Ecosystems*. Washington, DC: Island Press; pp. 23-47.
- Graedel, T. E.; Nassau, K.; Franey, J. P. (1987) Copper patinas formed in the atmosphere—I. Introduction. *Corros. Sci.* 27: 639-657.
- Grand Canyon Visibility Transport Commission. (1996) Report of the Grand Canyon Visibility Transport Commission to the United States Environmental Protection Agency: recommendations for improving western vistas. Denver, CO: Grand Canyon Visibility Transport Commission.
- Grant, L. D. (1989) Health effects issues associated with regional and global air pollution problems. In: *Proceedings of the changing atmosphere: implications for global security*; June, 1988; Toronto, Canada. Geneva, Switzerland: World Meteorological Organization; report no. WMO-No. 710.
- Grant, W. B. (2002) An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer* 6: 1867-1875.
- Grattan, S. R.; Maas, E. V.; Ogata, G. (1981) Foliar uptake and injury from saline aerosol. *J. Environ. Qual.* 10: 406-409.
- Gravenhorst, G.; Hoefken, K. D.; Georgii, H. W. (1983) Acidic input to a beech and spruce forest. In: Beilke, S.; Elshout, A. J., eds. *Acid deposition: proceedings of the CEC workshop organized as part of the concerted action "Physico-chemical behaviour of atmospheric pollutants"*; September 1982; Berlin, Federal Republic of Germany. Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 155-171.
- Greenberg, B. M.; Wilson, M. I.; Huang, X.-D.; Duxbury, C. L.; Gerhardt, K. E.; Gensemer, R. W. (1997) The effects of ultraviolet-B radiation on higher plants. In: Wang, W.; Gorsuch, J. W.; Hughes, J. S., eds. *Plants for Environmental Studies*. New York, NY: Lewis Publishers; pp. 1-35.
- Grennfelt, P. (1987) Deposition processes for acidifying compounds. *Environ. Technol. Lett.* 8: 515-527.
- Grimm, N. B.; Grove, J. M.; Pickett, S. T. A.; Redman, C. L. (2000) Integrated approaches to long-term studies of urban ecological systems. *BioScience* 50: 571-584.
- Gu, L. H.; Baldocchi, D.; Verma, S. B.; Black, T. A.; Vesala, T.; Falge, E. M.; Dowty, P. R. (2002) Advantages of diffuse radiation for terrestrial ecosystem productivity. *J. Geophys. Res. [Atmos.]* 107(D6): 10.1029/2001JD001242.
- Guderian, R. (1977) Accumulation of pollutants in plant organs. In: *Air pollution: phytotoxicity of acidic gases and its significance in air pollution control*. Berlin, Germany: Springer-Verlag; pp. 66-74. (Ecological studies: v. 22).
- Guderian, R., ed. (1985) *Air pollution by photochemical oxidants: formation, transport, control, and effects on plants*. New York, NY: Springer-Verlag. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. *Ecological studies: analysis and synthesis*: v. 52).
- Guderian, R. (1986) Terrestrial ecosystems: particulate deposition. In: Legge, A. H.; Krupa, S. V., eds. *Air pollutants and their effects on the terrestrial ecosystem*. New York, NY: John Wiley & Sons; pp. 339-363. (Advances in environmental science and technology: v. 18).

- Guderian, R.; Tingey, D. T.; Rabe, R. (1985) Effects of photochemical oxidants on plants. In: Guderian, R., ed. Air pollution by photochemical oxidants: formation, transport, control, and effects on plants. Berlin, Federal Republic of Germany: Springer-Verlag; pp. 127-333. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. Ecological studies: analysis and synthesis, v. 52).
- Gundersen, P.; Callesen, I.; De Vries, W. (1998a) Nitrate leaching in forest ecosystems is related to forest floor C/N ratios. *Environ. Pollut.* 102(suppl. 1): 403-407.
- Gundersen, P.; Emmett, B. A.; Kjonaas, O. J.; Koopmans, C. J.; Tietema, A. (1998b) Impact of nitrogen deposition on nitrogen cycling in forests: a synthesis of NITREX data. *For. Ecol. Manage.* 101: 37-55.
- Hamilton, R. S.; Revitt, D. M.; Vincent, K. J.; Butlin, R. N. (1995) Sulphur and nitrogen particulate pollutant deposition on to building surfaces. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. *Sci. Total Environ.* 167: 57-66.
- Hanchette, C. L.; Schwartz, G. G. (1992) Geographic patterns of prostate cancer mortality. *Cancer* 70: 2861-2869.
- Haneef, S. J.; Johnson, J. B.; Thompson, G. E.; Wood, G. C. (1993) Simulation of the degradation of limestones and dolomitic sandstone under dry deposition conditions. In: Corrosion control for low-cost reliability: proceedings [of the] 12th international corrosion congress. Volume 2: process industries plant operation. Houston, TX: National Association of Corrosion Engineers; pp. 700-710.
- Hanley, N.; Spash, C. L. (1993) Cost-benefit analysis and the environment. Hants, United Kingdom: Edward Elgar.
- Hansen, K.; Draaijers, G. P. J.; Ivens, W. P. M. F.; Gundersen, P.; van Leeuwen, N. F. M. (1994) Concentration variations in rain and canopy throughfall collected sequentially during individual rain events. *Atmos. Environ.* 28: 3195-3205.
- Hansen, J. E.; Sato, M.; Ruedy, R. (1997) Radiative forcing and climate response. *J. Geophys. Res.* 102: 6831-6864.
- Hansen, J.; Sato, M.; Ruedy, R.; Lacis, A.; Oinas, V. (2000) Global warming in the twenty-first century: an alternative scenario. *Proc. Natl. Acad. Sci. U. S. A.* 97: 9875-9880.
- Hanson, P. J.; Lindberg, S. E. (1991) Dry deposition of reactive nitrogen compounds: a review of leaf, canopy and non-foliar measurements. *Atmos. Environ. Part A* 25: 1615-1634.
- Hartge, P.; Devessa, S. S.; Graumen, D.; Fears, T. R.; Fraumeni, J. F. (1996) Non-Hodgkin's lymphoma and sunlight. *J. Natl. Cancer Inst.* 88: 298-300.
- Harwell, M. A.; Myers, V.; Young, T.; Bartuska, A.; Gassman, N.; Gentile, J. H.; Harwell, C. C.; Appelbaum, S.; Barko, J.; Causey, B.; Johnson, C.; McLean, A.; Smola, R.; Templet, P.; Tosini, S. (1999) A framework for an ecosystem integrity report card. *BioScience* 49: 543-556.
- Hauhs, M. (1989) Lange Bramke: an ecosystem study of a forested catchment. In: Adriano, D. C.; Havas, M., eds. Acidic precipitation: volume 1, case studies. New York, NY: Springer-Verlag; pp. 275-305. (Adriano, D. C.; Salomons, W., eds. Advances in environmental science).
- Haynie, F. H. (1976) Air pollution effects on stress induced intergranular corrosion of 7005-T53 aluminum alloy. In: Stress corrosion—new approaches. Philadelphia, PA: American Society for Testing and Materials; pp. 32-43. (ASTM special technical publication 610).
- Haynie, F. H. (1980) Theoretical air pollution and climate effects on materials confirmed by zinc corrosion data. In: Sereda, P. J.; Litvan, G. G., eds. Durability of building materials and components: proceedings of the 1st international conference; August 1978; Ottawa, ON, Canada. Philadelphia, PA: American Society for Testing and Materials; pp. 157-175. (ASTM special technical publication 691).
- Haynie, F. H. (1986) Environmental factors affecting corrosion of weathering steel. In: Baboian, R., ed. Materials degradation caused by acid rain: developed from the 20th state-of-the-art symposium of the American Chemical Society; June 1985; Arlington, VA. Washington, DC: American Chemical Society; pp. 163-171. (ACS symposium series 318).
- Haynie, F. H.; Lemmons, T. J. (1990) Particulate matter soiling of exterior paints at a rural site. *Aerosol Sci. Technol.* 13: 356-367.

- Haynie, F. H.; Upham, J. B. (1974) Correlation between corrosion behavior of steel and atmospheric pollution data. In: Coburn, S. K., ed. Corrosion in natural environments: presented at the 76th annual meeting American Society for Testing and Materials; June 1973; Philadelphia, PA. Philadelphia, PA: American Society for Testing and Materials; pp. 33-51. (ASTM special technical publication 558).
- Haywood, J. M.; Shine, K. P. (1995) The effect of anthropogenic sulfate and soot aerosol on the clear sky planetary radiation budget. *Geophys. Res. Lett.* 22: 603-606.
- Haywood, J. M.; Ramaswamy, V.; Soden, B. J. (1999) Tropospheric aerosol climate forcing in clear-sky satellite observations over the oceans. *Science (Washington, DC)* 283: 1299-1303.
- Heal, G. (2000) Valuing ecosystem services. *Ecosystems* 3: 24-30.
- Healey, K. D.; Rickert, K. G.; Hammer, G. L.; Bange, M. P. (1998) Radiation use efficiency increases when the diffuse component of incident radiation is enhanced under shade. *Aust. J. Agric. Res.* 49: 665-672.
- Heath, J. A.; Huebert, B. J. (1999) Cloudwater deposition as a source of fixed nitrogen in a Hawaiian montane forest. *Biogeochemistry* 44: 119-134.
- Hedin, L. O.; Granat, L.; Likens, G. E.; Buishand, T. A.; Galloway, J. N.; Butler, T. J.; Rodhe, H. (1994) Steep declines in atmospheric base cations in regions of Europe and North America. *Nature (London)* 367: 351-354.
- Heichel, G. H.; Hankin, L. (1972) Particles containing lead, chlorine, and bromine detected on trees with an electron microprobe. *Environ. Sci. Technol.* 6: 1121-1122.
- Heichel, G. H.; Hankin, L. (1976) Roadside coniferous windbreaks as sinks for vehicular lead emissions. *J. Air Pollut. Control Assoc.* 26: 767-770.
- Heinsdorf, D. (1993) The role of nitrogen in declining Scots pine forests (*Pinus sylvestris*) in the lowland of East Germany. *Water Air Soil Pollut.* 69: 21-35.
- Hemming, B. L.; Seinfeld, J. H. (2001) On the hygroscopic behavior of atmospheric organic aerosols. *Ind. Eng. Chem. Res.* 40: 4162-4171.
- Hermosin, B.; Saiz-Jimenez, C. (2000) Characterization of the organic compounds deposited on monument surfaces. *Fresenius' Environ. Bull.* 9: 322-327.
- Hess, A.; Iyer, H.; Malm, W. (2001) Linear trend analysis: a comparison of methods. *Atmos. Environ.* 35: 5211-5222.
- Hicks, B. B. (1986) Differences in wet and dry particle deposition parameters between North America and Europe. In: Lee, S. D.; Schneider, T.; Grant, L. D.; Verkerk, P. J., eds. *Aerosols: research, risk assessment and control strategies; proceedings of the second U.S.-Dutch international symposium; May 1985; Williamsburg, VA.* Chelsea, MI: Lewis Publishers; pp. 973-982.
- Hicks, B. B.; Matt, D. R.; McMillen, R. T.; Womack, J. D.; Shetter, R. E. (1984) Eddy fluxes of nitrogen oxides to a deciduous forest in complex terrain. In: Samson, P. J., ed. *Transactions: the meteorology of acid deposition, an APCA specialty conference; October 1983; Hartford, CT.* Pittsburgh, PA: Air Pollution Control Association; pp. 189-201.
- Hicks, B. B.; Baldocchi, D. D.; Meyers, T. P.; Hosker, R. P., Jr.; Matt, D. R. (1987) A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. *Water Air Soil Pollut.* 36: 311-330.
- Hicks, B. B.; Matt, D. R.; McMillen, R. T.; Womack, J. D.; Wesely, M. L.; Hart, R. L.; Cook, D. R.; Lindberg, S. E.; de Pena, R. G.; Thomson, D. W. (1989) A field investigation of sulfate fluxes to a deciduous forest. *J. Geophys. Res. [Atmos.]* 94: 13,003-13,011.
- Hidy, G. M.; Hales, J. M.; Roth, P. M.; Scheffe, R. (2000) Fine particles and oxidant pollution: developing an agenda for cooperative research. *J. Air Waste Manage. Assoc.* 50: 613-632.
- Hinckley, T.; Ford, D.; Segura, G.; Sprugel, D. (1992) Key processes from tree to stand level. In: Wall, G., ed. *Implications of climate change for Pacific Northwest forest management.* Waterloo, ON, Canada: University of Waterloo, Department of Geography; pp. 33-43. (Department of Geography publication series: occasional paper no. 15).
- Höfken, K. D.; Gravenhorst, G. (1982) Deposition of atmospheric aerosol particles to beech- and spruce forest. In: Georgii, H.-W.; Pankrath, J., eds. *Deposition of atmospheric pollutants: proceedings of a colloquium; November 1981; Oberursel/Taunus, Federal Republic of Germany.* Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 191-194.
- Höfken, K. D.; Meixner, F. X.; Ehhalt, D. H. (1983) Deposition of atmospheric trace constituents onto different natural surfaces. In: Pruppacher, H. R.; Semonin, R. G.; Slinn, W. G. N., eds. *Precipitation scavenging, dry deposition, and resuspension: v. 2, dry deposition and resuspension: proceedings of the fourth international conference; November-December 1982; Santa Monica, CA.* New York, NY: Elsevier; pp. 825-835.

- Hofstra, G.; Hall, R. (1971) Injury on roadside trees: leaf injury on pine and white cedar in relation to foliar levels of sodium and chloride. *Can. J. Bot.* 49: 613-622.
- Hogan, G. D.; Rennenberg, H.; Fink, S. (1998) Role and effect of sulfur in tree biology. In: Maynard, D. G., ed. *Sulfur in the environment*. New York, NY: Marcel Dekker, Inc.; pp. 173-217. (Books in soils, plants, and the environment: v. 67).
- Holbrow, G. L. (1962) Atmospheric pollution: its measurement and some effects on paint. *J. Oil Colour Chem. Assoc.* 45: 701-718.
- Holland, D. M.; Principe, P. P.; Sickles, J. E., II. (1999) Trends in atmospheric sulfur and nitrogen species in the eastern United States for 1989-1995. *Atmos. Environ.* 33: 37-49.
- Hollander, M.; Wolfe, D. A. (1973) *Nonparametric statistical methods*. New York: John Wiley & Sons.
- Holling, C. S. (1986) The resilience of terrestrial ecosystems: local surprise and global change. In: Clark, W. C.; Munn, R. E., eds. *Sustainable development of the biosphere*. Cambridge, UK: Cambridge University; pp. 292-317.
- Hooper, D. U.; Vitousek, P. M. (1997) The effects of plant composition and diversity on ecosystem processes. *Science* 277: 1302-1305.
- Hornung, M.; Langan, S. J. (1999) Nitrogen deposition, sources, impacts and responses in natural and semi-natural ecosystems. In: Bangan, S. J., ed. *Impact of nitrogen deposition on natural ecosystems and semi-natural ecosystems*. Dordrecht, Netherlands: Kluwer Academic Publishers; pp. 1-14. [Environmental Pollution, no.3].
- Horvath, H. (1993) Atmospheric light absorption—a review. *Atmos. Environ. Part A* 27: 293-317.
- Hosker, R. P., Jr.; Lindberg, S. E. (1982) Review: atmospheric deposition and plant assimilation of gases and particles. *Atmos. Environ.* 16: 889-910.
- Houghton, J. T.; Jenkins, G. J.; Ephraums, J. J., eds. (1990) *Climate change: the IPCC scientific assessment*. Cambridge, MA: Cambridge University Press; p. 55.
- Howarth, R. W.; Boyer, E. W.; Pabich, W. J.; Galloway, J. N. (2002) Nitrogen use in the United States from 1961-2000 and potential future trends. *Ambio* 31: 88-96.
- Hoyt, D. V. (1978) A model for the calculation of solar global insolation. *Sol. Energy* 21: 27-35.
- Huebert, B. J.; Luke, W. T.; Delany, A. C.; Brost, R. A. (1988) Measurements of concentrations and dry surface fluxes of atmospheric nitrates in the presence of ammonia. *J. Geophys. Res. [Atmos.]* 93: 7127-7136.
- Hughes, M. K. (1981) Cycling of trace metals in ecosystems. In: Lepp, N. W., ed. *Effect of heavy metal pollution on plants*. Volume 2: metals in the environment. London, United Kingdom: Applied Science Publishers; pp. 95-118.
- Humphreys, F. R.; Lambert, M. J.; Kelly, J. (1975) The occurrence of sulphur deficiency in forests. In: McLachlan, K. D., ed. *Sulphur in Australasian agriculture*. Sydney, Australia: Sydney University Press; pp. 154-162.
- Hunsaker, C.; Graham, R.; Turner, R. S.; Ringold, P. L.; Holdren, G. R., Jr.; Strickland, T. C. (1993) A national critical loads framework for atmospheric deposition effects assessment: II. Defining assessment end points, indicators, and functional subregions. *Environ. Manage. (N. Y.)* 17: 335-341.
- Huntington, T.G. (2000) The potential for calcium depletion in forest ecosystems of southeastern United States: review and analysis. *Global Biogeochem. Cycles* 14: 623-638.
- Husar, R. B.; Wilson, W. E. (1993) Haze and sulfur emission trends in the eastern United States. *Environ. Sci. Technol.* 27: 12-16.
- Husar, R. B.; Holloway, J. M.; Patterson, D. E.; Wilson, W. E. (1981) Spatial and temporal pattern of eastern U.S. haziness: a summary. *Atmos. Environ.* 15: 1919-1928.
- Husar, R. B.; Elkins, J. B.; Wilson, W. E. (1994) U.S. visibility trends, 1960-1992, regional and national. Presented at: 87th annual meeting of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-MP3.05.
- Hutchinson, A. J.; Johnson, J. B.; Thompson, G. E.; Wood, G. C.; Sage, P. W.; Cooke, M. J. (1992) The role of fly-ash particulate material and oxide catalysts in stone degradation. *Atmos. Environ. Part A* 26: 2795-2803.
- Huttunen, S. (1994) Effects of air pollutants on epicuticular wax structure. In: Percy, K. E.; Cape, J. N.; Jagels, R.; Simpson, C. J., eds. *Air pollutants and the leaf cuticle*. New York, NY: Springer-Verlag; pp. 81-96. (NATO ASI series).
- Hyer, K. E.; Webb, J. R.; Eshleman, K. N. (1995) Episodic acidification of three streams in Shenandoah National Park, Virginia, USA. *Water Air Soil Pollut.* 85: 523-528.
- Ibrahim, M.; Barrie, L. A.; Fanaki, F. (1983) An experimental and theoretical investigation of the dry deposition of particles to snow, pine trees and artificial collectors. *Atmos. Environ.* 17: 781-788.

- IMPROVE: interagency monitoring of protected visual environments [database]. (1998) [Data on gross chemical composition of PM_{2.5} particles in rural areas of the eastern and western United States as of 1998]. Fort Collins, CO: National Park Service (NPS); Cooperative Institute for Research in the Atmosphere (CIRA). Available at: <http://vista.cira.colostate.edu/improve/> [2001, January 26].
- Intergovernmental Panel on Climate Change (IPCC). (1998) The regional impacts of climate change: an assessment of vulnerability. Cambridge, United Kingdom: Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (2001a) Climate change 2001: the scientific basis. Contribution of working group I to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (2001b) Climate change 2001: impacts, adaptation, and vulnerability. Contribution of working group II to the third assessment report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom: Cambridge University Press.
- Iyer, H.; Patterson, P.; Malm, W. C. (2000) Trends in the extremes of sulfur concentration distributions. *J. Air Waste Manage. Assoc.* 50: 802-808.
- Jacobson, M. Z. (1998) Studying the effects of aerosols on vertical photolysis rate coefficient and temperature profiles over an urban airshed. *J. Geophys. Res. [Atmos.]* 103: 10,593-10,604.
- Jacobson, M. Z. (1999) Isolating nitrated and aromatic aerosols and nitrated aromatic gases as sources of ultraviolet light absorption. *J. Geophys. Res.* 104 (D3): 3527-3542.
- Jacobson, M. Z. (2000) A physically-based treatment of elemental carbon optics: implications for global direct forcing of aerosols. *Geophys. Res. Lett.* 27: 217-220.
- Jaworski, N. A.; Howarth, R. W.; Hetling, L. J. (1997) Atmospheric deposition of nitrogen oxides onto the landscape contributes to coastal eutrophication in the northeast United States. *Environ. Sci. Technol.* 31: 1995-2004.
- Jensen, V. (1974) Decomposition of angiosperm tree leaf litter. In: Dickinson, C. H.; Pugh, G. J. F., eds. *Biology of plant litter decomposition: v. I*. London, United Kingdom: Academic Press; pp. 69-104.
- John, W.; Wall, S. M.; Ondo, J. L. (1985) Dry acid deposition on materials and vegetation: concentrations in ambient air. Berkeley, CA: California Department of Health Services, Air and Industrial Hygiene Laboratory; report no. CA/DOH/AIHL/SP-34. Available from: NTIS, Springfield, VA; PB85-241206.
- Johnson, D. W. (1984) Sulfur cycling in forests. *Biogeochemistry* 1: 29-43.
- Johnson, D. W. (1992) Base cation distribution and cycling. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag, Inc.; pp. 275-333. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. *Ecological studies, analysis and synthesis: v. 91*).
- Johnson, F. R.; Desvousges, W. H. (1997) Estimating stated preferences with rated-pair data: environmental, health, and employment effects of energy programs. *J. Environ. Econ. Manage.* 34: 79-99.
- Johnson, D. W.; Lindberg, S. E., eds. (1992a) *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag, Inc. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. *Ecological studies: analysis and synthesis: v. 91*).
- Johnson, D. W.; Lindberg, S. E., eds. (1992b) Nitrogen chemistry, deposition, and cycling in forests. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag, Inc.; pp. 150-213. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. *Ecological studies: analysis and synthesis: v. 91*).
- Johnson, D. W.; Mitchell, M. J. (1998) Responses of forest ecosystems to changing sulfur inputs. In: Maynard, D. G., ed. *Sulfur in the environment*. New York, NY: Marcel Dekker, Inc.; pp. 219-262. [*Books in Soils, Plants, and the Environment*, v. 67].
- Johnson, D. W.; Taylor, G. E. (1989) Role of air pollution in forest decline in eastern North America. *Water Air Soil Pollut.* 48: 21-43.
- Johnson, D. W.; Todd, D. E., Jr. (1998) Harvesting effects on long-term changes in nutrient pools of mixed oak forest. *Soil Sci. Soc. Amer. J.* 62: 1725-1735.
- Johnson, D. W.; Henderson, G. S.; Huff, D. D.; Lindberg, S. E.; Richter, D. D.; Shriner, D. S.; Todd, D. E.; Turner, J. (1982a) Cycling of organic and inorganic sulphur in a chestnut oak forest. *Oecologia* 54: 141-148.
- Johnson, D. W.; Cole, D. W.; Bledsoe, C. S.; Cromack, K.; Edmonds, R. L.; Gessel, S. P.; Grier, C. C.; Richards, B. N.; Vogt, K. A. (1982b) Nutrient cycling in forests of the Pacific Northwest. In: Edmonds, R. L., ed. *Analysis of coniferous forest ecosystems in the Western United States*. Stroudsburg, PA: Hutchinson Ross Publishing Company; pp. 186-232. (*US/IBP synthesis series: v. 14*).
- Johnson, D. W.; Van Miegroet, H.; Lindberg, S. E.; Todd, D. E.; Harrison, R. B. (1991a) Nutrient cycling in red spruce forests of the Great Smoky Mountains. *Can. J. For. Res.* 21: 769-787.

- Johnson, D. W.; Cresser, M. S.; Nilsson, S. I.; Turner, J.; Ulrich, B.; Binkley, D.; Cole, D. W. (1991b) Soil changes in forest ecosystems: evidence for and probable causes. *Proc. R. Soc. Edinburgh Sect. B: Biol. Sci.* 97B: 81-116.
- Johnson, D. W.; Swank, W. T.; Vose, J. M. (1993) Simulated effects of atmospheric sulfur deposition on nutrient cycling in a mixed deciduous forest. *Biogeochemistry* 23: 169-196.
- Johnson, A. H.; Friedland, A. J.; Miller, E. K.; Siccama, T. G. (1994) Acid rain and soils of the Adirondacks. III. Rates of soil acidification in a montane spruce-fir forest at Whiteface Mountain, New York. *Can. J. For. Res.* 24: 663-669.
- Johnson, D.W.; Susfalk, R.B.; Brewer, P.,F.; Swank, W.T. (1999) Simulated effects of reduced sulfur, nitrogen, and base cation deposition on soils and solutions in southern Appalachian forests. *J. Environ. Qual* 28: 1336-1346.
- Joslin, J. D.; Wolfe, M. H. (1992) Red spruce soil chemistry and root distribution across a cloud water deposition gradient. *Can. J. For. Res.* 22: 893-904.
- Joslin, J. D.; Wolfe, M. H. (1994) Foliar deficiencies of mature Southern Appalachian red spruce determined from fertilizer trials. *Soil Sci. Soc. Am. J.* 58: 1572-1579.
- Joslin, J. D.; Kelly, J. M.; Van Miegroet, H. (1992) Soil chemistry and nutrition of North American spruce-fir stands: evidence for recent change. *J. Environ. Qual.* 21: 12-30.
- Justus, C. G.; Murphey, B. B. (1994) Temporal trends in surface irradiance at ultraviolet wavelengths. *J. Geophys. Res.* 99 (D1): 1389-1394.
- Kaminski, M. D.; Landsberger, S. (2000a) Heavy metals in urban soils of East St. Louis, IL, part I: total concentration of heavy metals in soils. *J. Air Waste Manage. Assoc.* 50: 1667-1679.
- Kaminski, M. D.; Landsberger, S. (2000b) Heavy metals in urban soils of East St. Louis, IL, part II: leaching characteristics and modeling. *J. Air Waste Manage. Assoc.* 50: 1680-1687.
- Kandeler, E.; Kampichler, C.; Horak, O. (1996) Influence of heavy metals on the functional diversity of soil microbial communities. *Biol. Fertil. Soils* 23: 299-306.
- Karmoker, J. L.; Clarkson, D. T.; Saker, L. R.; Rooney, J. M.; Purves, J. V. (1991) Sulphate deprivation depresses the transport of nitrogen to the xylem and the hydraulic conductivity of barley (*Hordeum vulgare L.*) roots. *Planta* 185: 269-278.
- Kaupp, H.; McLachlan, M. S. (1999) Atmospheric particle size distributions of polychlorinated dibenzo-*p*-dioxins and dibenzofurans (PCDD/Fs) and polycyclic aromatic hydrocarbons (PAHs) and their implications for wet and dry deposition. *Atmos. Environ.* 33: 85-95.
- Kelly, J.; Lambert, M. J. (1972) The relationship between sulphur and nitrogen in the foliage of *Pinus radiata*. *Plant Soil* 37: 395-407.
- Kelly, J. M.; Meagher, J. F. (1986) Nitrogen input/output relationships for three forested watersheds in eastern Tennessee. In: Correll, D. L., ed. *Watershed research perspectives*. Washington, DC: Smithsonian Institution Press; pp. 360-391.
- Kenk, G.; Fischer, H. (1988) Evidence from nitrogen fertilisation in the forests of Germany. *Environ. Pollut.* 54: 199-218.
- Kiehl, J. T.; Briegleb, B. P. (1993) The relative roles of sulfate aerosols and greenhouse gases in climate forcing. *Science* (Washington, DC) 260: 311-314.
- Kim, E.; Kalman, D.; Larson, T. (2000) Dry deposition of large airborne particles on a surrogate surface. *Atmos. Environ.* 34: 2387-2397.
- Kleeman, M. J.; Eldering, A.; Hall, J. R.; Cass, G. R. (2001) Effect of emissions control programs on visibility in southern California. *Environ. Sci. Technol.* 35: 4668-4674.
- Klepper, B.; Craig, D. K. (1975) Deposition of airborne particulates onto plant leaves. *J. Environ. Qual.* 4: 495-499.
- Kodama, Y.; Lee, S. D., eds. (1994) *The 13th UOEH international symposium and the 2nd pan Pacific cooperative symposium on impact of increased UV-B exposure on human health and ecosystem; October 1993; Kitakyushu, Japan.* Kitakyushu, Japan: University of Occupational and Environmental Health.
- Koslow, E. E.; Smith, W. H.; Staskawicz, B. J. (1977) Lead-containing particles on urban leaf surfaces. *Environ. Sci. Technol.* 11: 1019-1021.
- Krajčičková, A.; Mejstřík, V. (1984) The effect of fly-ash particles on the plugging of stomata. *Environ. Pollut.* 36: 83-93.
- Kripke, M. L. (1989) Health effects of stratospheric ozone depletion: an overview. In: Schneider, T.; Lee, S. D.; Wolters, G. J. R.; Grant, L. D. eds. *Atmospheric ozone research and its policy implications: proceedings of the Third US-Dutch International Symposium; May, 1988; Nijmegen, The Netherlands.* Amsterdam, The Netherlands: Elsevier, pp. 795-802. [Studies in Environmental Science 35].
- Krupa, S. V.; Legge, A. H. (1986) Single or joint effects of coarse and fine particulate sulfur aerosols and ozone on vegetation. In: Lee, S. D.; Schneider, T.; Grant, L. D.; Verkerk, P. J., eds. *Aerosols: research, risk assessment*

- and control strategies, proceedings of the second U.S.-Dutch international symposium; May 1985; Williamsburg, VA. Chelsea, MI: Lewis Publishers, Inc.; pp. 879-887.
- Krupa, S. V.; Legge, A. H. (1998) Sulphur dioxide, particulate sulphur and its impacts on a boreal forest ecosystem. In: Ambasht, R. S., ed. *Modern trends in ecology and environment*. Leiden, The Netherlands: Backhuys Publishers; pp. 285-306.
- Kylin, H.; Grimvall, E.; Östman, C. (1994) Environmental monitoring of polychlorinated biphenyls using pine needles as passive samplers. *Environ. Sci. Technol.* 28: 1320-1324.
- Lamaud, E.; Chapuis, A.; Fontan, J.; Serie, E. (1994) Measurements and parameterization of aerosol dry deposition in a semi-arid area. *Atmos. Environ.* 28: 2461-2471.
- Lamersdorf, N. P.; Meyer, M. (1993) Nutrient cycling and acidification of a northwest German forest site with high atmospheric nitrogen deposition. *For. Ecol. Manage.* 62: 323-354.
- Laurenroth, W. K.; Milchunas, D. G. (1985) SO₂ effects on plant community function. In: Winner, W. E.; Mooney, H. A.; Goldstein, R. A., eds. *Sulfur dioxide and vegetation: physiology, ecology, and policy issues*. Stanford, CA: Stanford University Press; pp. 454-477.
- Lawrence, G. B.; Huntington, T. G. (1999) Soil-Calcium depletion linked to acid rain and forest growth in the eastern United States. Reston, VA: U.S. Geological Survey. Available: <http://bqs.usgs.gov/acidrain/WRIR984267.pdf> [29 September, 2004].
- Lefkowitz, E. S.; Garland, C. F. (1994) Sunlight, vitamin D, and ovarian cancer mortality rates in U.S. women. *Int. J. Epidemiol.* 23: 1133-1136.
- Legge, A. H. (1980) Primary productivity, sulfur dioxide, and the forest ecosystem: an overview of a case study. In: Miller, P. R., ed. *Proceedings of symposium on effects of air pollutants on Mediterranean and temperate forest ecosystems*; June; Riverside, CA. Berkeley, CA: U.S. Department of Agriculture, Pacific Southwest Forest and Range Experiment Station; pp. 51-62; Forest Service general technical report no. PSW-43. Available from: NTIS, Springfield, VA; PB81-133720.
- Lerman, S. L.; Darley, E. F. (1975) Particulates. In: Mudd, J. B.; Kozlowski, T. T., eds. *Responses of plants to air pollution*. New York, NY: Academic Press; pp. 141-158.
- Leslie, M. (2001) Having a bad air day. *Science* (Washington, DC) 294: 2255.
- Levin, S. A. (1998) Ecosystems and the biosphere as complex adaptive systems. *Ecosystems* 1: 431-436.
- Li-Jones, X.; Prospero, J. M. (1998) Variations in the size distribution of non-sea-salt sulfate aerosol in the marine boundary layer at Barbados: impact of African dust. *J. Geophys. Res. [Atmos.]* 103: 16,073-16,084.
- Likens, G. E.; Driscoll, C. T.; Buso, D. C. (1996) Long-term effects of acid rain: response and recovery of a forest ecosystem. *Science* (Washington, DC) 272: 244-246.
- Likens, G. E.; Driscoll, C. T.; Buso, D. C.; Siccama, T. G.; Johnson, C. E.; Lovett, G. M.; Fahey, T. J.; Reiners, W. A.; Ryan, D. F.; Martin, C. W.; Bailey, S. W. (1998) The biogeochemistry of calcium at Hubbard Brook. *Biogeochemistry* 41: 89-173.
- Likens, G. E.; Butler, T. J.; Buso, D. C. (2001) Long- and short-term changes in sulfate deposition: effects of the 1990 Clean Air Act Amendments. *Biogeochemistry* 52: 1-11.
- Lin, J.-M.; Fang, G.-C.; Holsen, T. M.; Noll, K. E. (1993) A comparison of dry deposition modeled from size distribution data and measured with a smooth surface for total particle mass, lead and calcium in Chicago. *Atmos. Environ. Part A* 27: 1131-1138.
- Lindberg, S. E. (1992) Atmospheric deposition and canopy interactions of sulfur. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag, Inc.; pp. 74-90. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. *Ecological studies: analysis and synthesis: v. 91*).
- Lindberg, S. E.; Garten, C. T., Jr. (1988) Sources of sulphur in forest canopy throughfall. *Nature* (London) 336: 148-151.
- Lindberg, S. E.; Harriss, R. C. (1981) The role of atmospheric deposition in an eastern U.S. deciduous forest. *Water Air Soil Pollut.* 16: 13-31.
- Lindberg, S. E.; Lovett, G. M. (1985) Field measurements of particle dry deposition rates to foliage and inert surfaces in a forest canopy. *Environ. Sci. Technol.* 19: 238-244.
- Lindberg, S. E.; Lovett, G. M. (1992) Deposition and forest canopy interactions of airborne sulfur: results from the integrated forest study. *Atmos. Environ. Part A* 26: 1477-1492.

- Lindberg, S. E.; McLaughlin, S. B. (1986) Air pollutant interactions with vegetation: research needs in data acquisition and interpretation. In: Legge, A. H.; Krupa, S. V., eds. Air pollutants and their effects on the terrestrial ecosystem. New York, NY: John Wiley & Sons, Inc.; pp. 449-503. (Nriagu, J. O., ed. Advances in environmental science and technology: v. 18).
- Lindberg, S. E.; Owens, J. G. (1993) Throughfall studies of deposition to forest edges and gaps in montane ecosystems. *Biogeochemistry* 19: 173-194.
- Lindberg, S. E.; Harriss, R. C.; Turner, R. R. (1982) Atmospheric deposition of metals to forest vegetation. *Science* (Washington, DC) 215: 1609-1611.
- Lindberg, S. E.; Lovett, G. M.; Richter, D. D.; Johnson, D. W. (1986) Atmospheric deposition and canopy interactions of major ions in a forest. *Science* (Washington, DC) 231: 141-145.
- Lindberg, S. E.; Lovett, G. M.; Meiwes, K.-J. (1987) Deposition and forest canopy interactions of airborne nitrate. In: Hutchinson, T. C.; Meema, K. M., eds. Effects of atmospheric pollutants on forests, wetlands and agricultural ecosystems. Berlin, Federal Republic of Germany: Springer-Verlag; pp. 117-130. (NATO as 1 series: v. G16).
- Lindberg, S. E.; Lovett, G. M.; Schaefer, D. A.; Bredemeier, M. (1988) Dry deposition velocities and surface-to-canopy scaling factors for aerosol calcium from forest canopy throughfall. *J. Aerosol Sci.* 19: 1187-1190.
- Lindberg, S. E.; Bredemeier, M.; Schaefer, D. A.; Qi, L. (1990) Atmospheric concentrations and deposition of nitrogen and major ions in conifer forests in the United States and Federal Republic of Germany. *Atmos. Environ. Part A* 24: 2207-2220.
- Liousse, C.; Cachier, H.; Jennings, S. G. (1993) Optical and thermal measurements of black carbon aerosol content in different environments: variation of the specific attenuation cross-section, sigma (σ). *Atmos. Environ. Part A* 27: 1203-1211.
- Little, P. (1977) Deposition of 2.75, 5.0 and 8.5 μm particles to plant and soil surfaces. *Environ. Pollut.* 12: 293-305.
- Little, P.; Wiffen, R. D. (1977) Emission and deposition of petrol engine exhaust Pb--I. deposition of exhaust Pb to plant and soil surfaces. *Atmos. Environ.* 11: 437-447.
- Liu, S. C.; McKeen, S. A.; Madronich, S. (1991) Effect of anthropogenic aerosols on biologically active ultraviolet radiation. *Geophys. Res. Lett.* 18: 2265-2268.
- Lloyd, J. (2002) After years of smog, a shroud lifts. *The Christian Science Monitor*. June 4: 1(s. 1).
- Loehman, E.; Boldt, D.; Chaikin, K. (1985) Measuring the benefits of air quality improvements in the San Francisco Bay area: property value and contingent valuation studies, volume IV of methods development for environmental control benefits assessment. Washington, DC: U.S. Environmental Protection Agency; Office of Policy, Planning and Evaluation; report no. EPA-230-12-85-022. Available: <http://yosemite.epa.gov/EE/epa/eerm.nsf/vwRepNumLookup/EE-0278D?OpenDocument> (28 May 2003).
- Løkke, H.; Bak, J.; Falkengren-Grerup, U.; Finlay, R. D.; Ilvesniemi, H.; Nygaard, P. H.; Starr, M. (1996) Critical loads of acidic deposition for forest soils: is the current approach adequate. *Ambio* 25: 510-516.
- Lorente, J.; Redaño, A.; de Cabo, X. (1994) Influence of urban aerosol on spectral solar irradiance. *J. Appl. Meteorol.* 33: 406-415.
- Lorusso, S.; Marabelli, M.; Troili, M. (1997) Air pollution and the deterioration of historic monuments. *J. Environ. Pathol. Toxicol. Oncol.* 16: 171-173.
- Lovett, G. M. (1984) Atmospheric deposition to forests. In: Breece, L.; Hasbrouck, S., eds. Forest responses to acidic deposition: proceedings of the U. S.-Canadian conference; August 1983; Orono, ME. Orono, ME: University of Maine, Orono, Land and Water Resources Center; pp. 7-18.
- Lovett, G. M. (1992) Atmospheric deposition and canopy interactions of nitrogen. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer-Verlag, Inc.; pp. 152-166. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. Ecological studies: analysis and synthesis: v. 91).
- Lovett, G. M. (1994) Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. *Ecol. Appl.* 4: 629-650.
- Lovett, G. M.; Lindberg, S. E. (1984) Dry deposition and canopy exchange in a mixed oak forest as determined by analysis of throughfall. *J. Appl. Ecol.* 21: 1013-1027.
- Lovett, G. M.; Lindberg, S. E. (1986) Dry deposition of nitrate to a deciduous forest. *Biogeochemistry* 2: 137-148.
- Lovett, G. M.; Lindberg, S. E. (1992) Concentration and deposition of particles and vapors in a vertical profile through a forest canopy. *Atmos. Environ. Part A* 26: 1469-1476.
- Lovett, G. M.; Lindberg, S. E. (1993) Atmospheric deposition and canopy interactions of nitrogen in forests. *Can. J. For. Res.* 23: 1603-1616.

- Lovett, G. M.; Traynor, M. M.; Pouyat, R. V.; Carreiro, M. M.; Zhu, W.-X.; Baxter, J. W. (2000) Atmospheric deposition to oak forest along an urban-rural gradient. *Environ. Sci. Technol.* 34: 4294-4300.
- Lowenthal, D. H.; Rogers, C. F.; Saxena, P.; Watson, J. G.; Chow, J. C. (1995) Sensitivity of estimated light extinction coefficients to model assumptions and measurement errors. *Atmos. Environ.* 29: 751-766.
- Ludwig, J.; Marufu, L. T.; Huber, B.; Andreae, M. O.; Helas, G. (2003) Domestic combustion of biomass fuels in developing countries: a major source of atmospheric pollutants. *J. Atmos. Chem.* 44: 23-37.
- Madronich, S. (1987) Photodissociation in the atmosphere. 1. Actinic flux and the effects of ground reflections and clouds. *J. Geophys. Res. [Atmos.]* 92: 9740-9752.
- Magill, A. H.; Aber, J. D.; Berntson, G. M.; McDowell, W. H.; Nadelhoffer, K. J.; Melillo, J. M.; Steudler, P. (2000) Long-term nitrogen additions and nitrogen saturation in two temperate forests. *Ecosystems* 3: 238-253.
- Malm, W. C. (1999) Introduction to visibility. Fort Collins, CO: Colorado State University; Cooperative Institute for Research in the Atmosphere (CIRA); NPS Visibility Program. Available: <http://www.epa.gov/air/visibility/introvis.pdf> [2 April 2002].
- Malm, W. C. (2000) Spatial and seasonal patterns and temporal variability of haze and its constituents in the United States. Report III. Fort Collins, CO: Cooperative Institute for Research in the Atmosphere, Colorado State University. Available: <http://vista.cira.colostate.edu/improve/Publications/Reports/2000/2000.htm> [22 March, 2002].
- Malm, W. C.; Kreidenweis, S. M. (1997) The effects of models of aerosol hygroscopicity on the apportionment of extinction. *Atmos. Environ.* 31: 1965-1976.
- Malm, W. C.; Sisler, J. F.; Huffman, D.; Eldred, R. A.; Cahill, T. A. (1994) Spatial and seasonal trends in particle concentration and optical extinction in the United States. *J. Geophys. Res. [Atmos.]* 99: 1347-1370.
- Malm, W. C.; Molenaar, J. V.; Eldred, R. A.; Sisler, J. F. (1996) Examining the relationship among atmospheric aerosols and light scattering and extinction in the Grand Canyon area. *J. Geophys. Res. [Atmos.]* 101: 19,251-19,265.
- Malm, W. C.; Day, D.; Kreidenweis, S. M. (1997) Comparison of measured and reconstructed scattering during an intensive field study at Great Smoky Mountains National Park. Presented at: 90th annual meeting and exhibition of the Air & Waste Management Association; June; Toronto, ON, CA. Pittsburgh, PA: Air & Waste Management Association; 97-WA70.02.
- Malm, W. C.; Day, D. E.; Kreidenweis, S. M. (2000a) Light scattering characteristics of aerosols as a function of relative humidity: part I—a comparison of measured scattering and aerosol concentrations using the theoretical models. *J. Air Waste Manage. Assoc.* 50: 686-700.
- Malm, W. C.; Day, D. E.; Kreidenweis, S. M. (2000b) Light scattering characteristics of aerosols as a function of relative humidity: part II—a comparison of measured scattering and aerosol concentrations using the statistical models. *J. Air Waste Manage. Assoc.* 50: 701-709.
- Mansfeld, F. (1980) Regional air pollution study: effects of airborne sulfur pollutants on materials. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Sciences Research Laboratory; EPA report no. EPA-600/4-80-007. Available from: NTIS, Springfield, VA; PB81-126351.
- Marchwińska, E.; Kucharski, R. (1987) The combined influence of SO₂ and heavy metal-containing particulates on beans, carrots and parsley. *Environ. Monit. Assess.* 8: 11-25.
- Markewitz, D.; Richter, D.D.; Allen, L. H.; Urrego, J. B. (1998) Three decades of observed soil acidification in the Calhoun Experimental Forest: Has acid rain made a difference? *Soil Sci. Soc. Amer. J.* 62: 1428-1439.
- Marschner, H. (1995) Sulfur. In: Mineral nutrition of higher plants. 2nd ed. London, United Kingdom: Academic Press; pp. 255-265.
- Martin, M. H.; Coughtrey, P. J. (1981) Impact of metals on ecosystem function and productivity. In: Lepp, N. W., ed. Effect of heavy metal pollution on plants: volume 2, metals in the environment. Barking, United Kingdom: Applied Science Publishers; pp. 119-158. (Mellanby, K., ed. Pollution monitoring series).
- Martin, C. E.; Gravatt, D. A.; Loesch, V. S. (1992) Photosynthetic responses of three species to acute exposures of nitrate- and sulphate-containing aerosols. *Atmos. Environ. Part A* 26: 381-391.
- Mathai, C. V. (1995) The Grand Canyon visibility. *EM (December)*: 20-31.
- Matson, P. A.; Parton, W. J.; Power, A. G.; Swift, M. J. (1997) Agricultural intensification and ecosystem properties. *Science (Washington, DC)* 277: 504-509.
- May, P. F.; Till, A. R.; Cumming, M. J. (1972) Systems analysis of ³⁵sulphur kinetics in pastures grazed by sheep. *J. Appl. Ecol.* 9: 25-49.
- McCartney, H. A.; Aylor, D. E. (1987) Relative contributions of sedimentation and impaction to deposition of particles in a crop canopy. *Agric. For. Meteorol.* 40: 343-358.

- McClelland, G.; Schulze, W.; Waldman, D.; Schenk, D.; Irwin, J.; Stewart, T.; Deck, L.; Thayer, M. (1993) Valuing eastern visibility: a field test of the contingent valuation method. Washington, DC: U.S. Environmental Protection Agency, Office of Policy, Planning and Evaluation; cooperative agreement no. CR-815183.
- McCune, D. C.; Silberman, D. H.; Mandl, R. H.; Weinstein, L. H.; Freudenthal, P. C.; Giardina, P. A. (1977) Studies on the effects of saline aerosols of cooling tower origin on plants. *J. Air Pollut. Control Assoc.* 27: 319-324.
- McDonald, R. L.; Unni, C. K.; Duce, R. A. (1982) Estimation of atmospheric sea salt dry deposition: wind speed and particle size dependence. *J. Geophys. Res.* 87: 1246-1250.
- McDonnell, M. J.; Pickett, S. T. A.; Groffman, P.; Bohlen, P.; Pouyat, R. V.; Zipperer, W. C.; Parmelee, R. W.; Carreiro, M. M.; Medley, K. (1997) Ecosystem processes along an urban-to-rural gradient. *Urban Ecosyst.* 1: 21-36.
- McDow, S. R.; Vartiainen, M.; Sun, Q.; Hong, Y.; Yao, Y.; Kamens, R. M. (1995) Combustion aerosol water content and its effect on polycyclic aromatic hydrocarbon reactivity. *Atmos. Environ.* 29: 791-797.
- McGee, E. S.; Mossotti, V. G. (1992) Gypsum accumulation on carbonate stone. *Atmos. Environ. Part B* 26: 249-253.
- McGowan, T. F.; Lipinski, G. E.; Santoleri, J. J. (1993) New rules affect the handling of waste fuels. *Chem. Eng. (March)*: 122-128.
- McIntyre, N. E.; Knowles-Yáñez, K.; Hope, D. (2000) Urban ecology as an interdisciplinary field: differences in the use of "urban" between the social and natural sciences. *Urban Ecosyst.* 4: 5-24.
- McKay, W. A.; Garland, J. A.; Livesley, D.; Halliwell, C. M.; Walker, M. I. (1994) The characteristics of the shore-line sea spray aerosol and the landward transfer of radionuclides discharged to coastal sea water. *Atmos. Environ.* 28: 3299-3309.
- McLachlan, M. S. (1996a) Bioaccumulation of hydrophobic chemicals in agricultural food chains. *Environ. Sci. Technol.* 30: 252-259.
- McLachlan, M. S. (1996b) Biological uptake and transfer of polychlorinated dibenzo-*p*-dioxins and dibenzofurans. In: Hester, R. E.; Harrison, R. M., eds. Chlorinated organic micropollutants. Cambridge, United Kingdom: Royal Society of Chemistry; pp. 31-52. (Issues in environmental science and technology).
- McLachlan, M. S. (1999) Framework for the interpretation of measurements of SOCs in plants. *Environ. Sci. Technol.* 33: 1799-1804.
- McMahon, T. A.; Denison, P. J. (1979) Empirical atmospheric deposition parameters—a survey. *Atmos. Environ.* 13: 571-585.
- McMillen, R. T. (1988) An eddy correlation technique with extended applicability to non-simple terrain. *Boundary Layer Meteorol.* 43: 231-245.
- McMurry, P. H. (2000) A review of atmospheric aerosol measurements. *Atmos. Environ.* 34: 1959-1999.
- McMurry, P. H.; Stolzenburg, M. R. (1989) On the sensitivity of particle size to relative humidity for Los Angeles aerosols. *Atmos. Environ.* 23: 497-507.
- McMurry, P. H.; Zhang, X.; Lee, C.-T. (1996) Issues in aerosol measurement for optics assessments. *J. Geophys. Res. [Atmos.]* 101: 19,189-19,197.
- Meiwes, K. J.; Khanna, P. K. (1981) Distribution and cycling of sulphur in the vegetation of two forest ecosystems in an acid rain environment. *Plant Soil* 60: 369-375.
- Middleton, W. E. K. (1952) *Vision through the atmosphere*. Toronto, ON, Canada: University of Toronto Press.
- Midwest Regional Planning Organization. (2001) *Regional haze and visibility in the upper Midwest*. Des Plaines, IL: Midwest Regional Planning Organization.
- Milford, J. B.; Davidson, C. I. (1987) The sizes of particulate sulfate and nitrate in the atmosphere—a review. *JAPCA* 37: 125-134.
- Millar, C. S. (1974) Decomposition of coniferous leaf litter. In: Dickinson, C. H.; Pugh, G. J. F., eds. *Biology of plant litter decomposition: volume I*. New York, NY: Academic Press; pp. 105-128.
- Miller, P. R.; McBride, J. R., eds. (1999) *Oxidant air pollution impacts in the Montane forests of southern California: a case study of the San Bernardino Mountains*. New York, NY: Springer-Verlag. (Ecological Studies: v. 134.)
- Miller, W. C.; Fornes, R. E.; Gilbert, R. D.; Speer, A.; Spence, J. (1992) Removal of CaCO₃ extender in residential coatings by atmospheric acidic deposition. In: *Measurement of toxic and related air pollutants: proceedings of the 1992 U.S. EPA/A&WMA international symposium*. Pittsburgh, PA: Air & Waste Management Association; pp. 129-134. (A&WMA publication VIP-25).
- Miller, E. K.; Panek, J. A.; Friedland, A. J.; Kadlec, J.; Mohnen, V. A. (1993) Atmospheric deposition to a high-elevation forest at Whiteface Mountain, New York, USA. *Tellus Ser. B* 45: 209-227.
- Mims, F. M., III. (1996) Significant reduction of UVB caused by smoke from biomass burning in Brazil. *Photochem. Photobiol.* 64: 814-816.

- Mitchell, M. J.; Harrison, R. B.; Fitzgerald, J. W.; Johnson, D. W.; Lindberg, S. E.; Zhang, Y.; Autry, A. (1992a) Sulfur distribution and cycling in forest ecosystems. In: Johnson, D. W.; Lindberg, S. E., eds. *Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study*. New York, NY: Springer-Verlag; pp. 90-129. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Remmert, H., eds. *Ecological studies: analysis and synthesis: v. 91*).
- Mitchell, M. J.; Foster, N. W.; Shepard, J. P.; Morrison, I. K. (1992b) Nutrient cycling in Huntington Forest and Turkey Lakes deciduous stands: nitrogen and sulfur. *Can. J. For. Res.* 22: 457-464.
- Molenaar, J. V.; Malm, W. C.; Johnson, C. E. (1994) Visual air quality simulation techniques. *Atmos. Environ.* 28: 1055-1063.
- Molenaar, J. V. (2000) *Visibility science and trends in the Lake Tahoe Basin: 1989-1998*. Stateline, NV: Tahoe Regional Planning Agency; prepared by Air Resource Specialists, Inc.; February 15.
- Molenaar, J. V.; Malm, W. C.; Johnson, C. E. (1994) Visual air quality simulation techniques. *Atmos. Environ.* 28: 1055-1063.
- Monteith, J. L.; Unsworth, M. H. (1990) *Principles of environmental physics*. 2nd ed. London, United Kingdom: Edward Arnold; pp. 165-175.
- Moomaw, W. R. (2002) Energy, industry and nitrogen: strategies for decreasing reactive nitrogen emissions. *Ambio* 31: 184-189.
- Moosmüller, H.; Arnott, W. P.; Rogers, C. F. (1998) Photoacoustic and filter measurements related to aerosol light absorption during the Northern Front Range Air Quality Study (Colorado 1996/1997). *J. Geophys. Res. [Atmos.]* 103: 28,149-28,157.
- Moosmüller, H.; Arnott, W. P.; Rogers, C. F.; Bowen, J. L.; Gillies, J. A.; Pierson, W. R.; Collins, J. F.; Durbin, T. D.; Norbeck, J. M. (2001) Time-resolved characterization of diesel particulate emissions. 2. Instruments for elemental and organic carbon measurements. *Environ. Sci. Technol.* 35: 1935-1942.
- Moropoulou, A.; Bisbikou, K.; Torfs, K.; Van Grieken, R.; Zezza, F.; Macri, F. (1998) Origin and growth of weathering crusts on ancient marbles in industrial atmosphere. *Atmos. Environ.* 32: 967-982.
- Mueller, S. F. (1988) *Chemical deposition to high-elevation spruce-fir forests in the eastern United States*. Muscle Shoals, AL: Tennessee Valley Authority, Air Quality Branch.
- Naeem, S.; Thompson, L. J.; Lawler, S. P.; Lawton, J. H.; Woodfin, R. M. (1994) Declining biodiversity can alter the performance of ecosystems. *Nature (London)* 368: 734-737.
- Nash, T. H., III. (1975) Influence of effluents from a zinc factory on lichens. *Ecol. Monogr.* 45: 183-198.
- National Park Service. (1998) *IMPROVE program goals and objectives*. Washington, DC: National Park Service. Available: www2.nature.nps.gov/ard/impr/index.htm [24 November 1999].
- National Research Council (NRC). (1993) *Protecting visibility in national parks and wilderness areas*. Washington, DC: National Academy Press. 3v.
- National Science and Technology Council. (1998) *National acid precipitation assessment program biennial report to Congress: an integrated assessment; executive summary*. Silver Spring, MD: U.S. Department of Commerce, National Oceanic and Atmospheric Administration. Available: www.nnic.noaa.gov/CENR/NAPAP/NAPAP_96.htm [24 November 1999].
- National Weather Service. (1998) *Visibility*. Washington, DC: National Weather Service, National Oceanic and Atmospheric Administration. Available: www.nws.noaa.gov/asos/vsby.htm [24 November 1999].
- Nazaroff, W. W.; Cass, G. R. (1991) Protecting museum collections from soiling due to the deposition of airborne particles. *Atmos. Environ. Part A* 25: 841-852.
- Nelis, P. M.; Branford, D.; Unsworth, M. H. (1994) A model of the transfer of radioactivity from sea to land in sea spray. *Atmos. Environ.* 28: 3213-3223.
- Neuberger, H.; Hosler, C. L.; Kocmond, W. C. (1967) Vegetation as aerosol filter In: Tromp, S. W.; Weihe, W. H., eds. *Biometeorology, v. 2*. pp. 693-702.
- Neuroth, G.; Heisler, S. (2000) *The nature and sources of visibility impairment in Phoenix*. Presented at: Governor's Brown Cloud Summit; August; Phoenix, AZ. Phoenix, AZ: Arizona Department of Environmental Quality. Available: <http://www.adeq.state.az.us/environ/air/browncld/download/gary.pdf> (28 May 2003).
- Nicholson, K. W. (1988) The dry deposition of small particles: a review of experimental measurements. *Atmos. Environ.* 22: 2653-2666.
- Nicholson, K. W.; Davies, T. D. (1987) Field measurements of the dry deposition of particulate sulphate. *Atmos. Environ.* 21: 1561-1571.
- Nielsen, K. E.; Ladekarl, U. L.; Nornberg, P. (1999) Dynamic soil processes on heathland due to changes in vegetation to oak and Sitka spruce. *For. Ecol. Manage.* 114: 107-116.
- Nihlgård, B. (1985) The ammonium hypothesis—an additional explanation to the forest dieback in Europe. *Ambio* 14: 2-8.

- Niklińska, M.; Laskowski, R.; Maryański, M. (1998) Effect of heavy metals and storage time on two types of forest litter: basal respiration rate and exchangeable metals. *Ecotoxicol. Environ. Safety* 41: 8-18.
- Noble, I. R.; Dirzo, R. (1997) Forests as human-dominated ecosystems. *Science (Washington, DC)* 277: 522-525.
- Nodvin, S. C.; Van Miegroet, H.; Lindberg, S. E.; Nicholas, N. S.; Johnson, D. W. (1995) Acidic deposition, ecosystem processes, and nitrogen saturation in a high elevation southern Appalachian watershed. *Water Air Soil Pollut.* 85: 1647-1652.
- Noll, K. E.; Fang, K. Y. P. (1989) Development of a dry deposition model for atmospheric coarse particles. *Atmos. Environ.* 23: 585-594.
- Nord, A. G.; Ericsson, T. (1993) Chemical analysis of thin black layers on building stone. *Stud. Conserv.* 38: 25-35.
- Northeast States for Coordinated Air Use Management (NESCAUM). (2002) CAMNET: realtime air pollution and visibility monitoring. Boston, MA: NESCAUM. Available at: <http://www.hazecam.net>. (23 January 2002).
- Norton, R. A.; Childers, N. F. (1954) Experiments with urea sprays on the peach. *Proc. Am. Soc. Hortic. Sci.* 63: 23-31.
- Novakov, T.; Penner, J. E. (1993) Large contribution of organic aerosols to cloud-condensation-nuclei concentrations. *Nature (London)* 365: 823-826.
- Ockenden, W. A.; Stinnes, E.; Parker, C.; Jones, K. C. (1998) Observations on persistent organic pollutants in plants: implications for their use as passive air samplers and for POP cycling. *Environ. Sci. Technol.* 32: 2721-2726.
- Odum, E. P. (1969) The strategy of ecosystem development: an understanding of ecological succession provides a basis for resolving man's conflict with nature. *Science (Washington, DC)* 164: 262-270.
- Odum, E. P. (1985) Trends expected in stressed ecosystems. *BioScience* 35: 419-422.
- Odum, E. P. (1989) *Ecology and our endangered life-support systems*. Sunderland, MA: Sinauer Associates, Inc.; pp. 58-66, 108-118.
- Odum, E. P. (1993) The ecosystem. In: *Ecology and our endangered life-support systems*. 2nd ed. Sunderland, MA: Sinauer Associates, Inc.; pp. 38-67.
- Oehme, M.; Biseth, A.; Schlabach, M.; Wiig, Ø. (1995) Concentrations of polychlorinated dibenzo-*p*-dioxins, dibenzofurans and non-*ortho* substituted biphenyls in polar bear milk from Svalbard (Norway). *Environ. Pollut.* 90: 401-407.
- Olszyk, D. M.; Bytnerowicz, A.; Takemoto, B. K. (1989) Photochemical oxidant pollution and vegetation: effects of gases, fog, and particles. *Environ. Pollut.* 61: 11-29.
- Oosting, H. J.; Billings, W. D. (1942) Factors effecting vegetational zonation on coastal dunes. *Ecology* 23: 131-142.
- Ormrod, D. P. (1984) Impact of trace element pollution on plants. In: Treshow, M., ed. *Air pollution and plant life*. Chichester, United Kingdom: John Wiley & Sons; pp. 291-319.
- Ouimette, J. R.; Flagan, R. C. (1982) The extinction coefficient of multicomponent aerosols. *Atmos. Environ.* 16: 2405-2420.
- Padgett, P.; Allen, E. B. (1999) Differential responses to nitrogen fertilization in native shrubs and exotic annuals common to Mediterranean coastal sage scrub of California. *Plant Ecol.* 144: 93-101.
- Padgett, P. E.; Allen, E. B.; Bytnerowicz, A.; Minich, R. A. (1999) Changes in soil inorganic nitrogen as related to atmospheric nitrogenous pollutants in southern California. *Atmos. Environ.* 33: 769-781.
- Paerl, H. W.; Bales, J. D.; Ausley, L. W.; Buzzelli, C. P.; Crowder, L. B.; Eby, L. A.; Go, M.; Peierls, B. L.; Richardson, T. L.; Ramus, J. S. (2001) Ecosystem impacts of three sequential hurricanes (Dennis, Floyd, and Irene) on the United States' largest lagoonal estuary, Pamlico Sound, NC. *Proc. Natl. Acad. Sci. U. S. A.* 98: 5655-5611.
- Pan, W.; Tatang, M. A.; McRae, G. J.; Prinn, R. G. (1997) Uncertainty analysis of direct radiative forcing by anthropogenic sulfate aerosols. *J. Geophys. Res. [Atmos.]* 102: 21,915-21,924.
- Pan, W.; Tatang, M. A.; McRae, G. J.; Prinn, R. G. (1998) Uncertainty analysis of indirect radiative forcing by anthropogenic sulfate aerosols. *J. Geophys. Res. [Atmos.]* 103: 3815-3823.
- Pandis, S. N.; Seinfeld, J. H. (1989) Mathematical modeling of acid deposition due to radiation fog. *J. Geophys. Res. [Atmos.]* 94: 12,911-12,923.
- Pankow, J. F. (1987) Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere. *Atmos. Environ.* 21: 2275-2283.
- Parish, S. B. (1910) The effect of cement dust on citrus trees. *Plant World* 13: 288-291.
- Parker, A. (1955) The destructive effects of air pollution on materials. *Colliery Guardian* 191: 447-450.
- Parker, G. G. (1983) Throughfall and stemflow in the forest nutrient cycle. In: Macfadyen, A.; Ford, E. D., eds. *Advances in ecological research*, v. 13; pp. 57-120.

- Patz, J. A.; Engelberg, D.; Last, J. (2000a) The effects of changing weather on public health. *Ann. Rev. Public Health* 21: 271-307.
- Patz, J. A.; McGeehin, M. A.; Bernard, S. M.; Ebi, K. L.; Epstein, P. R.; Grambsch, A.; Gubler, D. J.; Reiter, P.; Romieu, I.; Rose, J. B.; Samet, J. M.; Trtanj, J. (2000b) The potential health impacts of climate variability and change for the United States: executive summary of the report of the health sector of the U.S. national assessment. *Environ. Health Perspect.* 108: 367-376.
- Pearcy, R. W.; Björkman, O.; Caldwell, M. M.; Keeley, J. E.; Monson, R. K.; Strain, B. R. (1987) Carbon gain by plants in natural environments. *BioScience* 37: 21-29.
- Penndorf, R. B. (1958) An approximation method to the Mie theory for colloidal spheres. *J. Phys. Chem.* 62: 1537-1542.
- Penner, J. E.; Charlson, R. J.; Hales, J. M.; Laulainen, N. S.; Leifer, R.; Novakov, T.; Ogren, J.; Radke, L. F.; Schwartz, S. E.; Travis, L. (1994) Quantifying and minimizing uncertainty of climate forcing by anthropogenic aerosols. *Bull. Am. Meteorol. Soc.* 75: 375-400.
- Percy, K. E. (1991) Effects of acid rain on forest vegetation: morphological and non-mensurational growth effects. In: Rennie, P. J.; Robitaille, G., eds. *Effects of Acid Rain on Forest Resources: Proceedings of the Conference*; June, 1983; Sainte-Foy, Quebec. Ottawa, Canada: Canadian Forestry Service; pp. 97-111.
- Perry, K. D.; Cahill, T. A.; Eldred, R. A.; Dutcher, D. D.; Gill, T. E. (1997) Long-range transport of North African dust to the eastern United States. *J. Geophys. Res. [Atmos]* 102: 11,225-11,238.
- Peterjohn, W. T.; Adams, M. B.; Gilliam, F. S. (1996) Symptoms of nitrogen saturation in two central Appalachian hardwood forest ecosystems. *Biogeochemistry* 35: 507-522.
- Peters, K.; Eiden, R. (1992) Modelling the dry deposition velocity of aerosol particles to a spruce forest. *Atmos. Environ. Part A* 26: 2555-2564.
- Peterson, J. T.; Flowers, E. C.; Rudisill, J. H. (1978) Urban-rural solar radiation and atmospheric turbidity measurements in the Los Angeles basin. *J. Appl. Meteorol.* 17: 1595-1609.
- Peterson, G.; Allen, C. R.; Holling, C. S. (1998) Ecological resilience, biodiversity, and scale. *Ecosystems* 1: 6-18.
- Petuskey, W. T.; Richardson, D. A.; Dolske, D. A. (1995) Aspects of the deterioration of sandstone masonry in Anasazi dwelling ruins at Mesa Verde National Park, Colorado, USA. *Sci. Total Environ.* 167: 145-159.
- Piatt, J. R.; Krause, P. D. (1974) Road and site characteristics that influence road salt distribution and damage to roadside aspen trees. *Water Air Soil Pollut.* 3: 301-304.
- Pickett, S. T. A.; Cadenasso, M. L.; Grove, J. M.; Nilon, C. Y.; Pouyat, R. V.; Zipperer, W. C.; Costanza, R. (2001) Urban ecological systems: linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas. *Annu. Rev. Ecol. Syst.* 32: 127-157.
- Pilinis, C.; Pandis, S. N.; Seinfeld, J. H. (1995) Sensitivity of direct climate forcing by atmospheric aerosols to aerosol size and composition. *J. Geophys. Res. [Atmos.]* 100: 18739-18754.
- Pimentel, D.; Wilson, C.; McCullum, C.; Huang, R.; Dwen, P.; Flack, J.; Tran, Q.; Saltman, T.; Cliff, B. (1997) Economic and environmental benefits of biodiversity. *BioScience* 47: 747-757.
- Pimm, S. L. (1984) The complexity and stability of ecosystems. *Nature (London)* 307: 321-326.
- Pitchford, M. L.; Malm, W. C. (1994) Development and applications of a standard visual index. *Atmos. Environ.* 28: 1049-1054.
- Pitchford, M. L.; McMurry, P. H. (1994) Relationship between measured water vapor growth and chemistry of atmospheric aerosol for Grand Canyon, Arizona, in winter 1990. *Atmos. Environ.* 28: 827-839.
- Pouyat, R. V.; McDonnell, M. J. (1991) Heavy metal accumulations in forest soils along an urban-rural gradient in southeastern New York, USA. *Water Air Soil Pollut.* 57/58: 797-808.
- Prescott, C. E.; Parkinson, D. (1985) Effects of sulphur pollution on rates of litter decomposition in a pine forest. *Can. J. Bot.* 63: 1436-1443.
- Pryor, S. C. (1996) Assessing public perception of visibility for standard setting exercises. *Atmos. Environ.* 30: 2705-2716.
- Pryor, S.; Steyn, D. (1994) Visibility and ambient aerosols in southwestern British Columbia during REVEAL (REgional Visibility Experimental Assessment in the Lower Fraser Valley): application of analytical techniques for source apportionment and related visibility impacts of the REVEAL fine particulate optical and scene data. Victoria, BC, Canada: British Columbia Ministry of Environment, Lands and Parks. Available from: NTIS, Springfield, VA; MIC-96-00314/INS.
- Pueschel, R. F. (1993) Potential climatic effects of anthropogenic aerosols. In: Jennings, S. G., ed. *Aerosol effects on climate*. Tucson, AZ: University of Arizona Press; pp. 110-132.
- Rabalais, N. N. (2002) Nitrogen in aquatic ecosystems. *Ambio* 31: 102-112.

- Rae, D. A. (1983) The value to visitors of improving visibility at Mesa Verde and Great Smoky National Parks. In: Rowe, R. D.; Chestnut, L. G., eds. Managing air quality and scenic resources at national parks and wilderness areas. Boulder, CO: Westview Press; pp. 217-234.
- Ragsdale, H. L.; Lindberg, S. E.; Lovett, G. M.; Schaefer, D. A. (1992) Atmospheric deposition and throughfall fluxes of base cations. In: Johnson, D. W.; Lindberg, S. E., eds. Atmospheric deposition and forest nutrient cycling: a synthesis of the integrated forest study. New York, NY: Springer Verlag; pp. 235-253. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. Ecological studies: analysis and synthesis: v. 91).
- Ramanathan, V.; Crutzen, P. J.; Kiehl, J. T.; Rosenfeld, D. (2001) Aerosols, climate and the hydrological cycle. Science (Washington, DC, U.S.) 294: 2119-2124.
- Rapport, D. J.; Whitford, W. G. (1999) How ecosystems respond to stress: common properties of arid and aquatic systems. BioScience 49: 193-203.
- Raynal, D. J.; Joslin, J. D.; Thornton, F. C.; Schaedle, M.; Henderson, G. S. (1990) Sensitivity of tree seedlings to aluminum: III. Red spruce and loblolly pine. J. Environ. Qual. 19: 180-187.
- Realini, M.; Negrotti, R.; Appollonia, L.; Vaudan, D. (1995) Deposition of particulate matter on stone surfaces: an experimental verification of its effects on Carrara marble. In: Saiz-Jimenez, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. Sci. Total Environ. 167: 67-72.
- Reiners, W. A.; Olson, R. K. (1984) Effects of canopy components on throughfall chemistry: an experimental analysis. Oecologia 63: 320-330.
- Reisinger, L. M. (1990) Analysis of airborne particles sampled in the southern Appalachian Mountains. Water Air Soil Pollut. 50: 149-162.
- Rennie, P. J. (1955) The uptake of nutrients by mature forest growth. Plant Soil 7: 49-95.
- Reuss, J. O. (1983) Implications of the calcium-aluminum exchange system for the effect of acid precipitation on soils. J. Environ. Qual. 12: 591-595.
- Reuss, J. O.; Johnson, D. W. (1986) Acid deposition and the acidification of soils and waters. New York, NY: Springer-Verlag. (Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. S.; Rimmert, H., eds. Ecological studies: analysis and synthesis: v. 59).
- Riggan, P. J.; Lockwood, R. N.; Lopez, E. N. (1985) Deposition and processing of airborne nitrogen pollutants in Mediterranean-type ecosystems of southern California. Environ. Sci. Technol. 19: 781-789.
- Rochette, P.; Desjardins, R. L.; Pattey, E.; Lessard, R. (1996) Instantaneous measurement of radiation and water use efficiencies of a maize crop. Agron. J. 88: 627-635.
- Rodney, D. R. (1952) The entrance of nitrogen compounds through the epidermis of apple leaves. Proc. Am. Soc. Hortic. Sci. 59: 99-102.
- Roelofs, J. G. M.; Kempers, A. J.; Houdijk, A. L. F. M.; Jansen, J. (1985) The effect of air-borne ammonium sulphate on *Pinus nigra* var. *maritima* in the Netherlands. Plant Soil 84: 45-56.
- Roelofs, J. G. M.; Boxman, A. W.; Van Dijk, H. F. G. (1987) Effects of airborne ammonium on natural vegetation and forests. In: Asman, W. A. H.; Diederik, H. S. M. A., eds. Ammonia and acidification: proceedings [of a] symposium of the European Association for the Science of Air Pollution (EURASAP); April; Bilthoven, The Netherlands. European Association for the Science of Air Pollution; pp. 266-276.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993a) Sources of fine organic aerosol. 4. Particulate abrasion products from leaf surfaces of urban plants. Environ. Sci. Technol. 27: 2700-2711.
- Rogge, W. F.; Hildemann, L. M.; Mazurek, M. A.; Cass, G. R.; Simoneit, B. R. T. (1993b) Sources of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust: roads as sources and sinks. Environ. Sci. Technol. 27: 1892-1904.
- Rosenberg, N. J.; Blad, B. L.; Verma, S. B. (1983) Microclimate: the biological environment. 2nd ed. New York, NY: John Wiley & Sons.
- Rosenfeld, D. (1999) TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall. Geophys. Res. Lett. 26: 3105-3108.
- Rovira, A. D.; Davey, C. B. (1974) Biology of the rhizosphere. In: Carson, E. W., ed. The plant root and its environment: proceedings of an institute; July, 1971; Blacksburg, VA. Charlottesville, VA: University Press of Virginia; pp. 153-204.
- Rueth, H.; Baron, J. S. (2002) Differences in Englemann spruce forest biogeochemistry east and west of the Continental Divide in Colorado, USA. Ecosystems 5: 45-57.
- Ruijgrok, W.; Tieben, H.; Eisinga, P. (1997) The dry deposition of particles to a forest canopy: a comparison of model and experimental results. Atmos. Environ. 31: 399-415.

- Sabbioni, C. (1995) Contribution of atmospheric deposition to the formation of damage layers. In: Jimenez-Saiz, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. *Sci. Total Environ.* 167: 49-56.
- Sabbioni, C.; Zappia, G.; Gobbi, G. (1992) Carbonaceous particles on carbonate building stones in a simulated system. In: Proceedings of the 1992 European aerosol conference; September; Oxford, United Kingdom. *J. Aerosol Sci.* 23(suppl. 1): S921-S924.
- Sabbioni, C.; Zappia, G.; Gobbi, G. (1996) Carbonaceous particles and stone damage in a laboratory exposure system. *J. Geophys. Res. [Atmos.]* 101: 19,621-19,627.
- Sabbioni, C.; Zappia, G.; Ghedini, N.; Gobbi, G.; Favoni, O. (1998) Black crusts on ancient mortars. *Atmos. Environ.* 32: 215-223.
- Saiz-Jimenez, C. (1993) Deposition of airborne organic pollutants on historic buildings. *Atmos. Environ. Part B* 27: 77-85.
- Salby, M. L. (1996) Fundamentals of atmospheric physics. New York, NY: Academic Press. (The International Geophysics Series, Textbooks and Monographs, v. 61).
- Samecka-Cymerman, A.; Kempers, A. J. (1999) Bioindication of heavy metals in the town Wroc/aw (Poland) with evergreen plants. *Atmos. Environ.* 33: 419-430.
- Samuels, H. J.; Twiss, S.; Wong, E. W. (1973) Visibility, light scattering and mass concentration of particulate matter: report of the California Tri-City Aerosol Sampling Project. Sacramento, CA: State of California, Air Resources Board.
- Sanyal, B.; Singhania, G. K. (1956) Atmospheric corrosion of metals: part I. *J. Sci. Ind. Res. Sect. B* 15: 448-455.
- Saunders, P. J. W.; Godzik, S. (1986) Terrestrial vegetation-air pollutant interactions: nongaseous air pollutants. In: Legge, A. H.; Krupa, S. V., eds. *Air pollutants and their effects on the terrestrial ecosystem*. New York, NY: John Wiley & Sons; pp. 389-394. (Advances in environmental science and technology: v. 18).
- Saxena, P.; Peterson, T. W. (1981) Thermodynamics of multicomponent electrolytic aerosols. *J. Colloid Interface Sci.* 79: 496-510.
- Saxena, P.; Mueller, P. K.; Kim, Y. P.; Seinfeld, J. H.; Koutrakis, P. (1993) Coupling thermodynamic theory with measurements to characterize acidity of atmospheric particles. *Aerosol Sci. Technol.* 19: 279-293.
- Saxena, P.; Hildemann, L. M.; McMurry, P. H.; Seinfeld, J. H. (1995) Organics alter hygroscopic behavior of atmospheric particles. *J. Geophys. Res. [Atmos.]* 100: 18,755-18,770.
- Schiavon, N.; Chiavari, G.; Schiavon, G.; Fabbri, D. (1995) Nature and decay effects of urban soiling on granitic building stones. In: Jimenez-Saiz, C., ed. The deterioration of monuments: proceedings of the 2nd international symposium on biodeterioration and biodegradation; January 1994; Sevilla, Spain. *Sci. Total Environ.* 167: 87-101.
- Schichtel, B. A.; Husar, R. B.; Falke, S. R.; Wilson, W. E. (2001) Haze trends over the United States, 1980-1995. *Atmos. Environ.* 35: 5205-5210.
- Schier, G. A.; Jensen, K. (1992) Atmospheric deposition effects on foliar injury and foliar leaching in red spruce. In: Eager, C.; Adams, M. B., eds. *Ecology and decline of red spruce in the eastern United States*. New York, NY: Springer-Verlag; pp. 271-294. [Billings, W. D.; Golley, F.; Lange, O. L.; Olson, J. D.; Remmert, H., eds. *Ecological Studies, Analysis and Synthesis Ecological Studies*, v. 96].
- Schimek, M. G. (1981) Smoothing and regression. New York, NY: Wiley.
- Schneider, S. H. (1992) Introduction to climate modeling. In: Trenberth, K. E. *Climate system modeling*. New York, NY: Cambridge University Press; pp. 1-26.
- Schnug, E. (1997) Significance of sulphur for the quality of domesticated plants. In: Cram, W. J.; De Kok, L. J.; Stulen, I.; Brunold, C.; Rennenberg, H., eds. *Sulphur metabolism in higher plants: molecular, ecophysiological and nutritional aspects*. Leiden, The Netherlands: Backhuys Publishers; pp. 109-130.
- Schönherr, J.; Huber, R. (1977) Plant cuticles are polyelectrolytes with isoelectric points around three. *Plant Physiol.* 59: 145-150.
- Schulze, E.-D. (1989) Air pollution and forest decline in a spruce (*Picea abies*) forest. *Science (Washington, DC)* 244: 776-783.
- Schuster, P. F.; Reddy, M. M.; Sherwood, S. I. (1994) Effects of acid rain and sulfur dioxide on marble dissolution. *Mater. Perform.* 33: 76-80.
- Schwar, M. J. R. (1998) Nuisance dust deposition and soiling rate measurements. *Environ. Technol.* 19: 223-229.
- Science Advisory Board. (2002) A framework for assessing and reporting on ecological condition: an SAB report. Young, T. F.; Sanzone, S., eds. Washington, DC: U.S. Environmental Protection Agency, Ecological Processes and Effects Committee; report no. EPA-SAB-EPEC-02-009. Available at: <http://www.epa.gov/science1/index.html> (27 January 2003).

- Seckmeyer, G.; McKenzie, R. L. (1992) Increased ultraviolet radiation in New Zealand (45 degrees S) relative to Germany (48 degrees N). *Nature (London)* 359: 135-137.
- Sehmel, G. A. (1980) Particle and gas dry deposition: a review. *Atmos. Environ.* 14: 983-1011.
- Sehmel, G. A.; Hodgson, W. H. (1976) Predicted dry deposition velocities. In: *Atmosphere-surface exchange of particulate and gaseous pollutants (1974): proceedings of a symposium; September 1974; Richland, WA. Oak Ridge, TN: Energy Research and Development Administration; pp. 399-422. (ERDA symposium series: 38). Available from: NTIS, Springfield, VA; CONF-740921.*
- Seinfeld, J. H. (1989) Urban air pollution: state of the science. *Science (Washington, DC)* 243: 745-752.
- Seinfeld, J. H.; Pandis, S. N. (1998) *Atmospheric chemistry and physics: from air pollution to climate change.* New York, NY: John Wiley & Sons, Inc.
- Sereda, P. J. (1974) Weather factors affecting corrosion of metals. In: *Corrosion in natural environments: three symposia presented at the seventy-sixth annual meeting [of the] American Society for Testing and Materials; June 1973; Philadelphia, PA. Philadelphia, PA: American Society for Testing and Materials; pp. 7-22. (ASTM special technical publication 558).*
- Shannon, J. D. (1999) Regional trends in wet deposition of sulfate in the United States and SO₂ emissions from 1980 through 1995. *Atmos. Environ.* 33: 807-816.
- Sharples, S.; Stewart, L.; Tregenza, P. R. (2001) Glazing daylight transmittances: a field survey of windows in urban areas. *Build. Environ.* 36: 503-509.
- Shinn, J. H. (1978) A critical survey of measurements of foliar deposition of airborne sulfates and nitrates. Presented at: 71st annual meeting of the Air Pollution Control Association; June; Houston, TX. Pittsburgh, PA: Air Pollution Control Association; paper no. 78-7.2.
- Shortle, W. C.; Bondietti, E. A. (1992) Timing, magnitude, and impact of acidic deposition on sensitive forest sites. *Water Air Soil Pollut.* 61: 253-267.
- Shortle, W. C.; Smith, K. T. (1988) Aluminum-induced calcium deficiency syndrome in declining red spruce. *Science (Washington, DC)* 240: 1017-1018.
- Shortle, W. C.; Smith, K. T.; Minocha, R.; Lawrence, G. B.; David, M. B. (1997) Acidic deposition, cation mobilization, and biochemical indicators of stress in healthy red spruce. *J. Environ. Qual.* 26: 871-876.
- Showak, W.; Dunbar, S. R. (1982) Effect of 1 percent copper addition on atmospheric corrosion of rolled zinc after 20 years' exposure. In: Dean, S. W.; Rhea, E. C., eds. *Atmospheric corrosion of metals: a symposium; May 1980; Denver, CO. Philadelphia, PA: American Society for Testing and Materials; pp. 135-162. (ASTM special technical publication 767).*
- Shriner, D. S.; Henderson, G. S. (1978) Sulfur distribution and cycling in a deciduous forest watershed. *J. Environ. Qual.* 7: 392-397.
- Sievering, H. (1987) Dynamics of sulfur exchange at the air/forest canopy interface: a review of throughfall inferred deposition rates. *Global Biogeochem. Cycles* 1: 233-249.
- Sievering, H.; Rusch, D.; Marquez, L. (1996) Nitric acid, particulate nitrate and ammonium in the continental free troposphere: nitrogen deposition to an alpine tundra ecosystem. *Atmos. Environ.* 30: 2527-2537.
- Simonich; S. L.; Hites, R. A. (1995) Organic pollutant accumulation in vegetation. *Environ. Sci. Technol.* 29: 2905-2914.
- Simpson, J. W.; Horrobin, P. J., eds. (1970) *The weathering and performance of building materials.* New York, NY: Wiley-Interscience.
- Sinclair, T. R.; Shiraiwa, T.; Hammer, G. L. (1992) Variation in crop radiation-use efficiency with increased diffuse-radiation. *Crop Sci.* 32: 1281-1284.
- Sisler, J. F. (1996) Spatial and seasonal patterns and long term variability of the composition of the haze in the United States: an analysis of data from the IMPROVE network. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere; EPA air docket A096-56, document no. VI-B-09-(ee).
- Sisler, J. F.; Malm, W. C. (2000) Interpretation of trends of PM_{2.5} and reconstructed visibility from the IMPROVE network. *J. Air Waste Manage. Assoc.* 50: 775-789.
- Sisler, J. F.; Huffman, D.; Lattimer, D. A.; Malm, W. C.; Pitchford, M. L. (1993) Spatial and temporal patterns and the chemical composition of the haze in the United States: an analysis of data from the IMPROVE network, 1988-1991. Fort Collins, CO: Colorado State University, Cooperative Institute for Research in the Atmosphere (CIRA).
- Skeffington, R. A.; Wilson, E. J. (1988) Excess nitrogen deposition: issues for consideration. *Environ. Pollut.* 54: 159-184.
- Skerry, B. S.; Alavi, A.; Lindgren, K. I. (1988) Environmental and electrochemical test methods for the evaluation of protective organic coatings. *J. Coat. Technol.* 60: 97-106.

- Slinn, W. G. N. (1977) Some approximations for the wet and dry removal of particles and gases from the atmosphere. *Water Air Soil Pollut.* 7: 513-543.
- Slinn, W. G. N. (1978) Parameterizations for resuspension and for wet and dry deposition of particles and gases for use in radiation dose calculations. *Nucl. Saf.* 19: 205-219.
- Slinn, W. G. N. (1982) Predictions for particle deposition to vegetative canopies. *Atmos. Environ.* 16: 1785-1794.
- Sloane, C. S. (1983) Optical properties of aerosols—comparison of measurements with model calculations. *Atmos. Environ.* 17: 409-416.
- Sloane, C. S. (1984) Optical properties of aerosols of mixed composition. *Atmos. Environ.* 18: 871-878.
- Sloane, C. S. (1986) Effect of composition on aerosol light scattering efficiencies. *Atmos. Environ.* 20: 1025-1037.
- Sloane, C. S.; Wolff, G. T. (1985) Prediction of ambient light scattering using a physical model responsive to relative humidity: validation with measurements from Detroit. *Atmos. Environ.* 19: 669-680.
- Smith, W. H. (1973) Metal contamination of urban woody plants. *Environ. Sci. Technol.* 7: 631-636.
- Smith, W. H. (1974) Air pollution - effects on the structure and function of the temperate forest ecosystem. *Environ. Pollut.* 6: 111-129.
- Smith, W. H. (1984) Pollutant uptake by plants. In: Treshow, M., ed. *Air pollution and plant life*. New York, NY: John Wiley & Sons, Ltd.; pp. 417-450.
- Smith, W. H. (1990a) Forests as sinks for air contaminants: soil compartment. In: *Air pollution and forests: interactions between air contaminants and forest ecosystems*. 2nd ed. New York, NY: Springer-Verlag; pp. 113-146. (Springer series on environmental management).
- Smith, W. H. (1990b) Forests as sinks for air contaminants: vegetative compartment. In: *Air pollution and forests: interactions between air contaminants and forest ecosystems*. 2nd ed. New York, NY: Springer-Verlag; pp. 147-180. (Springer series on environmental management).
- Smith, W. H. (1990c) Forest nutrient cycling: toxic ions. In: *Air pollution and forests: interactions between air contaminants and forest ecosystems*. 2nd ed. New York, NY: Springer-Verlag; pp. 225-268. (Springer series on environmental management).
- Smith, W. H. (1990d) Forest biotic agent stress: air pollutants and disease caused by microbial pathogens. In: *Air pollution and forests: interactions between air contaminants and forest ecosystems*. 2nd ed. New York, NY: Springer-Verlag; pp. 366-397. (Springer series on environmental management).
- Smith, W. H. (1991) Air pollution and forest damage. *Chem. Eng. News* 69(45): 30-43.
- Smith, K. E. C.; Jones, K. C. (2000) Particles and vegetation: implications for the transfer of particle-bound organic contaminants to vegetation. *Sci. Total Environ.* 246: 207-236.
- Smith, W. H.; Staskawicz, B. J. (1977) Removal of atmospheric particles by leaves and twigs of urban trees: some preliminary observations and assessment of research needs. *Environ. Manage. (N. Y.)* 1: 317-330.
- Soulen, P. E.; Frederick, J. E. (1999) Estimating biologically active UV irradiance from satellite radiance measurements: a sensitivity study. *J. Geophys. Res. [Atmos.]* 104: 4117-4126.
- Spence, J. W.; Haynie, F. H. (1972) Paint technology and air pollution: a survey and economic assessment. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Programs; publication no. AP-103. Available from: NTIS, Springfield, VA; PB-210736.
- Spence, J. W.; Haynie, F.; Upham, J. B. (1975) Effects of gaseous pollutants on paints: a chamber study. *J. Paint Technol.* 47: 57-63.
- Spence, J. W.; Haynie, F. H.; Lipfert, F. W.; Cramer, S. D.; McDonald, L. G. (1992) Atmospheric corrosion model for galvanized steel structures. *Corrosion* 48: 1009-1019.
- Spiker, E. C.; Comer, V. J.; Hosker, R. P.; Sherwood, S. I. (1992) Dry deposition of SO₂ on limestone and marble: the role of humidity. In: Rodrigues, J. D.; Henriques, F.; Jeremias, F. T., eds. *Proceedings of the 7th international congress on deterioration and conservation of stone*; June; Lisbon, Portugal. Lisbon, Portugal: Laboratório Nacional de Engenharia Civil; pp. 397-406.
- Spinka, J. (1971) Vliv znečištěného ovzduší na ovocné stromy a zeleninu [Effects of polluted air on fruit trees and legumes]. *Ziva* 19: 13-15.
- Stanhill, G.; Cohen, S. (2001) Global dimming: a review of the evidence for a widespread and significant reduction in global radiation with discussion of its probable causes and possible agricultural consequences. *Agric. For. Meteorol.* 107: 255-278.
- Steubing, L.; Klee, R. (1970) Vergleichende Untersuchungen zur Staubfilterwirkung von Laub- und Nadelgehölzen [Comparative investigations on the dust-filter effect of broad-leaved and coniferous trees]. *Angew. Bot.* 44: 73-85.
- Stevens, R. K.; Dzubay, T. G.; Lewis, C. W.; Shaw, R. W., Jr. (1984) Source apportionment methods applied to the determination of the origin of ambient aerosols that affect visibility in forested areas. *Atmos. Environ.* 18: 261-272.

- Stoddard, J. L. (1994) Long-term changes in watershed retention of nitrogen: its causes and aquatic consequences. In: Baker, L. A., ed. *Environmental chemistry of lakes and reservoirs*. Washington, DC: American Chemical Society; pp. 223-284. (Advances in chemistry series no. 237).
- Stoddard, J. L.; Murdoch, P. S. (1991) Catskill Mountains. In: Charles, D. F., ed. *Acidic deposition and aquatic ecosystems: regional case studies*. New York, NY: Springer-Verlag; pp. 237-271.
- Storm, G. L.; Fosmire, G. J.; Bellis, E. D. (1994) Persistence of metals in soil and selected vertebrates in the vicinity of the Palmerton zinc smelters. *J. Environ. Qual.* 23: 508-514.
- Strandberg, H.; Johansson, L.-G. (1997) The formation of black patina on copper in humid air containing traces of SO₂. *J. Electrochem. Soc.* 144: 81-89.
- Strickland, T. C.; Holdren, G. R., Jr.; Ringold, P. L.; Bernard, D.; Smythe, K.; Fallon, W. (1993) A national critical loads framework for atmospheric deposition effects assessment: I. method summary. *Environ. Manage.* (N. Y.) 17: 329-334.
- Swietlicki, E.; Zhou, J. C.; Berg, O. H.; Martinsson, B. G.; Frank, G.; Cederfelt, S. I.; Desek, U.; Berner, A.; Birmilli, W.; Wiedensohler, A.; Yuskiewicz, B.; Bower, K. N. (1999) A closure study of sub-micrometer aerosol particle hygroscopic behaviour. *Atmos. Res.* 50: 205-240.
- Swietlik, D.; Faust, M. (1984) Foliar nutrition of fruit crops. *Hortic. Rev.* 6: 287-355.
- Sydberger, T.; Ericsson, R. (1977) Laboratory testing of the atmospheric corrosion of steel. *Werkst. Korros.* 28: 154-158.
- Taback, H. J.; Brienza, A. R.; Macko, J.; Brunetz, N. (1979) Fine particle emissions from stationary and miscellaneous sources in the south coast air basin: appendix. Sacramento, CA: California Air Resources Board; contract no. A6-191-30.
- Talbot, J. J. (1979) A review of the potential biological impacts of cooling tower salt drift. *Atmos. Environ.* 13: 395-405.
- Tang, I. N. (1980) Deliquescence properties and particle size change of hygroscopic aerosols. In: Willeke, K., ed. *Generation of aerosols and facilities for exposure experiments*. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 153-167.
- Tang, I. N. (1997) Thermodynamic and optical properties of mixed-salt aerosols of atmospheric importance. *J. Geophys. Res. [Atmos.]* 102: 1883-1893.
- Tang, I. N.; Munkelwitz, H. R. (1994) Water activities, densities, and refractive indices of aqueous sulfates and sodium nitrate droplets of atmospheric importance. *J. Geophys. Res. [Atmos.]* 99: 18,801-18,808.
- Tanner, R. L. (1981) An ambient experimental study of phase equilibrium in the atmospheric system: aerosol H⁺, NH₄⁺, SO₄²⁻, NO₃⁻, NH₃(g), HNO₃(g). In: Fogler, H. S., ed. *Chemical reactors: based on a symposium sponsored by the Division of Industrial and Engineering Chemistry at the second chemical congress of the North American continent (180th ACS national meeting); August 1980; Las Vegas, NV*. Washington, DC: American Chemical Society; pp. 527-532. (ACS symposium series 168).
- Taylor, G. E., Jr.; Hanson, P. J.; Baldocchi, D. D. (1988) Pollutant deposition to individual leaves and plant canopies: sites of regulation and relationship to injury. In: Heck, W. W.; Taylor, O. C.; Tingey, D. T., eds. *Assessment of crop loss from air pollutants*. New York, NY: Elsevier Applied Science; pp. 227-257.
- Taylor, G. E., Jr.; Johnson, D. W.; Andersen, C. P. (1994) Air pollution and forest ecosystems: a regional to global perspective. *Ecol. Appl.* 4: 662-689.
- Tegen, I.; Lacis, A. A.; Fung, I. (1996) The influence on climate forcing of mineral aerosols from disturbed soils. *Nature (London)* 380: 419-422.
- Thornton, F. C.; Schaedle, M.; Raynal, D. J. (1987) Effects of aluminum on red spruce seedlings in solution culture. *Environ. Exp. Bot.* 27: 489-498.
- Tilman, D. (1996) Biodiversity: population versus ecosystem stability. *Ecology* 77: 350-363.
- Tilman, D. (2000) Causes, consequences and ethics of biodiversity. *Nature (London)* 405: 208-211.
- Tilman, D.; Downing, J. A. (1994) Biodiversity and stability in grasslands. *Nature (London)* 367: 363-365.
- Tjepkema, J. D.; Cartica, R. J.; Hemond, H. F. (1981) Atmospheric concentration of ammonia in Massachusetts and deposition on vegetation. *Nature (London)* 294: 445-446.
- Tolley, G. A.; Randall, A.; Blomquist, G.; Fabian, R.; Fishelson, G.; Frankel, A.; Hoehn, J.; Krumm, R.; Mensah, E.; Smith, T. (1986) Establishing and valuing the effects of improved visibility in eastern United States. Washington, DC: U.S. Environmental Protection Agency.
- Tombach, I.; Thurston, S. A. (1994) The quality of the SCENES measurements: the roles of data quality goals and evolving technology. In: *Aerosols and atmospheric optics: radiative balance and visual air quality: proceedings of the international specialty conference, volume A; September; Snowbird, UT*. Pittsburgh, PA: Air & Waste Management Association; pp. 21-32. (A&WMA publication VIP-41).

- Tong, S. T. Y. (1991) The retention of copper and lead particulate matter in plant foliage and forest soil. *Environ. Int.* 17: 31-37.
- Treshow, M. (1980) Pollution effects on plant distribution. *Environ. Conserv.* 7: 279-286.
- Trettin, C.C.; Johnson, D.W.; Todd, D.E., Jr. (1999) Forest nutrient and carbon pools at Walker Branch watershed: changes during a 21-year-period. *Soil Sci. Soc. Amer. J.* 63: 1436-1448.
- Trijonis, J.; Thayer, M.; Murdoch, J.; Hagemen, R. (1985) Air quality benefit analysis for Los Angeles and San Francisco based on housing values and visibility. Sacramento, CA: California Air Resources Board; ARB contract number A-2-088-32. Available: <ftp://ftp.arb.ca.gov/carbis/research/apr/past/a2-088-32.pdf> [29 April, 2004].
- Trijonis, J. C.; Pitchford, M.; McGown, M. (1987) Preliminary extinction budget results from the RESOLVE program. In: Bhardwaja, P. S., ed. *Visibility protection: research and policy aspects, an APCA international specialty conference; September 1986; Grand Teton National Park, WY.* Pittsburgh, PA: Air Pollution Control Association; pp. 872-883. (APCA transactions series no. TR-10).
- Trijonis, J. C.; Malm, W. C.; Pitchford, M.; White, W. H.; Charlson, R.; Husar, R. (1991) Visibility: existing and historical conditions—causes and effects. In: Irving, P. M., ed. *Acidic deposition: state of science and technology, volume III: terrestrial, materials, health and visibility effects.* Washington, DC: The U.S. National Acid Precipitation Assessment Program. (State of science and technology report no. 24).
- Turner, J.; Kelly, J. (1981) Relationships between soil nutrients and vegetation in a north coast forest, New South Wales. *Aust. For. Res.* 11: 201-208.
- Turner, J.; Lambert, M. J. (1980) Sulfur nutrition of forests. In: Shriner, D. S.; Richmond, C. R.; Lingberg, S. E., eds. *Atmospheric sulfur deposition: environmental impact and health effects, proceedings of the second life sciences symposium; October 1979; Gatlinburg, TN.* Ann Arbor, MI: Ann Arbor Science Publishers, Inc.; pp. 321-333.
- Turner, J.; Lambert, M. J.; Gessel, S. P. (1977) Use of foliage sulphate concentrations to predict response to urea application by Douglas-fir. *Can. J. For. Res.* 7: 476-480.
- Twomey, S. (1974) Pollution and the planetary albedo. *Atmos. Environ.* 8: 1251-1256.
- Tyler, G. (1972) Heavy metals pollute nature, may reduce productivity. *Ambio* 1: 52-59.
- U.S. Code. (1990) Clean Air Act, §169B, visibility. U. S. C. 42: §7492.
- U.S. Code. (1991) Clean Air Act, §108, air quality criteria and control techniques, §109, national ambient air quality standards. U. S. C. 42: §§7408-7409.
- U.S. Environmental Protection Agency. (1979) Protecting visibility: an EPA report to Congress. Research Triangle Park, NC: Office of Air Quality Planning and Standards; EPA report no. EPA-450/5-79-008. Available from: NTIS, Springfield, VA; PB80-220320.
- U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.
- U.S. Environmental Protection Agency. (1986) Air quality criteria for lead. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-83/028aF-dF. 4v. Available from: NTIS, Springfield, VA; PB87-142378.
- U.S. Environmental Protection Agency. (1987) Assessing the risks of trace gases that can modify the stratosphere. Washington, DC: U.S. Environmental Protection Agency, Office of Air Radiation; report no. EPA 400/1-87/001A-H.
- U.S. Environmental Protection Agency. (1993) Air quality criteria for oxides of nitrogen. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; report nos. EPA/600/8-91/049aF-cF. 3v. Available from: NTIS, Springfield, VA; PB95-124533, PB95-124525, and PB95-124517.
- U.S. Environmental Protection Agency. (1995a) Interim findings on the status of visibility research. Research Triangle Park, NC: Office of Research and Development; report no. EPA/600/R-95/021.
- U.S. Environmental Protection Agency. (1995b) Testing of meteorological and dispersion models for use in regional air quality modeling. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA-454/R-95-005. Available from: NTIS, Springfield, VA; PB95-215638.
- U.S. Environmental Protection Agency. (1996a) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1996b) Air quality criteria for ozone and related photochemical oxidants. Research Triangle Park, NC: Office of Research and Development; report nos. EPA/600/AP-93/004aF-cF. 3v. Available from: NTIS, Springfield, VA; PB96-185582, PB96-185590, and PB96-185608. Available: <http://cfpub2.epa.gov/ncea/>.

- U.S. Environmental Protection Agency. (1997a) Nitrogen oxides: impacts on public health and the environment. Washington, DC: Office of Air and Radiation; August. Available: www.epa.gov/ttncaaa1/t1/reports/noxrept.pdf [1999, November 24].
- U.S. Environmental Protection Agency. (1997b) Summary of the particulate monitoring program. Washington, DC: Office of Air and Radiation.
- U.S. Environmental Protection Agency. (1998) Monitoring PM_{2.5} in ambient air using designated reference of class I equivalent methods. In: Quality Assurance Handbook for Air Pollution Measurement Systems, V. II, Part II, Section 2.12. Research Triangle Park, NC: Research and Development, Environmental Monitoring Systems Laboratory.
- U.S. Environmental Protection Agency. (1999a) Protection of visibility. C. F. R. 40: §51.300-51.309.
- U.S. Environmental Protection Agency. (1999b) Visibility monitoring guidance. Research Triangle Park, NC: Emissions, Monitoring, and Analysis Division.
- U.S. Environmental Protection Agency. (2000a) Deposition of air pollutants to the great waters. Third report to Congress. [Executive Summary]. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-453/R-00-005.
- U.S. Environmental Protection Agency. (2000b) Latest findings on national air quality: 1999 status and trends. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards; report no. EPA-454/F-00-002.
- U.S. Environmental Protection Agency. (2001a) National air quality and emissions trends report, 1999. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA 454/R-01-004. Available: <http://www.epa.gov/oar/aqtrnd99/> [7 May 2003].
- U.S. Environmental Protection Agency. (2001b) Latest findings on national air quality: 2000 status and trends. Research Triangle Park, NC: Office of Air Quality Planning and Standards; report no. EPA-454/K-01-002. Available: <http://www.epa.gov/air/aqtrnd00/brochure/00brochure.pdf> (22 July 2003).
- Ulrich, B. (1983) Interaction of forest canopies with atmospheric constituents: SO₂, alkali and earth alkali cations and chloride. In: Ulrich, B.; Pankrath, J., eds. Effects of accumulation of air pollutants in forest ecosystems: proceedings of a workshop; May 1982; Göttingen, Federal Republic of Germany. Dordrecht, The Netherlands: D. Reidel Publishing Company; pp. 33-45.
- United Nations Environment Programme (UNEP). (1986) Report of the international conference on the assessment of the role of carbon dioxide and of other greenhouse gases in climate variations and associated impacts; October 1985; Villach, Austria. Geneva, Switzerland: World Meteorological Organization; WMO no. 661.
- United Nations Environment Programme (UNEP). (1998) Environmental effects of ozone depletion: 1998 assessment. *J. Photochem. Photobiol. B* 46: 1-4.
- United Nations Environment Programme (UNEP). (2000) Environmental effects of ozone depletion: interim summary. Available at: <http://www.gcio.org/ozone/unep2000summary.html> [9 April 2002].
- Unsworth, M. H. (1984) Evaporation from forests in cloud enhances the effects of acid deposition. *Nature (London)* 312: 262-264.
- Unsworth, M. H.; Wilshaw, J. C. (1989) Wet, occult and dry deposition of pollutants on forests. *Agric. For. Meteorol.* 47: 221-238.
- Van Aalst, R. M. (1982) Dry deposition of NO_x. In: Schneider, T.; Grant, L., eds. Air pollution by nitrogen oxides. Amsterdam, The Netherlands: Elsevier Scientific Publishing Company; pp. 263-270.
- Van Beelen, P.; Doelman, P. (1997) Significance and application of microbial toxicity tests in assessing ecotoxicological risks of contaminants in soil and sediment. *Chemosphere* 34: 455-499.
- Van Breemen, N.; Van Dijk, H. F. G. (1988) Ecosystem effects of atmospheric deposition of nitrogen in The Netherlands. In: Dempster, J. P.; Manning, W. J., eds. Excess nitrogen deposition. *Environ. Pollut.* 54: 249-274.
- Van Breemen, N.; Burrough, P. A.; Velthorst, E. J.; Van Dobben, H. F.; De Wit, T.; Ridder, T. B.; Reijnders, H. F. R. (1982) Soil acidification from atmospheric ammonium sulphate in forest canopy throughfall. *Nature (London)* 299: 548-550.
- Van Breemen, N.; Mulder, J.; Driscoll, C. T. (1983) Acidification and alkalization of soils. *Plant Soil* 75: 283-308.
- Van de Hulst, H. C. (1957, reprint 1981) Light scattering by small particles. New York, NY: Wiley. (Structure of matter series).
- Van der Leun, J. C.; Tang, X.; Tevini, M. (1995) Environmental effects of ozone depletion: 1994 assessment. *Ambio* 24: 138.
- Van der Leun, J. C.; Tang, X.; Tevini, M., eds. (1998) Environmental effects of ozone depletion: 1998 assessment. *J. Photochem. Photobiol. B* 46(1-3).

- Van Dijk, H. F. G.; Roelofs, J. G. M. (1988) Effects of excessive ammonium deposition on the nutritional status and condition of pine needles. *Physiol. Plant.* 73: 494-501.
- Van Dorland, R.; Dentener, F. J.; Lelieveld, J. (1997) Radiative forcing due to tropospheric ozone and sulfate aerosols. *J. Geophys. Res. [Atmos.]* 102: 28,079-28,100.
- Van Egmond, K.; Bresser, T.; Bouwman, L. (2002) The European nitrogen case. *Ambio* 31: 72-78.
- Van Miegroet, H.; Cole, D. W. (1984) The impact of nitrification on soil acidification and cation leaching in red alder ecosystem. *J. Environ. Qual.* 13: 586-590.
- Van Miegroet, H.; Johnson, D. W.; Todd, D. E. (1993) Foliar response of red spruce saplings to fertilization with Ca and Mg in the Great Smoky Mountains National Park. *Can. J. For. Res.* 23: 89-95.
- Venkatram, A.; Pleim, J. (1999) The electrical analogy does not apply to modeling dry deposition of particles. *Atmos. Environ.* 33: 3075-3076.
- Viles, H. A. (1990) The early stages of building stone decay in an urban environment. *Atmos. Environ. Part A* 24: 229-232.
- Virginia Climate Advisory. (2000) A closer look at visibility. Charlottesville, VA: University of Virginia, Virginia State Climatology Office. (Virginia Climate Advisory: 00/8). Available: <http://climate.virginia.edu/advisory/2000/ad00-8.htm> [29 September, 2004].
- Viskari, E.-L.; Kärenlampi, L. (2000) Roadside Scots pine as an indicator of deicing salt use - a comparative study from two consecutive winters. *Water Air Soil Pollut.* 122: 405-419.
- Vitousek, P. M.; Mooney, H. A.; Lubchenco, J.; Melillo, J. M. (1997) Human domination of Earth's ecosystems. *Science (Washington, DC)* 277: 494-499.
- Vogt, K. A.; Dahlgren, R.; Ugolini, F.; Zabowski, D.; Moore, E. E.; Zasoski, R. (1987a) Aluminum, Fe, Ca, Mg, K, Mn, Cu, Zn and P in above- and belowground biomass. I. *Abies amabilis* and *Tsuga mertensiana*. *Biogeochemistry* 4: 277-294.
- Vogt, K. A.; Dahlgren, R.; Ugolini, F.; Zabowski, D.; Moore, E. E.; Zasoski, R. (1987b) Aluminum, Fe, Ca, Mg, K, Mn, Cu, Zn and P in above- and belowground biomass. II. Pools, and circulation in a subalpine *Abies amabilis* stand. *Biogeochemistry* 4: 295-311.
- Von Glasow, R.; Bott, A. (1999) Interaction of radiation fog with tall vegetation. *Atmos. Environ.* 33: 1333-1346.
- Vora, A. B.; Bhatnagar, A. R. (1987) Comparative study of dust fall on the leaves in high pollution and low pollution area of Ahmedabad: V. Caused foliar injury. *J. Environ. Biol.* 8: 339-346.
- Vose, J. M.; Swank, W. T. (1990) Preliminary estimates of foliar absorption of ¹⁵N-labeled nitric acid vapor (HNO₃) by mature eastern white pine (*Pinus strobus*). *Can. J. For. Res.* 20: 857-860.
- Vuilleumier, L.; Harley, R. A.; Brown, N. J.; Slusser, J. R.; Kolinski, D.; Bigelow, D. S. (2001) Variability in ultraviolet total optical depth during the Southern California Ozone Study (SCOS97). *Atmos. Environ.* 35: 1111-1122.
- Waggoner, A. P.; Weiss, R. E. (1980) Comparison of fine particle mass concentration and light scattering extinction in ambient aerosol. *Atmos. Environ.* 14: 623-626.
- Waggoner, A. P.; Weiss, R. E.; Ahlquist, N. C.; Covert, D. S.; Will, S.; Charlson, R. J. (1981) Optical characteristics of atmospheric aerosols. In: White, W. H.; Moore, D. J.; Lodge, J. P., Jr., eds. *Plumes and visibility: measurements and model components: proceedings of the symposium; November 1980; Grand Canyon National Park, AZ.* *Atmos. Environ.* 15: 1891-1909.
- Wagrowski, D. M.; Hites, R. A. (1997) Polycyclic aromatic hydrocarbon accumulation in urban, suburban, and rural vegetation. *Environ. Sci. Technol.* 31: 279-282.
- Wall, D. H. (1999) Biodiversity and ecosystem functioning. *BioScience* 49: 107-108.
- Wall, D. H.; Moore, J. C. (1999) Interactions underground: soil biodiversity, mutualism, and ecosystem processes. *BioScience* 49: 109-117.
- Walton, J. R.; Johnson, J. B.; Wood, G. C. (1982) Atmospheric corrosion initiation by sulphur dioxide and particulate matter: II. characterisation and corrosivity of individual particulate atmospheric pollutants. *Br. Corros. J.* 17: 65-70.
- Wang, W. C.; Yung, Y. L.; Lacis, A. A.; Mo, T.; Hansen, J. E. (1976) Greenhouse effects due to man-made perturbations of trace gases. *Science (Washington, DC)* 194: 685-690.
- Waring, R. H. (1987) Nitrate pollution: a particular danger to boreal and subalpine coniferous forests. In: Fujimori, T.; Kimura, M., eds. *Human impacts and management of mountain forests: [proceedings of a symposium].* Ibaraki, Japan: Forestry and Forest Products Research Institute; pp. 93-105.
- Waring, R. H.; Schlesinger, W. H. (1985) The carbon balance of trees. In: *Forest ecosystems: concepts and management.* Orlando, FL: Academic Press, Inc.; pp. 7-37.
- Watson, J. G. (2002) Visibility: science and regulation. *J. Air Waste Manage. Assoc.* 52: 628-713.

- Watson, J. G.; Chow, J. C. (1993) Ambient air sampling. In: Willeke, K.; Baron, P. A., eds. *Aerosol measurement: principles, techniques, and applications*. New York, NY: Van Nostrand Reinhold; pp. 622-639.
- Watson, J. G.; Chow, J. C. (1994) Clear sky visibility as a challenge for society. *Annu. Rev. Energy Environ.* 19: 241-266.
- Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. *Environ. Software* 5: 38-49.
- Webb, A. H.; Bawden, R. J.; Busby, A. K.; Hopkins, J. N. (1992) Studies on the effects of air pollution on limestone degradation in Great Britain. *Atmos. Environ. Part B* 26: 165-181.
- Wedding, J. B.; Carlson, R. W.; Stukel, J. J.; Bazzaz, F. A. (1975) Aerosol deposition on plant leaves. *Environ. Sci. Technol.* 9: 151-153.
- Wedin, D. A.; Tilman, D. (1996) Influence of nitrogen loading and species composition on the carbon balance of grasslands. *Science (Washington, DC)* 274: 1720-1723.
- Weinbaum, S. A.; Neumann, P. M. (1977) Uptake and metabolism of ¹⁵N-labeled potassium nitrate by French prune (*Prunus domestica L.*) leaves and the effects of two surfactants. *J. Am. Soc. Hortic. Sci.* 102: 601-604.
- Wellburn, A. R. (1990) Tansley review no. 24: Why are atmospheric oxides of nitrogen usually phytotoxic and not alternative fertilizers? *New Phytol.* 115: 395-429.
- Wells, B. W. (1939) A new forest climax: the salt spray climax of Smith Island, North Carolina. *Torrey Bot. Club Bull.* 66: 629-634.
- Wells, B. W.; Shunk, I. V. (1937) Seaside shrubs: wind forms vs. spray forms. *Science* 85: 499.
- Wells, B. W.; Shunk, I. V. (1938) Salt spray: an important factor in coastal ecology. *Torrey Bot. Club Bull.* 65: 485-492.
- Wenny, B. N.; Schafer, J. S.; DeLuisi, J. J.; Saxena, V. K.; Barnard, W. F.; Petropavlovskikh, I. V.; Vergamini, A. J. (1998) A study of regional aerosol radiative properties and effects on ultraviolet-B radiation. *J. Geophys. Res. [Atmos.]* 103: 17,083-17,097.
- Wesely, M. L. (1989) Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models. *Atmos. Environ.* 23: 1293-1304.
- Wesely, M. L.; Hicks, B. B. (1977) Some factors that affect the deposition rates of sulfur dioxide and similar gases on vegetation. *J. Air Pollut. Control Assoc.* 27: 1110-1116.
- Wesely, M. L.; Hicks, B. B. (2000) A review of the current status of knowledge on dry deposition. *Atmos. Environ.* 34: 2261-2282.
- Wesely, M. L.; Eastman, J. A.; Stedman, D. H.; Yalvac, E. D. (1982) An eddy-correlation measurement of NO₂ flux to vegetation and comparison to O₃ flux. *Atmos. Environ.* 16: 815-820.
- Wesely, M. L.; Cook, D. R.; Hart, R. L. (1983) Fluxes of gases and particles above a deciduous forest in wintertime. *Boundary Layer Meteorol.* 27: 237-255.
- Wesely, M. L.; Cook, D. R.; Hart, R. L.; Speer, R. E. (1985) Measurements and parameterization of particulate sulfur dry deposition over grass. *J. Geophys. Res. [Atmos.]* 90: 2131-2143.
- Wesselink, L. G.; Meiwes, K.-J.; Matzner, E.; Stein, A. (1995) Long-term changes in water and soil chemistry in spruce and beech forests, Solling, Germany. *Environ. Sci. Technol.* 29: 51-58.
- Westman, W. E. (1977) How much are nature's services worth? Measuring the social benefits of ecosystem functioning is both controversial and illuminating. *Science (Washington, DC)* 197: 960-964.
- Whitby, K. T. (1978) The physical characteristics of sulfur aerosols. *Atmos. Environ.* 12: 135-159.
- White, W. H. (1991) Contributions to light extinction. In: Irving, P. M., ed. *Acidic deposition: state of science and technology, v. III: terrestrial, materials, health and visibility effects; report 24*. Washington, DC: National Acid Precipitation Assessment Program; pp. 85-102.
- White, W. H.; Macias, E. S. (1990) Light scattering by haze and dust at Spirit Mountain, Nevada. In: Mathai, C. V., ed. *Visibility and fine particles: an A&WMA/EPA international specialty conference; October 1989; Estes Park, CO*. Pittsburgh, PA: Air & Waste Management Association; pp. 914-922. (A&WMA transactions series no. TR-17).
- White, W. H.; Macias, E. S.; Nininger, R. C.; Schorran, D. (1994) Size-resolved measurements of light scattering by ambient particles in the southwestern U.S.A. *Atmos. Environ.* 28: 909-921.
- Williams, M. W.; Baron, J. S.; Caine, N.; Sommerfeld, R.; Sanford, R. (1996) Nitrogen saturation in the Rocky Mountains. *Environ. Sci. Technol.* 30: 640-646.
- Wilson, E. O. (1997) The creation of biodiversity. In: Raven, P. H., ed. *Nature and human society: the quest for a sustainable world: proceedings of the 1997 forum on biodiversity; October; sponsored by the National Research Council*. Washington, DC: National Academy Press.
- Wiman, B. (1981) Aerosol collection by Scots pine seedlings: design and application of a wind tunnel method. *Oikos* 36: 83-92.

- Wiman, B. L. B. (1985) A non-stationary model for sulphur aerosol depletion and deposition in a young spruce stand. *Tellus Ser. B* 37: 230-242.
- Wiman, B. L. B.; Ågren, G. I. (1985) Aerosol depletion and deposition in forests--a model analysis. *Atmos. Environ.* 19: 335-347.
- Wiman, B. L. B.; Lannefors, H. O. (1985) Aerosol characteristics in a mature coniferous forest--methodology, composition, sources and spatial concentration variations. *Atmos. Environ.* 19: 349-362.
- Wiman, B. L. B.; Ågren, G. I.; Lannefors, H. O. (1985) Aerosol concentration profiles within a mature coniferous forest--model versus field results. *Atmos. Environ.* 19: 363-367.
- Winner, W. E.; Atkinson, C. J. (1986) Absorption of air pollution by plants, and consequences for growth. *Trends Ecol. Evol.* 1: 15-18.
- Winner, W. E.; Bewley, J. D. (1978a) Terrestrial mosses as bioindicators of SO₂ pollution stress: synecological analysis and the index of atmospheric purity. *Oecologia* 35: 221-230.
- Winner, W. E.; Bewley, J. D. (1978b) Contrasts between bryophyte and vascular plant synecological responses in an SO₂-stressed white spruce association in central Alberta. *Oecologia* 33: 311-325.
- Witherspoon, J. P.; Taylor, F. G., Jr. (1971) Retention of 1-44 μ simulated fallout particles by soybean and sorghum plants. *Health Phys.* 21: 673-677.
- Wittenburg, C.; Dannecker, W. (1992) Dry deposition and deposition velocity of airborne acidic species upon different sandstones. In: *Proceedings of the 1992 European aerosol conference*; September, Oxford, United Kingdom. *J. Aerosol Sci.* 23(suppl. 1): S869-S872.
- Wolfe, A. H.; Patz, J. A. (2002) Reactive nitrogen and human health: acute and long-term implications. *Ambio* 31: 120-125.
- Wolff, G. T. (1984) On the nature of nitrate in coarse continental aerosols. *Atmos. Environ.* 18: 977-981.
- Wolff, G. T.; Collins, D. C.; Rodgers, W. R.; Verma, M. H.; Wong, C. A. (1990) Spotting of automotive finishes from the interactions between dry deposition of crustal material and wet deposition of sulfate. *J. Air Waste Manage. Assoc.* 40: 1638-1648.
- Woodcock, A. H. (1953) Salt nuclei in marine air as a function of altitude and wind force. *J. Meteorol.* 10: 362-371.
- Woodwell, G. M. (1970) Effects of pollution on the structure and physiology of ecosystems: changes in natural ecosystems caused by many different types of disturbances are similar and predictable. *Science (Washington, DC)* 168: 429-433.
- World Health Organization. (1997) *Nitrogen oxides*. 2nd ed. Geneva, Switzerland: World Health Organization. (Environmental health criteria 188).
- World Meteorological Organization. (1988) *Developing policies for responding to climatic change: a summary of the discussions and recommendations of workshops*; September-October 1987; Villach, Austria; and November 1987; Bellagio, Austria. Geneva, Switzerland: World Meteorological Organization; report no. WMO/TD; no. 225. [World Climate Impact Programme series report no. WCIP-1].
- World Meteorological Organization. (1999) *Scientific assessment of ozone depletion: 1998*. Geneva, Switzerland: World Meteorological Organization, Global Ozone and Monitoring Project; report no. 44.
- World Resources Institute. (2000) *World resources 2000-2001: people and ecosystems: the fraying web of life*. Washington, DC: World Resources Institute.
- Wu, P.-M.; Okada, K. (1994) Nature of coarse nitrate particles in the atmosphere--a single particle approach. *Atmos. Environ.* 28: 2053-2060.
- Wytenbach, A.; Tobler, L.; Bajo, S. (1989) Na, Cl and Br in needles of Norway spruce (*P. abies*) and in the aerosol adhering to the needles. *Toxicol. Environ. Chem.* 19: 25-33.
- Xu, J. R.; Balik, C. M. (1989) Infrared attenuated total reflectance study of latex paint films exposed to aqueous sulfur dioxide. *J. Appl. Polym. Sci.* 38: 173-183.
- Yanai, R. D.; Siccama, T. G.; Arthur, M. A.; Federer, C. A.; Friedland, A. J. (1999) Accumulation and depletion of base cations in forest floors in the northeastern United States. *Ecology* 80: 2774-2787.
- Yerrapragada, S. S.; Jaynes, J. H.; Chirra, S. R.; Gauri, K. L. (1994) Rate of weathering of marble due to dry deposition of ambient sulfur and nitrogen dioxides. *Anal. Chem.* 66: 655-659.
- Yocom, J. E.; Grappone, N. (1976) *Effects of power plant emissions on materials*. Palo Alto, CA: Electric Power Research Institute; report no. EPRI/EC-139. Available from: NTIS, Springfield, VA; PB-257 539.
- Yocom, J. E.; Upham, J. B. (1977) *Effects of economic materials and structures*. In: Stern, A. C., ed. *Air pollution*. 3rd ed. Volume II. New York, NY: Academic Press; pp. 93-94.
- Yoshida, L. C.; Allen, E. B. (2001) Response to ammonium and nitrate by a mycorrhizal annual invasive grass and a native shrub in southern California. *Am. J. Bot.* 88: 1430-1436.
- Young, P. (1996) Pollution-fueled "biodeterioration" threatens historic stone: researchers are probing the role microorganisms play in the decay of buildings and artwork. *Environ. Sci. Technol.* 30: 206A-208A.

- Young, J. R.; Ellis, E. C.; Hidy, G. M. (1988) Deposition of air-borne acidifiers in the western environment. *J. Environ. Qual.* 17: 1-26.
- Youngs, W. D.; Rutzke, M.; Gutenmann, W. H.; Lisk, D. J. (1993) Nickel and vanadium in foliage in the vicinity of an oil-fired power plant. *Chemosphere* 27: 1269-1272.
- Zappia, G.; Sabbioni, C.; Gobbi, G. (1993) Non-carbonate carbon content on black and white areas of damaged stone monuments. *Atmos. Environ. Part A* 27: 1117-1121.
- Zappia, G.; Sabbioni, C.; Pauri, M. G.; Gobbi, G. (1994) Mortar damage due to airborne sulfur compounds in a simulation chamber. *Mater. Struct.* 27: 469-473.
- Zappia, G.; Sabbioni, C.; Riontino, C.; Gobbi, G.; Favoni, O. (1998) Exposure tests of building materials in urban atmosphere. *Sci. Total Environ.* 224: 235-244.
- Zdanovskii, A. B. (1948) New methods of calculating solubilities of electrolytes in multicomponent systems. *Zh. Fiz. Khim.* 22: 1475-1485.
- Zepp, R. G.; Callaghan, T. V.; Erickson, D. J. (1998) Effects of enhanced solar ultraviolet radiation on biogeochemical cycles. *J. Photochem. Photobiol. B* 46: 69-82.
- Zhang, X. Q.; McMurry, P. H.; Hering, S. V.; Casuccio, G. S. (1993) Mixing characteristics and water content of submicron aerosols measured in Los Angeles and at the Grand Canyon. *Atmos. Environ. Part A* 27: 1593-1607.
- Zhang, L.; Gong, S.; Padro, J.; Barrie, L. (2001) A size-segregated particle dry deposition scheme for an atmospheric aerosol module. *Atmos. Environ.* 35: 549-560.
- Zipperer, W. C.; Wu, J.; Pouyat, R. V.; Pickett, S. T. A. (2000) The application of ecological principles to urban and urbanizing landscapes. *Ecol. Appl.* 10: 685-688.
- Zuburtikudis, I.; Triantafyllou, A. (2001) Environmental pollution and its impact on the monuments and works of art in Greece. *J. Environ. Prot. Ecol.* 2: 68-72.

Appendix 4A

Common and Latin Names

Alder, hazel	<i>Alnus serrulata</i> (Aiton) Willdenow
Alder, red	<i>Alnus rubra</i> Bong.
Bean, common	<i>Phaseolus vulgaris</i> L.
Beech	<i>Fagus sylvatica</i> L.
Birch, yellow	<i>Betula alleghaniensis</i> Britt.
Blackhaw	<i>Viburnum prunifolium</i> L.
Brush box	<i>Lophostemon confertus</i> (R. BR.) P.G. Wilson & Waterhouse
Ceanothus, hoaryleaf	<i>Ceanothus crassifolius</i> Torrey
Chaparral	<i>Ceanothus crassifolius</i>
Coachwood	<i>Ceratopetalum apetalum</i> , D. Don
Corn	<i>Zea mays</i> L.
Dogwood, flowering	<i>Cornus florida</i> L.
Elm	<i>Ulmus</i> spp.
Fir, balsam	<i>Abies balsamea</i> (L.) Mill.
Fir, Douglas	<i>Pseudotsuga menziesii</i> (Mirb.) Franco.
Fir, fraser	<i>Abies fraseri</i> (Pursh.) Poir
Grape	<i>Vitis</i> spp.
Grass, red brome	<i>Bromus rubens</i> L.
Grass, purple moor	<i>Molina caerulea</i> (L.) Moench.
Greenbriar	<i>Smilax</i> spp.
Gum, sweet	<i>Liquidambar styraciflua</i> L.
Haw, black	<i>Viburnum prunifolium</i> L.
Heather, Scottish	<i>Calluna vulgaris</i> Salisb.
Hickory	<i>Carya</i> spp.

Hophornbeam	<i>Ostrya virginiana</i> (Mill.) Koch
Ivy, English	<i>Hedera helix</i> L.
Laurel, mountain	<i>Kalmia latifolia</i> L.
Lichen, monks hood	<i>Hypogymnia physiodes</i>
Maize	<i>Zea mays</i> L.
Maple, red	<i>Acer rubrum</i> L.
Maple	<i>Acer</i> spp.
Maple, sugar	<i>Acer saccharum</i> Marsh.
Mustard, small podded	<i>Brassica geniculata</i> L.
Nettle, stinging	<i>Urtica dioica</i> L.
Oak, bur	<i>Quercus macrocarpa</i> Michx.
Oak, English	<i>Quercus rober</i> L.
Oak, chestnut	<i>Quercus prinus</i> = <i>Q. montana</i> Willd.
Oak, live	<i>Quercus virginiana</i> Mill.
Oak, northern red	<i>Quercus rubra</i> L.
Oak, turkey	<i>Quercus laevis</i> Walt.
Oak, white	<i>Quercus alba</i> L.
Oak	<i>Quercus</i> spp.
Oats, domestic	<i>Avena sativa</i> L.
Oats, wild	<i>Avena fatua</i> L.
Persimmon, common	<i>Diosporos virginiana</i> L.
Pine, eastern white	<i>Pinus strobus</i> L.
Pine, jack pine	<i>Pinus banksiana</i> Lamb.
Pine, loblolly	<i>Pinus taeda</i> L.
Pine, lodgepole	<i>Pinus contorta</i> Loud.
Pine, Scots (Scotch)	<i>Pinus sylvestris</i> L.
Pine, slash	<i>Pinus elliotti</i> Englem.
Pine, lodgepole x jack pine	<i>Pinus contorta</i> (Douglas ex Loud) x <i>P. banksiana</i> Lamb.

Poplar, black	<i>Populus nigra</i> L.
Poplar, white	<i>Populus alba</i> L.
Poplar, yellow or tulip	<i>Liriodendron tulipifera</i> L.
Privet	<i>Ligustrum</i> spp.
Purple Moor Grass	<i>Molina caerulea</i> (L.) Moench.
Ragweed	<i>Ambrosia</i> spp.
Rhododendron, Catawba	<i>Rhododendron catawbiense</i> Michx.
Rhododendron, rosebay	<i>Rhododendron maximum</i> L.
Sage, coastal	<i>Artemisia californica</i> Less.
Scottish Heather	<i>Calluna vulgaris</i> Salisb.
Soybean	<i>Glycine max</i> (L.) Merr.
Spruce, Norway	<i>Picea abies</i> (L.) Karst.
Spruce, red	<i>Picea rubens</i> Sarg.
Spruce, sitka	<i>Picea sitchensis</i> (Bong.) Carr.
Spruce, white	<i>Picea glauca</i> (Moench.) Voss.
Sunflower	<i>Helianthus annuus</i> L.
Sweetgum	<i>Liquidambar styraciflua</i> L.
Tulip poplar	<i>Liriodendron tulipifera</i> L.

EUMYCOTA-FUNGI

Zygomycota

Vesicular Arbuscular Mycorrhizae

Scutellospora

Gigaspora

Glomus agrigatum

Glomus leptototicum

Ascomycotina

Chaetomium sp.

Fungi Imperfecti

Aureobasidium pullulans

Cladosporium sp.

Epicoccum sp.

Pestalotiopsis

Phialophora verrucosa

Pleurophomella = *Sirodothis*

5. HUMAN EXPOSURE TO PARTICULATE MATTER AND ITS CONSTITUENTS

5.1 INTRODUCTION

5.1.1 Purpose

The U.S. Environmental Protection Agency's (EPA's) regulatory authority for particulate matter (PM) applies primarily to ambient air and those sources that contribute to ambient air PM concentrations. Most of the epidemiological studies discussed in Chapter 8 relate measured community levels of airborne pollutants to population-based health statistics. Of necessity, these studies rely on some simplifying assumptions regarding exposures. One such assumption is that air pollutant concentrations measured at a community (or population-oriented) monitoring site (or the average concentration of several such sites) can serve as a surrogate for the average personal exposure to ambient PM for the population. However, total personal exposure to PM includes both ambient and nonambient components; and both components may contribute to adverse health effects. Thus, a major emphasis must be to develop an understanding of exposure to PM from sources that contribute to ambient air pollution. Ultimately, it will be necessary to account for both ambient and nonambient components of personal exposure in order to fully understand the relationship between PM and health effects. In addition, knowledge of an individual's personal exposure to ambient, nonambient, and total PM would provide useful information for studies where health outcomes are tracked individually.

Exposure to environmental contaminants can be interpreted in a number of ways. For airborne PM, an individual's exposure is ideally based on measurements of PM concentrations in the air in the individual's breathing zone as the individual moves through space and time. However, epidemiological studies frequently use the ambient concentration as a surrogate for exposure. Therefore, understanding exposure is important because adverse health effects associated with elevated PM concentrations occur at the individual level. Human exposure data and models provide the link between ambient concentrations (from monitoring data or estimated with atmospheric transport models) and lung deposition and clearance models that enable estimates of the source-air concentration-exposure-dose relationship for input into dose-response

assessments for PM from ambient sources. Personal exposure includes contributions from many different types of particles, from many sources, and in many different environments.

The goal of this chapter is to provide current information on the development of human exposure data and models. This includes information on (a) relationships between PM measured at ambient sites and personal exposures to PM from both ambient and nonambient sources, and (b) factors that affect these relationships. The human exposure data and models presented in this chapter provide critical links between ambient monitoring data and PM dosimetry as well as between the toxicological studies and epidemiologic studies presented in other chapters. The specific objectives of this chapter are:

- (1) To provide an overall conceptual framework of exposure science as applied to PM, including the identification and evaluation of factors that determine personal PM exposure;
- (2) To provide a concise summary and review of recent data (since 1996) and findings from pertinent PM exposure studies;
- (3) To characterize quantitative relationships between ambient air quality measurements (mass, chemical components, number, etc. as determined at a community monitoring site) and total personal PM exposure and its ambient and nonambient components; and
- (4) To evaluate the implications of using ambient PM concentrations as a surrogate for personal exposure in epidemiologic studies of PM health effects.

5.1.2 Particulate Matter Mass and Constituents

Current EPA PM regulations are based on mass as a function of aerodynamic size. However, the EPA also measures the chemical composition of PM in both monitoring and research studies. The composition of PM is variable and, as discussed in Chapters 7 and 8, health effects may be related to PM characteristics other than mass. Because PM from ambient air and other microenvironments may have different physical and chemical characteristics, PM from such different sources may also have different health effects. Ultimately, to understand and control health effects caused by PM exposures from all sources, it is important to quantify and understand exposure to those chemical constituents from various sources that are responsible for adverse health effects.

The National Research Council (NRC) recognized the distinction between measuring exposure to PM mass and to chemical constituents in the document *Research Priorities for*

Airborne Particulate Matter I: Immediate Priorities and a Long-range Research Portfolio (National Research Council, 1998). Specifically, under Research Topic 1, Outdoor Measures versus Actual Human Exposures, the NRC recommended evaluation of “the relationships between concentrations of particulate matter and gaseous co-pollutants measured at stationary outdoor air monitoring sites and the contributions of these concentrations to actual personal exposures . . .” for PM mass. The NRC Research Topic 2 recommends evaluating exposures to biologically important constituents and specific characteristics of PM that cause responses in potentially susceptible subpopulations and the general population. It also was recognized by the NRC that “a more targeted set of studies under this research topic (#2) should await a better understanding of the physical, chemical, and biological properties of airborne particles associated with the reported mortality and morbidity outcomes” (National Research Council, 1999). The NRC also stated that the later studies “should be designed to determine the extent to which members of the population contact these biologically important constituents and size fractions of concern in outdoor air, outdoor air that has penetrated indoors, and air pollutants generated indoors” (National Research Council, 1999). Thus, exposure studies should include contributions from all sources. The emphasis in this chapter on PM mass reflects the current state of the science. Where available, data also have been provided on chemical constituents although in most cases the data are limited. As recognized by the NRC, a better understanding of exposures to PM chemical constituents from multiple sources will be required to more fully identify, understand, and control those sources of PM contributing to adverse health effects and to accurately define the relationship between PM exposure and health outcomes due to either short- or long-term exposures.

5.1.3 Relationship to Past Documents

Early versions of PM criteria documents did not emphasize total human exposure, but rather focused almost exclusively on outdoor air concentrations. For instance, the 1969 *Air Quality Criteria for Particulate Matter* (National Air Pollution Control Administration, 1969) did not discuss either *exposure* or *indoor concentrations*. The 1982 EPA PM Air Quality Criteria Document (1982 PM AQCD), however, provided some discussion of indoor PM concentrations reflecting an increase in microenvironmental and personal exposure studies (U.S. Environmental Protection Agency, 1982). The new data indicated that personal activities, along

with PM generated by personal and indoor sources (e.g., cigarette smoking), could lead to high indoor levels and high personal exposures to total PM. Some studies reported indoor concentrations that exceeded PM concentrations found in the air outside the monitored microenvironments or at nearby monitoring sites. Between 1982 and 1996, many more studies of personal and indoor PM exposure demonstrated that, in most inhabited domestic environments, indoor PM mass concentrations and personal PM exposures of the residents were greater than ambient PM mass concentrations measured simultaneously (e.g., Sexton et al., 1984; Spengler et al., 1985; Clayton et al., 1993). As a result, the National Research Council (1991) recognized the potential importance of indoor sources of contaminants (including PM) in causing health outcomes.

The 1996 PM AQCD (U.S. Environmental Protection Agency, 1996) reviewed the human PM exposure literature through early 1996, mainly to evaluate the use of ambient air monitors as surrogates for PM exposure in epidemiological studies. Many of the studies cited showed poor correlations between personal exposure or indoor measurements of PM mass and outdoor or ambient site measurements. Conversely, Janssen et al. (1995) and Tamura et al. (1996a) showed that in the absence of major nonambient sources, total PM exposures to individuals tracked through time were highly correlated with ambient PM concentrations. Analyses of these latter two studies led to the consideration of ambient and nonambient exposures as separate components of total personal exposure. As a result, the 1996 PM AQCD (U.S. Environmental Protection Agency, 1996) distinguished between ambient and nonambient PM personal exposure for the first time. This chapter builds on the work of the 1996 PM AQCD by further evaluating the ambient and nonambient components of PM and by reporting research that evaluates the relationship between ambient concentrations and total, ambient, and nonambient personal exposure.

5.1.4 Chapter Structure

The chapter is organized to provide information on the principles of exposure, to review the existing literature, and to summarize key findings and limitations in the information. Specific sections are described below:

- Section 5.2 discusses the basic concepts of exposure including definitions, methods for estimating exposure, and methods for estimating ambient air components of exposure.

- Section 5.3 presents PM mass data, including a description of the key available studies, the relationship of PM exposures with ambient concentrations, and factors that affect the relationship.
- Section 5.4 presents data on PM constituents, including a description of the key available studies, the relationship with ambient concentrations, and factors that affect the relationship.
- Section 5.5 then discusses the implications of using ambient PM concentrations in epidemiologic studies of PM health effects.
- Section 5.6 summarizes key findings and limitations of the information.

5.2 BASIC CONCEPTS OF EXPOSURE

5.2.1 The Concept of Exposure

“There is reasonable agreement that human exposure [to a substance] means contact with the chemical or agent.” However, contact can be either with “(a) the visible exterior of the person (skin and openings into the body such as mouth and nostrils), or (b) the so-called exchange boundaries where absorption takes place (skin, lung, gastrointestinal tract)” (Federal Register, 1986). In its 1992 Guidelines for Exposure Assessment (U.S. Environmental Protection Agency, 1992), EPA defined exposure as “taking place at the visible external boundary, as in (a) above, [concluding that this definition] is less ambiguous and more consistent with nomenclature in other scientific fields.” The 1992 Guidelines stated:

“Under this definition, it is helpful to think of the human body as having a hypothetical outer boundary separating inside the body from outside the body. This outer boundary of the body is the skin and the openings into the body such as the mouth, the nostrils, and punctures and lesions in the skin. As used in these Guidelines, exposure to a chemical is the contact of that chemical with the outer boundary. An exposure assessment is the quantitative or qualitative evaluation of that contact; it describes the intensity, frequency, and duration of contact, and often evaluates the rates at which the chemical crosses the boundary (chemical intake or uptake rates), the route by which it crosses the boundary (exposure route; e.g., dermal, oral, or respiratory), and the resulting amount of the chemical that actually crosses the boundary (a dose) and the amount absorbed (internal dose)” (U.S. Environmental Protection Agency, 1992).

When applied to PM exposure by the inhalation route, the concentration of interest is that of PM in the air that will enter the respiratory system, not the average concentration of inspired

and exhaled air as measured at the mouth or nostrils. Therefore, a measurement of inhalation exposure to PM is based on measurements of the PM concentration near the breathing zone, but not affected by exhaled air.

5.2.2 Components of Exposure

The total exposure of an individual over a discrete period of time includes exposures to many different types of particles from various sources while in different microenvironments. Duan (1982) defined a microenvironment as “a [portion] of air space with homogeneous pollutant concentration.” It also has been defined as a volume in space for a specific time interval during which the variance of concentration within the volume is significantly less than the variance between that microenvironment and surrounding ones (Mage, 1985). In general, people pass through a series of microenvironments including outdoor, in-vehicle, and indoor microenvironments as they go through space and time. Thus, total daily exposure for a single individual to PM must be expressed as the sum of various exposures for the microenvironments that the person occupies in the day (modified from National Research Council, 1991).

In a given microenvironment, particles may originate from a wide variety of sources. For example, in an indoor microenvironment, PM may be generated by (1) indoor activities, (2) outdoor PM entering indoors, (3) the chemical interaction of outdoor air pollutants and indoor air or indoor sources, (4) transport from another indoor microenvironment, or (5) personal activities. All of these disparate sources should be accounted for when estimating total human exposure to PM.

An analysis of personal exposure to PM mass (or to its constituent compounds) requires definition and discussion of several classes of particles and exposure. In this chapter, PM metrics will be described in terms of exposure or as an air concentration. Particulate matter also may be described according to both its source (i.e., ambient, nonambient) and the microenvironment in which exposure occurs. Table 5-1 provides a summary of the terms used in this chapter, the notations used for these terms, and their definitions. These terms are used throughout this chapter and provide the terminology for evaluating personal exposure to total PM and to PM from ambient and nonambient sources.

Exposures are significant only if they are associated with a biologically relevant duration of contact with a substance of concern (Lioy, 1990; National Research Council, 1991).

TABLE 5-1. TYPES OF PARTICULATE MATTER USED IN EXPOSURE AND CONCENTRATION VARIABLES

<i>General Definitions</i>		
Term	Notation	Definition
Concentration	C	General term for the average air concentration over some specified time period, used with subscript to indicate concentration of a specific type of PM, usually expressed in $\mu\text{g}/\text{m}^3$ units.
Personal Exposure	E	General term for the average personal exposure over some specified time period, used with subscript to indicate exposure to a specific type of PM, quantified as the concentration at the oral/nasal contact boundary.
<i>Types of Particulate Matter</i>		
Term	Subscript	Definition
Ambient PM	a	PM in the atmosphere measured at a community ambient monitoring site either emitted into the atmosphere directly (primary PM) or formed in it (secondary PM). Major sources of PM species are industry, motor vehicles, commerce, domestic emissions such as wood smoke, and natural wind-blown dust or soil. (C without subscript is used for C_a . E_a is used for exposure to ambient PM while outdoors.)
Ambient-Outdoor PM	o	Ambient PM in an outdoor microenvironment. (C_o is frequently considered to be equal to, or at least represented by, C.)
Indoor PM	i	All PM found indoors.
Ambient-Indoor PM	ai	Ambient PM that has infiltrated indoors (i.e., has penetrated indoors and remains suspended). E_{ai} is used for ambient exposure while indoors.
Indoor-generated PM	ig	PM generated indoors.
Indoor-reaction PM	ir	PM formed indoors by pollutants from outdoors reacting with indoor-generated pollutants.
Personal Cloud PM	pc	PM contributing to personal exposure but not contained in indoor or outdoor measurements of PM, usually related to personal activities.
Personal PM (of a subject)	s	PM at the oral/nasal contact zone as the subject moves through space and time.
<i>Concentration and Exposure Variable Used without Subscripts</i>		
Term	Notation	Definition
Ambient Concentration	C	Concentration measured at a community ambient air monitoring site (or the average of several such sites).
Total Personal Exposure	T	Total personal exposure as measured by a personal exposure monitor (PEM).
Ambient Exposure	A	Personal exposure to the ambient component of total personal exposure, i.e., personal exposure to that PM measured at an ambient air community monitoring site. Includes exposure to C and C_{ai} but not to resuspended ambient PM previously deposited indoors.
Nonambient Exposure	N	Personal exposure to nonambient PM.

TABLE 5-1 (cont'd). TYPES OF PARTICULATE MATTER USED IN EXPOSURE AND CONCENTRATION VARIABLES

Relationships among Concentration and Exposure Variables for a Two-Compartment Model (Indoors and Outdoors)

$$T = A + N$$

$$A = E_a + E_{ai}$$

A = yC + (y-1) C_{ai}, where y = fraction of time outside

$$N = E_{ig} + E_{ir} + E_{pc}$$

Application of this concept to PM exposure is complicated by a lack of understanding of the biological mechanisms of PM toxicity. It is not certain whether the relevant duration is the instantaneous exposure to a peak concentration, or hourly, daily, or long-term exposure for months or years (or possibly all of the above). Similarly, it is not certain as to how PM toxicity depends on particle size or particle composition; whether number, surface area, or mass is the appropriate metric; or how PM toxicity may be influenced by conditions that might increase susceptibility such as age, preexisting disease conditions (chronic obstructive pulmonary disease [COPD], asthma, diabetes, etc.), exposure to infectious agents, exposure to heat or cold, stress, etc. A person's exposure is clearly influenced by the distribution of many variables and parameters. A measurement at a single point in space and time along each distribution cannot adequately describe a person's exposure. Thus, it is important to think of exposure as a path function with the instantaneous exposure varying as the PM concentration and composition varies as the person moves through space and time.

The 1997 NAAQS were developed largely on the basis of evidence from epidemiologic studies that found relatively consistent associations between outdoor PM mass concentrations and observed health effects. Thus, an emphasis in this chapter is on the relationship between PM concentrations measured at ambient sites and personal exposures to the PM measured at those ambient sites (National Research Council, 1998), i.e., ambient PM exposure. Although this is an emphasis, it should be kept in mind that every particle that deposits in the lung becomes part of the dose delivered to an individual. It is likely that the nonambient component of total exposure also exerts health effects not necessarily detected by community time-series epidemiologic studies.

5.2.3 Quantification of Exposure

Quantification of inhalation exposure to PM or any other air pollutant starts with the concept of the variation in the concentration of the air pollutant in the breathing zone (but unperturbed by exhaled breath) as measured by a personal exposure monitor as a person moves through space and time. The relationships among the various types of exposure quantities can be easily seen in the hypothetical exposure time profile shown in Figure 5-1. The peak exposure, instantaneous exposure, and average exposure have units of concentration, for PM, usually $\mu\text{g}/\text{m}^3$. The integrated exposure has units of concentration \times time.




Figure 5-1. Hypothetical exposure time profile: pollutant exposure as a function of time showing how the averaged exposure, integrated exposure, and peak exposure relate to the instantaneous exposure. ($t_2 - t_1 = T$)

Source: Adapted from Duan (1989); Zartarian et al. (1997).

The integrated exposure may be written as:

$$\mathcal{E}(t_1, t_2) = \int_{t=t_1}^{t=t_2} C_s(t) dt, \quad (5-1)$$

where \mathcal{E} is the integrated personal exposure during the time period t_1 to t_2 and C_s is the instantaneous exposure of the subject as the subject moves through space and time (Lioy, 1990; National Research Council, 1991; Georgopoulos and Lioy, 1994). Because C_s is a function of both space and time, it cannot generally be correctly approximated by a measurement at a single point in space or at a single time.

For most of the discussion in this chapter, we will be interested in the average exposure given as:

$$E(t_1, t_2) = \frac{1}{t_2 - t_1} \int_{t=t_1}^{t=t_2} C_s(t) dt, \quad (5-2)$$

where E is the average exposure during the time period t_1 to t_2 . Most studies report 24-h averages, although some studies measure 12-h or 2- or 3-day averages.

Equations 5-1 and 5-2 apply to a specific individual moving through space and time on a specific path. When treating populations, the distribution of the values of the variables and parameters must be considered. Georgopoulos and Lioy (1994) showed how to modify these equations to consider the probability distributions.

5.2.4 Methods To Estimate Personal Exposure

Personal exposure may be estimated using either direct or indirect approaches. *Direct approaches* measure the contact of the person with the chemical concentration in the exposure media over an identified period of time. Direct measurement methods include personal exposure monitors (PEMs) for PM that are worn continuously by individuals as they encounter various microenvironments while performing their daily activities. *Indirect approaches* use models and available information on concentrations of chemicals in microenvironments, the time individuals spend in those microenvironments, and personal PM-generating activities to estimate personal

exposure. This section describes methods used to directly measure personal exposures and microenvironmental concentrations as well as the models used to estimate exposure. Several approaches to estimating personal exposure to ambient PM are also described.

5.2.4.1 Direct Measurement Methods

5.2.4.1.1 Personal Exposure Monitoring Methods

In theory, personal exposure to total PM is defined as the concentration of PM in inhaled air entering the nose or mouth. Practically, it is measured by sampling PM with a PEM worn by a person and sampling from a point near the breathing zone (but not affected by exhaled breath). Personal exposure monitors (PEMs) for PM use measurement techniques similar to those employed for measuring ambient PM. Most PEMs rely on filter-based mass measurement of a particle size fraction (PM₁₀ or PM_{2.5}) usually integrated over either a 24- or 12-h period at flow rates of 2 to 4 L/min using battery-operated pumps. Because PEMs must be worn by study participants, they must be quiet, compact, and battery-operated. These requirements limit the type of pumps and the total sample volume that can be collected. Generally, small sample volumes limit personal exposure measurements to PM mass and a few elements detected by X-ray fluorescence (XRF). In most studies, PM_{2.5} and PM₁₀ have not been collected concurrently; thus, very few data are available by which to estimate personal exposure to coarse thoracic PM (i.e., PM_{10-2.5}) exposures.

One of the earliest versions of a PEM was developed by the Harvard School of Public Health for use in the Particle Total Exposure Assessment Methodology (PTEAM) study (Clayton et al., 1993; Thomas et al., 1993; Özkaynak et al., 1996a). This 4 L/min sampler used a sharp cut impactor with 2.5 or 10 µm 50% cut points (Marple et al., 1987). A version of this device is manufactured by MSP, Inc. (Minneapolis, MN). Several exposure studies have used the MSP PEM (usually the 2.5 µm PEM version) with a flow rate of 2 L/min (Kamens et al., 1991; Williams et al., 2000a,b; Landis et al., 2001). Modifications to the MSP PEM, including a “scalper” nozzle to exclude particles > 4 µm, are described in Pellizzari et al. (1999). The penetration curve for the MSP scalper and the MSP 2.5 µm impactor (for 2 L/min flow rate) are given in Rodes et al. (2001).

More recently the Harvard School of Public Health has developed a personal particle sampler that can operate as a PM_{2.5} or a PM₁ sampler and that can be used in a personal

multipollutant sampler (Sioutas et al., 1996; Demokritou et al., 2001). The PM_{2.5} version (4 L/min) has been used in an exposure study (Liu et al., 2002). Evaluations of other PEMs have been reported (Williams et al., 1999; Lanki et al., 2002).

Other methods used for ambient PM have also been adapted for use as PEMs. For example, a personal nephelometer that measures light scattering has been worn by subjects and used in personal exposure studies to obtain real-time PM measurements (Quintana et al., 2000; Rea et al., 2001; Magari et al., 2002; Lanki et al., 2002). Light scattering instruments are most sensitive to particles in the accumulation-mode size range. Recent developments in light scattering devices have made possible the short-term measurement of personal and indoor exposures (Liu et al., 2002). These devices include the Radiance nephelometer (Radiance Research, Seattle, WA), the personal DataRAM or pDR (Thermo MIE Inc., Smyrna, GA), and the DustTrak aerosol monitor (TSI Inc., St. Paul, MN; Brauer, 1995; Howard-Reed et al., 2000; Quintana et al., 2000, 2001; Chang et al., 2000; Magari et al., 2002). The Radiance nephelometer is an active monitor with a 530-nm wavelength defined by an optical filter. The pDR uses a wavelength of 880 nm, making it more sensitive than the nephelometer to particles > 1 µm in diameter but less sensitive to particles < 0.3 µm diameter. The pDR (usually used in the passive mode, without a size fractionation inlet) has been widely used in exposure studies because of its light weight (no pump required), data logging capability, and easy handling (Quintana et al., 2000, 2001; Williams et al., 2000a; Muraleedharan and Radojevic, 2000; Rea et al., 2001). A portable condensation nuclei counter (with a lower size limit of 20 nm diameter) has also been used in exposure studies (Abraham et al., 2002).

Simultaneous measurements of both gravimetric mass and light scattering coefficients have provided assessment of both the nephelometer and pDR in outdoor and indoor air (Brauer, 1995; Quintana et al., 2000; Scheff and Wadden, 1979; Waggoner and Weiss, 1980; Sioutas et al., 2000). Less information is available to assess their use in personal environments. Liu et al. (2002) compared the pDR and the Radiance Nephelometer against gravimetric measurements from both the Harvard impactor for PM_{2.5} (HI_{2.5}) and the Harvard personal environmental monitor (HPEM_{2.5}) in indoor, outdoor, and personal settings at residences and on the person of elderly subjects living across the metropolitan Seattle area.

Ten-minute averages for collocated indoor nephelometers and pDRs at diverse residential sites have produced varied coefficients of determination across homes ($R^2 = 0.75$ to 0.96). The

differences between the nephelometer and the pDR increased during cooking hours. When regressed against 24-h gravimetric measurements made with the HI_{2.5}, the Radiance nephelometer showed higher coefficients of determination indoors ($R^2 = 0.81$ [non-cooking days] and 0.93 [days with cooking]) and outdoors ($R^2 = 0.88$ [outdoors]) than the pDR indoors ($R^2 = 0.84$ [non-cooking days] and 0.77 [days with cooking]). A comparison of the pDR versus the HPEM_{2.5}, when both were carried by subjects, gave lower coefficients of determination ($R^2 = 0.44$ [non-cooking days] and 0.60 [days with cooking]).

Liu et al. (2002) also report:

“The pDR measurements generally overestimated gravimetric PM_{2.5} measurements, with a mean indoor pDR/HI_{2.5} ratio of 1.56 ± 0.07 . As the pDR uses a calibration dust density of 2.6 g/cm^3 , overestimating gravimetric measurements by a factor of 1.56 gives an estimated indoor particle density of 1.67 g/cm^3 . This estimated indoor particle density is within the range for sulfates (1.76), nitrate (2.1), organic carbon (1.4), and elemental carbon (2.0) (Ames et al., 2000). Somewhat different results were obtained for the comparison with the personal gravimetric measurements. When both the pDR and HPEM_{2.5} were carried by subjects, the mean personal pDR/HPEM_{2.5} ratio was 1.27 ± 0.08 . This overestimation factor leads to an estimated density of the personal aerosol of 2.05 g/cm^3 . This larger density seem reasonable if the personal aerosol is largely suspended crustal material, as has been found by Ozkaynak et al. (1996a). These overestimation factors were similar but slightly lower than the mean indoor pDR/HI_{2.5} ratio 1.41 reported by Quintana et al. (2000) and the mean personal pDR/PEM ratio 1.49 by Howard-Reed et al. (2000).”

Howard-Reed et al. (2000) compared pDR readings, averaged over 24 h with corresponding gravimetric measurements taken with MSP and PM_{2.5} PEMs (2 L/min). Both personal samplers were worn by participants in a Baltimore exposure study. Howard-Reed et al. (2000) reported a coefficient of determination of $R^2 = 0.66$ and a regression equation of $\text{pDR } (\mu\text{g/m}^3) = 1.10 \times \text{PEM } (\mu\text{g/m}^3) + 5.84$ for 34 24-h measurements. Lanki et al. (2002), using the pDR in a flow-through mode and a PM_{2.5} PEM of their own design, found $R^2 = 0.66$ and $\text{pDR} = 1.85 \times \text{PEM} + 4.52$. Sioutas et al. (2000) have also evaluated the pDR in a flow-through mode of operation (2 L/min, with a diffusion dryer in the inlet). They found a significant increase in pDR to gravimetric mass for relative humidity (RH) above 60%. The ratio was 2:1 at RH = 80%. They subsequently used a diffusion drying tube in the inlet. With this system, they were able to study the sensitivity of the pDR to particle size by comparing the pDR reading to the mass measured by the multi-orifice, uniform deposit detector (MOUDI) as a function of the

particle mass mean diameter (MMD) as determined by the MOUDI. As the MMD increases from 0.3 to 1.1 μm , the pDR-to-MOUDI ratio increases from approximately 0.7 to about 1.6. As the MMD increases to 1.5 μm , the ratio decreases to about 1.0.

Chang et al. (2001) evaluated the DustTrak against a MSP PEM in a roll-around system. Particles were passed through at 2.5 μm PEM size-selective inlet and a Nafion diffusion dryer before entering the DustTrak. Hourly $\text{PM}_{2.5}$ concentrations measured by the DustTrak were averaged and compared to 12-h integrated MSP PEM $\text{PM}_{2.5}$ measurements. The DustTrak concentrations were higher than the PEM by a factor of about 2. The use of a DustTrak correction factor (2.07 in summer and 2.02 in winter) resulted in good agreement between the two methods, with $R^2 = 0.87$ in the summer (35 pairs) and $R^2 = 0.81$ in the winter (42 pairs).

5.2.4.1.2 Microenvironmental Monitoring Methods

Direct measurements of microenvironmental PM concentrations, which are used with models to estimate personal exposure to PM, also employ methods similar to those used for ambient PM. These methods differ from PEMs in that they are stationary with respect to the microenvironment (i.e., a stationary PEM). Microenvironmental monitoring methods include filter-based mass measurements of particle size fractions (PM_{10} , $\text{PM}_{2.5}$) usually integrated over either a 24- or 12-h period such as the Federal Reference Methods (FRMs) for $\text{PM}_{2.5}$ and PM_{10} . Flow rates vary between various devices from 4 to 20 L/min. Larger sample volumes allow more extensive chemical characterization to be conducted on microenvironmental samples. Because more than one pumping system can be used in a microenvironment, $\text{PM}_{2.5}$ and PM_{10} can be collected simultaneously.

The Harvard School of Public Health has developed a 10 L/min sampler, known as the Harvard Impactor ($\text{HI}_{2.5}$, Air Diagnostics and Engineering, Inc., Naples, ME), which has been used as an indoor or outdoor monitor in exposure studies (Liu et al., 2002). Babich et al. (2000) reported that the $\text{HI}_{2.5}$ showed excellent agreement ($R^2 = 0.99$) with the $\text{PM}_{2.5}$ Federal Reference Method (FRM) for 81 24-h samples in Riverside and Bakersfield, CA. The HI also showed good precision (4.8%) for 243 24-h collocated samples over eight studies. Other continuous ambient PM measurement methods that have been utilized for microenvironmental monitoring include the Tapered Element Oscillating Microbalance (TEOM) and nephelometers. Various continuous

techniques for counting particles by size (Climet, LASX, SMPS, APS) also have also been used. Measurement techniques are discussed in Chapter 2.

5.2.4.2 Indirect or Modeling Methods

5.2.4.2.1 Personal Exposure Models

As a relatively new field, exposure modeling for PM mass (PM_{2.5} and PM_{10-2.5}) and PM chemical constituents faces significant methodological challenges and input data limitations. Exposure models typically use one of two general approaches: (1) a time-series approach that estimates individuals' microenvironmental exposures sequentially over time or (2) a time-averaged approach that estimates individuals' microenvironmental exposures using average microenvironmental concentrations and the total time spent in each microenvironment. Although the time-series approach to modeling personal exposures provides the appropriate structure for accurately estimating personal exposures (Esmen and Hall, 2000; Mihlan et al., 2000), a time-averaged approach typically is used when the input data needed to support a time-series model are not available. However, the time-varying dose profile of an exposed individual can only be modeled by using the time-series approach (McCurdy, 1997, 2000).

Even though the processes that lead to exposure are nonlinear in nature, personal exposure models are often used to combine microenvironmental concentration data with human activity pattern data to estimate personal exposures. Time-averaged models can be used to estimate personal exposure for an individual or for a defined population. Total personal exposure models estimate exposures for all of the different microenvironments in which a person spends time, and total average personal exposure is calculated from the sum of these microenvironmental exposures:

$$E = \sum_j E_j = \frac{1}{T} \left(\sum_j C_j t_j \right), \quad (5-3)$$

where E_j is the personal exposure in each microenvironment, j (Duan, 1982). Example microenvironments include outdoors, indoors at home, indoors at work, and in transit. Each microenvironmental exposure, E_j , is calculated from the average concentration in microenvironment j , C_j , weighted by the time spent in microenvironment j , t_j . T is the sum

of t_j over all j . This model has been applied to concentration data in a number of studies (Ott, 1984; Ott et al., 1988, 1992; Miller et al., 1998; Klepeis et al., 1994; Lachenmyer and Hidy, 2000).

Many exposure studies employ 24-h average measurements of concentration indoors and outdoors and use these concentrations with time spent indoors and outdoors, as shown in Equation 5-3. It is important to note that although measurement data may be an average concentration over some time period (i.e., 24 h), significant variations in PM concentrations can occur during that time period. Thus, an error may be introduced if real-time concentrations are highly variable and an average concentration for a microenvironment is used to estimate exposure when the individual is in that microenvironment for only a fraction of the total time. This may create large errors if the indoor (e.g., in a house) 24-h average includes significant time periods when there are no people in the house, because the indoor concentrations are increased by the activities of people. In an effort to overcome these errors, the Air Pollution Exposure Distribution within Urban Populations in Europe (EXPOLIS) study (Kousa et al., 2002) turned outdoor samplers off when the subject was indoors and the indoor sampler off with the subject was outdoors. This provides a better estimate of E_a and E_i to compare with the PEM measurement and allows a better calculation of E_{pc} . However, it does not provide data that can be used to regress C or C_o with C_i . This problem could be overcome with continuous monitoring, and the extent of the error could be calculated. Also, shorter time interval information might be useful in examining relationships between acute effects and short-term exposures.

Microenvironmental concentrations used in the exposure models can be measured directly or estimated by one or more microenvironmental models. Microenvironmental models vary in complexity from a simple indoor/outdoor (I/O) ratio to a multi-compartmental mass-balance model. A discussion of microenvironmental models is presented in Section 5.3.4.2.2.

On the individual level, the time spent in the various microenvironments is obtained from time-activity diaries that are completed by the individual. For population-based estimates, the time spent in various microenvironments is obtained from human activity databases. Many of the largest human activity databases have been consolidated by the EPA's National Exposure Research Laboratory (NERL) into one comprehensive database called the Consolidated Human Activity Database (CHAD). CHAD contains over 22,000 person-days of 24-h activity data from

11 different human activity-pattern studies (McCurdy et al., 2000). Population cohorts with diverse characteristics can be constructed from the activity data in CHAD and used for exposure analysis and modeling (McCurdy, 2000). These databases can also be used to estimate inhalation rates based on activity levels, age, gender, and weight for dosimetry calculations. A human activity database may contain information on human location, activity, and exertion rate as well as information such as the presence of combustion sources (e.g., wood fireplaces, smokers). However, in exposure studies, “activity” usually refers to a person’s location in space, i.e., in what microenvironment at what times. In dosimetry, “activity” is used as an indication of the level of physical exertion and is used to estimate breathing rate and the extent of mouth, nose, or combined breathing. Table 5-2 is a summary of the human activity studies in CHAD.

Methodologically, personal exposure models can be divided into three general types: (1) statistical models based on empirical data obtained from one or more personal monitoring studies, (2) simulation models based upon known or assumed physical relationships, and (3) physical-stochastic models that include Monte Carlo or other techniques to explicitly address variability and uncertainty in model structure and input data (Ryan, 1991; MacIntosh et al., 1995). The attributes, strengths, and weaknesses of these model types are discussed by Ryan (1991), National Research Council (1991), Frey and Rhodes (1996), and Ramachandran and Vincent (1999). A review of the logic of exposure modeling is found in Klepeis (1999).

Personal exposure models that have been developed for PM are summarized in Table 5-3. The regression-based models (Johnson et al., 2000; Janssen et al., 1997; Janssen et al., 1998a) were developed specifically to account for the observed difference between personal exposure and microenvironmental measurements and are based on data from a single study, limiting their utility for broader purposes. Other types of models in Table 5-3 were limited by a lack of data for the various model inputs. For example, ambient PM monitoring data are not generally of adequate spatial and temporal resolution for these models. Lurmann and Korc (1994) assumed a constant relationship between coefficient of haze (COH) and PM_{10} and used site-specific COH information to stochastically develop a time series of 1-h PM_{10} data from every sixth day 24-h PM_{10} measurements. A mass-balance model typically was used for indoor microenvironments when sufficient data were available, such as for a residence. For most other microenvironments, I/O ratios were used because of the lack of data required for a mass-balance model. In addition, only the deterministic model, PMEX, included estimation of inhaled dose

TABLE 5-2. ACTIVITY PATTERN STUDIES INCLUDED IN THE CONSOLIDATED HUMAN ACTIVITY DATABASE (CHAD)

Study Name	Calendar Time Period of the Study	Age ¹	Days ²	Diary		Rate ⁵	Documentation or Reference	Notes
				Type ³	Time ⁴			
Baltimore, MD	Jan-Feb 1997 Jul-Aug 1998	65+	391	Diary; 15-min blocks	24-h Standard	No	Williams et al. (2000a,b)	Multiple days, varying from 5-15; part of a PM _{2.5} PEM study
CARB: Adolescents and Adults	Oct 1987-Sept 1988	12-94	1762	Retrospective	24-h Standard	No	Robinson et al. (1991) Wiley et al. (1991a)	
CARB: Children	Apr 1989-Feb 1990	0-11	1200	Retrospective	24-h Standard	No	Wiley et al. (1991b)	
Cincinnati (EPRI)	Mar-Apr and Aug 1985	0-86	2614	Diary	24 h; nominal 7 pm-7 am	Yes	Johnson (1989)	3 consecutive days; 186 P-D removed ⁷
Denver (EPA)	Nov 1982-Feb 1983	18-70	805	Diary	24 h; nominal 7 pm-7 am	No	Akland et al. (1985) Johnson (1984)	Part of CO PEM ⁶ study; 2 consec. days; 55 P-D removed ⁷
Los Angeles: Elem. School Children	Oct 1989	10-12	51	Diary	24-h Standard	Yes	Spier et al. (1992)	7 P-D removed ⁷
Los Angeles: High School Adoles.	Sept-Oct 1990	13-17	43	Diary	24-h Standard	Yes	Spier et al. (1992)	23 P-D removed ⁷
National: NHAPS-A ⁸	Sept 1992-Oct 1994	0-93	4723	Retrospective	24-h Standard	No ⁹	Klepeis et al. (1995) Tsang and Klepeis (1996)	A national random-probability survey
National: NHAPS-B ⁸	As above	0-93	4663	Retrospective	24-h Standard	No ⁹	As above	As above
U. Michigan: Children	Feb-Dec1997	0-13	5616	Retrospective	24-h Standard	No	Institute for Social Research (1997)	2 days of data: one is a weekend day
Valdez, AK	Nov 1990-Oct 1991	11-71	401	Retrospective	Varying 24-h period	No	Goldstein et al. (1992)	4 P-D removed ⁷
Washington, DC (EPA)	Nov 1982-Feb 1983	18-98	699	Diary	24 h; nominal 7 pm-7 am	No	Akland et al. (1985) Hartwell et al. (1984)	Part of a CO PEM ⁶ study; 6 P-D removed ⁷

¹ All studies included both genders. The age range depicted is for the subjects actually included; in most cases, there was not an upper limit for the adult studies. Ages are inclusive.

Age 0 = babies < 1 year old.

² The actual number of person-days of data in CHAD after the “flagging” and removal of questionable data. See the text for a discussion of these procedures.

³ Retrospective: a “what did you do yesterday” type of survey; also known as an ex-post survey. Diary: a “real-time” paper diary that a subject carried as he or she went through the day.

⁴ Standard = midnight to midnight.

⁵ Was activity-specific breathing rate data collected?

⁶ PEM = a personal monitoring study. In addition to the diary, a subject carried a small CO or PM_{2.5} monitor throughout the sampling period.

⁷ P-D removed = The number of person-days of activity pattern data removed from consolidated CHAD because of missing activity *and* location information; completeness criteria are listed in the text.

⁸ National Human Activity Pattern Study; A = the air version; B = the water version. The activity data obtained on the two versions are identical.

⁹ A question was asked regarding which activities (within each 6-h time block in the day) involved “heavy breathing,” lifting heavy objects, and running hard.

TABLE 5-3. PERSONAL EXPOSURE MODELS FOR PARTICULATE MATTER

Study Citation	Model Name	Model Type	Microenvironments or Predictors	Output	Notes
Time-series Models					
Hayes and Marshall (1999)	PMEX	Deterministic	Indoors: residential, work, school Outdoors: near roadway, other Motor vehicle	Inhaled dose of PM ₁₀ Hourly for 24 h By age/gender groups Source contributions	Used IAQM Used human activity data with activity-specific breathing rate info
Johnson et al. (2000)	None	Regression-based	Auto travel, roadside, ETS, food prep., grilling, high ambient PM	PM _{2.5} exposure 24-h average	Developed from scripted activity study (Chang et al., 2000)
Klepeis et al. (1994)	None	Stochastic	ETS, cooking, cleaning, attached garage, wood burning	Respirable particle (PM _{3.5}) exposure	
Lurmann and Korc (1994)	REHEX-II	Stochastic	12 residential with different sources, restaurant/bar, nonresidential indoors, in transit, outdoors	Distribution of PM ₁₀ exposure for population Three averaging times (1 h, 24 h, season)	Fixed I/O ratio of 0.7 for indoors w/o sources and 1.2 for in transit Reduced form mass balance model for indoors with PM sources
Koontz and Niang (1998)	CPIEM	Stochastic	Indoors: residence, office, industrial plant, school, public building, restaurant/lounge, other. Outdoors, in vehicle	Distribution of PM ₁₀ exposure for population	Used California activity pattern and breathing rate data Used either a mass balance model or I/O ratio distribution for indoor microenvironments Indoor sources included
Time-averaged Models					
Clayton et al. (1999a)	SIM	Stochastic		Distribution of annual PM _{2.5} exposures	Based on 3-day ambient measurements
Janssen et al. (1997)	None	Regression-based	Smoking parent, ETS exposure, outdoor physical activity	Accounts for difference between personal and microenvironmental PM ₁₀	Children only
Janssen et al. (1998a)	None	Regression-based	Number of cigarettes smoked, hours of ETS exposure, residence on busy road, time in vehicle	Accounts for difference between personal and microenvironmental PM ₁₀	Adults only

TABLE 5-3 (cont'd). PERSONAL EXPOSURE MODELS FOR PARTICULATE MATTER

Study Citation	Model Name	Model Type	Microenvironments or Predictors	Output	Notes
Time-averaged Models (cont'd)					
Ott et al. (2000)	RCS	Statistical	Not separated	Distribution of PM ₁₀ exposure for population	A random-component superposition (RCS) model that uses distribution of ambient PM ₁₀ and estimated nonambient PM ₁₀ concentrations Results for Ontario, Canada not corrected for 72-h compared to 24-h averaging time in Riverside, CA and Phillipsburg, NJ
Burke et al. (2001)	SHEDS-PM	Stochastic	Outdoors, indoors: residence, office, stores, school, in vehicle, restaurant/lounge	PM _{2.5} exposure distributions for population by age, gender, smoking and employment status; PM _{2.5} exposure uncertainty predictions. Percent contribution from PM of ambient origin to total personal exposures	A 2-stage Monte-Carlo simulation model for predicting population distribution of daily-average personal exposures to PM. Model has been applied to Philadelphia using spatially and temporally interpolated PM _{2.5} ambient measurements from 1992-1993 and 1990 census data. Does not consider PM _{2.5} exposure from active smoking or exposure in subways
Chao and Tung (2001)	None	Mass balance with empirical corrections	Indoors in unoccupied residences in Hong Kong	Predictions of ambient PM in indoor microenvironments	Model makes corrections for nonideal mixing (residence with multiple compartments with limited intermixing)

from activity-specific breathing rate information. Data from recent PM personal exposure and microenvironmental measurement studies should help in the future to facilitate the development of improved personal exposure models for PM.

An integrated human exposure source-to-dose modeling system that will include exposure models to predict population exposures to environmental pollutants, such as PM, currently is being developed by the EPA/NERL. A first-generation population exposure model for PM, called the Stochastic Human Exposure and Dose Simulation (SHEDS-PM) model, recently has been developed. The SHEDS-PM model uses a 2-stage Monte Carlo sampling technique previously applied by MacIntosh et al. (1995) for benzene exposures. This technique allows for separate characterization of variability and uncertainty in the model predictions to predict the distribution of total exposure to PM for the population of an urban/metropolitan area and to estimate the contribution of ambient PM to total PM exposure. Results from a case study using data from Philadelphia, PA have been reported (Burke et al., 2001). Work is underway to link exposure modeling with dosimetry so as to provide estimates of integrated PM doses for different regions of the lung. In the future, both exposure and dose metrics generated for various subgroups of concern should aid in evaluation of PM health effects.

5.2.4.2.2 Microenvironmental Models

The mass-balance model has been used extensively in exposure analysis to estimate PM concentrations in indoor microenvironments (Calder, 1957; Sexton and Ryan, 1988; Duan, 1982, 1991; McCurdy, 1995; Johnson, 1995; Klepeis et al., 1995; Dockery and Spengler, 1981; Ott, 1984; Ott et al., 1988, 1992, 2000; Miller et al., 1998; Mage et al., 1999; Wilson et al., 2000). The mass balance model describes the infiltration of particles from outdoors into the indoor microenvironment, the removal of particles in indoor microenvironments, and the generation of particles from indoor sources:

$$V dC_i / dt = vPC - vC_i - kVC_i + Q_i, \quad (5-4)$$

- where
- V = volume of the well-mixed indoor air (m³);
 - C_i = concentration of indoor PM (μg/m³);
 - v = volumetric air exchange rate between indoors and outdoors (m³/h);

- P = penetration ratio, the fraction of ambient (outdoor) PM that is not removed from ambient air during its entry into the indoor volume;
- C = concentration of PM in the ambient air ($\mu\text{g}/\text{m}^3$);
- k = removal rate (h^{-1}); and
- Q_i = rate of generation of particles from indoor sources ($\mu\text{g}/\text{h}$).

Q_i contains a variety of indoor, particle-generating sources, including combustion or mechanical processes; condensation of vapors formed by combustion or chemical reaction; suspension from bulk material; and resuspension of previously deposited PM. The removal rate, k , includes dry deposition to interior surfaces by diffusion, impaction, electrostatic forces, and gravitational fallout. It may include other removal processes, such as filtration by forced air heating, ventilation, or air conditioning (HVAC) or by independent air cleaners. All parameters except V are functions of time. P and k also are functions of particle aerodynamic diameter, D_a , air exchange rate, v , and house characteristics such as the surface to volume ratio, type of surface, etc. All variables in Equation 5-3 will have distributions within the population and, in some cases, may vary by a factor of 5 to 10. It is important to determine the distribution of these variables. Sensitivity and uncertainty analyses are necessary when attempting to explain model results.

In addition to the mass-balance model, some single-source or single-microenvironment models exist. However, most are used to estimate personal exposures to environmental tobacco smoke (ETS). These models include both empirically based statistical models and physical models based on first principles: some are time-averaged, whereas others are time-series. These models evaluate the contribution of ETS to total PM exposure in an enclosed microenvironment and can be applied as activity-specific components of total personal exposure models. Examples of ETS-oriented personal exposure models include Klepeis (1999), Klepeis et al. (1996, 2000), Mage and Ott (1996), Ott (1999), Ott et al. (1992, 1995), and Robinson et al. (1994).

5.2.4.3 Methods for Estimating Personal Exposure to Ambient Particulate Matter

In keeping with the various components of PM exposure described in Section 5.3.2, personal exposure to PM can be expressed as the sum of exposure to particles from different sources summed over all microenvironments in which exposure occurs. Total personal exposure may be expressed as:

$$T = E_a + E_{ai} + E_{ig} + E_{ir} + E_{pc} \quad (5-5)$$

or

$$T = A + N, \quad (5-6)$$

where T is the total personal exposure to ambient and nonambient PM, E_a is personal exposure to ambient PM while outdoors, E_{ai} is personal exposure to ambient PM that has infiltrated indoors while indoors, E_{ig} is personal exposure to indoor-generated PM, E_{ir} is exposure to indoor-reaction PM, and E_{pc} is personal exposure to PM from personal activity (personal cloud). T can also be expressed as A + N where A is ambient PM exposure ($E_a + E_{ai}$) and N is nonambient PM exposure ($E_{ig} + E_{ir} + E_{pc}$). Although personal exposure to ambient and nonambient PM cannot be measured directly, they can be calculated or estimated from other measurement data.

Approaches for estimating these components of PM exposure are described in the next section.

5.2.4.3.1 Mass Balance Approach

Ambient-Indoor Concentrations of Particulate Matter

The mass balance model described above (Equation 5-4) has been used to estimate PM concentrations in indoor microenvironments. This model also may be used to estimate ambient-indoor (C_{ai}) and indoor-generated (C_{ig}) PM concentrations. The mass balance model can be solved for C_{ai} and C_{ig} assuming equilibrium conditions, i.e., all variables remain constant (Ott et al., 2000; Dockery and Spengler, 1981; Koutrakis et al., 1992) and no indoor reaction PM (C_{ir}). By substituting $a = v/V$, where a = the number of air exchanges per hour substituting, $dC_{ai} + dC_{ig}$ for dC_i in Equation 5-4, and assuming that dC_{ai} and $dC_{ig} = 0$, i.e., ambient-indoor PM (C_{ai}) and indoor-generated PM (C_{ig}) are at equilibrium, C_{ai} and C_{ig} are given by Equations 5-7 and 5-8.

$$C_{ai} = C \left[\frac{Pa}{(a + k)} \right] \quad (5-7)$$

$$C_{ig} = \frac{Q_i}{V(a + k)} \quad (5-8)$$

Equations 5-7 and 5-8 assume equilibrium conditions and, therefore, are valid only when the parameters P , k , a , C , and Q_i are not changing rapidly and when the C s are averaged over several hours. It should be understood that equilibrium is a simplification of indoor microenvironments that are occupied by residents. This assumption of equilibrium may only represent a virtual set of individuals or populations at risk. Under certain conditions (e.g., air-conditioned homes, homes with HVAC or air cleaners that cycle on and off, or ambient pollutants with rapidly varying concentrations), nonequilibrium versions of the mass-balance model (Ott et al., 2000; Freijer and Bloemen, 2000; Isukapalli and Georgopoulos, 2000) are likely to provide a more accurate estimate of C_{ai} and C_{ig} . However, the equilibrium model provides a useful, although simplified, example of the basic relationships (Ott et al., 2000).

Equation 5-7 may be rearranged further to give C_{ai}/C , the equilibrium fraction of ambient PM that is found indoors, defined as the infiltration factor, F_{INF} , (Dockery and Spengler, 1981).

$$F_{INF} = \frac{C_{ai}}{C} = \frac{Pa}{a+k} \quad (5-9)$$

The penetration ratio, P , and the decay rate, k , can be estimated using a variety of techniques. A discussion of these variables and estimation techniques is given in Section 5.4.3.2.2. Both P and k are a function of particle aerodynamic diameter, air exchange rate, and housing characteristics. F_{INF} will also be a function of these parameters; and, as a result, F_{INF} may vary substantially within a population. Distributions of this parameter should be estimated to understand the uncertainty and variability associated with estimating exposure to PM of ambient origin. The distribution of daytime F_{INF} , as estimated from PTEAM data, is shown in Figure 5-2a (Wilson et al., 2000).

Personal Exposure to Ambient Particulate Matter

Personal exposure to ambient PM, A , may be estimated using ambient-indoor PM concentration, C_{ai} , from the mass balance model, ambient outdoor PM concentrations, C , and information on the time an individual spent in the various microenvironments. For a two-compartment model, A may be expressed as





Figure 5-2. Distribution of F_{INF} (a) and α (b) for daytime as estimated from PTEAM study data.

Source: Wilson et al. (2000).

$$A = yC + (1 - y)C_{ai} = yC + (1 - y)C \left[\frac{Pa}{(a + k)} \right], \quad (5-10)$$

where y is the fraction of time that an individual spent outdoors, and $(1 - y)$ is the fraction of time spent indoors.

It is convenient to express personal exposure to ambient PM, A , as the product of the ambient PM concentration, C , and a personal exposure or attenuation factor. Following the usage in several recent papers (Zeger et al., 2000; Dominici et al., 2000; Ott et al., 2000), the symbol α will be used for this attenuation factor. Equation 5-10 can be rearranged to obtain an expression for α :

$$\alpha = \frac{T}{C} = y + (1 - y) \left[\frac{Pa}{a + k} \right] \quad (5-11)$$

Substituting equation 5-9 in equation 5-11 gives a relationship for α in terms of the infiltration factor F_{INF} and the fraction of time spent in the various microenvironments:

$$\alpha = y + (1 - y)F_{INF}. \quad (5-12)$$

Thus, personal exposures to ambient PM, A , may be calculated from measurable quantities:

$$A = \alpha C. \quad (5-13)$$

The factor α can be measured directly or calculated from measured or estimated values of the parameters a , k , and P and the time spent in various microenvironments from activity pattern diaries (Wilson et al., 2000). Because α depends on housing and lifestyle factors, air exchange rate, and PM deposition rate, it could vary to a certain extent from region to region and from season to season. Consequently, predicted exposures based on these physical modeling concepts provide exposure distributions derived conceptually as resulting from housing, lifestyles, and meteorological considerations. For any given population the coefficient α may represent

substantial intra- and inter-personal variability based on personal activities, housing characteristics, particle size, and composition. The distribution of daytime α as estimated from PTEAM data is shown in Figure 5-2b. Note that the distribution of α is shifted to higher values compared to F_{INF} because of the inclusion of time outdoors in α . Distributions of α should be determined using population studies in order to evaluate the uncertainty and variability associated with model exposures.

The mass-balance model has been used to separate indoor concentrations into ambient and nonambient components. This approach, based on Equation 5-5 as given in Duan (1982) and called superposition of component concentrations, has been applied using multiple microenvironments to estimate exposures to carbon monoxide (Ott, 1984; Ott et al., 1988, 1992), volatile organic compounds (Miller et al., 1998), and particles (Koutrakis et al., 1992; Klepeis et al., 1994). However, in these studies and in most of the exposure literature, the ambient and nonambient components are added to yield a personal exposure from all sources of the pollutant. The use of the mass-balance model, ambient concentrations, and exposure parameters to estimate exposure to ambient PM and exposure to indoor-generated PM separately as different classes of exposure has been discussed in Wilson and Suh (1997) and in Wilson et al. (2000).

5.2.4.3.2 The Sulfate Ratio Technique for Estimating Ambient PM Exposure

The ratio of personal exposure to ambient concentration for sulfate has been recommended as a technique to estimate α (Wilson et al., 2000). If sulfate has no indoor sources, then $A^S = T^S$. (Superscript S refers to sulfate; superscript 2.5 to $PM_{2.5}$.) As can be seen in Equation 5-11, P and k depend on particle size, but a and y do not. It has been known since the mid-1970s that sulfate and accumulation-mode mass have similar size distributions (Whitby, 1978). Therefore, if the coarse mode contribution to $PM_{2.5}$ is small, so that sulfate and $PM_{2.5}$ would have similar size distributions, then $T^S/C^S = A^{2.5}/C^{2.5} = \alpha^S = \alpha^{2.5}$. Sulfate is formed in the ambient air via photochemical oxidation of gaseous sulfur dioxide arising from the primary emissions from the combustion of fossil fuels containing sulfur. It also arises from the direct emissions of sulfur-containing particles from nonanthropogenic sources (e.g., volcanic activity, windblown soil). In the indoor environment, the only common sources of sulfate may be resuspension by human activity of deposited PM containing ammonium sulfates or soil sulfates that were tracked into the home. However, resuspended PM will be mostly larger than $PM_{2.5}$. In some homes, an

unvented kerosene heater using a high-sulfur fuel may be a major contributor during winter (Leaderer et al., 1999). The use of matches to light cigarettes or gas stoves can also be a source of sulfates.

Studies that have used the sulfate ratio technique to estimate α and ambient PM exposures are discussed in Section 5.4.3.1. When there are no indoor sources of accumulation-mode sulfates, one may deduce that the ambient-to-personal relationship found for sulfates probably would be the same as that for other PM with the same size range and physical/chemical properties. This assumption has been validated for several homes in Boston (Sarnat et al., 2002). For particle sizes within the accumulation-mode size range, the ratio C_{air}/C was similar for sulfate and $\text{PM}_{2.5}$ as estimated from SMPS measurements. However, ambient PM with different physical or chemical characteristics than sulfate will not behave similarly to sulfate. Sulfate has been used as a marker of outdoor air in the indoor microenvironments (Jones et al., 2000; Ebel et al., 2000). However, the personal exposure of sulfate ($T^s = A^s$) should not be taken as an indicator or surrogate for ambient $\text{PM}_{2.5}$ exposure ($A^{2.5}$) unless it has been previously determined that $\text{PM}_{2.5}$ and sulfate concentrations are highly correlated. This may be the case in some air sheds with high sulfate concentrations, but will not be true in general.

5.2.4.3.3 Source-Apportionment Techniques

Source apportionment techniques provide a method for determining personal exposure to PM from specific sources. If a sufficient number of samples are analyzed with sufficient compositional detail, it is possible to use statistical techniques to derive source category signatures, identify indoor and outdoor source categories, and estimate their contribution to indoor and personal PM. Daily contributions from sources that have no indoor component can be used as tracers to generate exposure to ambient PM of similar aerodynamic size or directly as exposure surrogates in epidemiological analyses. Studies that have used source-apportionment are discussed in Section 5.4.3.3 (i.e., Özkaynak and Thurston, 1987; Yakovleva et al., 1999; Mar et al. 2000; Laden et al., 2000).

5.2.4.3.4 Comparison and Validation of Methods

There are as yet no published comparisons or validations of the various methods for estimating personal exposure to ambient PM on an average or individual daily basis.

5.3 SUMMARY OF PARTICULATE MATTER MASS DATA

5.3.1 Types of Particulate Matter Exposure Measurement Studies

A variety of field measurement studies have been conducted to quantify personal exposure to PM mass, to measure microenvironmental concentrations of PM, to evaluate relationships between personal exposure to PM and PM air concentrations measured at ambient sites, and to evaluate factors that affect exposure. In general, exposure measurement studies are of two types depending on how the participants are selected for the study. In a *probability* study, participants are selected using a probability sampling design where every member of the defined population has a known, positive probability of being included in the sample. Probability study results can be used to make statistical inferences about the target population. In a *purposeful* or *nonprobability* design, any convenient method may be used to enlist participants; and the probability of any individual in the population being included in the sample is unknown (National Research Council, 1991). Participants in purposeful samples may not have the same characteristics that would lead to exposure as the rest of the unsampled population. Thus, results of purposeful studies apply only to the subjects sampled on the days that they were sampled and not to other subjects or other periods of time. Although such studies may report significant differences, confidence intervals, and *p* values, they do not have inferential validity (Lessler and Kalsbeek, 1992). Purposeful studies, however, may have generalizability (external validity). The extent of generalizability is a matter of judgement based on study participant characteristics. Purposeful studies of PM personal exposure can provide data with which to develop relationships based on important exposure factors and can provide useful information for developing and evaluating either statistical or physical/chemical human exposure models.

Regardless of the sampling design (probability or purposeful), there are three general categories of study design that can be used to measure personal exposure to PM and evaluate the relationship between personal PM exposure levels and ambient PM concentrations measured simultaneously: (1) longitudinal, in which each subject is measured for many days; (2) pooled, in which each subject is measured for only one or two days, different days for different subjects; and (3) daily-average, in which many subjects are measured on the same day. Only one study, in which 14 subjects were measured for 14 days, provided sufficient data for a comparison of longitudinal and daily-average data (Lioy et al., 1990). Longitudinal and pooled studies are discussed in Section 5.4.3.1.1.

5.3.2 Available Data

5.3.2.1 Personal Exposure Data

Table 5-4 gives an overview of the personal exposure studies that have been reported since the 1996 PM AQCD. In addition, major studies that were reported before that time also have been included to provide a comprehensive evaluation of data in this area. Table 5-4 gives information on the sampling and study designs, the study population, the season, number of participants, PM exposure metric, and the PM size fraction measured.

Although there are a number of studies listed in the table, the data available with which to evaluate longitudinal relationships and the factors that influence these are limited. Few studies are based on probability sampling designs that allow study results to be inferred to the general population and to develop distributional data on exposures and the factors that affect exposure. Unfortunately, none of these probability studies used a longitudinal study design. This limits our ability to provide population estimates and distributional data on the relationship between personal PM exposures and ambient site measurements. In addition, most of the probability studies of PM exposure were conducted during a single season; thus, variations in ambient concentrations, air exchange rates, and personal activities are not accounted for across seasons. In these cases, study results are only applicable to a specific time period. Longitudinal studies, on the other hand, generally have small sample sizes and use a purposeful sampling design. Some studies did not include ambient site measurements to allow comparisons with the exposure data. Approximately half of these studies monitored $PM_{2.5}$. Only one or two studies measured both PM_{10} and $PM_{2.5}$ to provide information on $PM_{10-2.5}$.

Four large-scale probability studies that quantify personal exposure to PM under normal ambient source conditions have been reported in the literature. These include the EPA's PTEAM study (Clayton et al., 1993; Özkaynak et al., 1996a,b); the Toronto, Ontario study (Clayton et al., 1999a; Pellizzari et al., 1999); the EXPOLIS study (Jantunen et al., 1998, 2000; Oglesby, et al., 2000; Gotshi et al., 2002; Kousa et al., 2002); and a study of a small, highly polluted area in Mexico City (Santos-Burgoa et al., 1998). A fifth study conducted in Kuwait during the last days of the oil-well fires (Al-Raheem et al., 2000) is not reported here because the ambient PM levels were not representative of normal ambient source conditions.

TABLE 5-4. SUMMARY OF RECENT PM PERSONAL EXPOSURE STUDIES

Study Design	Study Location and Population	No. of Subjects	Study Period	Age (Years)	Days per Subject	PM Exposure ^a Metrics	PM Size Measured ^b	Co-pollutant Metrics	Reference
Probability Studies									
Pooled ^c	Indianapolis, IN	240	1996	16-?	One 72-h sample/ subject	P, I, A, O	PM _{2.5} , PM ₁₀	Mn, Al, Ca	Pellizzari et al. (2001)
Pooled	Riverside, CA, PTEAM	178	Fall 1990	10-70	1 (12 h)	P, I, O, A	PM ₁₀		Clayton et al. (1993) Özkaynak et al. (1996a,b)
Pooled	Basel, Switzerland, EXPOLIS	50	1997		1 (48 h)	P, I, P	PM _{2.5}	VOC, CO, NO ₂ , S, K, Pb, Br, Ca	Oglesby et al. (2000) Jantunen et al. (1998)
Pooled	Toronto, Canada	732	Sept 1995- Aug 1996	16+	3	P, I, O, A	PM _{2.5} (12 mo) PM ₁₀ (3 mo)		Clayton et al. (1999a), Pellizzari et al. (1999)
Pooled	Mexico City	66	1992	< 65	1	P, I, O	PM ₁₀		Santos-Burgoa et al. (1998)
Pooled	Baltimore, MD nonsmoke exposed adults, children, COPD patients	20 Adults 21 Children 15 COPD	Summer 1998, winter 1999	A 75 Ch 9-13 COPD 65	12-days 8-days for children in summer	P, A	PM _{2.5} PM ₁₀	O ₃ , NO ₂ , SO ₂ , VOCs, ED/OC, CO	Sarnat et al. (2001)
Purposeful Studies									
Longitudinal ^d	Wageningen, Netherlands, school children	13	1995	10-12	6	P, A, School	PM _{2.5} , PM ₁₀		Janssen et al. (1999a)
Longitudinal	Amsterdam (Am), Helsinki (Hls), elderly angina or coronary heart disease	41 (Am) 49 (Hls)	Winter 1998, spring 1999	50-84	22 (Am) 27 (Hls)	P, I, O	PM _{2.5}		Janssen et al. (2000)
Longitudinal	Baltimore, elderly healthy and COPD	21	July-Aug 1998	72-93	5-22	P, I, O, A	PM _{2.5} , PM ₁₀	CO, O ₃ , NO ₂ , SO ₂	Williams et al. (2000a,b)
Longitudinal	Fresno I Fresno II (elderly)	516	Feb 1999, Apr-May 1999	60+	24 24	P, I, O, A P, I, O, A	PM _{2.5} , PM ₁₀ PM _{2.5} , PM ₁₀	CO, O ₃	Evans et al. (2000)

TABLE 5-4 (cont'd). SUMMARY OF RECENT PM PERSONAL EXPOSURE STUDIES

Study Design	Study Location and Population	No. of Subjects	Study Period	Age (Years)	Days per Subject	PM Exposure ^a Metrics	PM Size Measured ^b	Co-pollutant Metrics	Reference
Purposeful Studies (cont'd)									
Longitudinal	Los Angeles, elderly COPD subjects	30	Summer/fall 1996	56-83	4	P, I, O	PM _{2.5}		Linn et al. (1999)
Longitudinal	Boston, COPD subjects	18	Winter 1996-1997, summer 1996		12	P, I, O, A	PM _{2.5} , PM ₁₀		Rojas-Bracho et al. (2000)
Longitudinal	Vancouver, British Columbia, COPD	16	April-Sept 1998	54-86	7	P, A	PM _{2.5} , PM ₁₀		Ebelt et al. (2000)
Longitudinal	Amsterdam and Wageningen, Neth., school children	45	1994-1995	10-12	4-8	P, A, School	PM ₁₀		Janssen et al. (1997)
Longitudinal	Amsterdam, adults	37	1994	51-70	5-8	P, I, A	PM ₁₀		Janssen et al. (1998a)
Longitudinal	Baltimore, elderly	15	Summer 1998, spring 1999	75 ± 6.8	12	P	PM _{2.5} , PM ₁₀	O ₃ , NO ₂ , SO ₂ , VOCs	Sarnat et al. (2000)
Longitudinal	Baltimore, elderly, COPD, children	56	Summer 1998, winter 1999	Adults: 75 ± 6.8 Children: 9-13 COPD: 65 ± 6.6	12	P, I, O, A	PM _{2.5}	O ₃ , NO ₂ , SO ₂ , CO, EC, OC, VOC	Sarnat et al. (2000)
Longitudinal	Tokyo, Japan, elderly housewives	18	1992		3	P, I, O, A	SPM	NO ₂	Tamura et al. (1996a)
Longitudinal	Osaka, Japan	26	Fall 1990-1995		Multiple days	P, I, O	PM ₂ , PM ₂₋₁₀ , PM ₋₁₀		Tamura et al. (1996b)
Pooled	Milan, Italy, office workers	100	Spring/summer and winter, year not stated		1	P, Home, Office, Commuting	PM ₁₀	NO ₂ , CO, VOCs	Carrer et al. (1998)
Pooled	Banská Bystrica, Slovakia	49	1997-1998	15-59	1	P, I, O, A	PM ₁₀ , PM _{2.5}	SO ₄ ²⁻ , nicotine	Brauer et al. (2000)
Longitudinal	Wageningen, NL	13	Mar-June 1995	12-14	5-8	P, A, I at school	PM _{2.5} , PM ₁₀	None	Janssen et al. (1999a)
Longitudinal μE diary	Mpala, Kenya	252	1996-1998	5-75	2 years	I	Undefined Optical MIE	CO	Ezzati and Kammen (2001)

TABLE 5-4 (cont'd). SUMMARY OF RECENT PM PERSONAL EXPOSURE STUDIES

Study Design	Study Location and Population	No. of Subjects	Study Period	Age (Years)	Days per Subject	PM Exposure ^a Metrics	PM Size Measured ^b	Co-pollutant Metrics	Reference
Longitudinal	London, UK	10	1997	9-11	5 day/season 3 seasons	P, I, O	PM _{2.5} , PM ₁₀		Wheeler et al. (2000)
Pooled	Zurich, CH	10	1998	Adults	12 h/day for 3 days	P, I, O	Pollen		Riediker et al. (2000)
Pooled	Minneapolis/St. Paul, MN Volunteers	32	Spring, summer, fall 1999	24-64	2-15	P, I, O	PM _{2.5}		Adgate et al. (2002)
Pooled	Birmingham, UK - healthy adults, children and susceptibles	11 healthy adults, 18 susceptible	Season and year not given	Adults, Adults > 65 Child < 10	Adults and children 10, susceptibles 5, daytime only	P, I, O	PM ₁₀	CO, NO ₂	Harrison et al. (2002)
Pooled	Santiago, Chile children	8 in 1998 20 in 1999	Winters 1998 & 1999	10-12	5	P, I, O	PM _{2.5} , PM ₁₀	NO ₂ , O ₃	Rojas-Bracho et al. (2002)
Pooled	Copenhagen, DK non-smoking students	68 subjects	Winter 1999, spring, summer, fall 2000	20-33	2	P, A	PM _{2.5} , BS (from PM _{2.5} filter)		Sørensen et al. (2003)

^a All based on gravimetric measurements.

^b P = personal, I = indoors, O = outdoors, A = ambient.

^c Pooled; data from many subjects with only a few days per subject.

^d Longitudinal; one subject measured for many days.

Recent longitudinal exposure studies have focused on potentially susceptible subpopulations such as the young and elderly with preexisting respiratory and heart diseases (hypertension, COPD, and congestive heart disease). This is in keeping with epidemiological studies that indicate mortality associated with high levels of ambient PM_{2.5} is greatest for elderly people with cardiopulmonary disease (U.S. Environmental Protection Agency, 1996). Longitudinal studies were conducted in the Netherlands by Janssen (1998) and Janssen et al. (1997, 1998a,b, 1999b,c) on purposefully selected samples of adults (50 to 70 years old) and children (10 to 12 years old). School children have also been studied in Chile (Rojas-Bracho et al., 2002). Several additional studies have focused on nonsmoking elderly populations in Amsterdam and Helsinki (Janssen et al., 2000), Tokyo (Tamura et al., 1996a), Baltimore, MD (Liao et al., 1999; Williams et al., 2000a,b,c), and Fresno, CA (Evans et al., 2000). These cohorts were selected because of the low incidence of indoor sources of PM (such as combustion or cooking). This should allow an examination of the relationship between personal and ambient PM concentrations without the large influences caused by smoking, cooking, and other indoor particle-generating activities. The EPA has a research program focused on understanding PM exposure characteristics and relationships. Within the program, longitudinal studies are being conducted on elderly participants with underlying heart and lung disease (COPD, patients with cardiac defibrillator, and myocardial infarction), an elderly environmental justice cohort, and asthmatics. These studies are being conducted in several cities throughout the United States and over several seasons (Rodes et al., 2001; Conner et al., 2001; Landis et al., 2001; Rea et al., 2001).

A series of studies by Phillips et al. (1994, 1996, 1997a,b, 1998a,b, 1999) examined personal ETS exposure in several European cities. Participants varied by age and occupation. Respirable suspended particulate (RSP) concentrations were reported. These studies are not included in Table 5-4 because of their focus on ETS exposure (which is not the focus of this chapter). A small personal exposure study in Zurich, Switzerland was reported by Monn et al. (1997) for PM₁₀. This study also is not listed in Table 5-4 because indoor and outdoor measurements were not taken simultaneously with the personal measurements and other details of the study were not published.

5.3.2.2 Microenvironmental Data

Usually, personal PM monitoring is conducted using integrated measurements over a 12- or 24-h period. As such, total PM exposure estimates based on PEM measurements do not capture data from individual microenvironments. Recent studies have examined PM concentrations in various microenvironments using a number of different types of instruments ranging from filter-based to continuous particle monitors. Details on the instruments used, measurements collected, and findings of these studies according to microenvironment (residential indoor, nonresidential indoor, and traffic-related) are summarized in Table 5-5. Those studies which collected microenvironmental data as part of a personal exposure monitoring study are summarized in Table 5-4 above. In general, the studies listed in Table 5-5 are relatively small, purposeful studies designed to provide specific data on the factors that affect microenvironmental concentration of PM from both ambient and nonambient sources.

Recently published studies have used various types of continuous monitors to examine particle concentrations in specific microenvironments and resulting from specific activities. Continuous particle monitors such as the scanning mobility particle sizer (SMPS), aerodynamic particle sizer (APS), and Climet have been used to measure particle size distributions in residential microenvironments (Abt et al., 2000a; Long et al., 2000, 2001a; Wallace et al., 1997; Wallace, 2000a; McBride et al., 1999; Vette et al., 2001; Wallace and Howard-Reed, 2002). These studies have been able to assess the infiltration factor for ambient particles to indoor microenvironments, as well as penetration factors and deposition rates. Continuous instruments are also a valuable tool for assessing the impact of particle resuspension caused by human activity. A semi-quantitative estimate of PM exposure can be obtained using personal nephelometers that measure PM using light-scattering techniques. Recent PM exposure studies have used condensation nuclei counters (1-s averaging time) and personal nephelometers (1-min averaging time) to measure PM continuously (Howard-Reed et al., 2000; Quintana et al., 2000; Magari et al., 2002; Lanki et al., 2002) in various microenvironments. These data have been used to identify the most important ambient and nonambient sources of PM, to provide an estimate of source strength, and to compare modeled time activity data and PEM 24-h mass data to nephelometer measurements (Rea et al., 2001).

TABLE 5-5. SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Residential Indoor: Nonsmoking Homes					
Abt et al. (2000a) Boston, MA	2 homes, 2 seasons, 6 days	SMPS APS		Detailed traces of PM in various size classes for different air exchange rates (< 1h ⁻¹ to > 2 h ⁻¹).	Major indoor sources of PM: cooking, cleaning, human activity.
Long et al. (2000a) Boston, MA	9 homes, 2 seasons	SMPS APS	0.02-10 µm	Continuous PM distributions and size distributions obtained for indoor and outdoor air using SMPS and APS monitors.	Sources of fine particles: cooking and outdoor particles. Sources of coarse particles: cooking, cleaning, indoor activities. 50% of particles generated by indoor events were ultrafine particles by volume.
Anuszewski et al. (1998) Seattle, WA	9 homes, 18 days	Nephelometer (Radiance)	PM ₁	Simultaneous indoor and outdoor PM measured continuously; 1-h avg time, I/O = 0.98; air exchange rate: 0.7-1.7 h ⁻¹ .	Homes contained asthmatic children, heavy wood burning. Dominant source of fine particles was outdoor air.
Leaderer et al. (1999) Southwest VA	58 homes, summer		PM ₁₀ PM _{2.5}	24 h mean: Regional air 26.0 ± 11.5 µg/m ³ (n = 47); Outdoor homes 28.0 ± 17.7 µg/m ³ (n = 43); Indoor w/ AC 28.9 ± 18.7 µg/m ³ (n = 49); Indoor w/o AC 33.3 ± 14.2 µg/m ³ (n = 8). 24 h mean: Regional air 20.2 ± 9.9 µg/m ³ (n = 50); Outdoor homes 21.8 ± 14.8 µg/m ³ (n = 43); Indoor w/ AC 18.7 ± 13.2 µg/m ³ (n = 49); Indoor w/o AC 21.1 ± 7.5 µg/m ³ (n = 9).	Epidemiologic study of maternal and infant health effects associated with indoor air pollution. Indoor PM concentrations were lower for homes with air conditioning (AC) than non-air-conditioned homes.
Wallace et al. (1997); Wallace (2000b) Reston, VA	1 home, 4 years	SMPS Climet PAHs Black carbon	6 size bins; 100 size channels 0.01-0.4 µm	Time activity data, whole-house air exchange rates. Continuous carbon monoxide; descriptive data for monitored pollutants; size profiles for six indoor particle sources.	0.3- to 0.5-µm particles linked to outdoor concentrations, frying, broiling; 0.5- to 2.5-µm particles related to cooking events; > 2.5-µm particles influenced by physical movement.
Howard-Reed et al. (2000) Fresno, CA Baltimore, MD	15 participants	Nephelometer (personal MIE) PEM	0.1-10 µm PM _{2.5}	Continuous (15-min avg) PM and time activity data; 24-h PM mass; participants from Baltimore and Fresno PM panel studies. Descriptive statistics from each study for five microenvironments.	Time-series plots of personal nephelometer data showed that each participant's PM exposure consisted of a series of short-term peaks, imposed on a background caused by ambient PM concentrations.

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Residential Indoor: Nonsmoking Homes (cont'd)					
Rea et al. (2001) Baltimore, MD Fresno, CA	15 participants	Nephelometer (personal MIE) PEM	0.1-10 µm PM _{2.5} and PM ₁₀	54 ± 31% of average daily PM _{2.5} exposure occurred indoor residences, where participants spent 83 ± 10% of their time. A significant portion of PM _{2.5} exposure occurred where participants spent 4-13% of their time.	Continuous (15-min avg) PM and time activity data; 24-h PM mass; modeled PM mass and time activity data to apportion time spent in a location. Good comparison with nephelometer mass (6-20%).
Quintana et al. (2000) San Diego, CA	Asthmatic children indoor and outdoor 9 homes	Nephelometer (personal MIE) Harvard impactors TEOM	0.1-10 µm PM _{2.5} and PM ₁₀	Nephelometer correlates best with PM _{2.5} : vs. Indoor PM _{2.5} r = 0.66, vs. indoor PM ₁₀ r = 0.13, vs. outdoor PM _{2.5} r = 0.42, vs. outdoor PM ₁₀ r = 0.20.	Indoor and outdoor measurements collected using passive, active, and active heated nephelometers for comparison to PM mass measurements.
Chang et al. (2000) Baltimore, MD	1 person performing predetermined activities	“Roll-around” monitor (RAS) (PM _{2.5} , CO, VOC, O ₃ , NO ₂ , SO ₂)	PM _{2.5}	One-hour personal O ₃ exposures were significantly lower in indoor than outdoor microenvironments. One-hour personal CO exposures were highest in vehicles. Personal and ambient PM _{2.5} correlations were strongest for outdoor microenvironments and those with high air exchange rates (i.e., vehicles).	One-hour personal exposures measured simultaneously. Personal and ambient concentrations were compared.
Liroy et al. (1999)	10 vacuum cleaners		0.3-0.5 µm	Vacuum cleaners ranged in collection efficiency from 29-99%. Substantial fine particle emissions from motors with emission rates from 0.028-128.8 µg/min.	
Ezzati and Kammen (2001) Mpala, Kenya	55 Native huts 2-years	MiniRam (MIE)	Not specified. Optical device detects particles 1-10 µm, but it is not PM ₁₀	Measured PM surrounding wood fires in unvented huts. PM measures were up to 8000 µg/m ³ , but uncalibrated against wood smoke.	Exposures were related to ARI.
Chao and Tung (2001) Hong Kong	5 unoccupied homes measured indoors and outdoors, along with air exchange rates	DustTrak (TSI)	PM _{2.5} real time, calibrated against an Andersen Mark II	In the limit as air exchange goes to zero, there appears to be a residual source, perhaps from drafts or thermal effects. Above air exchange rates of 4.5/hr penetration goes to 1, but indoor turbulence resuspends previously settled PM _{2.5} .	Developed an excellent model for ambient PM infiltration in the absence of anthropogenic indoor sources.

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Residential Indoor: Nonsmoking Homes (cont'd)					
Fischer et al. (2000) Amsterdam, NL	Measured traffic related differences of PM and VOCs, indoor/outdoor in 18 paired homes at varying distances from traffic	Harvard Impactors	PM _{2.5} and PM ₁₀ . EC was measured by reflectance of the PM _{2.5} filters. PAH also measured as indicator of diesel traffic.	Outdoor PM ₁₀ and PM _{2.5} were approximately 15-20% higher at higher traffic streets than at the quiet streets on the same days. However, much larger differences were found for PAH and EC which are traffic specific.	"This [study] supports the use [of] traffic related pollution mapping as an exposure proxy in large-scale epidemiologic studies into health effects of motorized traffic emissions."
Kingham et al. (2000) Huddersfield, UK	Measured PM at ten homes of non-smokers, < 50 m and > 300 m from traffic	Harvard Impactors	PM _{2.5} and PM ₁₀ and PAH. EC measured by filter reflectance.	Median ratio = 1 (no indoor combustion sources).	Found an absence of a spatial gradient, perhaps due to wind direction effects (e.g., sometimes upwind and sometimes downwind of traffic).
Morawska et al. (2001) Brisbane, Australia	Measured PM indoors and outdoors at 16 homes while residents were absent. Air exchange rate estimated, not measured	Scanning mobility particle sizer, aerodynamic particle sizer, and a TSI Dust Trak	Submicron PM, supramicron PM, PM _{2.5}	For supra- and submicron particles, indoor = outdoor for normal ventilation conditions of > 2 air changes/hour.	Average outdoor PM concentrations are good estimates of average indoor concentrations of PM of ambient origin for air exchange rates of > 0.5/h.
Abraham et al. (2002)	Measured ultrafine PM in various home and traffic microenvironments	TSI 8525	< 1 µm optical diameter	1-second readings collected while noting presence of sources.	The TSI 8525 is a useful instrument for screening UFPM in microenvironments.
Geller et al. (2002)	Comparison of indoor and outdoor PM in homes in Coachella Valley, CA	USC Personal PM 5 Lpm	PM _{2.5} and PM _{10-2.5}	I2.5 = 4.3 + 0.74 O I10-2.5 = 3 + 0.3 O I EC = 0.84 O EC I OC > O OC.	Thirteen volunteers had two to four 23-h measurements I and O analyzed for elemental carbon (EC) and organic carbon (OC). Some unoccupied homes measured.
Gotschi et al. (2002)	Measured Black Smoke (BS) by reflection from PM _{2.5} filters for 4 European cities, indoor and outdoor.	XPOLIS PM monitor and EEL 43 reflectometer.	PM _{2.5}	PM measured I and O for two nights while subject was home (5 pm - 8 am). 183 sampled PM _{2.5} and BS were compared for these samples.	BS is a supplementary measurement that can be made on PM _{2.5} filters as an indication of EC.
Sarnat et al. (2002)	Measured PM _{2.5} for sulfate analysis in and out of 6 Boston homes	Harvard Impactors with SMPS and APS particle counters	PM _{2.5}	The ratio of I/O PM _{2.5} , sulfate and size sub-fractions are developed as a function of season and air exchange rate.	Sulfur is primarily of outdoor origin and can be used to track ambient PM of similar (0.06-0.5 µm) AD, but different relations exist for ultrafine PM and coarse-mode PM < 2.5 µm AD.
Wallace and Howard-Reed (2002)	Measured ultrafine, fine and coarse PM in one Reston, VA tri-level home for 18 months with air exchange and meteorologic data	TSI 3071 SMPS TSI 3320 APS Climet 500-I MIE pDR-1000	10 nm to > 10 µm	Analysis of particle counts as a function of air exchange rate and meteorological variables.	Wind speed has little influence on AER. The home sometimes acts as 1 compartment and sometimes as multiple compartments.

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Residential Indoor: Other Home Types					
Brauer et al. (1996) Mexico	22 rural Mexican homes (smoking and nonsmoking)	Inertial impactor Radiance nephelometer	PM ₁₀ PM _{2.5}	Indoor PM _{2.5} : 132-555 µg/m ³ , PM ₁₀ : 282-768 µg/m ³ . Outdoor PM _{2.5} : 37 µg/m ³ . PM ₁₀ : 68 µg/m ³ ; I/O PM _{2.5} : 1.8-12.4; PM ₁₀ : 4.7-10.0.	Variety of cooking fuels used. Nephelometer data were highly correlated with PM _{2.5} and PM ₁₀ indoors (r = ~0.87-0.95).
Jenkins et al. (1996 a,b) 16 U.S. Cities	Smoking and nonsmoking homes	Fluoropore membrane filters	Particle phase ETS markers	Mean PM _{3.5} concentrations were 17-20 µg/m ³ in smoking homes over nonsmoking homes.	
McBride et al. (1999) NA	Combustion source (incense) and walking (1 room, carpeted)	Met-One laser particle counter		Ratios of particle counts at 1.0 and 5.7 m from the combustion source/activity were obtained.	Proximity to source may help explain the existence of a personal cloud.
Vette et al. (2001) Fresno, CA	Detached semioccupied residence	SMPS LASX	0.01-2.5 µm	Temporal relationships between indoor and outdoor aerosol concentrations evaluated; penetration factors and deposition rates estimated. Fresno panel study empty residence.	Diurnally variable indoor/outdoor aerosol concentration ratios because of resuspension from daytime activities. Penetration factors ranged from 0.5 to 0.9.
Douce et al. (2001)	ETS measured by 4 methods in smokers homes and offices	37 mm filter and XAD-4 resin backup	PM ₅	Samples extracted and analyzed for UV absorption, fluorescence, solanesol, and scopoletin.	Solanesol is best method of 4, but needs study of solanesol stability on filter.
Nonresidential Microenvironments					
Bohadana et al. (2000)	Manufacturing plant, woodworkers		Not given	443 personal time-weighted average occupations samples of airborne dust.	
Donham et al. (2000) San Francisco, CA	34 poultry workers	NIOSH Method 0600 monitors probed respirators	PM ₅	Total dust sampled indoor respiratory masks. Personal monitoring: 630 ± 980 µg/m ³ (n = 210) ranging from 10-7730 µg/m ³ .	Respirable dust constituted about 10% of total dust measured.
Klepeis et al. (1996) San Francisco, CA	Airport lounge, ETS	TSI 8510 piezobalance	PM _{3.5}	Estimated cigarette emission rate of 1.43 mg/min/cigarette.	Personal exposures to ETS can be modeled in these types of microenvironments.
Nieuwenhuijsen et al. (1999)	Agricultural activities		PM ₄	Average respirable fraction: 4.5 mg/m ³ .	
Teschke et al. (1999)	Wood production, wood finishing, wood construction workers		PM ≈50	1,632 observations from 1979-1997. Arithmetic mean exposure: 7.93 mg/m ³ . Geometric mean exposure: 1.86 mg/m ³ .	

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (μm)	Summary of Measurements	Notes
Nonresidential Microenvironments (cont'd)					
Baek et al. (1997) Korea	Indoor and outdoor smoking restaurants		PM _{3,5}	Indoor concentrations: 33-475 $\mu\text{g}/\text{m}^3$ Outdoor concentrations: 12-172 $\mu\text{g}/\text{m}^3$ I/O: 2.4.	No significant correlation between indoor and outdoor measurements.
Ott et al. (1996) California	Bar before and after smoking prohibited	Piezobalance	PM _{3,5}	Smoking permitted: indoor 26.3-182 $\mu\text{g}/\text{m}^3$; outdoor < 5-67 $\mu\text{g}/\text{m}^3$ Smoking prohibited: indoor 4-82 $\mu\text{g}/\text{m}^3$; outdoor 2-67 $\mu\text{g}/\text{m}^3$.	I/O nonsmoking: 2.2. I/O smoking: 3.4.
Houseman et al. (2002) Boston, MA	Indoor and outdoor restaurants, stores	TSI DustTrak		Indoor restaurants: 14-278 $\mu\text{g}/\text{m}^3$ Outdoor restaurants: 7-281 $\mu\text{g}/\text{m}^3$ Indoor stores: 12-206 $\mu\text{g}/\text{m}^3$ Outdoor stores: 7-281 $\mu\text{g}/\text{m}^3$.	Avg I/O for restaurants: 2.3. Not known if the restaurants allowed smoking. In stores, indoor and outdoor measurements were correlated, avg I/O: 0.83.
Brauer and Mannetje (1998) Vancouver, BC	Indoor restaurants, various smoking policies		PM _{2,5} PM ₁₀	Nonsmoking: PM _{2,5} 7-65 $\mu\text{g}/\text{m}^3$, PM ₁₀ < 10-74 $\mu\text{g}/\text{m}^3$ Restricted smoking (> 40% nonsmoking) PM _{2,5} 11-163 $\mu\text{g}/\text{m}^3$; PM ₁₀ 24-89 $\mu\text{g}/\text{m}^3$ Unrestricted smoking: PM _{2,5} 47-253 $\mu\text{g}/\text{m}^3$; PM ₁₀ 51-268 $\mu\text{g}/\text{m}^3$.	
Lee and Chang (1999) Hong Kong	Indoor and outdoor 5 classrooms		PM ₁₀	Indoor PM ₁₀ : 30-470 $\mu\text{g}/\text{m}^3$ Outdoor PM ₁₀ : 20-617 $\mu\text{g}/\text{m}^3$	
Chan (2002) Hong Kong	A Hong Kong Office monitored indoor and outdoor	TEOM 1400a	PM _{2,5}	Daily 9 a.m - 6 p.m./weekday for 9 months.	Temp, RH and solar intensity influence In/Out. Wind speed has no effect.
Levy et al. (2002) Boston, MA	Arbitrary choice of one library, coffee shop, urban shopping mall, food court, apartment, hospital, subway, diesel bus in Boston, MA	TSI DustTrak 8520 TSI P-Trak 8525	PM counter calibrated to PM _{2,5} < 1 μm OD	Measured inside and outside the various microenvironments tested, weekdays morning and afternoon. At least 3 visits, several 10-min averages.	10-min outdoor averages are poor predictors of indoor values.

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Traffic-Related Microenvironments (TRMs)					
Praml and Schierl (2000) Munich, Germany	Trams and buses, rural and urban	Continuous millipore polycarbonate filter	PM ₁₀	n = 201 4-h trips, mean concentration 155 µg/m ³ range: 13-686 µg/m ³ I/O: 2.8.	Tram > circular bus route > radial bus route. Day > night.
Monn et al. (1997) Switzerland	Spatial scale from a city street	Harvard microenvironment monitor	PM ₁₀	48- or 72-h avg times; horizontal distance from street: 0, 15, 50, and 80 m; vertical distance from street: 20 m. Mean PM ₁₀ 27.3 ± 3.0 µg/m ³ .	No vertical gradient (0-20 m) and horizontal gradient (0-80 m) in distance from road, each about 13%. No significant differences between wet and dry periods.
Rodes et al. (1998) Saevanenso, Los Angeles, CA	In-vehicle, various road types, 2-h trips		PM _{2.5} PM ₁₀	Vehicles in front of the monitored vehicle accounted for most of the in-vehicle commuting exposure; average I/O: 0.6-0.8 h ⁻¹ for PM _{2.5} ; carpool lane concentrations were 30-60% lower than noncarpool lane concentrations.	Air exchange rates measured at various ventilation settings and speeds. Monitoring vehicle followed a diesel bus or truck.
Roorda-Knape et al. (1998) van Vliet et al. (1997) Netherlands	Gradient in distance from roadway	Harvard impactor	PM _{2.5} PM ₁₀ Black smoke	PM monitoring at 50, 100, 150, and 300 m from roadway; 1-week avg time.	No concentration gradient with increasing distance from the roadways for PM _{2.5} and PM ₁₀ ; concentration gradient did exist for black smoke, also found an effect with wind direction.
Houseman et al. (2002) Boston, MA	Indoor and outdoor vehicles buses, subways	TSI DustTrak	PM ₁₀	Vehicle concentrations ranged from 33-170 µg/m ³ . Outdoor vehicle concentrations ranged from 40-144 µg/m ³ . Bus concentrations: 17-268 µg/m ³ ; outdoor 10-203 µg/m ³ . Subway: 28-174 µg/m ³ ; outdoor 8-203 µg/m ³ .	The average in-vehicle to outdoor ratio was 0.99. Average I/O: 3; subway values were correlated with outdoor concentrations.
Brauer et al. (1999) Vancouver, BC	Commuting environments	APC-1000		PM < 5: greatest concentrations by combustion powered vehicles. PM > 5: greatest concentrations by bicycling and buses.	
Janssen et al. (1997) Netherlands	Background and roadway		PM _{2.5} PM ₁₀	PM _{2.5} background: 21-35 µg/m ³ ; roadway 23-43 µg/m ³ . PM ₁₀ background: 13-32 and 29-62 µg/m ³ ; roadway 16-56 and 30-75 µg/m ³ .	Average roadway/background ratio: 3 for PM _{2.5} and PM ₁₀ . Average increase in concentration at the roadway 7.2-12.7 µg/m ³ .

TABLE 5-5 (cont'd). SUMMARY OF RECENT MICROENVIRONMENTAL PM MEASUREMENT STUDIES

Reference	Study Description	Instrument(s)	Size Fraction (µm)	Summary of Measurements	Notes
Traffic-Related Microenvironments (TRMs) (cont'd)					
Adams et al. (2001) London, UK	PM by volunteers in TRM in London, UK	16 L/min personal monitor. Porous foam size selector	PM _{2.5}	Volunteers rode/cycled along fixed routes repetitively.	Personal exposures were generally double those at fixed-site ambient station. Subway exposures are maximal.
Alm et al. (1999) Kuopio, Finland	9-km commuter route, rush hours 1/mo	Climet	6 channels	Windows closed, vents open air exchange rate 36-47 h ⁻¹ .	Morning commutes were generally higher than afternoon commutes; relationships determined between PM and wind speed and vehicle speed.
Chan et al. (2002) Hong Kong	PM measured in TRM in Hong Kong	TSI 8520 DustTrac Calibrated to Partisol PM _{2.5} HiVol PM ₁₀	PM _{2.5} and PM ₁₀	Repetitive sampling over 8 fixed routes.	Highly variable by mode. Tram exposures are maximal.
Hoek et al. (2001) Hoek et al. (2002)	BS in Netherlands interpolated to outdoor locations relative to traffic at subject's homes	Black Smoke measured from filters	Not Recorded	GIS used to interpolate background and regional ambient BS with added increment for distance to traffic < 50 m and < 100 m.	May be useful technique. Estimates not validated with BS measurements at interpolated loci.
Jinsart et al. (2002) Bangkok	PM Exposure of traffic police at intersection post in Bangkok, Thailand	Sibata personal single nozzle 2.5 L/min	PM _{2.5} and PM ₁₀	Sampled 12-h while on duty at post at intersection.	Exposures of both PM _{2.5} and PM ₁₀ higher than ambient PM measured at station with β gauge. Not comparable to U.S. conditions.
Lena et al. (2002) New York, NY	PM _{2.5} and EC in the Bronx, NY in area of high Diesel truck traffic	PM _{2.5} @ 4 L/min 3 Lpm quartz filter for EC EEL for BS	PM _{2.5}	Sampled 10-h at sidewalk locations while counting cars and trucks for 3-weeks in summer of 1999.	EC is a large component of diesel PM _{2.5} and varies with truck traffic.
Zhu et al. (2002) Los Angeles, CA	Ultrafine near Interstate Highway in Los Angeles, CA with heavy Diesel traffic	CPC TSI 3022A SMPS TSI 3936 BC aethalometer	6 nm-220 nm	200 m upwind and 17, 20, 30, 90, 150, and 300 m downwind.	Ultrafine PM decreased exponentially from the freeway and was equal to upwind at 300 m.

5.3.2.3 Traffic-Related Microenvironments

There has been increasing interest in the possible role of traffic-related pollutants. Distance to roadways has been used as a surrogate for exposure to traffic-related pollutants (Hoek et al., 2001), and this exposure indicator was subsequently used in an epidemiological study (Hoek et al., 2002). A traffic model, using traffic volume, direct exhaust emissions rate, and a reentrainment rate has been used to estimate concentrations of traffic-related emissions at several schools in east Los Angeles (Korenstein and Piazza, 2002). Personal exposure studies have been made in a variety of commuting situations including vehicle traffic (Adams et al., 2001; Chan et al., 2002). Other studies have measured various indicators of traffic near roadways (Lena et al., 2002), inside vehicles in traffic (Abraham et al., 2002), and in several types of traffic-related microenvironments (Levy et al., 2002). Table 5-5 above provides a brief description of these studies, instruments used, measurements made, and key findings.

5.3.2.4 Reanalyses of Previously-Reported Particulate Matter Exposure Data

Papers that have reanalyzed and interpreted the data collected in previous PM exposure studies are summarized in Table 5-6. These reanalyses are directed toward understanding the personal cloud, the variability in total PM exposure, and the personal exposure-to-ambient concentration relationships for PM. Brown and Paxton (1998) determined that the high variability in personal exposure to PM makes the personal-to-ambient PM relationship difficult to predict. Wallace (2000b) used data from a number of studies to test two hypotheses: elderly COPD patients have (1) smaller personal clouds and (2) higher correlations between personal exposure and ambient concentrations compared to healthy elderly, children, and the general population. The analysis by Wallace (2000a) and three subsequent longitudinal studies (Williams 2000a,b,c; Ebelt et al., 2000; Sarnat et al., 2000) supported hypothesis 1 but not hypothesis 2. Özkaynak and Spengler (1996) showed that at least 50% of personal PM_{10} exposure for the general population comes from ambient particles. Wilson and Suh (1997) concluded that fine and coarse particles should be treated as separate classes of pollutants because of differences in characteristics and potential health effects. Wilson et al. (2000) gave a review of what they call the “exposure paradox” and determined that personal PM needs to be divided into different classes according to source type and that correlations between personal and ambient PM will be higher when nonambient sources of PM are removed from the personal PM

TABLE 5-6. PAPERS REPORTING REANALYSES OF PARTICULATE MATTER EXPOSURE STUDIES

Reference	Study Cited	Objectives/Hypotheses	Findings
Wallace (2000a)	PTEAM (Özkaynak et al., 1990; Spengler et al., 1989; Wiener 1988, 1989; Wiener et al., 1990) THEES (Liroy et al., 1990) Amsterdam COPD (Janssen et al., 1997, 1998a) Boston COPD (Rojas-Bracho et al., 2000)	Examines the differences between pooled and longitudinal correlations in personal and ambient (or outdoor) data for PM _{2.5} and PM ₁₀ . Discusses the personal cloud for PM _{2.5} and PM ₁₀ . Hypothesizes that COPD patients have (1) smaller personal clouds (supported) and (2) higher correlations of personal exposure with outdoor concentrations because of reduced mobility (not supported).	Median longitudinal correlation coefficient is much higher than the pooled correlation coefficient for the same data sets. Personal cloud for PM ₁₀ : 3-67 µg/m ³ ; PM _{2.5} 6-27 µg/m ³ . Personal cloud for elderly COPD was much smaller (PM ₁₀ : 6-11 µg/m ³ ; PM _{2.5} ≈ 6 g/m ³) than for other healthy populations (PM ₁₀ : 27-56 µg/m ³ ; PM _{2.5} : 11-27 µg/m ³) of elderly, children, and the general population. However, correlations of personal exposure with ambient concentrations were not higher for elderly COPD than for other groups.
Özkaynak and Spengler (1996)	Dockery and Spengler (1981) PTEAM (Özkaynak et al., 1996a,b) Netherlands (Janssen et al., 1995)	Uses statistical modeling techniques to examine the relationship between ambient PM concentrations and personal exposures. Data analysis involves use of air exchange rates, penetration factors, and June 25, 2004 ratios, as well as examining exposure in various microenvironments (traveling, working, outdoors, indoors) activities (exposure to smoke, cooking), and source strengths.	The important components of personal exposures are received during contact with indoor sources, mainly in homes and work places. Ambient aerosols contribute about 50% or more to the personal PM ₁₀ exposures of the general population. The contribution of ambient aerosols to the total toxicity of inhaled particles is significant.
Brown and Paxton (1998)	THEES (Liroy et al., 1990) PTEAM pilot (Wallace, 1996) Boston and Nashville COPD (Rojas-Bracho et al., 2000); Bahadori et al., 1998)	Cross-sectional and longitudinal regression analysis on data sets.	Individual personal PM exposure is subject to high variability, which makes the personal-to-ambient PM relationship difficult to predict.
Wilson and Suh (1997)	Philadelphia (Burton et al., 1996; Suggs and Burton, 1983) EPA AIRS database	Determines the utility of fine and coarse PM concentrations as indicators of time-series epidemiology with regard to day-to-day variability, area uniformity, and June 25, 2004 PM ratios. Necessary to treat personal exposure to ambient PM and personal exposure to nonambient PM as separate components of total personal PM exposure.	Fine and coarse particles should be considered separate classes of pollutants. Fixed-site ambient fine-particle measurements likely give a reasonable indication of the variability in the concentration of ambient fine particles across the community. Coarse-particle measurements most likely will not.
Wilson et al. (2000)	New Jersey (Liroy et al., 1990) Japan (Tamura et al., 1996a) PTEAM (Clayton et al., 1993; Özkaynak et al., 1996a,b) Netherlands (Janssen, 1998a; Suh et al., 1992)	Synoptic review of the “exposure paradox”: low correlations between personal exposure and ambient PM concentrations in spite of the existence of statistical association between ambient PM and epidemiologic health effects. Uses personal exposure equation, mass balance, regression analysis, and deductive logic.	Personal PM exposure needs to be divided into different classes according to source type: exposure to ambient PM (outdoor and indoors) and exposure to nonambient PM (indoor source and personal activity). Correlations are higher between personal exposure and ambient PM concentrations when PM exposures from nonambient sources are removed.

TABLE 5-6 (cont'd). PAPERS REPORTING REANALYSES OF PARTICULATE MATTER EXPOSURE STUDIES

Reference	Study Cited	Objectives/Hypotheses	Findings
Mage et al. (1999)	Japan (Tamura et al., 1996a) State College (Suh et al., 1995) Netherlands (Janssen et al., 1997, 1998a, 1999a) New Jersey (Lioy et al., 1990) PTEAM (Clayton et al., 1993; Özkaynak et al., 1996a,b)	Examines the influence of nonambient PM on total PM concentrations and how it may confound the outdoor/personal PM relationship. Missing data and outlier values created using an algorithm. Linear regression analysis of subsequent data sets.	Variation in daily personal exposure for subjects with similar lifestyles and no ETS exposure are driven by variations in ambient PM concentrations. Exposure to ambient PM is highly correlated in time with ambient PM concentrations measured at a community site. Indoor PM does not confound the relationship between daily mortality and ambient PM.
Mage (1998)	PTEAM (Clayton et al., 1993; Özkaynak et al., 1993, 1996a,b)	Uses a reduced-form mass-balance model to predict the average fraction of ambient PM to which the average person is exposed.	On average, a person is exposed to > 75% of ambient PM _{2.5} and > 64% of ambient PM ₁₀ measured by the community monitor.
Monn (2001)	Multiple Literature Review	Objective review of literature published since 1996 as an implicit update to the 1996 U.S. EPA PM AQCD. Emphasis on European studies.	“It is important to note that a personal measurement does not <i>a priori</i> provide more valid data than a stationary ambient measurement, i.e. a personal sample in a study investigating effects from outdoor combustion particles is often influenced by sources other than outdoor sources and may thus confound the exposure-effect outcome.” “Despite some lack of correlation between personal (PM ₁₀) and outdoor values, outdoor fine particle concentrations were strongly associated with mortality and morbidity indicating that outdoor sources (e.g. vehicular emissions) emit the toxic entity” (Dockery et al., 1993; Schwartz et al., 1996).
Rotko et al. (2000a)	Jantunen et al. (1998) Carrer et al. (1997) Koistinen et al. (1999)	Compares exposure relationships between the six EXPOLIS European cities (Athens, Basel, Grenoble, Helsinki, Milan, Prague).	Demographic bias exists because women and more-educated individuals are more likely to respond to survey. Socioeconomic bias exists in low SES subjects less likely to participate in diary keeping and exposure monitoring. Weighting is required for inter-city comparisons. Selection bias is not a problem for characterizing physical factors influencing personal exposure.
Rotko et al. (2000b)	Rotko et al. (2000a), Jantunen et al. (1998)	Determines sociodemographic influences of exposure in Helsinki.	Distinct male vs. female differences: males had higher exposures to PM _{2.5} , related to ETS, and a larger variance between sociodemographic groupings. No sociodemographic differences existed in outdoor PM _{2.5} concentrations. Lower occupational status contributed to greater PM _{2.5} exposures than higher (professional) occupational status.
Rodes et al. (2001)	EPA Baltimore and Fresno 1 and 2	Investigates relationships between the different retirement centers and identify most likely factors influencing personal and indoor concentrations.	Mean personal exposure PM _{2.5} was higher than their apartment concentrations. Personal cloud of 3 µg/m ³ for PM _{2.5} was negligible but cloud for PM ₁₀ was 20 µg /m ³ . Indoor PM _{2.5} data were less than ambient concentrations.

concentration. Mage (1998) conducted analysis using the PTEAM data and showed that the average person in PTEAM (Riverside, CA in the fall) was exposed to > 75% of ambient PM_{2.5} and > 64% of ambient PM₁₀. Mage et al. (1999) used an algorithm to fill in missing data and outliers to analyze data sets and showed that variation in daily personal exposures for subjects with similar activity patterns and no ETS exposure are driven by variation in ambient PM concentrations.

5.3.3 Factors Influencing and Key Findings on Particulate Matter Exposures

5.3.3.1 Relationship of Personal/Microenvironmental Particulate Matter with Ambient Particulate Matter

Understanding the relationship between ambient site measurements and personal exposure to PM is important for several reasons. First, it allows us to examine the extent to which ambient measurements for PM and various PM constituents can serve as valid surrogates for exposure to ambient PM or ambient constituents of PM in epidemiological studies. Second, it provides information that may improve surrogate exposure measurements and, hence, increase the power of epidemiologic studies. Finally, because compliance with the NAAQS is based on ambient monitoring, it can be used to understand the effect of regulation on exposures to PM and its constituents and can, therefore, help link the effect of regulations to health outcomes. Many of the studies summarized in Table 5-4 have analyzed this relationship using measurements of personal PM exposures and ambient PM concentrations. Of main interest are the PM concentrations measured in ambient, indoor, and outdoor air; personal exposure measurements; the statistical correlations between measurements; and the attenuation and/or infiltration factors developed for personal exposure and indoor microenvironments. Attenuation and infiltration factors are discussed in Section 5.3.4.3.1. Information on correlation analysis is provided below.

5.3.3.1.1 Types of Correlations

The three types of correlation data that will be discussed in this section are longitudinal, “pooled,” and daily-average correlations. *Longitudinal correlations* are calculated when data from a study includes measurements over multiple days for each subject (longitudinal study design). Longitudinal correlations describe the temporal relationship between daily personal PM exposure or microenvironment concentration and daily ambient PM concentration for each

individual subject. The longitudinal correlation coefficient, r , may differ for each subject. An analysis of the variability in r across subjects can be performed with this type of data. Typically, the median r is reported along with the range across subjects in the study. *Pooled correlations* are calculated when a study involves one or only a few measurements per subject and when different subjects are studied on subsequent days. Pooled correlations combine individual subject/individual day data for the correlation calculation. Pooled correlations describe the relationship between daily personal PM exposure and daily ambient PM concentration across all subjects in the study. For some studies, the multiple days of measurements for each subject were assumed to be independent (after autocorrelation and sensitivity analysis) and combined together in the correlation calculation (Ebelt et al., 2000). *Daily-average correlations* are calculated by averaging exposure across subjects for each day. Daily-average correlations then describe the relationship between the daily average exposure and daily ambient PM concentration. The term cross-sectional is used to refer to both pooled and daily average correlations, so the meaning of this term must be determined from context.

Pooled correlations have been simulated from longitudinal data by using a random-sampling procedure to select a random day from each subject's measurements for use in the correlation. This procedure was repeated many times, and statistics (such as the mean and standard deviation of the pooled correlation coefficient) were reported (Janssen et al., 1997, 1998a, 1999c).

The type of correlation analysis can have a substantial effect on the resultant r value. Mage et al. (1999) mathematically demonstrated that very low correlations between personal exposure and ambient concentrations could be obtained when people with very different nonambient exposures are pooled, even though their individual longitudinal correlations are high. The longitudinal studies conducted by Tamura et al. (1996a) and Janssen et al. (1997, 1998a, 1999c) determined that the longitudinal correlations between personal exposure and ambient PM concentrations were higher than the correlations obtained from a pooled data set. Wallace (2000a) reviewed a number of longitudinal studies and found that the median longitudinal correlation coefficient was higher than the pooled correlation coefficient for the same data (see Tables 1 and 2, Wallace, 2000a).

Mage et al. (1999) examined three longitudinal exposure data sets where several subjects were measured each day. They showed that by averaging daily exposures across subjects, daily-

average correlations could be obtained. These were all higher than the median longitudinal correlations. Williams et al. (2000a,b) and Evans et al. (2000) have also reported higher correlation coefficients for daily-average correlations compared to longitudinal correlations. The higher correlations found between daily-average personal exposures and ambient PM concentrations, as opposed to lower correlations found between individual exposures and ambient PM levels, have been attributed to the statistical process of averaging (Ott et al., 2000). Personal exposures include contributions from nonambient as well as ambient PM concentrations. When several subjects are measured on the same day, the mean variability due to variations in nonambient exposures is reduced due to averaging. Therefore, the correlation between personal exposure and ambient concentrations increases as the number of subjects measured daily increases. Ott et al. (2000), using the theory on which their Random Component Superposition (RCS) model is based, predict expected correlations above 0.9 for the PTEAM study and above 0.70 for the New Jersey study (Lioy et al., 1990) if 25 subjects had been measured daily in each study.

5.3.3.1.2 Correlation Data from Personal Exposure Studies

Measurement data and correlation coefficients for the personal exposure studies described in Section 5.4.2.1 are summarized in Table 5-7. All data are based on mass measurements. The studies are grouped by the type of study design, longitudinal or pooled. For each study in Table 5-7, summary statistics for the total personal PM exposure measurements are presented as well as statistics for residential indoor, residential outdoor, and ambient PM concentrations when available. The correlation coefficients, r , between total personal PM exposures and ambient PM concentrations also are presented and classified as longitudinal or pooled correlations. When reported, p -values for the correlation coefficients are included. Correlation coefficients between personal, indoor, outdoor, and ambient also are reported when available.

5.3.3.1.3 Correlations Between Personal Exposures, Indoor, Outdoor, and Ambient Measurements

Longitudinal and pooled correlations between personal exposure and ambient or outdoor PM concentrations varied considerably between study and study subjects. Most studies report longitudinal correlation coefficients that range from < 0 to ≈ 1 , indicating that an individual's

TABLE 5-7. PERSONAL MONITORING STUDIES FOR PARTICULATE MATTER: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

Size Fraction	Avg. Time	Statistic	Sample Size ¹	Measured Concentration Levels ($\mu\text{g}/\text{m}^3$)				Personal-Ambient ² Correlation Coefficients (r)			Other Correlation Coefficients (r)	
				Personal	Residential Indoor	Residential Outdoor	Ambient	Type ³	Value (Range)	Notes	Type ³	Value (Range)
Longitudinal Studies												
<i>Ebelt et al. (2000) – Vancouver, BC</i>												
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	106	18.2 \pm 14.6 2-91			11.4 \pm 4.1 4-29	Median L P	0.48 (-0.68-0.83) 0.15	n = 16 COPD subjects		
<i>Evans et al. (2000) – Fresno, CA</i>												
PM _{2.5}	24 h	\bar{x} Range	24	13.3 1-24	9.7 4-17	20.5 4-52	21.7 6-37	P	0.414	Fresno-1 study	P _{p-i} P _{p-o}	0.81 ⁴ 0.80 ⁴
PM _{2.5}	24 h	\bar{x} Range	12	11.1 7-16	8.0 4-12	10.1 5-20	8.6 4-16	P	0.844	Fresno-2 study	P _{p-i} P _{p-o}	0.95 ⁴ 0.80 ⁴
<i>Janssen et al. (1997) – Netherlands</i>												
PM ₁₀	24 h	$\bar{x} \pm \text{SD}$ Range	301	105.2 \pm 28.7 57-195			38.5 \pm 5.6 25-56	Median L Median L Median L Mean P Mean P Mean P	0.63 (0.1-0.9) 0.63 0.59 0.28 (0.12) ⁵ 0.45 (0.16) ⁵ 0.20 (0.19) ⁵	n = 45 school children With nonsmoking parents With smoking parents All With nonsmoking parents With smoking parents		
<i>Janssen et al. (1998a) – Netherlands</i>												
PM ₁₀	24 h	$\bar{x} \pm \text{SD}$ Range	262	61.7 \pm 18.3 38-113	35.0 \pm 9.4 19-65		41.5 \pm 4.3 32-50	Median L Median P Median P	0.50 (-0.41-0.92) 0.50 (0.07-0.83) ⁵ 0.34 (-0.09-0.67) ⁵	n = 37 adults No ETS exposure All	Med. L _{p-i} Med. L _{i-a}	0.72 (-0.10-0.98) 0.73 (-0.88-0.95)
<i>Janssen et al. (1999c) – Netherlands</i>												
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	77	28.3 \pm 11.3 19-60			17.1 \pm 2.8 14-22	Median L Median P	0.86 (-0.11-0.99) 0.41 (-0.28-0.93) ⁵	n = 13 school children		
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	55	24.4 \pm 4.9 19-33			17.1 \pm 2.6 15-22	Median L Median P	0.92 0.82 ⁵	With nonsmoking parents		
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	22	37.0 \pm 17.4 21-60			17.1 \pm 3.7 14-21			With smoking parents		

TABLE 5-7 (cont'd). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

Size Fraction	Avg. Time	Statistic	Sample Size ¹	Measured Concentration Levels ($\mu\text{g}/\text{m}^3$)				Personal-Ambient ² Correlation Coefficients (r)			Other Correlation Coefficients (r)		
				Personal	Residential Indoor	Residential Outdoor	Ambient	Type ³	Value (Range)	Notes	Type ³	Value (Range)	
<i>Janssen et al. (2000) – Netherlands</i>													
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	338	24.3 \pm 25.7 9-134	28.6 \pm 41.8 9-239		20.6 \pm 4.0	Median L	0.79 (-0.41-0.98)	n = 36 elderly w/CV disease No ETS exposures	Med.	0.91	
							13-31	Median L	0.85		L _{p-i}	(-0.28-1.0)	
											Med.	0.84	
											L _{i-a}	(-0.00-0.98)	
<i>Janssen et al. (2000) – Finland</i>													
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	336	10.8 \pm 4.4 4-33	11.0 \pm 4.0 3-27		12.6 \pm 2.0	Median L	0.76 (-0.12-0.97)	n = 46 elderly w/CV disease	Med.	0.89	
							10-18				L _{p-i}	(0.14-1.0)	
											Med.	0.70	
											L _{i-a}	(-0.15-0.94)	
<i>Linn et al. (1999) – Los Angeles</i>													
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ Range	60	23.8 \pm 15.1 4-65	23.5 \pm 15.3 4-92	24.8 \pm 14.5 4-63						P _{i-a}	0.26 ⁶
												P _{o-a}	0.47 ⁶
PM ₁₀	24 h	$\bar{x} \pm \text{SD}$ Range	59	34.8 \pm 14.8 5-85	32.6 \pm 15.6 9-105	39.8 \pm 18.3 7-97	33 \pm 15 9-??					P _{i-a}	0.32 ⁶
												P _{o-a}	0.66 ⁶
<i>Rojas-Bracho et al. (2000) – Boston</i>													
PM _{2.5}	12 h	$\bar{x} \pm \text{SD}$ Range	224	21.6 \pm 13.6 1-128	17.5 \pm 14.1 2-73	14.2 \pm 11.2 1-57					n = 17 adults	Med.	0.87 ⁶
												L _{p-i}	0.74 ⁶
												Med.	0.71 ⁶
												L _{p-i}	0.50 ⁶
												L _{i-o}	
PM _{10-2.5}	12 h	$\bar{x} \pm \text{SD}$ Range	222	15.6 \pm 14.6 -11-103	14.5 \pm 9.2 -3-255	8.1 \pm 6.8 -2-64						Med.	0.42 ⁶
												L _{p-i}	0.20 ⁶
												Med.	
												L _{i-o}	

TABLE 5-7 (cont'd). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

Size Fraction	Avg. Time	Statistic	Sample Size ¹	Measured Concentration Levels ($\mu\text{g}/\text{m}^3$)				Personal-Ambient ² Correlation Coefficients (r)			Other Correlation Coefficients (r)	
				Personal	Residential Indoor	Residential Outdoor	Ambient	Type ³	Value (Range)	Notes	Type ³	Value (Range)
<i>Sarnat et al. (2000) – Baltimore</i>												
PM _{2.5}	24 h	$\bar{x} \pm \text{SD}$ $\bar{x} \pm \text{SD}$	3736	26.7 \pm 13.7 18.5 \pm 11.2			25.2 \pm 11.5	Median L	0.76 (-0.21-0.95) ⁷	n = 15 adults, summer n = 15 adults, winter Ventilation: High summer Med. summer Low summer Winter		
							5.6 \pm 49.0	Median L	0.25 (-0.38-0.81) ⁷			
				P	0.89 ⁸							
				P	0.75 ⁸							
							P	0.50 ⁸				
							P	0.44 ⁸				
PM ₁₀	24 h	$\bar{x} \pm \text{SD}$ $\bar{x} \pm \text{SD}$	3736	33.9 \pm 11.7 28.0 \pm 16.5			34.0 \pm 12.8	Median L	0.64 (0.08-0.86) ⁷	Summer Winter		
							7.5 \pm 73.2	Median L	0.53 (-0.79-0.89) ⁷			
PM _{10-2.5}	24 h	$\bar{x} \pm \text{SD}$ $\bar{x} \pm \text{SD}$	3736	7.2 \pm 4.0 9.6 \pm 7.9			8.4 \pm 2.3	Median L	0.11 (-0.60-0.64) ⁷	Summer Winter		
							-1.3 \pm 24.2	Median L	0.32 (-0.48-0.68) ⁷			
<i>Tamura et al. (1996a) – Tokyo</i>												
PM ₁₀	48 h							P	0.83	n = 7 elderly adults		
<i>Williams et al. (2000a,b) – Baltimore</i>												
PM _{2.5}	24 h	\bar{x} Range	23	13.0 7-25	9.4 4-19	22.0 7-52	22.0 8-59	Median L	0.80 (0.38-0.98) ⁶	n = 21 elderly adults		
								P	0.89 ⁴			
											P _{p-i}	0.90 ⁴
											P _{p-o}	0.95 ⁴
										P _{i-o}	0.94 ⁴	
											P _{i-a}	0.87 ⁴
											P _{o-a}	0.96 ⁴
PM ₁₀	24 h	\bar{x} Range	28	11.0 4-23	30.0 13-66	29.9 13-74					P _{i-o}	0.82 ⁴
											P _{i-a}	0.81 ⁴
											P _{o-a}	0.94 ⁴
PM _{10-2.5}	24 h	\bar{x} Range	26	1.0 -3-5	8.0 -2-16	8.0 1-15					P _{i-o}	0.18 ⁴
											P _{i-a}	0.08 ⁴
											P _{o-a}	0.45 ⁴

TABLE 5-7 (cont'd). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

Size Fraction	Avg. Time	Statistic	Sample Size ¹	Measured Concentration Levels ($\mu\text{g}/\text{m}^3$)				Personal-Ambient ² Correlation Coefficients (r)			Other Correlation Coefficients (r)	
				Personal	Residential Indoor	Residential Outdoor	Ambient	Type ³	Value (Range)	Notes	Type ³	Value (Range)
<i>Williams et al. (2000a,b) – Baltimore (cont'd)</i>												
Keeler et al. (2002)		Mean (Std)	20 asthmatic children	68.4 (39.2)	52.2 (30.6)		25.8 (11.8)					
Detroit, MI PM ₁₀ 24-h					34.4 (21.7)		15.6 (8.2)					
Landis et al. (2001)		Mean	10 elderly retirees	12.8	10.2	21.0		Median L	r = 0.82 (0.51-0.92)		Lp-i r = 0.60 (0.41-0.85)	
Baltimore, MD PM _{2.5} 24-h				4.5	4.0	10.2			r = 0.95 (0.74-0.97)		r = 0.95 (0.73-0.97)	
Sulfate24-h												
Pooled Studies												
<i>Bahadori (1998) – Nashville</i>												
PM _{2.5}	12 h	$\bar{x} \pm$ SD Range	30	21.7 \pm 10.5 10-67	15.5 \pm 6.6 5-40	23.4 \pm 6.8 3-61		P	0.09	n = 10 COPD subjects; daytime	P _{p-i} 0.72 P _{i-o} 0.31	
PM ₁₀	12 h	$\bar{x} \pm$ SD Range	30	33.0 \pm 16.9 5-88	21.6 \pm 10.7 9-77	32.5 \pm 8.1 7-76		P	-0.08	n = 10 COPD subjects; daytime	P _{p-i} 0.43 P _{i-o} 0.06	
<i>Pellizzari et al. (1999) – Toronto</i>												
PM _{2.5}	3 d	\bar{x}	922	28.4	21.1	15.1		P	0.23	n = 178; n for indoor, outdoor lower than personal	P _{p-i} 0.79 P _{i-o} 0.33	
PM ₁₀	3 d	\bar{x}	141	67.9	29.8	24.3				No correlations reported		
<i>Oglesby et al. (2000) – EXPOLIS Basel</i>												
PM _{2.5}	48 h	$\bar{x} \pm$ SD	4420	23.7 \pm 17.1 17.5 \pm 13.0		19.0 \pm 11.7 17.7 \pm 7.1		P P	0.07 0.21	All No ETS exposure		
<i>Santos-Burgoa et al. (1998) – Mexico City</i>												
PM ₁₀	24 h	$\bar{x} \pm$ SD	66	97 \pm 44	99 \pm 50			P	0.26		P _{p-i} 0.47 P _{i-a} 0.23	
<i>Tamura et al. (1996b) – Osaka</i>												
PM ₂	48 h							P	0.74			
PM ₁₀	48 h							P	0.67			

TABLE 5-7 (cont'd). PERSONAL MONITORING STUDIES FOR PM: MEASURED CONCENTRATIONS AND CORRELATION COEFFICIENTS

Size Fraction	Avg. Time	Statistic	Sample Size ¹	Measured Concentration Levels (µg/m ³)				Personal-Ambient ² Correlation Coefficients (r)			Other Correlation Coefficients (r)	
				Personal	Residential Indoor	Residential Outdoor	Ambient	Type ³	Value (Range)	Notes	Type ³	Value (Range)
<i>Pellizzari et al. (2001) – Indianapolis</i>												
PM _{2.5}	72 h	Median	250	23	18	18	18	P	0.102	Between the Logarithms of concentrations	P _{p-o}	0.138
											P _{p-i}	0.923
<i>Brauer et al. (2000) Banska Bystrica</i>												
PM _{2.5} PM ₁₀	24 h	Mean	PM ₁₀ summer PM ₁₀ winter PM _{2.5} summer PM _{2.5} winter SO ₄ winter	1.2212e + 10	796655535		354522326	P	PM ₁₀ R ² = 0.17	Multivariate model with PM ₁₀ and nicotine	P _{p-i}	R ² = 0.15
											P _{p-o}	R ² = 0.23
<i>Kousa et al. (2002)</i>												
Helsinki + 3 cities PM _{2.5} 48 h								P	0.69	Leisure time, non-ETS exposed, Helsinki, log-transformed	P _{p-o} P _{p-i}	0.65 0.83

Abbreviations used:

Avg. = Averaging (time)
 Conc. = Concentration
 CV = Cardiovascular
 d = Day
 ETS = Environmental tobacco smoke

h = Hour
 i-a = Indoor-ambient correlation
 i-o = Indoor-outdoor correlation
 L = Longitudinal correlation
 Med. = Median
 o-a = Outdoor-ambient correlation

P = Pooled correlation
 p-i = Personal-indoor correlation
 p-o = Personal-outdoor correlation
 SD = Standard deviation
 Stat. = Statistic
 \bar{x} = Mean

Notes:

- ¹ Sample size is for personal concentrations; indoor, outdoor and ambient sample sizes may differ.
- ² Correlation coefficient is for personal-residential outdoor if no ambient concentration data reported.
- ³ See text for description of types of correlations.
- ⁴ Daily-averaged correlation (values for individual subjects averaged for each day).
- ⁵ Pooled correlations estimated using a Monte Carlo sampling procedure, n = 1000. If mean P is shown, then SD given; if median P is shown, then range is given.
- ⁶ Obtained from a regression equation; $r = \sqrt{R^2}$.
- ⁷ Spearman rank correlations.
- ⁸ Calculated, $r = \sqrt{R^2}$, from R² from a mixed model regression.

activities and residence type may have a significant effect on total personal exposure to PM. General population studies tend to show lower correlations because of the higher variation in the levels of PM-generating activities. In contrast, the absence of indoor sources for the populations in several of the longitudinal studies resulted in high correlations between personal exposure and ambient PM within subjects over time for these populations. But even for these studies, correlations varied by individual depending on their activities and the microenvironments that they occupied.

Probability Studies

In the Toronto study (Pellizzari et al., 1999), pooled correlations were derived for personal, indoor, outdoor, and fixed-site ambient measurements. This study involved a probability sample of 732 participants who represented the general population of people 16 years and older. The study included between 185 and 203 monitoring periods with usable PM data for personal, residential indoor, and outdoor measurements. For PM₁₀ measurements, the mean concentrations were 67.9 µg/m³ for personal, 29.8 µg/m³ for indoor, and 24.3 µg/m³ for outdoor air samples. For PM_{2.5}, the mean concentrations were 28.4 µg/m³ for personal, 21.1 µg/m³ for indoor air, and 15.1 µg/m³ for outdoor air samples. A low, but significant, correlation ($r = 0.23$, $p < 0.01$) was reported between personal exposure and ambient measurements. The correlations between indoor concentrations and the various outdoor measurements of PM_{2.5} ranged from 0.21 to 0.33. The highest correlations were for outdoor measurements at the residences with the ambient measurements made at the roof site (0.88) and the other fixed site (0.82). Pellizzari et al. (1999) state that much of the difference among the data for personal/indoor/outdoor PM:

“... can be attributed to tobacco smoking, since all variables reflecting smoking ... were found to be highly correlated with the personal (and indoor) particulate matter levels, relative to other variables that were measured ... none of the outdoor concentration data types (residential or otherwise) can adequately predict personal exposures to particulate matter.” (p. 729)

Using a Random Component Superposition (RCS) statistical model, Ott et al. (2000) calculated an attenuation factor of 0.61 for personal exposure for PM₁₀ for the Toronto study. The mean nonambient exposure component for PM₁₀ was estimated as 53 µg/m³ with a standard

deviation of $85 \mu\text{g}/\text{m}^3$. Although the data were available for $\text{PM}_{2.5}$, similar calculations were not made.

PM_{10} data from the PTEAM study were analyzed using the same approach (Ott et al., 2000). For PTEAM, an attenuation factor of 0.55 was calculated for personal exposure. Infiltration factors were calculated for each residence with an average of 0.56 and a standard deviation of 0.15. Values ranged from a minimum of 0.19 to a maximum of 0.87 showing the substantial variability that can be seen between homes depending upon the housing characteristics and operation of the HVAC system. The mean \pm SD nonambient exposure component for PM_{10} was estimated as $59 \pm 46 \mu\text{g}/\text{m}^3$.

Santos-Burgoa et al. (1998) described a 1992 study of personal exposures and indoor concentrations to a randomly sampled population near Mexico City. The sample of 66 monitored subjects included children, students, office and industrial workers, and housewives. None of the people monitored were more than 65 years old. The mean 24-h personal exposure and indoor concentrations were 97 ± 44 and $99 \pm 50 \mu\text{g}/\text{m}^3$, respectively, with an $r_{\text{Personal}/\text{Ambient}} = 0.26$ ($p = 0.099$). Other correlations of interest were $r_{\text{Personal}/\text{Indoor}} = 0.47$ ($p = 0.002$) and $r_{\text{Indoor}/\text{Ambient}} = 0.23$ ($p = 0.158$). A strong statistical association was found between personal exposure and socioeconomic class ($p = 0.047$) and a composite index of indoor sources at the home ($p = 0.039$).

Correlation analysis for personal exposure has not yet been reported for EXPOLIS. Some preliminary results (Jantunen et al., 2000) show that in Basel and Helsinki a single ambient monitoring station was sufficient to characterize the ambient $\text{PM}_{2.5}$ concentration in each city. Using microenvironmental concentration data collected while the subjects were at home, at work, and outdoors, they calculated the sum of the time-weighted-averages of these data and found the results closely matched the personal $\text{PM}_{2.5}$ exposure data collected by the monitors carried by most of the subjects, although a few subjects (mostly smokers) were noticeable exceptions.

Longitudinal Studies

A number of longitudinal studies using a purposeful sampling design have been conducted and reported in the literature since 1996. Several of these studies (Janssen et al., 1998a, 1999b, 2000; Williams et al., 2000b; Evans et al., 2000) support the previous work by Janssen et al.

(1995) and Tamura et al. (1996a) and demonstrate that, for individuals with little exposure to nonambient sources of PM, correlations between total PM exposure and ambient PM measurements are high. Other studies (Ebelt et al., 2000; Sarnat et al., 2000) show strong correlations for the SO_4^{2-} component of $\text{PM}_{2.5}$ but poorer correlations for $\text{PM}_{2.5}$ mass. Still other studies show only weak correlations (Rojas-Bracho et al., 2000; Linn et al., 1999). Even when strong longitudinal correlations are demonstrated for individuals in a study, the variety of living conditions may lead to variations in attenuating factors or the fraction of ambient PM contributing to personal exposure. Groups with similar living conditions, especially if measurements are conducted during one season, may have similar α and, therefore, very high correlations between personal exposure and ambient concentrations, even for pooled correlations. However, when studies contain subjects with homes of very different ventilation characteristics or cover more than one season, variations in α can be high across subjects, thus, showing poor pooled correlations even in the absence of indoor sources.

Elderly Subjects. Janssen et al. (2000) continued their longitudinal studies with measurements of personal, indoor, and outdoor concentrations of $\text{PM}_{2.5}$ for elderly subjects with doctor-diagnosed angina pectoris or coronary heart disease. Studies were conducted in Amsterdam, the Netherlands and Helsinki, Finland in the winter and spring of 1998 and 1999. In the Amsterdam study with 338 to 417 observations, mean $\text{PM}_{2.5}$ concentrations were 24.3, 28.6, and 20.6 $\mu\text{g}/\text{m}^3$ for personal, indoor, and outdoor samples, respectively. If the measurements with ETS in the home were excluded, the mean indoor concentration dropped to 16 $\mu\text{g}/\text{m}^3$, which was lower than outdoor concentrations. In the Helsinki study, the mean $\text{PM}_{2.5}$ concentrations were 10.8 $\mu\text{g}/\text{m}^3$ for personal, 11.0 $\mu\text{g}/\text{m}^3$ for indoor air, and 12.6 $\mu\text{g}/\text{m}^3$ outdoor air samples. The authors noted that for this group of subjects, personal exposure, indoor concentrations, and ambient concentrations of $\text{PM}_{2.5}$ were highly correlated within subjects over time. Median Pearson's correlation coefficients between personal exposure and outdoor concentrations were 0.79 in Amsterdam and 0.76 in Helsinki. The median Pearson's r for the indoor/outdoor relationship was 0.85 for the Amsterdam study when homes with ETS were excluded. The correlation for indoors versus outdoors was 0.70 for all homes.

Results from the correlation analysis can be used to estimate infiltration factors and penetration factors for these two groups of subjects. In Amsterdam, the attenuation factor was

0.43 and the infiltration factor was 0.47. Very similar results were seen in Helsinki for the attenuation factor (0.45) and the infiltration factor (0.51).

A series of three PM personal monitoring studies involving elderly subjects was conducted in Baltimore County, MD, and Fresno, CA. The first study was a 17-day pilot (January to February 1997) to investigate daily personal and indoor $PM_{1.5}$ concentrations, as well as outdoor $PM_{2.5}$ and $PM_{10-2.5}$ concentrations experienced by nonsmoking elderly residents of a retirement community located near Baltimore (Liao et al., 1999; Williams et al., 2000c). The 26 residents were aged 65 to 89 (mean = 81) years, and 69% of them reported a medical condition such as hypertension or coronary heart disease. In addition, they were quite sedentary: on average, less than 5 h/day were spent on ambulatory activities. Because most of the residents ate meals in a communal dining area, the average daily cooking time in the individual apartments was only 0.5 h (range = 0 to 4.5 h). About 96% of the residents' time was spent indoors (Williams et al., 2000c). Personal monitoring, conducted for five subjects, yielded longitudinal correlation coefficients between ambient concentrations and personal exposure ranging from 0.00 to 0.90.

The main Baltimore study and the Fresno study were conducted using similar monitoring techniques and study design. Concentrations measured in these studies are summarized in Table 5-8. For $PM_{2.5}$, personal exposure and indoor air concentrations are similar for all three studies, even though outdoor air concentrations for Fresno in the spring are only half of those measured for Fresno in the winter and for Baltimore. This difference is presumably due to high penetration efficiencies in the spring in Fresno when the weather was warm and participants tended to keep the windows and doors of their homes open. These data also show that, even when correlations are high, the use of an ambient monitor as a surrogate for exposure in epidemiologic studies can bias the strength of the health effect found due to differing exposure levels.

Calculated correlation coefficients are summarized in Tables 5-9 and 5-10. In Table 5-9, the Baltimore results show high daily average correlations for both $PM_{2.5}$ and PM_{10} . These results primarily represent the behavior of fine-particle regional sulfate for a group of participants who have few indoor or personal sulfate sources. However, even for this group, there was a wide range of individual correlation coefficients. The Fresno data, on the other hand, shows much lower daily average correlations. The residential site may have been

TABLE 5-8. MEAN CONCENTRATION FOR PM MASS REPORTED FOR THE BALTIMORE (Williams et al., 2000a,b,c) AND THE FRESNO (Evans et al., 2000) STUDIES

Study	PM _{2.5} Concentration (µg/m ³)			PM ₁₀ Concentration (µg/m ³)		
	Personal	Indoors	Outdoors	Personal	Indoors	Outdoors
Baltimore	13.0 ± 4.2	10.5 ± 4.9	22.0 ± 12.0	—	13.5 ± 6.3	30.0 ± 13.7
Fresno-Winter	13.3 ± 5.9	9.7 ± 5.0	20.5 ± 13.4	—	15.1 ± 4.1	28.2 ± 15.9
Fresno-Spring	11.1 ± 2.8	8.0 ± 1.8	10.1 ± 3.2	37.3	16.7 ± 3.1	28.7 ± 6.6

TABLE 5-9. DAILY-AVERAGE CORRELATION COEFFICIENTS REPORTED FOR THE BALTIMORE (Williams et al., 2000a,b,c) AND THE FRESNO (Evans et al., 2000) STUDIES

Study	PM _{2.5} R ²			PM ₁₀ R ²
	Ambient/Outdoor	Personal/Ambient	Personal/Indoors	Ambient/Outdoor
Baltimore	0.92	0.80 (0.14-0.80) ^a	0.98 (0.20-0.99) ^a	0.89
Fresno-Winter	0.48	—	—	0.48
Fresno-Spring	0.53	0.7	0.77	0.61

^aRange for individual participants.

TABLE 5-10. REGRESSION ANALYSIS REPORTED FOR INDOOR/OUTDOOR RELATIONSHIPS FOR PM_{2.5} IN THE BALTIMORE (Williams et al., 2000a,b,c) AND FRESNO (Evans et al., 2000) STUDIES

Study	Daily Average			Individual		
	R ²	Slope	Intercept (µg/m ³)	R ²	Slope	Intercept (µg/m ³)
Baltimore	0.92	0.39	1.5	0.73 ± 0.16	0.43 ± 0.15	0.9 ± 2.6
Fresno-Winter	0.86	nr	nr	0.55 ± 0.25	0.25 ± 0.17	4.4 ± 3.2
Fresno-Spring	0.56	nr	nr	0.39 ± 0.21	0.49 ± 0.38	3.0 ± 3.7

nr = not reported.

influenced by highway traffic which could account for the low correlations between ambient and outdoor residential monitors.

In addition, the composition of particles is not the same for these two locations. In Fresno, ammonium nitrate represents a much larger fraction of fine-particle mass. Because this compound is in equilibrium with its precursor gases, ammonia and nitric acid, its partitioning between the gaseous and particulate phase can be easily disturbed while outdoor particles infiltrate indoors. Thus, differences in June 25, 2004 ratios and correlations between Fresno and Baltimore could be a function of differences in ventilation, indoor sources, and sampling artifacts.

The correlation analysis in Table 5-10 shows correlation coefficients as well as the slope (infiltration factor) and the intercept (indoor concentration due to nonambient sources) for the Baltimore and Fresno studies. These data show strongest correlations for Baltimore, where there are very low indoor concentrations from nonambient sources. Correlations are not as strong for Fresno, where there are higher concentrations from nonambient sources. The infiltration factors for Baltimore and Fresno-spring are very similar at approximately 0.5. The infiltration factors for Fresno-winter are considerably lower.

Subjects with COPD. Linn et al. (1999) describe a 4-day longitudinal assessment of personal $PM_{2.5}$ and PM_{10} exposures (on alternate days) in 30 COPD subjects aged 56 to 83 years. Concurrent indoor and outdoor monitoring was conducted at their residences in the Los Angeles area during summer/autumn of 1996. PM_{10} data from the nearest fixed-site monitoring station to each residence was also obtained. Pooled correlations for personal exposure to outdoor measurements gave R^2 values of 0.26 and 0.22 for $PM_{2.5}$ and PM_{10} , respectively. Correlations of day-to-day changes in $PM_{2.5}$ and PM_{10} measured outside the homes and correlated with concurrent PM_{10} measurements at the nearest ambient monitoring location gave R^2 values of 0.22 and 0.44, respectively. Correlations of day-to-day changes in PM mass measured indoors correlated with outdoor measurements at the homes gave R^2 values of 0.27 and 0.19 for PM_{10} and $PM_{2.5}$, respectively.

Personal, indoor, and outdoor $PM_{2.5}$, PM_{10} , and $PM_{2.5-10}$ correlations were reported by Rojas-Bracho et al. (2000) for a study conducted in Boston, MA on 18 individuals with COPD. Both the mean and median personal exposure concentrations were higher than the indoor

concentrations, which were higher than outdoor concentrations for all three PM measurement parameters. June 25, 2004 geometric mean ratios were 1.4 ± 1.9 for PM_{10} , 1.3 ± 1.8 for $PM_{2.5}$, and 1.5 ± 2.7 for $PM_{10-2.5}$. Median longitudinal R^2 s between personal exposure and ambient PM measurements were 0.12 for PM_{10} , 0.37 for $PM_{2.5}$ and 0.07 for $PM_{10-2.5}$. The relationship between the indoor and outdoor concentrations was strongest for $PM_{2.5}$, with a median R^2 of 0.55 and with significant R^2 values for 11 homes. For PM_{10} , the median R^2 value was 0.25, with significant values for eight homes. Only five homes had significant indoor/outdoor associations for $PM_{10-2.5}$, with an insignificant median R^2 value of 0.04. The poor correlations for $PM_{10-2.5}$ are a result of poorer penetration efficiencies, higher decay rates, and spatial inhomogeneities.

Bahadori et al. (1998) reported a pilot study of PM exposure of 10 nonrandomly chosen COPD patients in Nashville, TN during the summer of 1995. Each subject alternately carried a personal $PM_{2.5}$ or PM_{10} monitor for a 12-h daytime period (8:00 am to 8:00 pm) for 6 consecutive days. These same pollutants were monitored simultaneously indoors and outdoors at their homes. All of the homes were air-conditioned and had low air exchange rates (mean = 0.57/h), which may have contributed to the finding that mean indoor $PM_{2.5}$ was 66% of the mean ambient $PM_{2.5}$. This can be contrasted with the PTEAM study in Riverside, CA, where no air conditioners were in use and the mean indoor $PM_{2.5}$ was 98% of the mean ambient $PM_{2.5}$ (Clayton et al., 1993). Data sets were pooled for correlation analysis. Resulting pooled correlations between personal and outdoor concentrations were $r = 0.09$ for $PM_{2.5}$ and -0.08 for PM_{10} .

5.3.3.1.4 Personal Exposure to Sulfate Compared to Personal Exposure to Ambient Particulate Matter

A study conducted in Vancouver involving 16 COPD patients aged 54 to 86 years reported low median longitudinal ($r = 0.48$) and pooled ($r = 0.15$) correlation coefficients between personal exposures and ambient concentrations of $PM_{2.5}$ (Ebelt et al., 2000). However, the mean correlation between personal exposures to sulfate and ambient concentrations of sulfate was much higher ($r = 0.96$). Because there are typically minimal indoor sources of sulfate, the relationship between ambient concentrations and personal exposures to sulfate would not be weakened by variability in an indoor-generated sulfate component as, for example, in the case of $PM_{2.5}$ for which there are many primary indoor sources as well as some secondary indoor

sources. Correlations of ambient concentrations versus personal exposures for PM_{2.5} and sulfate are compared in Figure 5-3.




Figure 5-3. Comparison of correlation coefficients for longitudinal analyses of personal exposure versus ambient concentrations for individual subjects for PM_{2.5} and sulfate.

Another study, conducted in Baltimore, MD, involved 15 nonsmoking adult subjects (> 64 years old) who were monitored for 12 days during summer 1998 and winter 1999 (Sarnat et al., 2000). All subjects (nonrandom selection) were retired, physically healthy, and lived in nonsmoking private residences. Each residence, except one, was equipped with central air-conditioning; however, not all residences used air-conditioning throughout the summer. The average age of the subjects was 75 years (± 6.8 years). Sarnat et al. (2000) reported higher longitudinal and pooled correlations for PM_{2.5} during summer than winter. Similar to Ebelt et al. (2000), Sarnat et al. (2000) reported stronger associations between personal exposure to SO₄²⁻

and ambient concentrations of SO_4^{2-} than for total personal $\text{PM}_{2.5}$ exposure and ambient $\text{PM}_{2.5}$ concentrations. The ranges of correlations are shown in Figure 5-3, along with similar data from Ebel et al. (2000).

The higher correlation coefficients and the narrower range of the correlation coefficient for sulfate suggest that removing indoor-generated and personal-activity PM from total personal PM would result in a higher correlation with ambient concentrations. If, as discussed in 5.3.4.3.1, there are no indoor sources, a personal exposure measurement for sulfate gives the ambient exposure of sulfate; the ratio of personal sulfate to ambient sulfate gives the attenuation coefficient on an individual, daily basis; and the attenuation coefficient times the ambient $\text{PM}_{2.5}$ concentration gives the individual, daily values of ambient $\text{PM}_{2.5}$ exposures (Wilson et al., 2000). This technique applies only to the nonvolatile components of fine PM, as measured by $\text{PM}_{2.5}$. It requires that the sulfate concentration be large enough so that it can be measured with reasonable accuracy. It does not require that sulfate be correlated with $\text{PM}_{2.5}$ or the non-sulfate components of $\text{PM}_{2.5}$, because the sulfate data is used to estimate the attenuation coefficient, not $\text{PM}_{2.5}$. The technique does require that there be minimal indoor sources of sulfate, as indicated by a near-zero intercept for the regression, and that the size distribution of $\text{PM}_{2.5}$ and sulfate be similar.

Sarnat et al. (2001) subsequently extended the Baltimore study to include 20 older adults, 21 children, and 15 individuals with COPD for a total of 56 subjects. In both studies (Sarnat et al., 2000, 2001), personal and ambient sulfate data were used to estimate the ambient $\text{PM}_{2.5}$ exposure. They used this information in mixed-model analysis (mixed models account for differences among individual subjects), but did not report correlations between ambient $\text{PM}_{2.5}$ exposure and ambient $\text{PM}_{2.5}$ concentrations based on the pooled data set. However, Sarnat et al. (2001) did report slopes from the mixed model analyses. The t-statistic for the slope of ambient exposure versus ambient concentration, as compared to total personal exposure versus ambient concentration, increased from 9.96 to 11.12 (total exposure versus ambient concentration) for the summer period and from 4.36 to 19.88 (ambient exposure versus ambient concentration) for the winter period.

The study conducted by Sarnat et al. (2000) also illustrates the importance of ventilation on personal exposure to PM. During the summer, subjects recorded the ventilation status of every visited indoor location (e.g., windows open, air-conditioning use). As a surrogate for the air

exchange rate, personal exposures were classified by the fraction of time the windows were open while a subject was in an indoor environment (F_v). Sarnat et al. (2000) reported regression analyses of personal exposure on ambient concentration for $PM_{2.5}$ and for sulfate for each of the three ventilation conditions (Figure 5-4). The correlation between personal exposure and ambient concentration is higher for sulfate than for $PM_{2.5}$, presumably because $PM_{2.5}$ has indoor sources as well as ambient sources but sulfate has only ambient sources. As expected, the improvement is better for the lower ventilation conditions, because under these conditions the ambient concentration is larger. For the lowest ventilation condition, R^2 improves from 0.25 to 0.72.

5.3.3.1.5 Personal Exposure to Ambient and Nonambient Particulate Matter

The utility of treating personal exposure to ambient PM, A, and personal exposure to nonambient PM, N, as separate and distinct components of total personal exposure to PM, T, was pointed out by Wilson and Suh (1997). The PTEAM study measured, in addition to indoor, outdoor, and personal PM_{10} , the air exchange rate for each home and collected information on the time spent in various indoor and outdoor microenvironments. This information is available for 147 12-h daytime periods. With this information and statistically estimated values of P and k , it is possible to estimate the daytime A and N as described in Section 5.2.4.3. Various examples of this information have been reported (Mage et al., 1999; Wilson et al., 2000). Graphs showing the relationships between ambient concentration and the various components of personal exposure (T, A, and N) are shown in Figure 5-5. The correlation coefficient for the pooled data set improves from $r = 0.377$ for T versus C (Figure 5-5a) to $r = 0.856$ for A versus C (Figure 5-5b) because of the removal of the N, which, as shown in Figure 5-5c, is highly variable and independent of C. The correlation between A and C is less than 1.0 because of the day-to-day variation in the α of each individual. The regression of T on C gives $\bar{\alpha} = 0.711$ and $\bar{N} = 81.6 \mu\text{g}/\text{m}^3$. The regression of A on C gives $\bar{\alpha} = 0.625$. The regression of N on C gives $\bar{N} = 79.2 \mu\text{g}/\text{m}^3$. The finite intercept in the regression with A must be attributed to bias or error in some of the measurements. No reported studies, other than PTEAM, have provided the quantity of data on individual, daily values of T, C, C_i , and a that are required to conduct an analysis comparable to that shown in Figure 5-5. It should be noted that the PTEAM study was conducted in southern California in the fall, when houses were open and air exchange rates were




Figure 5-4. Personal exposure versus ambient concentrations for PM_{2.5} and sulfate for well, moderately, and poorly ventilated indoor environments. (Slope estimated from mixed models).

Source: Sarnat et al. (2000).




Figure 5-5. Regression analyses of aspects of daytime personal exposure to PM_{10} estimated using data from the PTEAM study: (a) total personal exposure to PM, T, regressed on ambient concentration, C; (b) personal exposure to ambient PM, A, regressed on C; and (c) personal exposure to nonambient PM, N, regressed on C.

Source: Data taken from Clayton et al. (1993). Adapted from Mage et al. (1999) and Wilson et al. (2000).

high and relatively uniform. These are best case conditions for showing high correlations between ambient site measurements and personal correlations. Such high correlations are not usually found and would not be expected with lower and more variable air exchange rates.

The RCS model introduced by Ott et al. (2000) presents a modeling framework to determine the average contribution of ambient PM₁₀ and indoor-generated PM₁₀ to average personal exposures in large urban metropolitan areas. The model has been tested using personal, indoor, and outdoor PM₁₀ data from three urban areas (Riverside, CA; Toronto; and Phillipsburg, NJ). Results suggest that it is possible to separate the average ambient and nonambient PM contributions to personal exposures on a community-wide basis. However, as discussed in the paper, the authors make some assumptions that require individual consideration in each city-specific application of the model for exposure or health effects investigations. Primarily, housing factors, air-conditioning, seasonal differences, and complexities in time-activity profiles specific to the cohort being studied have to be taken into account prior to adopting the model to a given situation. Finally, the exposure-based analyses presented here do not predict the relative contribution of indoor and outdoor PM to particle mass burden to the lung as a function of human activities and different microenvironmental sources and concentrations of PM and its co-pollutants.

5.3.3.2 Factors That Affect Relationships Between Personal Exposures and Ambient PM

A number of factors can affect the relationship between personal PM exposures and PM measured at ambient-site community monitors. Spatial variability in outdoor microenvironmental concentrations of PM and variations in penetration of PM into indoor microenvironments influence the relationship of ambient PM concentration to ambient PM exposure. Air exchange rates and decay rates in indoor microenvironments influence the relationship of ambient PM concentrations to both ambient and total PM exposure, whereas personal activities that generate particles influence the relationship to total, but not ambient PM exposure. Information on these effects is presented in detail in the following sections.

5.3.3.2.1 Spatial Variability and Correlations Over Time

Chapter 3, Section 3.2.3, presents information on the spatial variability of PM mass and chemical components at fixed-site ambient monitors; for purposes of this chapter, this spatial

variability is called an “ambient gradient.” The data presented in Section 3.2.3 indicate that ambient gradients of PM and its constituents exist to a greater or lesser degree in urban areas. These and any other gradients that may exist between a fixed-site monitor and the outdoor microenvironments near where people live, work, and play obviously affect their exposure. The purpose of this section is to review the available data on ambient monitor-to-outdoor microenvironmental concentration gradients or relationships that have been measured by researchers since 1996. These analyses, presented below, are generally consistent with earlier studies covered in the 1996 PM AQCD. A few outdoor-to-outdoor monitoring studies also are included to highlight relationships among important microenvironment categories. To assess spatial variability or gradients, the spatial correlations in the data are usually analyzed. However, it should be noted that high temporal correlation between two monitoring locations does not imply low spatial variability or low ambient gradients. A high temporal correlation between two sites indicates that changes in concentrations at one site may be estimated from data collected at another site.

In a paper on the EXPOLIS-EAS study, Oglesby et al. (2000) concluded that in Basel, Switzerland little spatial variability exists between PM levels measured at fixed site monitors and the participants’ outdoor microenvironments. The authors reported a high correlation between home outdoor PM_{2.5} levels (48-h measurements beginning and ending at 8:00 a.m.) and the corresponding 24-h average PM₄ (time-weighted values calculated from midnight to midnight) measured at a fixed monitoring station (n = 38, $r_{sp} = 0.96$, $p < 0.001$). They considered each home outdoor monitor as a temporary fixed monitor and concluded that “the PM_{2.5} level measured at home outdoors . . . represents the fine particle level prevailing in the city of Basel during the 48-h measuring period.”

In a study conducted in Helsinki, Finland, Buzorius et al. (1999) concluded that a single monitor may be used to adequately describe the temporal variations in concentration across the metropolitan area. Particle size distributions were measured using a differential mobility particle sizer (DMPS; Wintlmayer) coupled with a condensation particle counter (CPC TSI 3010, 3022) at four locations, including the official air monitoring station, which represented a “background” site. The monitoring period varied between 2 weeks and 6 mo for the sites, and data were reported for 10-min and 1-, 8-, and 24-h averages. As expected, temporal variation decreased as the averaging time increased. The authors reported that particle number concentration varied in

magnitude with local traffic intensity. Linear correlation coefficients computed for all possible site-pairs and averaging times showed that the correlation coefficient improved with increasing averaging time. Using wind speed and direction vectors, lagged correlations were calculated and were generally higher than the “raw” data correlations. The authors noted that weekday correlations were higher than weekend correlations as “traffic provides relatively uniform spatial distribution of particulate matter” and concluded that, even for time periods of 10 min and 1 h, sampling at one station can describe temporal variations across relatively large areas of the city with $r > 0.7$.

Dubowsky et al. (1999) pointed out that, although the variation of $PM_{2.5}$ mass concentration across a community may be small, there may be significant spatial variations of specific components of the total mass on a local scale. An example is given of a study of concentrations of polycyclic aromatic hydrocarbons (PAHs) at three indoor locations in a community: (1) an urban site, (2) a semi-urban site 1.6 km away, and (3) a suburban site located further away. The authors found the geometric mean PAH concentrations at these three locations varied respectively as 31:19:8 ng/m^3 and suggested that the local variations in traffic density were responsible for this gradient. Note that these concentrations are 1,000 times lower than the total PM mass concentration and that such a small gradient would not be detectable for total $PM_{2.5}$ mass measurements on the order of 25 $\mu g/m^3$.

The Total Human Environmental Exposure Study (THEES) reported by Waldman et al. (1991) measured indoor, outdoor, and personal benzo(a)pyrene (BaP) levels and found that the outdoor BaP was the same at all outdoor sites across the three sampling periods. This study showed seasonal differences for BaP levels as well as for exposures due to indoor and outdoor sources and individual activities.

Leaderer et al. (1999) monitored 24-h PM_{10} , $PM_{2.5}$, and sulfates during the summers of 1995 and 1996 at a regional site in Vinton, VA (6 km from Roanoke, VA). One similar 24-h measurement was made outdoors at residences in the surrounding area at distances ranging from 1 to > 175 km from the Vinton site, at an average separation distance of 96 km. The authors reported significant correlations for $PM_{2.5}$ and sulfates between the residential outdoor values and those measured at Vinton on the same day. In addition, the mean values of the regional site and residential site $PM_{2.5}$ and sulfates showed no significant differences in spite of the large distance separations and mountainous terrain intervening in most directions. However, for the

concentrations of coarse particles estimated as PM_{10} minus $PM_{2.5}$, no significant correlations among these sites were found ($n = 30$; $r = -0.20$).

Lillquist et al. (1998) found no significant gradient in PM_{10} concentrations in Salt Lake City, UT when levels were low, but a gradient existed when levels were high. Outdoor PM_{10} levels were measured for a period of about 5 mo at three hospitals using a Minivol 4.01 sampler (Airmetrics, Inc.) operating at 5 L/min and at the Utah Department of Air Quality (DAQ) ambient monitoring station located 3 to 13 km from the hospitals.

Pope et al. (1999) monitored ambient PM_{10} concentrations in Provo, UT (Utah Valley) during the same time frame the following year and reported nearly identical concentrations at three sites separated by 4 to 12 km. Pearson correlation coefficients for the data were between 0.92 and 0.96. The greater degree of variability in the Salt Lake City PM_{10} data relative to the Provo data may be related to the higher incidence of windblown crustal material in Salt Lake City.

Väkevä et al. (1999) found significant vertical gradients in submicron particles existed in an urban street canyon of Lahti, Finland. Particle number concentrations were measured using a TSI screen diffusion battery and a condensation particle counter at 1.5 and 25 m above the street at rooftop level. The authors found a five-fold decrease in concentration between the two sampling heights and attributed the vertical gradient to dilution and dispersion of pollutants emitted at street level.

White (1998) suggested that the higher random measurement error for the coarse PM fraction compared to the error for the fine PM fraction may be responsible for a major portion of the apparent greater spatial variability of coarse ambient PM concentration compared to fine ambient PM concentration in a community (e.g., Burton et al., 1996; Leaderer et al., 1999). When $PM_{2.5}$ and PM_{10} are collected independently and the coarse fraction is obtained by difference ($PM_{10-2.5} = PM_{10} - PM_{2.5}$), the expected variance in the coarse fraction is influenced by the variances of the PM_{10} and $PM_{2.5}$ measurements. When a dichotomous sampler collects $PM_{2.5}$ and $PM_{10-2.5}$ on two separate filters, the coarse fraction is expected to have a larger error than the fine fraction. There is a possible error caused by loss of mass below the cut-point size and a gain of mass above the cut-point size that is created by the asymmetry of the product of the penetration times PM concentration about the cut-point size. Because a dichotomous PM

sampler collects coarse mass using an upper and lower cut point, it is expected to have a larger variance than for fine mass collected using only one cut point.

Wilson and Suh (1997) concluded that $PM_{2.5}$ and PM_{10} concentrations are correlated more highly across Philadelphia than are $PM_{10-2.5}$ concentrations. Ambient monitoring data from 1992 to 1993 was reviewed for $PM_{2.5}$, $PM_{10-2.5}$, and PM_{10} , as well as for $PM_{2.5}$ and $PM_{10-2.5}$ dichotomous data for 212 site-years of information contained in the AIRS database (U.S. Environmental Protection Agency, 2000). The authors also observed that PM_{10} was frequently correlated more highly with $PM_{2.5}$ than with $PM_{10-2.5}$. The authors noted that $PM_{2.5}$ constitutes a large fraction of PM_{10} and that this is the likely reason for the strong agreement between $PM_{2.5}$ and PM_{10} . Similar observations were made by Keywood et al. (1999) in six Australian cities. The authors reported that PM_{10} was more highly correlated with $PM_{2.5}$ than with coarse PM ($PM_{10-2.5}$) suggesting that “variability in PM_{10} is dominated by variability in $PM_{2.5}$.”

Lippmann et al. (2000) examined the site-to-site temporal correlations in Detroit (1981 to 1994) and found the ranking of median site-to-site correlations was O_3 (0.83), PM_{10} (0.78), TSP (0.71), NO_2 (0.70), CO (0.50), and SO_2 (0.49). The authors explain that O_3 and a fraction of TSP and PM_{10} (e.g., sulfate) are secondary pollutants that would tend to be distributed more uniformly spatially within the city than primary pollutants such as CO and SO_2 , which are more likely to be influenced by local emission sources. Lippman et al. (2000) concluded “. . . spatial uniformity of pollutants may be due to area-wide sources, or to transport (e.g., advection) of fairly stable pollutants into the urban area from upwind sources. Relative spatial uniformity of pollutants would therefore vary from city to city or region to region.”

Goswami et al. (2002) used data collected at outdoor monitors of homes in a large exposure study in Seattle, WA to analyze the spatial variability of outdoor $PM_{2.5}$ concentrations. The day-to-day variability between sites was 10 times higher than the spatial variability between sites. However, differences between sites was sufficient to potentially contribute to measurement error. An examination of the spatial characteristics of the monitoring sites showed that the most representative monitoring sites were located at elevations of 80 to 120 m above sea level and at distances of 100 to 300 m from the nearest arterial road.

5.3.3.2.2 *Physical Factors Affecting Indoor Microenvironmental Particulate Matter Concentrations*

Several physical factors can affect ambient particle concentrations in the indoor microenvironment, including air exchange, penetration, and particle deposition. Combined, these factors are critical variables that describe ambient particle dynamics in the indoor microenvironments and, to a large degree, significantly affect an individual's personal exposure to ambient particles while indoors. The relationship between the concentrations of ambient particles outdoors, C , and ambient particles that have infiltrated indoors, C_{ai} , is given by

$$\frac{C_{ai}}{C} = \frac{Pa}{a + k} \quad (5-14)$$

where P is the penetration factor; a is the air exchange rate; and k is the particle deposition rate. (As discussed in Section 5.3.2.3.1, use of this model assumes equilibrium conditions and assumes that all variables remain constant.) Particle penetration is a dimensionless quantity that describes the fraction of ambient particles that effectively penetrates the building shell. "Air exchange" is a term used to describe the rate at which the indoor air in a building or residence is replaced by outdoor air. The dominant processes governing particle penetration are air exchange and deposition of particles as they traverse through cracks and crevices and other routes of entry into the building. Although air exchange rates have been measured in numerous studies, very few field data existed prior to 1996 to determine size-dependent penetration factors and particle deposition rates. All three parameters (P , a , and k) vary substantially depending on building type, region of the country, and season. In the past several years, researchers have made significant advancements in understanding the relationship between particle size and penetration factors and particle deposition rates. This section highlights studies that have been conducted to better understand physical factors affecting indoor particle dynamics.

Air Exchange Rates

The air exchange rate, a , in a residence varies depending on a variety of factors, including geographical location, age of the building, the extent to which window and doors are open, and season. Murray and Burmaster (1995) used measured values of a from households throughout the United States to describe empirical distributions and to estimate univariate parametric

probability distributions of air exchange rates. Figure 5-6 shows the results classified by season and region. In general, a is highest in the warmest region and increases from the coldest to the warmest region during all seasons. Air exchange rates also are quite variable within and between seasons, as well as between regions (Figure 5-6). Data from the warmest region in summer should be viewed cautiously, because many of the measurements were made in southern California in July when windows were more likely to be open than in other areas of the country where air-conditioning is used. Use of air-conditioning generally results in lowering air exchange rates. In a separate analysis of these data, Koontz and Rector (1995) suggested that a conservative estimate for air exchange in residential settings would be 0.18 h^{-1} (10th percentile) and a typical air exchange would be 0.45 h^{-1} (50th percentile).




Figure 5-6. Air exchange rates measured in homes throughout the United States. Climatic regions are based on heating-degree days: coldest region ≥ 7000 , colder region = 5500 to 6999, warmer region = 2500 to 4999, and warmest region ≤ 2500 heating-degree days.

Source: Adapted from Murray and Burmaster (1995).

These data provide reasonable experimental evidence that a varies by season in locations with distinct seasons. As a result, infiltration of ambient particles may be more efficient during warmer seasons when windows are likely to be opened more frequently and air exchange rates are higher. This suggests that the fraction of ambient particles present in the indoor microenvironment would be greater during warmer seasons than colder seasons. For example, in a study conducted in Boston, MA, participants living in non-air-conditioned homes kept the windows closed except during the summer (Long et al., 2000). This resulted in higher and more variable air exchange rates in summer than during any other season (Figure 5-7). During nighttime periods when indoor sources are negligible, the indoor/outdoor concentration ratio or infiltration factor may be used to determine the relative contribution of ambient particles in the indoor microenvironment. Particle data collected during this study were used to determine the indoor/outdoor concentration ratios by particle size (Figure 5-8). For these nine homes in Boston the fraction of ambient particles penetrating indoors was higher during summer when air exchange rates were higher than during fall when air exchange rates were lower (Long et al., 2001a).




Figure 5-7. Box plots of hourly air exchange rates stratified by season in Boston, MA during 1998.

Source: Long et al. (2000).




Figure 5-8. Geometric mean infiltration factor (indoor/outdoor ratio) for hourly nighttime, nonsource data for two seasons. Box plots of air exchange rates are shown as inserts for each plot. (Boston, 1998)

Source: Long et al. (2001a).

Wallace et al. (2002) conducted a year-long investigation of air rates in an occupied house to establish the effects of temperature, wind velocity, use of exhaust fans, and window-opening behavior. Air exchange rates were calculated by periodically injecting a tracer gas (SF_6) into the return air duct and measuring the concentration in 10 indoor locations sequentially every minute by a gas chromatograph equipped with an electron capture detector. Temperature and relative humidity were measured outdoors and in multiple indoor locations. Wind speed and direction in the horizontal plane were also measured. Use of the thermostat-controlled attic fan was recorded automatically. Indoor temperatures increased from 21 °C in winter to 27 °C in summer. Windows were open only a few percent of the time in winter but more than half the time in summer. About 4,600 hour-long average air exchange rates were calculated from the measured tracer gas decay rates. The mean (\pm SD) rate was 0.65 (0.56) h^{-1} . Tracer gas decay rates in different rooms were very similar, ranging only from 0.62 to 0.67 h^{-1} , suggesting that the air in

the house was well mixed throughout the year. The strongest influence on air exchange rates was opening windows, which could increase the rate to as much as 2 h⁻¹ for extended periods, and up to 3 h⁻¹ for short periods of a few hours. The use of the attic fan also increased air exchange rates by amounts up to 1 h⁻¹. Use of the furnace fan had no effect on air exchange rates (ducts were all inside the house). A clear effect of indoor-outdoor temperature difference, ΔT , could be discerned. However, wind speed and direction were found to have very little influence on air exchange rates at the house where the measurements were made.

The air exchange rate, a , is a critical parameter in determining the fraction of ambient PM found indoors and the extent of build-up of indoor PM due to indoor sources. Wallace et al. (2002) provided a brief review of the existing literature on the theory and measurement of air exchange rates. Open windows and frequent opening of doors lead to higher values of a . However, many homes are kept closed for heating in winter and air-conditioning in summer. Windows may or may not be opened during moderate weather conditions. In some areas, on the other hand, heating or air-conditioning may not be required and ventilation by open windows may be more frequent. Thus, a may vary geographically with climate. However, wind speed and direction were found to have very little influence on air exchange rates at the house where measurements were made. The variation of a with ΔT , as shown in Figure 5-9, is given by $a \text{ (h}^{-1}\text{)} = 0.176 + 0.0162 \Delta T \text{ (}^\circ\text{C)}$. Thus, an increase of 10 °C in ΔT would lead to an increase in a of 0.164 h⁻¹, or almost a doubling of the value of a for no indoor/outdoor temperature difference and no open windows or forced ventilation.

The observation of a minimal effect of wind speed on a is an important finding. If a were strongly dependent on wind speed, especially at low wind speeds, there might be a correlation between the indoor-generated PM found indoors and the ambient concentration outdoors. Such a correlation could lead to a confounding of the relationship of ambient PM with health outcomes by indoor-generated PM. Wallace et al. (2002) suggested that “the generally tighter construction of homes and the use of vapor barriers may have reduced the effect of wind speed and direction on residential air change rates compared to earlier studies.”

Wind speed might be expected to have a larger effect on a in a home with open windows. Under conditions of large a , the ambient infiltrated indoors PM concentration will be a large fraction of the outdoor PM and the two concentrations will be highly correlated. However, the indoor-generated PM concentration will be kept low by the high a , making a significant




Figure 5-9. Regression of air exchange rate on absolute indoor-outdoor temperature difference. Overnight values (midnight-8 a.m.) in winter (January-March 2000). a (h^{-1}) = 0.176 (0.011 SE) + 0.0164 (0.0005) ΔT $^{\circ}\text{C}$ ($N = 183$, $R^2 = 0.82$).

correlation between ambient PM concentrations and indoor-generated PM concentrations unlikely. The observed lack of a strong wind effect on a in closed houses (Wallace et al., 2002; Howard-Reed et al., 2002) and the observed lack of correlation between ambient PM concentrations and indoor-generated PM concentrations indicate that the possibility of correlation of ambient and indoor-generated PM concentrations can be discounted.

Particle Deposition Rates and Penetration Factors


Physical factors affecting indoor particle concentrations including particle deposition rates, k , and penetration factors, P , are possibly the most uncertain and variable quantities. Although k can be modeled with some success, direct measurements are difficult, and results often vary from study to study. Particle deposition rates vary considerably depending on particle size because of the viscous drag of air on the particles hindering their movement to varying degrees. The nature and composition of particles also affect deposition rates. Surface properties of

particles, such as their electrostatic properties, can have a significant influence on deposition rates. In addition, thermophoresis can also affect k , but probably to a lesser degree in the indoor microenvironment, because temperatures generally vary over a small range. Combined, these effects can produce order of magnitude variations in k between particles of different size and, in the case of electrophoresis and thermophoresis, particles of the same size.

Particle penetration efficiency into indoor microenvironments depends on particle size and air exchange rates. Penetration varies with particle size because of the size-dependent deposition of particles caused by impaction, interception, and diffusion of particles onto surfaces as they traverse through cracks and crevices. Penetration also is affected by air exchange rates. When air exchange rates are high, P approaches unity because the majority of ambient particles have less interaction with the building shell. In contrast, when air exchange rates are low, P is governed by particle deposition as particles travel through cracks and crevices.

Significant advancements have been made in the past few years to better characterize particle deposition rates and penetration factors. Several new studies, including two in which semi-continuous measurements of size distributions were measured indoors and outdoors, have produced new information on these quantities, which are key to understanding the contributions of ambient PM to indoor PM concentrations (Equation 5-7).

Studies involving semi-continuous measurements of indoor and outdoor particle size distributions have been used to estimate k and P as a function of particle size (Vette et al., 2001; Long et al., 2001a; Abt et al., 2000b). These studies each demonstrated that the indoor/outdoor concentration ratios (C_{ai}/C in Equation 5-9) were highest for accumulation mode particles and lowest for ultrafine and coarse-mode particles. Various approaches were used to estimate size-specific values for k and P . Vette et al. (2001) and Abt et al. (2000b) estimated k by measuring the decay of particles at times when indoor levels were significantly elevated. Vette et al. (2001) estimated P using measured values of k and indoor/outdoor particle measurements during nonsource nighttime periods. Long et al. (2001a) used a physical-statistical model, based on Equation 5-12, to estimate k and P during nonsource nighttime periods. The results for k reported by Long et al. (2001a) and Abt et al. (2000b) are compared with other studies in Figure 5-10. Although not shown in Figure 5-10, the results for k obtained by Vette et al. (2001) were similar to the values of k reported by Abt et al. (2000b) for particle sizes up to 1 μm . Results for P by Long et al. (2001a) showed that penetration was highest for accumulation-mode



- ^aDecay Rates represent summary estimates from the four houses examined.
- ^bDecay rates are based on sulfate and are presented as < 2.5 μm . Estimates were computed using a surface-to-volume ratio of 2 m^{-1} (Koutrakis *et al.*, 1992).
- ^cData represents $\text{PM}_{2.5}$.
- ^dParticle sizes are the midpoint of the ranges examined.
- ^eDecay rates presented are estimates of k for nightly average data from all nine study homes.
- ^fDecay rates are theoretically modeled deposition values for smooth indoor surfaces and homogeneous and isotropically turbulent air flow. Presented curves assume typical room dimensions (3 m x 4 m x 5 m) and a friction velocity of 1.0 cm/s.

Figure 5-10. Comparison of deposition rates from Long et al. (2001a) with literature values (from Abt et al., 2000b). Error bars represent standard deviations for same-study estimates.

Source: Adapted from Abt et al. (2000b); Long et al. (2001a).

particles and decreased substantially for coarse-mode particles (Figure 5-11). The results for P reported by Vette et al. (2001) show similar trends, but were lower than those reported by Long et al. (2001a). This is likely because of the lower air exchange rates in the one Fresno, CA test home ($a \approx 0.5 \text{ h}^{-1}$; Vette et al., 2001) than in the nine test homes in Boston, MA study ($a > 1 \text{ h}^{-1}$; Long et al., 2001a). These data for P and k illustrate the role that the building shell may provide in increasing the concentration of particles because of indoor sources and reducing




Figure 5-11. Penetration efficiencies and deposition rates from models of nightly average data. Error bars represent standard errors.

Source: Boston (1998).

the concentration of indoor particles from ambient sources, especially for homes with low air exchange rates.

Several other studies have investigated particle loss as a function of particle size. The penetration of particles across building envelopes has been modeled for several sizes of idealized rectangular cracks (Liu and Nazaroff, 2001). Particles of 0.1 to 1.0 μm diameter had penetration efficiencies near 1.0. Supermicron and ultrafine particles were removed to a greater degree by gravitational settling and Brownian diffusion. Thatcher et al. (2002) conducted an experimental study of the effects of room furnishings and air speed on particle deposition rates indoors. The deposition loss rate (k) increased by as much as a factor of 2.6 in going from a bare room (35 m^2 surface area) to a fully furnished room (12 m^2 additional surface area) with the greatest increase seen for the smallest particles. Air speed increases from < 5 to 19 cm/s enhanced the deposition rates by factors of 1.3 to 2.4, with greater effects on large particles than small particles. The

authors stated that, “The significant effect of particle size and room conditions on deposition loss rates argues against using a single first-order loss-rate coefficient to represent deposition for integrated mass measurements (PM_{2.5} or PM₁₀).” Riley et al. (2002) have modeled the infiltration of particles into two building types: offices and residences. They developed representative values of P , k , and ventilation-system filter efficiencies for particle sizes from 0.001 to 10 μm . They then used a typical rural and urban outdoor size distribution and calculated predicted indoor concentrations for number, surface area, and volume distributions.

Compositional Differences Between Indoor-Generated and Ambient Particulate Matter

Wilson et al. (2000) discussed the differences in composition between particles from indoor and outdoor sources. They noted that, because of the difficulty in separating indoor PM into ambient and nonambient PM, there is little direct experimental information on the composition differences between the two. Although experimental data are limited, Wilson et al. (2000) suggested the following:

“Photochemistry is significantly reduced indoors; therefore, most secondary sulfate [H_2SO_4 , NH_4HSO_4 , and $(\text{NH}_4)_2\text{SO}_4$] and nitrate (NH_4NO_3) found indoors come from ambient sources. Primary organic emissions from incomplete combustion may be similar, regardless of the source. However, atmospheric reactions of polyaromatic hydrocarbons and other organic compounds produce highly oxygenated and nitrated products, so these species are also of ambient origin. Gasoline, diesel fuel, and vehicle lubricating oil all contain naturally present metals or metal additives. Coal and heavy fuel oil also contain more metals and nonmetals, such as selenium and arsenic, than do materials such as wood or kerosene burned inside homes. Environmental tobacco smoke (ETS), however, with its many toxic components, is primarily an indoor-generated pollutant.”

Particles generated indoors may have different chemical and physical properties than those generated by anthropogenic ambient sources. Siegmann et al. (1999) have demonstrated that elemental carbon in soot particles generated indoors has different properties than in those generated outdoors by automotive or diesel engines. In the United States, combustion-product PM in the ambient/outdoor air generally is produced by burning fossil fuels (e.g., coal, gasoline, fuel oil) and wood, whereas combustion-product PM from indoor sources is mainly produced by biomass burning (e.g., tobacco, wood, foods). However, some indoor sources of PM (such as cigarette smoking, meat cooking, and coal burning) occur both indoors and outdoors and may

constitute an identifiable portion of measured ambient PM (Cha et al., 1996; Kleeman and Cass, 1998).

Indoor Air Chemistry

Gas- and aerosol-phase chemical reactions in the indoor microenvironment are responsible for secondary particle formation and modification of existing particles. This process could be complex and may influence the interpretation of exposures to indoor-generated particles in instances when particles are generated by outdoor gases reacting with gases generated or released indoors to produce fresh particles. For example, homogeneous gas phase reactions involving ozone and terpenes (specifically *d*-limonene, α -terpinene, and α -pinene) have been identified as an important source of submicron particles (Weschler and Shields, 1999). Terpenes are present in several commonly available household cleaning products, and *d*-limonene has been identified in more than 50% of the buildings monitored in the Building Assessment and Survey Evaluation (BASE) study (Hadwen et al., 1997). Long et al. (2000) found that when PineSol (primary ingredient is α -pinene) was used indoors, indoor PM_{2.5} mass concentrations increased from 11 to 32 $\mu\text{g}/\text{m}^3$ (indoor O₃ concentrations unknown, but ambient O₃ levels were 44 to 48 ppb). Similarly, a 10-fold increase in number counts of 0.1 to 0.2 μm particles was observed in an experimental office containing supplemental *d*-limonene and normally encountered indoor O₃ concentrations (< 5 to 45 ppb), resulting in an average increase in particle mass concentration of 2.5 to 5.5 $\mu\text{g}/\text{m}^3$ (Weschler and Shields, 1999). Ozone appears to be the limiting reagent, because particle number concentration varied proportionally to O₃ concentrations (Weschler and Shields, 1999). Other studies showed similar results (e.g., Jang and Kamens, 1999; Wainman et al., 2000). Such particles, if toxic, would represent an increased health risk due to ambient air pollution. However, the concentration would depend on the ambient O₃ concentration, the O₃ infiltration factor, and the indoor generation rate of terpenes. It seems unlikely that the concentration of the resulting particles would be correlated with ambient PM on an individual or population basis. Therefore, it seems more appropriate to consider indoor-reaction particles as part of nonambient exposure. However, since ambient O₃ and ambient PM_{2.5} are correlated in the summertime, indoor-reaction particles due to O₃-terpene reactions might be correlated with outdoor PM_{2.5} mass.

Indoor Sources of Particles

The major sources of indoor PM in nonsmoking residences and buildings include suspension of PM from bulk material, cooking, cleaning, and the use of combustion devices such as stoves and kerosene heaters. Human and pet activities also lead to PM detritus production (from tracked-in soil, fabrics, skin and hair, home furnishings, etc.) that is found ubiquitously in house dust deposited on floors and other interior surfaces. House dust and lint particles may be resuspended indoors by agitation (cleaning) and turbulence (HVAC systems, human activities, etc.). Ambient particles that have infiltrated into the indoor microenvironments also may be resuspended after deposition to indoor surfaces. Typically, resuspension of particles from any source involves coarse particles ($> 1 \mu\text{m}$); particles with smaller diameters are not resuspended efficiently. On the other hand, cooking produces both fine- and coarse-mode particles, whereas combustion sources typically produce ultrafine particles.

Environmental tobacco smoke (ETS) is also a major indoor source of PM. It is, however, beyond the scope of this chapter to review the extensive literature on ETS. A number of articles provide source-strength information for cigarette or cigar smoking (e.g., Daisey et al. [1998] and Nelson et al. [1998]).

A study conducted on two homes in the Boston metropolitan area (Abt et al., 2000a) showed that indoor PM sources predominate when air exchange rates were $< 1 \text{ h}^{-1}$, and outdoor sources predominate when air exchange rates were $\geq 2 \text{ h}^{-1}$. The authors attributed this to the fact that when air exchange rates were low ($< 1 \text{ h}^{-1}$), particles released from indoor sources tend to accumulate, because particle deposition is the mechanism governing particle decay and not air exchange. Particle deposition rates are generally $< 1 \text{ h}^{-1}$, especially for accumulation-mode particles. When air exchange rates were higher ($\geq 2 \text{ h}^{-1}$), infiltration of ambient aerosols and exfiltration of indoor-generated aerosols occur more rapidly, reducing the effect of indoor sources on indoor particle levels. The study also confirmed previous findings that the major indoor sources of PM are cooking, cleaning, and human activity. The authors discussed the size characteristics of these ubiquitous sources as follows:

“The size of the particles generated by these activities reflected their formation processes. Combustion processes (oven cooking, toasting, and barbecuing) produced fine particles and mechanical processes (sauteing, frying, cleaning, and movement of people) generated coarse particles. These activities increased particle concentrations by many orders of magnitude higher than outdoor levels and altered indoor size distributions.” (Abt et al., 2000a; p. 43)

They also noted that variability in indoor PM for all size fractions was greater (2 to 33 times higher) than for outdoor PM, especially for short averaging times.

In a separate study conducted in nine nonsmoking homes in the Boston area, Long et al. (2000) concluded that the predominant source of indoor fine particles was infiltration of outdoor particles and that cooking activities were the only other significant source of fine particles. Coarse particles, however, had several indoor sources such as cooking, cleaning, and various indoor activities. This study also concluded that more than 50% of the particles (by volume) generated during indoor events were ultrafine particles. Events that elevated indoor particle levels were found to be brief, intermittent, and highly variable, thus requiring the use of continuous instrumentation for their characterization. Because the concentration of ultrafine particles is greater near the source (they grow in size into the accumulation mode as they age), the personal cloud for ultrafine particles may be higher than for accumulation mode particles if the person is nearer the source than the indoor monitor. Table 5-11 provides information on the mean volume mean diameter (VMD) for various types of indoor particle sources. The differences in mean VMD confirm the clear separation of source types and suggest that there is very little resuspension of accumulation-mode PM. In addition, measurements of OC and EC indicated that OC had significant indoor sources whereas EC was primarily of ambient origin.

Vette et al. (2001) found that resuspension was a significant indoor source of particles $> 1 \mu\text{m}$. Concentrations of fine particles were not affected by resuspension. Figure 5-12 shows the diurnal variability in the indoor/outdoor aerosol concentration ratio from an unoccupied residence in Fresno, CA. The study was conducted in the absence of common indoor particle sources such as cooking and cleaning. The data in Figure 5-12 show the mean indoor/outdoor concentration ratio for particles $> 1 \mu\text{m}$ increased dramatically during daytime hours. This pattern was consistent with indoor human activity levels. In contrast, the mean indoor/outdoor concentration ratio for particles $< 1 \mu\text{m}$ (ultrafine and accumulation-mode particles) remain fairly constant during both day and night.

Wallace and Howard-Reed (2002) used three instruments (SMPS, APS, and Climet) to measure ultrafine, fine, and coarse particles in an inhabited residence for 18 months. They confirmed the observations of Abt et al. (2000a) and Long et al. (2000) that indoor sources primarily generate ultrafine and coarse particles. Wallace and Howard-Reed reported that “Indoor sources affecting ultrafine particle concentrations were observed 22% of the time, and

TABLE 5-11. VOLUME MEAN DIAMETER (VMD) AND MAXIMUM PM_{2.5} CONCENTRATIONS OF INDOOR PARTICLE SOURCES^{a,b}

Particle Source	n	Size Statistics		PM _{2.5}	
		Indoor Activity Mean VMD (µm)	Background ^{a,c} Mean VMD (µm)	Maximum Concentration ^{c,d} Mean (µg/m ³)	SD (µg/m ³)
Cooking					
Baking (Electric)	8	0.189 ^f	0.221 ^f	14.8	7.4
Baking (Gas)	24	0.107 ^f	0.224 ^f	101.2	184.9
Toasting	23	0.138 ^f	0.222 ^f	54.9	119.7
Broiling	4	0.114 ^f	0.236 ^f	29.3	43.4
Sautéing	13	0.184 ^f , 3.48 ^g	0.223 ^f , 2.93 ^g	65.6	95.4
Stir-Frying	3	0.135 ^f	0.277 ^f	37.2	31.4
Frying	20	0.173 ^f	0.223 ^f	40.5	43.2
Barbecuing	2	0.159 ^f	0.205 ^f	14.8	5.2
Cleaning					
Dusting	11	5.38 ^g	3.53 ^g	22.6	22.6
Vacuuming	10	3.86 ^g	2.79 ^g	6.5	3.9
Cleaning with Pine Sol	5	0.097 ^f	0.238 ^f	11	10.2
General Activities					
Walking Vigorously (w/Carpet)	15	3.96 ^g	3.18 ^g	12	9.1
Sampling w/Carpet	52	4.25 ^g	2.63 ^g	8	6.6
Sampling w/o Carpet	26	4.28 ^g	2.93 ^g	4.8	3
Burning Candles	7	0.311 ^f	0.224 ^f	28	18

Notes:

^a All concentration data corrected for background particle levels.

^b Includes only individual particle events that were unique for a given time period and could be detected above background particle levels.

^c PM concentrations in µg/m³.

^d Maximum concentrations computed from 5-min data for each activity.

^e Background data are for time periods immediately prior to the indoor event.

^f Size statistics calculated for PV_{0.02-0.5} using SMPS data.

^g Size statistics calculated for PV_{0.7-10} using APS data.

Source: Long et al. (2000).




Figure 5-12. Mean hourly indoor/outdoor particle concentration ratio from an unoccupied residence in Fresno, CA during spring 1999.

Source: Vette et al. (2001).

sources affecting fine and coarse particle concentrations were observed 12 and 15% of the time, respectively . . . Indoor sources, such as cooking with natural gas, and simple physical activities, such as walking, accounted for a majority (50-90%) of the ultrafine and coarse particle concentrations, whereas outdoor sources were more important for accumulation-mode particles between 0.1 and 1 μm in diameter.”

5.3.3.2.3 Time-Activity Patterns

Total exposure to PM is the sum of various microenvironmental exposures that an individual encounters during the day and will depend on the microenvironments occupied. As discussed previously, PM exposure in each microenvironment is the sum of exposures from ambient PM (in outdoor and indoor microenvironments), indoor-generated PM, and indoor-reaction PM. In addition, there is exposure to PM generated by personal activities that is determined by the specific personal activities that the individual conducts while in those

microenvironments. As mentioned before, PM exposures and their components are variable across the population and, thus, each are distributions rather than point estimates. A thorough analysis of these distributions would require a comprehensive sensitivity and uncertainty analysis.

Determining microenvironments and activities that contribute significantly to human exposure begins with establishing human activity pattern information for the general population as well as for subpopulations. Personal exposure and time-activity pattern studies have shown that different populations have varying time-activity patterns and, accordingly, different personal PM exposures. Both characteristics will vary greatly as a function of age, health status, ethnic group, socioeconomic status, season, and region of the country. Collecting detailed time-activity data can be very burdensome on participants but is clearly valuable in assessing human exposure and microenvironments. For modeling purposes, human activity data frequently come from the general databases that are discussed below.

The gathering of human activity information, often called “time-budget” data, started in the 1920s; however, their use for exposure assessment purposes only began to be emphasized in the 1980s. As noted earlier, many of the largest U.S. human activity databases have been consolidated by the EPA’s National Exposure Research Laboratory (NERL) into one comprehensive database (CHAD) containing more than 22,000 person-days of 24-h activity (Glen et al., 1997; McCurdy et al., 2000). The information in CHAD is accessible for constructing population cohorts of people with diverse characteristics that are useful for analysis and modeling (McCurdy, 2000). See Table 5-2 for a summary listing of human activity studies in CHAD. Most of the databases in CHAD are also available elsewhere, including the National Human Activity Pattern Survey (NHAPS), California’s Air Resources Board (CARB), and the University of Michigan’s Institute for Survey Research data sets.

Although CHAD provides a very valuable resource for time and location data, there is little information on PM-generating personal activities. In addition, very few of the time-activity studies have collected longitudinal data within a season or over multiple seasons. Such longitudinal data are important in understanding potential variability in activities and how they affect correlations between PM exposure and ambient site measurements for both total PM and PM of ambient origin.

5.3.3.3 Effect of Ambient Sources on Exposures to Particulate Matter

Different sources may generate ambient PM with different aerodynamic and chemical characteristics, which may, in turn, result in different health responses. Thus, to fully understand the relationship between PM exposure and health outcomes, exposure from different sources should be identified and quantified. Source apportionment techniques provide a method for determining personal exposure to PM from specific sources. Daily contributions from sources that have no indoor component can be used as tracers to generate exposure estimates for ambient PM of similar aerodynamic size or used directly as exposure surrogates in epidemiological analyses. The most recent EPA PM Research Needs Document (U. S. Environmental Protection Agency, 1998) recommended using source apportionment techniques to determine daily time series of source categories for use in community, time-series epidemiology.

A number of epidemiologic studies (discussed more fully in Chapter 8) have evaluated relationships between health outcomes and sources of PM determined from measurements at a community monitor. These studies suggest the importance of examining sources and constituents of indoor, outdoor, and personal PM. For example, Özkaynak and Thurston (1987) evaluated the relationship between PM sources and mortality in 36 Standard Metropolitan Statistical Areas (SMSAs). Particulate matter samples from the EPA's Inhalable Particle (IP) Network were analyzed for SO_4^{2-} and NO_3^- by automated colorimetry, and elemental composition was determined with X-ray fluorescence (XRF). Mass concentrations from five PM source categories were determined from multiple regression of absolute factor scores on the mass concentration: (1) resuspended soil, (2) auto exhaust, (3) oil combustion, (4) metals, and (5) coal combustion.

In another study, Mar et al. (2000) applied factor analysis to evaluate the relationship between PM composition (and gaseous pollutants) in Phoenix, AZ. In addition to daily averages of $\text{PM}_{2.5}$ elements from XRF analysis, they included in their analyses OC and EC in $\text{PM}_{2.5}$ and gaseous species emitted by combustion sources (CO , NO_2 , and SO_2). They identified five source factors classified as (1) motor vehicles, (2) resuspended soil, (3) vegetative burning, (4) local SO_2 , and (5) regional sulfate. Additionally, Laden et al. (2000) applied specific rotation factor analysis to XRF PM composition data from six eastern cities (Ferris et al., 1979). Fine PM was regressed on the recentered scores to determine the daily source contributions. Three main

sources were identified: (1) resuspended soil (Si), (2) motor vehicle (Pb), and (3) coal combustion (Se).

Source apportionment or receptor modeling has been applied to the personal exposure data to understand the relationship between personal and ambient sources of PM. Application of source apportionment to ambient, indoor, and personal PM composition data is especially useful in sorting out the effects of particle size and composition. If a sufficient number of samples are analyzed with sufficient compositional detail, it is possible to use statistical techniques to derive source category signatures, to identify indoor and outdoor source categories, and to estimate their contribution to indoor and personal PM.

Positive Matrix Factorization (PMF) has been applied to the PTEAM database by Yakovleva et al. (1999). The authors utilized mass and XRF elemental composition data from indoor and outdoor PM_{2.5} and personal, indoor, and outdoor PM₁₀ samples. PMF is an advance over ordinary factor analysis, because it allows measurements below the quantifiable limit to be used by weighting them according to their uncertainty. This effectively increases the number of species that can be used in the model. The factors used by the authors correspond to general source categories of PM, such as outdoor soil, resuspended indoor soil, indoor soil, personal activities, sea salt, motor vehicles, nonferrous metal smelters, and secondary sulfates. PMF, by identifying the various source factors and apportioning them among the different monitor locations (personal, indoor, and outdoor), was able to estimate the contribution of resuspended indoor dust to the personal cloud (15% from indoor soil and 30% from resuspended indoor soil). Factor scores for these items then were used in a regression analysis to estimate personal exposures (Yakovleva et al., 1999).

The most important contributors to PM₁₀ personal exposure were indoor soil, resuspended indoor soil, and personal activities; these accounted for approximately 60% of the mass (Yakovleva et al., 1999). Collectively, they include personal-cloud PM, smoking, cooking, and vacuuming. For both PM_{2.5} and PM₁₀, secondary sulfate and nonferrous metal operations accounted for another 25% of PM mass. Motor vehicle exhausts, especially from vehicles started inside attached garages, accounted for another 10% of PM mass. The PTEAM study was conducted in Riverside, CA in the fall of 1990. Yakovleva et al. (1999) cautioned that their results may not apply to other geographic areas, seasons of the year, or weather conditions.

Simultaneous measurement of personal (PM_{10}) and outdoor measurements ($PM_{2.5}$ and PM_{10}) were evaluated as a three-way problem with PMF, which allowed for differentiation of source categories based on their variation in time and type of sample, as well as their variation in composition. By using this technique, three sources of coarse-mode, soil-type PM were identified. One was associated with ambient soil, one with indoor soil dispersed throughout the house, and one with soil resulting from the personal activity of the subject.

Two other source apportionment models have been applied to ambient measurement data and can be used for the personal exposure studies. The effective variance weighted Chemical Mass Balance (CMB) receptor model (Watson et al., 1984, 1990, 1991) solves a set of linear equations that incorporate the uncertainty in the sample and source composition. CMB requires the composition of each potential source of PM and the uncertainty for the sources and ambient measurements. Source apportionment with CMB can be conducted on individual samples; however, composition of each of the sources of PM must be known. An additional source apportionment model, UNMIX (Henry et al., 1994) is a multivariate source apportionment model. UNMIX is similar to PMF, but does not explicitly use the measurement uncertainties. Because measurement uncertainties are not used, only species above the detection limit are evaluated in the model. UNMIX provides the number of sources and source contributions and requires a similar number of observations as PMF.

The Yakovleva et al. (1999) study demonstrated that source apportionment techniques also could be very useful in determining parameters needed for exposure models and for determining exposure to ambient PM. Exposure information, similar to that obtained in the PTEAM study, but including other PM components useful for definition of other source categories (e.g., OC, EC; organic tracers for EC from diesel vehicle exhaust, gasoline vehicle exhaust, and wood combustion; nitrate; Na, Mg, and other metal tracers; and gas-phase pollutants) would be useful as demonstrated in the use of EC/OC and gas-phase pollutants by Mar et al. (2000).

5.3.3.4 Correlations of Particulate Matter with Other Pollutants

Correlations between ambient concentrations and between ambient concentrations and personal exposures for PM and other pollutants are of importance in understanding possible confounding in epidemiological studies and are discussed more fully in Chapter 9. Available information from exposure studies is presented in this section. Several epidemiologic studies

have included the gaseous pollutants CO, NO₂, SO₂, and O₃ along with PM₁₀ or PM_{2.5} in the analysis of the statistical association of health responses with pollutants. In a recent study, the personal exposure to O₃ and NO₂ was determined, as well as that to PM_{2.5} and PM_{10-2.5} for a cohort of 15 elderly subjects in Baltimore, MD, although measured personal exposures to O₃, NO₂, and SO₂ were below their respective level of detection (LOD) for 70% of the samples. Spearman correlations for 14 subjects in summer and 14 subjects in winter are given in Table 5-12 for relationships between personal PM_{2.5} and ambient concentrations of PM_{2.5}, PM_{10-2.5}, O₃, and NO₂. In contrast to ambient concentrations, neither personal exposure to total PM_{2.5} nor to ambient PM_{2.5} was correlated significantly with personal exposures to the co-pollutants, nonambient PM_{2.5} and ambient PM_{10-2.5}, O₃, NO₂, and SO₂. Personal-ambient associations for PM_{10-2.5}, O₃, NO₂, and SO₂ were similarly weak and insignificant. Based on these results, Sarnat et al. (2000) concluded that the potential for confounding of PM_{2.5} by O₃, NO₂, or PM_{10-2.5} appears to be limited, because, despite significant correlations observed among ambient pollutant concentrations, the correlations among personal exposures were low.

Sarnat et al. (2001) further evaluated the role of gaseous pollutants in PM epidemiology by extending the measurements taken on the earlier adult cohort of 20 individuals in Baltimore, MD through inclusion of additional PM and gaseous pollutant measurements that were collected during the same 1998 to 1999 period from 15 individuals with COPD and from 21 children. Twenty-four-hour average personal exposures for PM_{2.5}, O₃, SO₂ and NO₂, and corresponding ambient concentrations for PM_{2.5}, O₃, SO₂, NO₂ and CO for all 56 subjects were collected over 12 consecutive days. Results from correlation and regression analysis of the personal and ambient data showed that personal PM_{2.5} and personal gaseous pollutant exposures were generally not correlated. The analysis also showed that ambient PM_{2.5} concentrations had significant associations with personal PM_{2.5} exposures in both seasons. On the other hand, ambient gaseous pollutant concentrations were not correlated with their corresponding personal exposure concentrations. However, ambient gaseous concentrations were found to be strongly associated with personal PM_{2.5} exposures, suggesting that ambient gaseous concentrations for O₃, NO₂, and SO₂ are acting as surrogates, as opposed to confounders, of PM_{2.5} in the estimation of PM health effects based on multipollutant models. This study did not measure personal CO and did not find a significant association between summertime ambient CO and personal PM_{2.5} (a significant wintertime association, however, was found).

TABLE 5-12. CORRELATIONS BETWEEN PERSONAL PM_{2.5} AND AMBIENT POLLUTANT CONCENTRATIONS¹

	Subject	Personal PM _{2.5} vs. Ambient				Personal PM _{2.5} of Ambient Origin vs. Ambient		
		PM _{2.5}	O ₃	NO ₂	PM _{2.5-10}	O ₃	NO ₂	PM _{10-2.5}
SUMMER	SA1	0.55	0.15	0.38	-0.12	0.27	<i>0.71</i>	0.15
	SA2	<i>0.85</i>	0.31	<i>0.66</i>	0.57	0.21	0.64	0.68
	SA5	<i>0.89</i>	0.18	<i>0.82</i>	0.64	0.33	<i>0.81</i>	0.79
	SB1	<i>0.65</i>	0.40	-0.15	0.38	<i>0.89</i>	<i>-0.74</i>	-0.03
	SB2	-0.21	-0.62	<i>0.81</i>	0.15	0.26	0.08	0.33
	SB3	<i>0.82</i>	0.55	-0.14	-0.04	0.52	-0.20	0.00
	SB4	<i>0.73</i>	<i>0.62</i>	-0.34	-0.12	0.45	-0.29	-0.14
	SB5	<i>0.73</i>	0.45	-0.42	0.23	0.36	-0.48	0.33
	SB6	0.53	0.15	-0.38	0.12	-0.03	<i>-0.57</i>	0.32
	SC1	<i>0.95</i>	<i>0.78</i>	<i>0.66</i>	<i>0.65</i>	<i>0.83</i>	<i>0.63</i>	0.57
	SC2	<i>0.78</i>	<i>0.68</i>	0.36	0.51	<i>0.66</i>	<i>0.65</i>	<i>0.76</i>
	SC3	<i>0.85</i>	<i>0.78</i>	<i>0.73</i>	<i>0.68</i>	<i>0.69</i>	<i>0.71</i>	<i>0.80</i>
	SC4	<i>0.78</i>	<i>0.66</i>	0.59	<i>0.70</i>	0.50	0.50	0.51
	SC5	0.55	0.51	0.32	0.43	0.34	0.33	0.27
	WINTER	WA1	0.22	-0.18	-0.26	-0.05	<i>-0.78</i>	-0.04
WA2		-0.38	-0.07	-0.36	-0.70	-0.15	-0.15	0.02
WA4		-0.18	<i>0.67</i>	-0.22	-0.29	-0.33	0.20	0.00
WA5		0.22	-0.43	<i>0.61</i>	0.50	<i>-0.72</i>	-0.09	0.40
WB1		<i>0.80</i>	-0.84	<i>0.77</i>	0.41	<i>-0.87</i>	0.53	<i>0.66</i>
WB2		<i>0.62</i>	-0.32	<i>0.59</i>	0.09	<i>-0.76</i>	<i>0.59</i>	<i>0.59</i>
WB3		0.55	-0.45	<i>0.62</i>	0.04	<i>-0.77</i>	0.56	0.60
WB4		-0.12	-0.01	0.34	-0.10	<i>-0.80</i>	<i>0.68</i>	0.48
WC1		<i>0.74</i>	-0.62	-0.15	0.44	<i>-0.64</i>	0.02	0.69
WC2		<i>0.79</i>	-0.88	0.17	<i>0.77</i>	<i>-0.87</i>	0.25	<i>0.71</i>
WC3		0.28	-0.42	0.03	0.57	<i>-0.77</i>	0.30	-0.45
WC4		0.19	-0.84	0.50	0.45	<i>-0.72</i>	0.22	0.67
WC5		<i>0.81</i>	-0.62	0.08	<i>0.81</i>	<i>-0.76</i>	0.05	0.42
WC6		0.01	-0.03	<i>0.65</i>	0.37	<i>-0.75</i>	0.19	-0.45
Median	Summer	0.76	0.48	0.37	0.41	0.41	0.42	0.33
Median	Winter	0.25	-0.43	0.26	0.39	-0.76	0.21	0.45

¹Correlations represent Spearman's r values; italicized values indicate significance at the p < 0.05 level.

Source: Samat et al. (2000).

Personal EC and SO_4^{2-} were also measured during the winter for the cohort of COPD patients only. The analysis of this subset of the data showed that personal SO_4^{2-} was significantly and negatively associated with ambient O_3 and SO_2 , and that personal EC was significantly associated with ambient O_3 , NO_2 , and CO. The authors interpret these findings as suggesting that O_3 is primarily a surrogate for secondary particle exposures, whereas ambient CO and NO_2 are primarily surrogates for particles from traffic. Sarnat et al. (2001) cautioned that these findings were from only one location and various physical and personal factors, such as ventilation, time spent outdoors, and household characteristics, could affect the strength of the reported associations for certain individuals and cohorts even though the qualitative results found were unlikely to change.

A newly developed Roll-Around System (RAS) was used to evaluate the hourly relationship between gaseous pollutants (CO, O_3 , NO_2 , SO_2 , and VOCs) and PM (Chang et al., 2000). Exposures were characterized over a 15-day period for the summer and winter in Baltimore, MD based on scripted activities to simulate activities performed by older adults (65+ years of age). Spearman rank correlations were reported for $\text{PM}_{2.5}$, O_3 , CO, and toluene for both the summer and winter. The correlations are given for each microenvironment in Table 5-13: indoor residence, indoor other, outdoor near roadway, outdoor away from road, and in vehicle. No significant relationships ($p < 0.05$) were found between hourly $\text{PM}_{2.5}$ and O_3 . Significant relationships were found between hourly $\text{PM}_{2.5}$ and CO in: indoor residence, winter; indoor other, summer and winter; and outdoor away from roadway, summer. Significant relationships also were found between hourly $\text{PM}_{2.5}$ and toluene in: indoor residence, winter; indoor other, winter; and in vehicle, winter. The significant relationships between CO and $\text{PM}_{2.5}$ in the winter may be caused by reduced air exchange rates that could allow them to accumulate (Chang et al., 2000). Note that, although no significant correlation was found between in-vehicle $\text{PM}_{2.5}$ and CO, another significant component of vehicle exhaust, toluene (Conner et al., 1995), was significantly correlated to $\text{PM}_{2.5}$ in the winter.

Carrer et al. (1998) presented data on the correlations among microenvironmental and personal PM_{10} exposures and concentrations and selected environmental chemicals monitored simultaneously (using methods not described). These chemicals were nitrogen oxides (NO_x), carbon monoxide (CO), and total volatile organic compounds (TVOC), benzene, toluene, xylene,

TABLE 5-13. CORRELATIONS BETWEEN HOURLY PERSONAL PM_{2.5} AND GASEOUS POLLUTANTS

	Indoor Residence		Indoor Other		Outdoor Near Roadway		Outdoor Away from road		In Vehicle	
	N	r _s	N	r _s	N	r _s	N	r _s	N	r _s
PM_{2.5} vs. O₃										
Summer	35	0.29	16	-0.14	10	0.05	12	0.45	37	0.21
Winter	56	0.05	37	-0.06	11	-0.28	7	0.04	34	-0.10
PM_{2.5} vs. CO										
Summer	41	0.25	19	0.59 ^a	13	0.14	12	0.62	46	0.23
Winter	59	0.43 ^a	39	0.62 ^a	13	0.37	8	0.41	37	0.1
PM_{2.5} vs. Toluene										
Summer	46	0.23	21	-0.14	14	0.26	14	0.02	48	0.12
Winter	66	0.38 ^a	47	0.44 ^a	17	0.4	8	0.48	42	0.43 ^a

^aCorrelations represent Spearman's r values; italicized values indicate significance at the p < 0.05 level.

Source: Chang et al. (2000).

and formaldehyde. The Kendall τ correlation coefficient was used; only results significant at p < 0.05 are mentioned here. Significant associations were found only between the following pairs of substances (τ shown in parentheses): personal PM₁₀ (24 h) and NO_x (0.34), CO (0.34), TVOC (0.18), toluene (0.19), and xylene (0.26); office PM₁₀ and NO_x (0.31); home PM₁₀ and NO_x (0.24), CO (0.24), toluene (0.17), and xylene (0.25). Even though most of these chemicals are associated with motor vehicular emissions, there was no significant correlation between commuting PM₁₀ and any of the substances.

5.4 SUMMARY OF PARTICULATE MATTER CONSTITUENT DATA

5.4.1 Introduction

Atmospheric PM contains a number of chemical constituents that may be of significance to human exposure and health effects. These constituents may be either components of the ambient particles or bound to the surface of particles. They may be elements, inorganic species, or organic compounds. A limited number of studies have collected data on concentrations of elements, acidic aerosols, and PAHs in ambient, personal, and microenvironmental PM samples. However, there have not been extensive analyses of the constituents of PM in personal or microenvironmental samples. Data from relevant studies are summarized in this section. The summary does not address bacteria, bioaerosols, viruses, or fungi (e.g., Owen et al., 1992; Ren et al., 1999).

5.4.2 Monitoring Studies That Address Particulate Matter Constituents

Relevant studies published in recent years that have measured the constituents of PM in personal or microenvironmental samples are summarized in Tables 5-11 and 5-12 for personal exposure measurements of PM and microenvironmental samples, respectively. Studies that measured both personal and microenvironmental samples are included in Table 5-11.

The largest database on personal, microenvironmental, and outdoor measurements of PM elemental concentrations is the PTEAM study (Özkaynak et al., 1996b). The results are highlighted in the table and discussed below. The table shows that a number of studies have measured concentrations of elements (by XRF), organic carbon (OC), various indicators of elemental carbon (EC), aerosol acidity, sulfate, ammonia, and nitrate. Additionally, a number of studies have measured PAHs, both indoors and outdoors. Other than the PAHs, there are few data on the organic constituents of PM.

5.4.3 Key Findings

5.4.3.1 Correlations of Personal and Indoor Concentrations with Ambient Concentrations of Particulate Matter Constituents

The elemental composition of PM₁₀ in personal samples was measured in the PTEAM study, the first probability-based study of personal exposure to particles. A number of important observations made from the PTEAM data collected in Riverside, CA are summarized by

Özkaynak et al. (1996b). Population-weighted daytime personal exposures averaged $150 \pm 9 \mu\text{g}/\text{m}^3$ compared to concurrent indoor and outdoor concentrations of $95 \pm 6 \mu\text{g}/\text{m}^3$. The personal exposure measurements suggested that there was a “personal cloud” of particles associated with personal activities. Daytime personal exposures to 14 of the 15 elements measured in the samples were considerably greater than concurrent indoor or outdoor concentrations; sulfur was the only exception.

The PTEAM data also showed good agreement between the concentrations of the elements measured outdoors in the backyard of the residences with the concentrations measured at the central site in the community. The agreement was excellent for sulfur. Although the particle and element mass concentrations were higher in personal samples than for indoor or outdoor samples, a nonlinear mass-balance method showed that the penetration factor was nearly 1 for all particles and elements.

Measurements of element concentrations in NHEXAS are similar to the PTEAM results in that they show elevated concentrations of arsenic (As) and lead (Pb) in personal samples relative to indoor and outdoor samples (Clayton et al., 1999b). The elevated concentrations of As and Pb were consistent with elevated levels of PM_{50} in personal samples (median particle exposure of $101 \mu\text{g}/\text{m}^3$) as compared to indoor concentrations ($34.4 \mu\text{g}/\text{m}^3$). There was a strong association between personal and indoor concentrations and between indoor and outdoor concentrations for both As and Pb. However, there were no central site ambient measurements for comparison to the outdoor or indoor measurements at the residences.

Manganese (Mn) concentrations were measured in $\text{PM}_{2.5}$ samples collected in Toronto (Crump, 2000). The mean $\text{PM}_{2.5}$ Mn concentrations were higher outdoors than indoors. However, the outdoor concentrations measured at the participants’ homes were lower than those measured at two fixed locations. Crump (2000) suggested that the difference in the concentrations may have been because the fixed locations were likely closer to high-traffic areas than were the participants’ homes.

Studies of acidic aerosols and gases typically measure strong acidity (H^+), SO_4^{2-} , NH_4^+ , and NO_3^- . The relationship between the concentrations of these ions and the relationship between indoor and outdoor concentrations have been addressed in a number of studies during which personal, microenvironmental, and outdoor samples have been collected, as shown in Tables 5-14 and 5-15. Key findings from these studies include the following:

TABLE 5-14. STUDIES THAT HAVE MEASURED PARTICULATE MATTER CONSTITUENTS IN PERSONAL EXPOSURE SAMPLES

PM Constituent	Study Name/Reference	Study Location	Population Size/No. of Samples	Summary of Results
Elements	PTEAM/Özkaynak et al. (1996b)	Riverside, CA	178 adults	Outdoor air was the major source for most elements indoors, providing 70 to 100% of the observed indoor concentrations for 12 of the 15 elements. Correlation coefficients for central monitoring site vs. outdoor at the residences were 0.98 for S and 0.5 to 0.9 for other elements (except Cu).
As and Pb	NHEXAS/Clayton et al. (1999b)	EPA Region 5	167 samples	Personal As and Pb levels higher than indoor or outdoor levels. No community ambient site for comparison.
Mn	Pellizzari et al. (1998, 1999); Clayton et al. (1999a); Crump (2000)	Toronto	925 personal samples	Mean PM _{2.5} Mn higher outdoors than indoors. But PM _{2.5} Mn concentrations higher at two fixed locations than at participants' homes.
Acid Aerosol Constituents	Sarnat et al. (2000)	Baltimore, MD	20 adults	High correlations between personal and ambient sulfate measurements in summer and winter.
	Brauer et al. (1989)	Boston, MA	—	Personal exposures to aerosol strong acidity slightly lower than concentrations measured at stationary site.
	Suh et al. (1992)	Uniontown, PA	24 children for 2 days	Personal exposures to H ⁺ and SO ₄ ²⁻ lower than outdoor levels, but higher than indoor microenvironmental levels; personal NH ₄ ⁺ and NO ₃ ⁻ higher than indoor or outdoor levels.
	Suh et al. (1993a,b)	State College, PA	47 children	Results similar to Uniontown, PA study.
	Suh et al. (1994)			Results indicate strong neutralization of acidity indoors.
	Waldman and Liang (1993); Waldman et al. (1990)	Georgia and New Jersey	Hospital, day cares	Indoor sulfate levels were 70 to 100% of outdoor levels. Indoor ammonia levels 5- to 50-times higher than outdoors. Indoors, acid aerosols were largely neutralized.
PAHs	Zmirou et al. (2000)	Grenoble, France	38 adults	Ambient air concentrations close to traffic emissions were 1.1- to 3.5-times higher than personal exposure concentrations.
Individual Particle Analyses by CCSEM	Conner et al. (2001)	Baltimore, MD Same study locations for both	3 sets of indoor-outdoor-personal filters, > 2000 particles/filter	CCSEM was used to identify individual particles and classify them by most likely sources.
Trace Elemental analyses by XRF	Landis et al. (2001)		19 days with P, I, A for 10 elderly retirees	P and A highly correlated for PM _{2.5} and sulfates. P and A not significantly correlated for soil and trace element oxides.
Elemental analysis by HR-ICP-MS	Kinney et al. (2002)	New York City, NY	46 student volunteers, 1 week in summer and winter, PM _{2.5}	29 PM constituents were measured. Most were similar for P, I, and O, suggesting outdoor sources dominant.

TABLE 5-15. STUDIES THAT HAVE MEASURED PARTICULATE MATTER CONSTITUENTS IN MICROENVIRONMENTAL SAMPLES

PM Constituent	Study Name/Reference	Study Location	Population Size/No. of Samples	Summary of Results
Acid Aerosol Constituents	Jones et al. (2000)	Birmingham, England	12 residences	Sulfate I/O ratios ranged from 0.7 to 0.9 for three PM size fractions.
	Patterson and Eatough (2000)	Lindon, UT	1 school	Ambient sulfate, SO ₂ , nitrate, soot, and total particle number showed strong correlations with indoor exposure although ambient PM _{2.5} mass was not a good indicator of total PM _{2.5} exposure.
	Leaderer et al. (1999)	Virginia and Connecticut	232 homes	The regional ambient air monitoring site provided a reasonable estimate of indoor and outdoor sulfate at nonsmokers homes. I/O sulfate ratio of 0.74 during summer. Ammonia concentrations were an order of magnitude higher indoors than outdoors. Nitrous acid levels higher indoors than outdoors.
	Brauer et al. (1990)	Boston, MA	11 homes	Outdoor levels of H ⁺ , SO ₂ , HNO ₃ , and SO ₄ ²⁻ exceeded indoor levels in winter and summer. I/O ratios of H ⁺ lower than I/O ratios of SO ₄ ²⁻ indicated neutralization of the acidity by ammonia.
PAHs	Chuang et al. (1999)	Durham, NC	24 homes	Measurements with continuous monitor; PAH levels generally higher indoors than outdoors.
	Dubowsky et al. (1999)	Boston, MA	3 buildings	PAHs indoors attributable to traffic, cooking, and candle-burning.
	Sheldon et al. (1993a,b)	Placerville and Roseville, CA	280 homes	Mass balance model used to estimate source strengths for PAH sources such as smoking, wood-burning and cooking.
PAHs and phthalates	PTEAM/Özkaynak et al. (1996b), Sheldon et al. (1993c)	Riverside, CA	120 homes	12-h I/O ratios for particulate-phase PAHs ranged from 1.1 to 1.4 during the day and 0.64 to 0.85 during night. The concentrations of phthalates and the number of samples with detectable phthalates were higher indoors than outdoors.

5-97

Abbreviations used:
 I/O = Indoor/Outdoor.
 A = ambient.
 I = indoor.
 O = outdoor.
 P = personal.

- Acid aerosol concentrations measured at the residences in the Uniontown, PA study were significantly different from those measured at a fixed ambient site located 16 km from the community. However, Leaderer et al. (1999) reported that the regional ambient air monitoring site in Vinton, VA provided a reasonable estimate of indoor and outdoor sulfate measurements during the summer at homes without tobacco combustion.
- Approximately 75% of the fine aerosol indoors during the summer was associated with outdoor sources based on I/O sulfate ratios measured in the Leaderer et al. (1999) study.
- Personal exposures to strong acidity (H^+) were lower than corresponding outdoor levels measured in studies by Brauer et al. (1989, 1990) and Suh et al. (1992). However, the personal exposure levels measured by Suh et al. (1992) were higher than the indoor microenvironmental levels.
- Personal exposures to NH_4^+ and NO_3^- were reported by Suh et al. (1992) to be higher than either indoor or outdoor levels.

Personal exposures to SO_4^{2-} were also lower than corresponding outdoor levels, but higher than the indoor microenvironmental levels (Suh et al., 1992; 1993a,b) as shown in Table 5-16.

The fact that the personal and indoor H^+ concentrations were substantially lower than outdoor concentrations suggests that a large fraction of aerosol strong acidity is neutralized by ammonia. Ammonia is emitted in relatively high concentrations in exhaled breath and sweat. The difference between indoor and outdoor H^+ concentrations in the Suh et al. (1992, 1993a,b) studies was also much higher than the difference for indoor and outdoor SO_4^{2-} , indicative of neutralization of the H^+ . Results of the Suh et al. (1992, 1993a,b) studies also showed substantial interpersonal variability of H^+ concentrations that could not be explained by variation in outdoor concentrations.

Similar results for ammonia were reported by Waldman and Liang (1993). They reported that levels of ammonia in monitored institutional settings were 10- to 50-times higher than outdoors and that acid aerosols were largely neutralized. Leaderer et al. (1999) reported that ammonia concentrations during both winter and summer in residences were an order of magnitude higher indoors than outdoors, consistent with results of other studies and the presence of sources of ammonia indoors.

Sulfate aerosols appear to penetrate indoors effectively. Waldman et al. (1990) reported I/O ratios of 0.7 to 0.9 in two nursing care facilities and a day-care center. Sulfate I/O ratios were measured for three particle size fractions in 12 residences in Birmingham, England by Jones et al. (2000). The sulfate I/O ratios were 0.7 to 0.9 for $PM < 1.1 \mu m$, 0.6 to 0.8 for PM

TABLE 5-16. SUMMARY STATISTICS FOR PERSONAL, INDOOR, AND OUTDOOR CONCENTRATIONS OF SELECTED AEROSOL COMPONENTS IN TWO PENNSYLVANIA COMMUNITIES

Aerosol	Home Type	Number of Samples (In/Out) ^a	Concentration (nmol m ⁻³)		
			Indoor (12 h) GM ± GSD ^b	Outdoor (24 h) GM ± GSD ^b	Personal (12 h) GM ± GSD ^b
<i>State College</i>					
NO ⁻³	A/C Homes ^c	53/71	2.1 ± 2.7	1.4 ± 2.1	—
	Non-A/C	254/71	3.2 ± 2.3	1.4 ± 2.1	—
SO ₄ ²⁻	A/C Homes	56/75	61.8 ± 2.5	109.4 ± 2.4	—
	Non-A/C	259/75	96.7 ± 2.5	109.4 ± 2.4	—
	All Homes ^d	214/76	69.1 ± 2.6	91.0 ± 2.5	71.5 ± 2.4
NH ₄ ⁺	All Homes	314/155	154.7 ± 2.8	104.4 ± 2.3	—
H ⁺	A/C Homes	28/74	4.2 ± 4.3	82.5 ± 2.6	—
	Non-A/C	230/74	11.2 ± 3.1	82.5 ± 2.6	—
	All Homes ^e	163/75	9.1 ± 3.5	72.4 ± 2.9	18.4 ± 3.0
<i>Uniontown</i>					
SO ₄ ²⁻	All Homes ^e	91/46	87.8 ± 2.1	124.9 ± 1.9	110.3 ± 1.8
NH ₄ ⁺	All Homes ^e	91/44	157.2 ± 2.8	139.4 ± 2.1	167.0 ± 2.0
H ⁺	All Homes ^e	91/46	13.7 ± 2.5	76.6 ± 2.7	42.8 ± 2.2

^aIn/Out = Indoor samples/outdoor samples.

^bGM ± GSD = Geometric mean ± geometric standard deviation.

^cA/C Homes = Homes that had air-conditioning (A/C); this does not imply that it was on during the entire sampling period.

Non-A/C = Homes without air conditioning.

^dThe sample size, n, for the personal monitoring = 209.

^en = 174 for personal monitoring.

Source: Suh et al. (1992, 1993a,b).

1.1 to 2.1 μm, and 0.7 to 0.8 for PM 2.1 to 10 μm. Suh et al. (1993b) reported that personal and outdoor sulfate concentrations were highly correlated, as depicted in Figure 5-13.

Patterson and Eatough (2000) reported indoor/outdoor relationships for a number of PM_{2.5} components and related species in Lindon, UT during January and February of 1997. Outdoor samples were collected at the Utah State Air Quality monitoring site. Indoor samples were collected in the adjacent Lindon Elementary School. The infiltration factors, C_{ai}/C, given by the




Figure 5-13. Personal versus outdoor SO_4^{2-} in State College, PA. Open circles represent children living in air-conditioned homes; the solid line is the 1:1 line.

Source: Suh et al. (1993b).

slope of the regression lines (Table 5-17), were low (0.27 for sulfate and 0.12 for $\text{PM}_{2.5}$) possibly because of removal of particles in the air heating and ventilation system. The authors concluded that indoor $\text{PM}_{2.5}$ mass may not always be a good indicator of exposure to ambient combustion material due to the influence of indoor particle sources. Presumably this occurs because the concentrations of indoor-generated particles are not well correlated with the concentrations of ambient combustion particles. However, ambient sulfate, SO_2 , nitrate, soot, and total particle number displayed strong correlations with indoor exposure, presumably because these species have few indoor sources in the absence of indoor combustion. Ambient $\text{PM}_{2.5}$ mass was not a good indicator of indoor $\text{PM}_{2.5}$ mass exposure, presumably due to uncorrelated indoor sources of $\text{PM}_{2.5}$ mass.

TABLE 5-17. REGRESSION ANALYSIS OF INDOOR VERSUS OUTDOOR CONCENTRATIONS (units are nmol/m³, except for soot and metals, which are µg/m³ and absorption units/m³, respectively.)^a

Species	Slope = F _{INF}	Intercept = \bar{C}_{ig}	R ²	Average Outdoors, C _o
SO ₂ All Samples	0.0272 ± 0.0023	0.34 ± 0.13	0.73	38
SO ₂ Day Samples	0.0233 ± 0.0037	0.75 ± 0.26	0.62	56
SO ₂ Night Samples	0.0297 ± 0.0029	0.099 ± 0.075	0.82	20
Sulfate All Samples	0.267 ± 0.024	-0.14 ± 0.48	0.7	16
Sulfate Day Samples	0.261 ± 0.034	0.40 ± 0.66	0.71	16
Sulfate Night Samples	0.282 ± 0.035	-0.84 ± 0.68	0.7	16
Nitrate All Samples	0.0639 ± 0.0096	0.9 ± 1.5	0.54	134
Nitrate Day Samples	0.097 ± 0.0096	-0.4 ± 1.4	0.88	126
Nitrate Night Samples	0.047 ± 0.011	1.5 ± 1.8	0.44	139
Soot Day Samples	0.43 ± 0.25	3.5 ± 1.7	0.43	6
Soot Night Samples	0.33 ± 0.13	0.00 ± 0.55	0.69	4
Total Acidity All Samples	0.04 ± 0.73	0.42 ± 0.23	0	0.2
Metals All Samples	0.10 ± 0.30	0.0014 ± 0.0042	0.01	0.0042

^aLindon Elementary School, Lindon, UT January and February 1997.

Source: Patterson and Eatough (2000).

Oglesby et al. (2000) conducted a study to evaluate the validity of fixed-site fine particle concentration measurements as exposure surrogates for air pollution epidemiology. Using 48-h EXPOLIS data from Basel, Switzerland, they investigated the personal exposure/outdoor concentration relationships for four indicator groups: (1) PM_{2.5} mass, (2) sulfur and potassium for regional air pollution, (3) lead and bromine for traffic-related particles, and (4) calcium for crustal particles. The authors reported that personal exposures to PM_{2.5} mass were not correlated to corresponding home outdoor levels (n = 44, r = 0.07). In the study group reporting neither relevant indoor sources nor relevant activities, personal exposures and home outdoor levels of sulfur were highly correlated (n = 40, r = 0.85). These results are consistent with spatially homogeneous regional pollution and higher spatial variability of traffic and crustal materials.

Indoor, outdoor, and personal concentrations of $PM_{2.5}$ mass and a variety of PM constituents were measured at an 18-story retirement facility in Towson, MD for 28 24-hour monitoring periods during July and August 1998 (Landis et al., 2001). Indoor and outdoor measurements were made with a Versatile Air Pollutant Sampler (VAPS). Indoor, outdoor, and personal samples were made with a Personal Exposure Monitor (PEM). The VAPS (a dichotomous sampler) collected $PM_{2.5}$ (15 L/min) and $PM_{10-2.5}$ (2 L/min) while the PEM collected $PM_{2.5}$ (2 L/min). A comparison of the VAPS and the PEM indicated that the indoor PEM collected much higher mass and more soil components than the indoor VAPS although the differences between outdoor results were smaller and not significant. These differences were attributed to the presence of a larger coarse particle concentration inside (and perhaps larger diameter particles) and either more particle bounce or a higher 50% cut point for the PEM. In their analysis, Landis et al. (2001) compared indoor and outdoor VAPS data, as well as outdoor and personal PEM data.

As shown in Table 5-18 (PEM) and Table 5-19 (VAPS), higher correlations were found for fine-particle components of $PM_{2.5}$ and lower correlations for coarse-particle components. Like Patterson and Eatough (2000), Landis et al. (2001) found low infiltration factors for nitrate along with a reasonable correlation suggesting that fine-mode ammonium nitrate may be evaporating after it penetrates indoors. Neither sulfate nor nitrate had indoor sources.

Indoor and outdoor $PM_{2.5}$ and PM_{10} mass and chemical composition were measured in 13 homes (2 to 4 days for each home) in the Coachella Valley, a unique desert area in southern California during the winter and spring of 2000 (Geller et al., 2002). Maximum infiltration of ambient PM would be expected during this period because the mild climate minimizes the use of heating or air conditioning. Regression analysis was used to estimate \bar{F}_{INF} and \bar{C}_{ig} . Results are shown in Table 5-20. The Coachella PM is generally considered to be rich in coarse PM and epidemiological studies have associated PM_{10} (Ostro et al., 1999) and estimated $PM_{10-2.5}$ (Ostro et al., 2000) with mortality. However, the results of Geller et al. (2002) indicate that even during periods of high air exchange rates, indoor exposures would be dominated by $PM_{2.5}$. Geller et al. (2002) also reported results for some chemical components of $PM_{2.5}$ and $PM_{10-2.5}$ as well as EC and OC in $PM_{2.5}$. For EC, $F_{INF}^{EC} = 0.74$, in good agreement with $F_{INF}^{2.5} = 0.74$. Indoor concentrations of OC were much higher than outdoor concentrations; the average I/O ratio was 1.77.

TABLE 5-18. MIXED MODEL ANALYSIS OF PERSONAL VERSUS OUTDOOR CONCENTRATIONS^a

	\bar{R}^2	$\bar{\alpha}$	\bar{C}	\bar{A}	\bar{N}	\bar{T}	$\bar{A} + \bar{N}$
PM _{2.5} (μg/m ³)	0.82	0.46 ± 0.04	21	9.66	3.57 ± 0.80	12.8	13.2
Sulfate ^b (μg/m ³)	0.95	0.40 ± 0.02	10.2	4.08	0.1 ± 0.04	4.5	4.2
Soil ^c (ng/m ³)	0.03	0.05 ± 0.12	458	93	544 ± 95	553	637
TEO ^d (ng/m ³)	0.39	0.43 ± 0.11	165	71	99 ± 23	170	170

^a $\bar{\alpha}$ (slope) and \bar{N} (intercept) taken from mixed model results. \bar{A} calculated as $\bar{\alpha} \cdot \bar{C}$; \bar{T} is measured value.

Note good agreement between measured \bar{T} and \bar{T} estimated from $\bar{A} + \bar{N}$ except for soil.

^b as (NH₄)₂SO₄.

^c Soil = sum of oxides Si, Ca, Fe, and Ti (due to coarse PM in PM_{2.5}).

^d TEO = trace element oxides (mostly fine PM components).

Source: Landis et al. (2001).

TABLE 5-19. REGRESSION ANALYSIS OF INDOOR VERSUS OUTDOOR CONCENTRATIONS^a

	R^2	\bar{F}_{INF}	\bar{C}	\bar{C}_{ai}	\bar{C}_{ig}	\bar{C}_i	$\bar{C}_{ai} + \bar{C}_{ig}$
PM _{2.5} (μg/m ³)	0.74	0.35	18.9	6.6	0.32	6.7	6.9
Sulfate ^b (μg/m ³)	0.98	0.41	10.4	4.4	-0.24	4	4.2
OC • 1.4 (μg/m ³)	0.44	0.3	5.4	1.6	8	9.7	9.6
EC ^c (μg/m ³)	0.58	0.32	0.5	0.16	0.09	0.4	0.25
NaCl (ng/m ³)	0.32	0.06	231	14	37	48	53
Soil ^d (ng/m ³)	0.3	0.07	363	25	51	74	76
TEO ^e (ng/m ³)	0.38	0.35	94	33	7	39	40
Nitrate (ng/m ³)	0.83	0.09	372	33	12	68	45

^a \bar{F}_{INF} (slope) and \bar{C}_{ig} (intercept) taken from regression equation. C_{ai} calculated as $\bar{C} \cdot \bar{F}_{INF}$; \bar{C}_i is measured value.

Note reasonable agreement between measured \bar{C}_i and C_i estimated from $\bar{C}_{ai} + \bar{C}_{ig}$ except for EC and nitrate.

^b as (NH₄)₂SO₄.

^c revised regression with outliers omitted.

^d Soil = sum of oxides Si, Ca, Fe, and Ti (due to coarse PM in PM_{2.5}).

^e TEO = trace element oxides (mostly fine PM components).

Source: Landis et al. (2001).

TABLE 5-20. REGRESSION ANALYSIS OF INDOOR VERSUS OUTDOOR CONCENTRATIONS^a

	R^2	\bar{F}_{INF}	\bar{C}	\bar{C}_{ai}	\bar{C}_{ig}	\bar{C}_i
Fine	0.37	0.74	15	11.1	4.3	15.4
Coarse	0.35	0.3	8.6	2.6	3	5.6

^a \bar{F}_{INF} (slope) and \bar{C}_{ig} (intercept) taken from regression equation. \bar{C}_{ai} calculated as $\bar{C} \cdot \bar{F}_{INF}$. \bar{C}_i is measured value. Note excellent agreement between measured \bar{C}_i and \bar{C}_i estimated from $\bar{C}_{ai} + \bar{C}_{ig}$.

Source: Geller et al. (2002).

PAHs have been measured in studies by EPA and the California Air Resources Board. PAH results from a probability sample of 125 homes in Riverside are discussed in reports by Sheldon et al. (1992a,b) and Özkaynak et al. (1996b). Data for two sequential 12-h samples were reported for PAHs by ring size (3 to 7) and for individual phthalates. The results can be summarized as follows.

- The particulate-phase 5- to 7-ring species had lower relative concentrations than the more volatile 3- to 4-ring species.
- The 12-h I/O ratios for the 5- to 7-ring species ranged from 1.1 to 1.4 during the day and from 0.64 to 0.85 during the night (Sheldon et al., 1993a).
- An indoor air model used to calculate indoor “source strengths” for the PAHs showed that smoking had the strongest effect on indoor concentrations.

Results from a larger PAH probability study in 280 homes in Placerville and Roseville, CA (Sheldon et al., 1993a,b) were similar to the 125-home study. The higher-ring, particle-bound PAHs had lower indoor and outdoor concentrations than the lower-ring species. For most PAHs, the I/O ratio was greater than 1 for smoking and smoking/fireplace homes and less than 1 for fireplace-only, wood stove, wood stove/gas heat, gas heat, and “no source” homes.

A study of PAHs in indoor and outdoor air was conducted in 14 inner-city and 10 rural low-income homes near Durham, NC in two seasons (winter and summer) in 1995 (Chuang et al., 1999). Fine-particle-bound PAH concentrations measured with a real-time monitor were usually higher indoors than outdoors (2.47 ± 1.90 versus $0.53 \pm 0.58 \mu\text{g}/\text{m}^3$). Higher indoor

levels were seen in smokers' homes compared with nonsmokers' homes, and higher outdoor and indoor PAH levels were seen in urban areas compared with rural areas.

In a study reported by Dubowsky et al. (1999), the weekday indoor PAH concentrations attributable to traffic (indoor source contributions were removed) were $39 \pm 25 \text{ ng/m}^3$ in a dormitory that had a high air exchange rate because of open windows and doors; $26 \pm 25 \text{ ng/m}^3$ in an apartment; and $9 \pm 6 \text{ ng/m}^3$ in a suburban home. The study showed that both indoor and outdoor sources (especially motor vehicular traffic) contributed to indoor PAH concentrations. BaP concentrations were measured in the THEES study (Waldman et al., 1991). A comprehensive analysis of the data showed considerable seasonal variability of indoor and outdoor sources and resultant changes in personal exposures to BaP.

The indoor and outdoor concentrations of 30 PAHs were measured in 55 nonsmoking residences in Los Angeles, CA; Houston, TX; and Elizabeth, NJ (Naumova et al., 2002). A comparison of I/O ratios of low molecular weight PAHs (3 to 4 rings) and higher molecular weight PAHs (5 to 7 rings) indicated that indoor sources had a significant effect on indoor concentrations of 3-ring PAHs and a smaller effect on 4-ring PAHs, while outdoor sources dominated the indoor concentrations of 5- to 7-ring PAHs.

5.4.4 Factors Affecting Correlations Between Ambient Measurements and Personal or Microenvironmental Measurements of Particulate Matter Constituents

The primary factors affecting correlations between personal exposure and ambient air PM measurements are discussed in Section 4.3.2. These include air exchange rates, particle penetration factors, decay rates, removal mechanisms, indoor air chemistry, indoor sources, and freshly-generated particles indoors. The importance of these factors varies for different PM constituents. For acid aerosols, indoor air chemistry is particularly important as indicated by the discussion of the neutralization of the acidity by ammonia which is present at higher concentrations indoors because of the presence of indoor sources. For SVOCs, including PAHs and phthalates, the presence of indoor sources will substantially affect the correlation between indoor and ambient concentrations (Özkaynak et al., 1996b; Sheldon et al., 1993b). Penetration factors for PM will affect correlations between indoors and outdoors for most elements, except Pb, which may have significant indoor sources in older homes. Indoor air chemistry, decay

rates, and removal mechanisms may affect soot and OC. Furthermore, reactions between indoor or outdoor gases and particles may produce freshly generated aerosols indoors. These factors must be fully evaluated when attempting to correlate concentrations of PM constituents in ambient, personal, or indoor samplers.

5.4.5 Limitations of Available Data

The previous discussion demonstrates that there is limited data available that can be used to compare personal, microenvironmental, and ambient air concentrations of PM constituents. Because of resource limitations, PM constituents have not been measured in many studies of PM exposure. There are little data on freshly generated indoor aerosols. Although there are some data on acid aerosols, the comparisons between the personal and indoor data generally have been with outdoor measurements at the participants' residences, not with community ambient air measurement sites. The relationship between personal exposure and indoor levels of acid aerosols is not clear because of the limited database, although there is some evidence that acidity will be neutralized by indoor ammonia (Suh et al., 1994). The exception is total sulfate, for which there appears to be a strong correlation between indoor and ambient concentrations.

With the exception of PAHs, there are practically no data available with which to relate personal or indoor concentrations to outdoor or ambient site concentrations of SVOCs that may be generated from a variety of combustion and industrial sources. The relationship between exposure and ambient concentrations of particles from specific sources, such as diesel engines, has not been determined.

Although there is an increasing amount of research being performed to measure PM constituents in different PM size fractions, the current data are inadequate to adequately assess the relationship between indoor and ambient concentrations of most PM constituents. Additional information is also needed on PM exposures that result from outdoor gases reacting with indoor gases. This is a source that could also vary with outdoor PM, e.g., when the outdoor gas is O₃.

5.5 IMPLICATIONS OF USING AMBIENT PM CONCENTRATIONS IN TOXICOLOGICAL AND EPIDEMIOLOGICAL STUDIES OF PM HEALTH EFFECTS

5.5.1 Toxicology

Most studies of PM toxicity have used either pure chemicals or ambient PM. Indoor-generated PM differs somewhat from ambient PM in terms of sources, size, and composition. It is possible, therefore, that indoor-generated PM could have different toxicological properties. Environmental tobacco smoke (ETS) has been studied extensively; however, little toxicological information exists on PM from other indoor sources. Long et al. (2001b) have assessed the in vitro toxicity of 14 paired indoor and outdoor PM_{2.5} samples collected in nine Boston-area homes. Bioassays using rat alveolar macrophages (AMs) and measuring tumor necrosis factor (TNF) were used to assess particle-induced proinflammatory responses. TNF production was found to be significantly higher in AMs exposed to indoor PM than to those exposed to outdoor PM. This result held even after normalization for endotoxin concentrations, which were higher in indoor samples. The authors concluded that their results “suggest that indoor-generated particles may be more bioactive than ambient particles.” PM, in its various forms, produces many types of biological effects. It seems possible that indoor PM could be more active than ambient PM for some effects and less for others.

5.5.2 Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiological Analyses

In this section, the exposure issues that relate to the interpretation of the findings from epidemiologic studies of PM health effects are examined. This section examines the errors that may be associated with using ambient PM concentrations in epidemiologic analyses of PM health effects. First, implications of associations found between personal exposure and ambient PM concentrations are reviewed. This is discussed separately in the context of either community time-series studies or long-term, cross-sectional studies of chronic effects. Next, the role of compositional and spatial differences in PM concentrations are discussed and how these may influence the interpretation of findings from PM epidemiology. Finally, using statistical methods, an evaluation of the influence of exposure measurement errors on PM epidemiologic studies is presented.

Measurement studies of personal exposures to PM are still few in number and limited in spatial, temporal, and demographic coverage. Consequently, with the exception of a few longitudinal panel studies, most epidemiologic studies of PM health effects rely on ambient community monitoring data giving 24-h average PM concentration measurements. Moreover, because of limited sampling for PM_{2.5}, many of these epidemiologic studies used available PM₁₀ or in some instances relied on historic data for other PM measures or indicators, such as TSP, SO₄²⁻, IP₁₅ (inhalable particles with an upper cut of 15 μm), RSP (respirable suspended particles), COH (coefficient of haze), etc. A critical question often raised in the interpretation of results from acute or chronic epidemiological community-based studies of PM is whether the use of ambient stationary site PM concentration data influenced or biased the findings from those studies.

If it is assumed that total personal PM exposure is responsible for observed effects, use of ambient concentrations could lead to misclassification of individual exposures and to errors in the epidemiologic analysis of pollution and health data depending on the pollutant and on the mobility and lifestyles of the population studied. Ambient monitoring stations can be some distance away from the individuals and may represent only a fraction of all likely outdoor microenvironments that individuals come in contact with during the course of their daily lives. Furthermore, most individuals are quite mobile and move through multiple microenvironments (e.g., home, school, office, commuting, shopping) and engage in diverse personal activities at home (e.g., cooking, gardening, cleaning, smoking). Some of these microenvironments and activities may have different sources of PM and result in distinctly different concentrations of PM than that monitored by the fixed-site ambient monitors. Consequently, exposures of some individuals could be classified incorrectly if only ambient monitoring data are used to estimate individual-level total personal PM exposures. Thus, improper assessment of exposures using data routinely collected by the neighborhood monitoring stations could conceivably lead to a bias or increase in the standard error in epidemiologic analyses. Except for extremely unlikely situations, however, the bias would be expected to reduce the estimated health risk coefficient.

Many individuals are typically exposed to particles in a multitude of indoor and outdoor microenvironments during the course of a day. Thus, concern about possible error introduced in the estimation of PM risk coefficients using ambient, as opposed to personal, PM measurements has received considerable attention recently from exposure analysts, epidemiologists, and

biostatisticians. Some exposure analysts contend that, for community time-series epidemiology to yield information on the statistical association of a pollutant with a health response, there must be an association between personal exposure to a pollutant and the ambient concentration of that pollutant because people tend to spend around 90% time indoors and are exposed to both indoor-generated and ambient-infiltrated PM (cf. Wallace, 2000b; Brown and Paxton, 1998; Ebel et al., 2000). Consequently, numerous findings reported in the epidemiologic literature on significant associations between ambient PM concentrations and various morbidity and mortality health indices, in spite of the low correlations between ambient PM and concentrations and measures of personal exposure, have been described by some exposure analysts as an exposure paradox (Lachenmyer and Hidy, 2000; Wilson et al., 2000).

To resolve that exposure paradox, two types of analyses must be considered. The first type of analysis is to examine the correlations between ambient PM concentrations and personal exposures that are relevant to most of the existing PM epidemiologic studies using either pooled, daily-average, or longitudinal exposure data. The second approach is to study the degree of correlations between the two key components of personal PM exposures (i.e., exposures caused by ambient PM and exposures caused by nonambient PM) with ambient or outdoor PM concentrations, for each of the three types of exposure study designs. Yet, even with these two approaches, it may still be difficult to examine complex synergisms which, in some situations, may preclude simple decoupling of indoor and outdoor particles either in terms of exposure or total dose delivered to the lung. In addition, several factors that influence either the exposure or health response characterization of the subjects have to be addressed. Such factors include:

- spatial variability of PM components,
- health or sensitivity status of subjects,
- variations of PM with other co-pollutants,
- co-generation of fine and ultrafine particles from outdoor air and indoor gaseous pollutants,
- formal evaluation of exposure errors in the analysis of health data, and
- how the results may depend on the variations in epidemiological study design.

To facilitate the discussion of these topics, a brief review of concepts pertinent to exposure analysis issues in epidemiology is presented.

5.5.2.1 Associations Between Personal Exposures and Ambient Particulate Matter Concentrations

As defined in Sections 5.3 and 5.4, personal exposures to PM result from an individual's exposures to PM in many different types of microenvironments (e.g., outdoors near home, outdoors away from home, indoors at home, indoors at office or school, commuting, restaurants, malls, other public places). Total personal exposures that occur in these indoor and outdoor microenvironments can be classified as those resulting from ambient PM (ambient PM exposure includes exposure to ambient PM while outdoors and exposure while indoors to ambient PM that has infiltrated indoors) and those primarily generated by indoor sources, indoor reaction, and personal activities (nonambient PM exposure). The associations between personal exposures and ambient PM concentrations that have been reported from various personal exposure monitoring studies under three broad categories of study design, (1) longitudinal, (2) daily-average, or (3) pooled exposure studies, and are summarized below.

In Sections 5.4.3.1.2 and 5.4.3.1.3, recent studies conducted mainly in the United States and involving children, the elderly, and subjects with COPD were reviewed, and they indicated that both intra- and interindividual variability in the relationships between personal exposures and ambient PM concentrations were observed. A variety of different physical, chemical, and personal or behavioral factors were identified by the original investigators that seem to influence the magnitude and the strength of the associations reported.

For cohort studies in which individual daily health responses are obtained, individual longitudinal PM personal exposure data (including ambient and nonambient components) may provide the appropriate indicators. In this case, health responses of each individual can be associated with the total personal exposure, the ambient exposure, or the nonambient exposure of each individual. Additionally, the relationships of personal exposure indicators with ambient concentration can be investigated. In the case of community time-series epidemiology, however, it is not feasible to obtain experimental measurements of personal exposure for the millions of people over the periods of years that are needed to investigate the relationship between air pollution and infrequent health responses such as deaths or even hospital admissions. The epidemiologist must work with the aggregate number of health responses occurring each day and a measure of the ambient concentration that is presumed to be representative of the entire community. The relationship of PM exposures of the potentially susceptible groups to monitored ambient PM concentrations depends on their activity pattern and level, residential

building and HVAC factors (which influence the infiltration factor), status of exposure to ETS, amount of cooking or cleaning indoors, and seasonal factors, among others. For these special subgroups, average personal exposures to ambient PM correlate well with ambient PM concentrations regardless of individual variation in the absence of major microenvironmental sources.

Even though both ambient and nonambient PM exposure contribute to daily baseline PM dose received by the lung, there seem to be clear differences in the relationships of ambient and nonambient PM exposure with ambient PM concentration. Various researchers have shown that nonambient PM exposure is independent of ambient PM concentration, but that ambient PM exposure is a function of ambient PM concentration. Wilson et al. (2000) explained the difference based on different temporal patterns that affect PM concentrations, “Concentrations of ambient PM are driven by meteorology and by changes in the emission rates and locations of emission sources, while concentrations of nonambient PM are driven by the daily activities of people.” Still, although nonambient PM exposure may not correlate with ambient PM concentration or ambient PM exposure, it will nevertheless add to the daily baseline dose received by the lung. An important concern, for which there is little information, is the relative biological activity of ambient and indoor-generated PM both in terms of the type of toxic effect and the relative potency for a given effect.

Ott et al. (2000) also discussed the reasons for assuming that nonambient PM exposure is independent of ambient PM exposure and ambient PM concentration. They showed that the nonambient component of total personal exposure is uncorrelated with the outdoor concentration data. Ott et al. (2000) demonstrated that the ambient PM exposure was similar for three population-based exposure studies: two large probability-based studies (the PTEAM study conducted in Riverside [Clayton et al., 1993; Thomas et al., 1993; Özkaynak et al., 1996a,b] and a study in Toronto [Pelizzarri et al., 1999; Clayton et al., 1999a]), and a nonprobability-based study conducted in Phillipsburg (Lioy et al., 1990). Based on these three studies, they concluded that the average nonambient PM exposure and the distribution of individual, daily values of nonambient PM exposure can be treated as constant from city to city.

Dominici et al. (2000) examined a larger database consisting of five different PM exposure studies and also concluded that nonambient PM exposure can be treated as relatively constant from city to city, although their data show greater variability than the data reported by Ott

(2000). However, the constancy of nonambient PM exposure remains an open question. If daily values of nonambient PM exposure were constant, this would imply a zero correlation with ambient PM concentration. However, this hypothesis of constant individual, daily nonambient PM exposure has not been fully established, since only a few studies have produced the data needed to estimate daily, individual values of nonambient PM exposure. Although nonambient PM exposure is independent of ambient PM concentration, it may not be independent of the attenuation factor (ambient PM exposure/ambient PM concentration). Sarnat et al. (2000) showed that nonambient PM exposure increases as the ventilation rate (and attenuation factor) decreases. By comparing winter and summer regression equations Lachenmeyer and Hidy (2000) also showed that as the slope, which gives the attenuation factor, decreases, the intercept, which gives the nonambient PM exposure, increases. If ambient and nonambient PM_{2,5} are not correlated, as seems likely, statistical analyses using ambient concentrations will not be able to demonstrate health effects, or the lack of health effects, due to indoor-generated PM. However, indoor-generated PM could raise the overall exposure level, contribute to the observed health effects, and make it more difficult to observe a threshold level for ambient PM.

Mage et al. (1999) assumed that the PM₁₀ concentration component from indoor sources (e.g., smoking, cooking, cleaning, burning candles) is not correlated with the outdoor concentration. They indicated that this lack of correlation is to be expected because people are unaware of ambient concentrations and do not necessarily change their smoking or cooking activities as outdoor PM₁₀ concentrations vary, an assumption supported by other empirical analyses of personal exposure data. For the PTEAM data set, Mage et al. (1999) have shown that individual, daily exposures to indoor-generated PM and daily ambient PM concentrations have a correlation coefficient near zero ($R^2 = 0.005$). Wilson et al. (2000) have shown that individual, daily values of concentration of ambient PM which has infiltrated indoors and indoor-generated PM concentrations also have a near zero correlation ($R^2 = 0.03$). Figure 5-14 shows the relationship of estimated values of nonambient PM exposure with ambient PM concentrations (calculated by EPA, daily individual values from PTEAM and daily average values for the cohort from THEES).

Based on these results, it is reasonable to assume that nonambient PM exposure will ordinarily have little correlation with ambient PM concentration. A possible exception could be caused by indoor-reaction PM, PM formed when an ambient gas infiltrates indoors and reacts




Figure 5-14. Plots of nonambient exposure to PM₁₀ showing (a) daily individual daytime values from PTEAM data and (b) daily-average values from THEES data.

Source: Data taken from (a) Clayton et al. (1993) and (b) Liroy et al. (1990).

with an indoor-generated gas, e.g., the reaction of O₃ with terpenes from air cleaners. However, ambient O₃ does not appear to be highly correlated with ambient or personal PM_{2.5} (see Table 5-12, Sarnat et al., [2000]). Additionally, not every home will use air fresheners or have the same level of terpene emissions. Hence, indoor-reaction PM concentrations would not be expected to correlate with ambient PM concentrations on a community basis. Therefore, in linear nonthreshold models of PM health effects, nonambient PM exposure is not expected to contribute to the relative risk determined in a regression of health responses on ambient PM concentration. Furthermore, in time-series analysis of pooled or daily health data, it is expected that ambient PM exposure, rather than total personal PM exposure, will have the stronger association with ambient PM concentration.

5.5.2.2 Role of Compositional Differences in Exposure Characterization for Epidemiology

The majority of the available data on PM exposures and relationships with ambient PM have come from a few large-scale studies, such as PTEAM, or longitudinal studies on selected populations. Consequently, for most analyses, exposure scientists and statisticians had to rely on PM₁₀ or PM_{2.5} mass data, instead of elemental or chemical compositional information on individual or microenvironmental samples. In a few cases, researchers have examined the factors influencing I/O ratios or penetration and deposition coefficients using elemental mass data on personal, indoor, and outdoor PM data (e.g., Özkaynak et al., 1996a,b; Yakovleva et al., 1999). These results have been informative in terms of understanding relative infiltration of different classes of particle sizes and sources into residences (e.g., fossil fuel combustion, mobile source emissions, soil-derived). Clearly, accumulation-mode particles, associated with stationary or mobile combustion sources, have greater potential for penetration into homes and other microenvironments than crustal material. There will be variability in the chemical composition of these broad categories of source classes as well as probable variations in relative toxicity. Moreover, when particles and reactive gases are present indoors in the presence of other pollutants or household chemicals, they may react to form additional or different compounds and particles with yet unknown physical, chemical, and toxic compositions (Wainman et al., 2000). Thus, if indoor-generated and ambient PM were responsible for different types of health effects or had significantly different toxicities on a per unit mass basis, it would then be important that ambient and nonambient exposure should be separated and

treated as different components, much like the current separation of PM_{10} into $PM_{2.5}$ and $PM_{10-2.5}$. These complexities in personal exposure profiles may introduce nonlinearities and other statistical challenges in the selection and fitting of concentration-response models. Unfortunately, PM health effects models have not yet been able to meaningfully consider such complexities.

It is important to also note that individuals spend time in places other than their homes and outdoors. Many of the interpretations reported in the published literature on factors influencing personal PM_{10} exposures, as well as in this chapter, come from the PTEAM study. The PTEAM study was conducted 10 years ago in one geographic location in California, during one season, and most residences had very high and relatively uniform air exchange rates. Non-home indoor microenvironments were not monitored directly during the PTEAM study. Commuting exposures from traffic or exposures in a variety of different public places or office buildings could not be assessed directly. Nonresidential buildings may have lower or higher ambient infiltration rates depending on the use and type of the mechanical ventilation systems employed. Because the source and chemical composition of PM affecting personal exposures in different microenvironments vary by season, day-of-the-week, and time of day, it is conceivable that some degree of misclassification of exposures to toxic PM agents of concern could be introduced when health-effects models use only daily-average mass measures such as PM_{10} or $PM_{2.5}$. However, because of the paucity of currently available data on many of these factors, it is not now possible to ascertain the potential magnitude or severity of any such complex exposure misclassification problems or their potential implications for interpretation of results from PM epidemiology.

5.5.2.3 Role of Spatial Variability in Exposure Characterization for Epidemiology

Chapter 3, Section 3.2.3 and Chapter 5, Section 5.3 present information on the spatial variability of PM mass and chemical components at fixed-site ambient monitors; for purposes of this chapter, this spatial variability is called an “ambient gradient.” Any gradient that may exist between a fixed-site monitor and the outdoor microenvironments near where people live, work, and play will obviously affect the concentration profile actually experienced by people as they go about their daily lives.

However, the evidence so far indicates that PM concentrations, especially fine PM (mass and sulfate), generally are uniformly distributed in most metropolitan areas. This reduces the potential for exposure misclassification because of outdoor spatial gradients when a limited number of ambient PM monitors are used to represent population average ambient exposures in time-series or cross-sectional epidemiological studies of PM. This topic is further discussed in Section 5.6.5. However, as discussed earlier, the same assumption is not necessarily true for different components of PM such as PM_{10-2.5}, because source-specific and other spatially nonuniform pollutant emissions could alter the spatial profile of individual PM components in a community. For example, particulate and gaseous pollutants emitted from motor vehicles tend to be higher near roadways and inside cars. Likewise, acidic and organic PM species may be location- and time-dependent. Furthermore, human activities are complex. If outdoor PM constituent concentration profiles are either spatially or temporally variable, it is likely that exposure misclassification errors could be introduced in the analysis of PM air pollution and health data.

5.5.3 Analysis of Exposure Measurement Error Issues in Particulate Matter Epidemiology

The effects of exposure misclassification on the relative risk estimates of disease using the classical 2×2 contingency design (i.e., exposed/nonexposed versus diseased/nondiseased) have been studied extensively in the epidemiologic literature. It has been shown that the magnitude of the exposure-disease association (e.g., relative risk), because of either misclassification of exposure or disease alone (i.e., nondifferential misclassification), biases the effect results toward the null; and differential misclassification (i.e., different magnitudes of disease misclassification in exposed and nonexposed populations) can bias the effect measure toward or away from the null value relative to the true measure of association (Shy et al., 1978; Gladen and Rogan, 1979; Copeland et al., 1977; Özkaynak et al., 1986). However, the extension of these results from contingency analysis design to multivariate models (e.g., log-linear regression, Poisson, logit) typically used in recent PM epidemiology has been more complicated.

5.5.3.1 Time-Series Analyses

Researchers have investigated the appropriateness of using ambient PM concentration as an exposure metric and have developed a framework for analyzing measurement errors typically encountered in the analysis of time-series mortality and morbidity effects from exposures to ambient PM (cf. Zeger et al., 2000; Dominici et al., 2000; Samet et al., 2000). Use of this framework, discussed more extensively in Chapters 8 and 9, leads to the following conclusions: the deviation of an individual's personal exposure from the risk-weighted average exposure due to variations in ambient concentrations, infiltration rates, and indoor-generated PM concentrations is a Berkson error and will not bias the estimated regression coefficient (β , the increase in risk per unit increase in PM) in a time-series analysis of mortality as a function of ambient PM concentrations. However, the difference between the average personal exposure and the true ambient concentration will bias β . If the daily, individual values of ambient PM exposure and nonambient PM exposure are poorly correlated and the attenuation coefficient, α , equal to the ambient PM concentration/ambient PM exposure were constant, the bias would be given by $\beta_C = \alpha\beta_A$ where β_C is calculated using the ambient PM concentration, C, and β_A is calculated using the ambient PM exposure, A; i.e., the risk determined from an analysis using the daily ambient PM concentrations would be lower than the risk obtained using ambient PM exposure by the factor α . However, β_C provides the correct information on the change in health risks that would be produced by a change in ambient concentrations. However, if the daily, individual values of ambient PM exposure and nonambient PM exposure are highly correlated and the nonambient PM is toxic for the effect being studied, nonambient PM exposure will act as a confounder and could introduce substantial bias into β_C . Only one study has reported the association between the daily, individual values of ambient PM exposure and nonambient PM exposure. Wilson et al. (2000), in a further analysis of the PTEAM data set, found a coefficient of determination, $R^2 = 0.03$, suggesting that the daily, individual values of ambient PM exposure and nonambient PM exposure are independent, meaning that nonambient PM exposure will not confound β_C .

5.5.3.2 Studies of Chronic Effects

The Six Cities (Dockery et al., 1993) and American Cancer Society (ACS) (Pope et al., 1995) Studies have played an important role in assessing the health effects from long-term

exposures to particulate pollution. Even though these studies often have been considered as chronic epidemiologic studies, it is not easy to differentiate the role of historic exposures from those of recent exposures on chronic disease mortality. In the Six Cities study, fine particles and sulfates were measured at the community level, and the final analysis of the database used six city-wide average ambient concentration measurements. This limitation also applies to the ACS study but has less impact because of the larger number of cities considered in that study. In a HEI-sponsored reanalysis of the Six Cities and the ACS data sets, Krewski et al. (2000) examined some of the exposure misclassification issues either analytically or through sensitivity analysis of the aerometric and health data. The HEI reanalysis project also addressed exposure measurement error issues related to the Six Cities study. For example, the inability to account for exposures prior to the enrollment of the cohort hampered accurate interpretation of the relative risk estimates in terms of acute versus chronic causes. Although the results seem to suggest that past exposures are more strongly associated with mortality than recent exposures, the measurement error for long-term averages could be higher, influencing these interpretations. As another example, Krewski et al. (2000), using the individual mobility data available for the Six Cities cohort, analyzed the mover and nonmover groups separately. The relative risk of fine particle effects on all-cause mortality was shown to be higher for the nonmover group than for the mover group, suggesting the possibility of higher exposure misclassification biases for the movers. The issue of using selected ambient monitors in the epidemiological analyses was also investigated by the ACS and Six Cities studies reanalysis team. Krewski et al. (2000) described the sensitivity of results to choices made in selecting stationary or mobile-source-oriented monitors. For the ACS study, reanalysis of the sulfate data using only those monitors designated as residential or urban and excluding sites designated as industrial, agricultural, or mobile did not appreciably change the risk estimates. On the other hand, application of spatial analytic methods designed to control confounding at larger geographic scales (i.e., between cities) caused changes in the particle and sulfate risk coefficients. Spatial adjustment may account for differences in pollution mix or PM composition, but many other cohort-dependent risk factors will vary across regions or cities in the United States. Therefore, it is difficult to interpret these findings solely in terms of spatial differences in pollution composition or relative PM toxicity until further research is concluded.

The influence of measurement errors in air pollution exposure and health effects assessments has also been examined by Navidi et al. (1999). This study developed techniques to incorporate exposure measurement errors encountered in long-term air pollution health-effects studies and tested them on the data from the University of Southern California Children's Health Study conducted in 12 communities in California. These investigators developed separate error analysis models for direct (i.e., personal sampling) and indirect (i.e., microenvironmental) personal exposure assessment methods. These models were generic to most air pollutants, but a specific application was performed using a simulated data set for studying ozone health effects on lung function decline in children. Because the assumptions made in their microenvironmental simulation modeling framework were similar to those made in estimating personal PM exposures, it is useful to consider the conclusions from Navidi et al. (1999). According to Navidi et al. (1999), neither the microenvironmental nor the personal sampler method produces reliable estimates of the exposure-response slope for O₃ when measurement error is uncorrected. Because of nondifferential measurement error, the bias was toward zero under the assumptions made in Navidi et al. (1999) but could be away from zero if the measurement error was correlated with the health response. A simulation analysis indicated that the standard error of the estimate of a health effect increases as the errors in exposure assessment increase (Navidi et al., 1999). According to Navidi et al. (1999), when a fraction of the ambient level in a microenvironment is estimated with a standard error of 30%, the standard error of the estimate is 50% higher than it would be if the true exposures were known. It appears that errors in estimating ambient PM indoor/ambient PM outdoor ratios have much more influence on the accuracy of the microenvironmental approach than do errors in estimating time spent in these microenvironments.

Epidemiologic studies of chronic effects that use long-term average ambient PM concentrations as the exposure metric generally do not address the nonambient component of personal exposure (N). However, if N contributes to the health effect being studied and the average N is different in different cities, the correlation between the average ambient PM concentration and the health effect could be reduced. In an analysis of the effect of nonambient exposure on time-series epidemiology, Dominici et al. (2000) examined nonambient exposure data from several cities and concluded that the average nonambient PM exposure varies little

among cities in developed countries. Also, Ott et al. (2000) examined estimated average and daily, individual values of nonambient exposure from three studies and concluded that both the average and the distribution of daily, individual values were similar for the three studies.

5.5.4 Conclusions from Analysis of Exposure Measurement Errors on Particulate Matter Epidemiology

Personal exposures to PM are influenced by a number of factors and sources of PM located in both indoor and outdoor microenvironments. However, PM resulting from ambient sources penetrate into indoor environments such as residences, offices, public buildings, etc., in which individuals spend a large portion of their daily lives. The correlations between total personal exposures and ambient or outdoor PM concentrations can vary depending on the relative contributions of indoor PM sources to total personal exposures. Panel studies of both adult and young subjects have shown that, in fact, individual correlations of personal exposures with ambient PM concentrations could vary from person to person, and even day to day, depending on the specific activities of each person. Separation of PM exposures into two components, ambient PM and nonambient PM, would reduce uncertainties in the analysis and interpretation of PM health effects data. Nevertheless, because ambient PM is an integral component of total personal exposures to PM, statistical analyses of cohort-average exposures are strongly correlated with ambient PM concentrations when a large underlying population is studied. Using the PTEAM study data, analysis of exposure measurement errors, in the context of time-series epidemiology, has also shown that errors or uncertainties introduced by using surrogate exposure variables, such as ambient PM concentrations, could lead to biases in the estimation of health risk coefficients. These then would need to be corrected by suitable calibration of the PM health risk coefficients. Correlations between the PM exposure variables and other covariates (e.g., gaseous co-pollutants, weather variables) also could influence the degree of bias in the estimated PM regression coefficients. However, most time-series regression models employ seasonal or temporal detrending of the variables, thus reducing the magnitude of this cross-correlation problem (Özkaynak and Spengler, 1996).

Ordinarily, exposure measurement errors are not expected to influence the interpretation of findings from either the chronic or time-series epidemiologic studies that have used ambient concentration data if they include sufficient adjustments for seasonality and key confounders.

There is no question that better estimates of exposures to components of PM of health concern are beneficial. Composition of PM may vary in different geographic locations and different exposure microenvironments. Compositional and spatial variations could lead to further errors in using ambient PM measures as surrogates for exposures to PM. Even though the spatial variability of PM (PM_{2.5} in particular) mass concentrations in urban environments seems to be small, the same conclusions drawn above regarding the influence of measurement errors may not necessarily hold for all of the toxic PM components. Again, the expectation based on statistical modeling considerations is that these exposure measurement errors or uncertainties will most likely reduce the statistical power of the PM health effects analysis, making it difficult to detect a true underlying association between the correct exposure metric and the health outcome studied. However, until more data on exposures to specific toxic components of PM become available, existing studies on PM exposure measurement errors must be relied on; thus, at this time, the use of ambient PM concentrations as a surrogate for exposures is not expected to change the principal conclusions from PM epidemiologic studies that use community average health and pollution data.

5.6 SUMMARY OF OBSERVATIONS AND LIMITATIONS

Exposure Definitions and Components

- Personal exposure to PM mass or its constituents results when individuals come in contact with particulate pollutant concentrations in locations or microenvironments that they frequent during a specific period of time. Various PM exposure metrics can be defined according to its source (i.e., ambient, nonambient) and the microenvironment where exposure occurs.
- Personal exposure to PM results from an individual's exposure to PM in many different types of microenvironments (e.g., outdoors near home, outdoors away from home, indoors at home, indoors at office or school, commuting, restaurants, malls, other public places). Thus, total daily exposure to PM for a single individual can be expressed as the sum of various microenvironmental exposures that the person encounters during the course of a day.
- In a given microenvironment, particles may originate from a wide variety of sources. In an indoor microenvironment, PM may originate from PM-generating activities (e.g., cooking, cleaning, smoking, resuspending PM from PM resulting from both indoor and outdoor sources that had settled out), from outside (outdoor PM entering through cracks and

openings in the structure), and from the chemical interaction of pollutants from outdoor air with indoor-generated pollutants.

- The total daily exposure to PM for a single individual can also be expressed as the sum of contributions of ambient and nonambient PM. Nonambient PM exposure is due to PM generated by indoor sources, personal activities, and chemical reactions in indoor air. Ambient PM exposure includes exposure to ambient PM while outdoors, and ambient PM that has infiltrated indoors while indoors. However, within a large population, there will be distributions of total personal exposure and its components due to variations in human activities and microenvironmental concentrations and sources each individual encounters.
- Exposure models are useful tools for examining the importance of sources, micro-environments, and physical and behavioral factors that influence personal exposures to PM. However, the development and evaluation of population exposure models for PM and its components have been limited; and improved modeling methodologies and new model input data are needed.

Factors Affecting Concentrations and Exposures to Particulate Matter

- Concentrations of PM indoors are affected by several factors and mechanisms: ambient concentrations outdoors, air exchange rates, particle penetration factors, particle production from indoor sources and indoor air chemistry, and indoor particle decay rates and removal mechanisms caused by physical processes or resulting from mechanical filtration, ventilation or air-conditioning devices.
- Average personal exposures to PM mass and its constituents are influenced by microenvironmental PM concentrations and by how much time each individual spends in these various indoor and outdoor microenvironments. Nationwide, individuals, on average, spend nearly 90% of their time indoors (at home and in other indoor locations) and about 6% of their time outdoors.
- Personal exposures are associated with both indoor as well as outdoor sources. The personal exposure/outdoor concentration ratios present substantial intra- and interpersonal variability. This variability is due to both the presence of personal and microenvironmental sources and the varying effect of the outdoor particles on indoor environments.
- Home characteristics may be the most important factor that affects the relationship between the average population exposures and ambient concentrations. Air exchange rate seems to be an important home characteristic surrogate that can explain a large fraction of the observed inter- and intrapersonal variability. One reason why longitudinal studies (many repeated measurements per person) provide stronger correlations between personal exposure and outdoor concentrations than pooled studies (few repeated measurements per individual) may be because home characteristics remain the same.

- Because home characteristics constitute the most important factor affecting personal exposures, one would expect that correlations between average population exposures and outdoor concentrations will vary by season and geography.
- The relative size of personal exposure to ambient PM relative to nonambient PM depends on the ambient concentration, the infiltration rate of outdoor PM into indoor microenvironments, the amount of PM generated indoors (e.g., ETS, cooking and cleaning emissions), and the amount of PM generated by personal activity sources. Infiltration rates primarily depend on air exchange rate, size-dependent particle penetration across the building membrane, and size-dependent removal rates. All of these factors vary over time and across subjects and building types.
- The relationship between PM exposure, dose, and health outcome could depend on the concentration, composition, and toxicity of PM or PM components originating from different sources. Application of source apportionment techniques to indoor and outdoor PM_{2.5} and personal, indoor, and outdoor PM₁₀ composition data have identified the following general source categories: outside soil, resuspended indoor soil, indoor soil, personal activities, sea salt, motor vehicles, nonferrous metal smelters, and secondary sulfates.
- Only a limited number of studies have measured the physical and chemical constituents of PM in personal or microenvironmental samples. Available data on PM constituents indicate the following:
 - personal and indoor sulfate measurements often are correlated highly with outdoor and ambient sulfate concentration measurements;
 - for acid aerosols, indoor air chemistry is particularly important because of the neutralization of the acidity by ammonia, which is present at higher concentrations indoors because of the presence of indoor sources of ammonia;
 - for SVOCs, including PAHs and phthalates, the presence of indoor sources will substantially affect the relation between indoor and ambient concentrations;
 - penetration and decay rates are functions of size and will cause variations in the attenuation factors as a function of particle size (infiltration rates will be higher for PM₁ and PM_{2.5} than for PM₁₀, PM_{10-2.5} or ultrafine particles); and
 - indoor air chemistry may increase indoor concentrations of organic PM.
- Even though there is an increasing amount of research being performed to measure PM constituents in different PM size fractions, with few exceptions (i.e., sulfur or sulfates) the current data are inadequate to adequately assess the relationship between personal, indoor, and ambient concentrations of most PM constituents.

Correlations Between Personal Exposures, Indoor, Outdoor, and Ambient Measurements

- Most of the available personal data on PM measurements and information on the relationships between personal and ambient PM come from a few large-scale studies (e.g., the PTEAM study) or the longitudinal panel studies that have been conducted on selected populations, such as the elderly.
- Panel and cohort studies that have measured PM exposures and concentrations typically have reported their results in terms of three types of correlations: (1) longitudinal, (2) pooled, and (3) daily-average correlations between personal and ambient or outdoor PM.
- The type of correlation analysis performed can have a substantial effect on the resulting correlation coefficient. Low correlations with ambient concentrations could result when people with very different nonambient exposures are pooled even though their individual personal exposures may temporally be highly correlated with ambient concentrations.
- Longitudinal and pooled correlations between personal exposure and ambient or outdoor PM concentrations reported by various investigators varied considerably among the different studies and in each study between the study subjects. Most studies report longitudinal correlation coefficients that range from close to 0 to near 1.0, indicating that individual's activities and residence type may have a significant effect on total personal exposures to PM.
- Longitudinal studies that measured sulfate found high correlations between personal and ambient sulfate.
- In general, probability-based population studies tend to show low pooled correlations because of the high differences in levels of nonambient PM-generating activities from one subject to another. In contrast, the absence of indoor sources for the populations in several of the longitudinal panel studies resulted in high correlations between personal exposure and ambient PM within subjects over time for these populations. However, even for these studies, correlations varied by individual depending on their activities and the microenvironments that they occupied.

Potential Sources of Error Resulting from Using Ambient Particulate Matter Concentrations in Epidemiological Analyses

As yet, there is no clear consensus among exposure analysts as to how well community monitor measurements of ambient air PM concentrations represent a surrogate for personal exposure to total PM or to ambient PM.

- Measurement studies of personal exposures to PM are still few in number and limited in spatial, temporal, and demographic coverage. Consequently, with the exception of a few longitudinal panel studies, most epidemiological studies on PM health effects have relied on daily-average PM concentration measurements obtained from ambient community monitoring data as a surrogate for the exposure variable.

- Because individuals are exposed to particles in a multitude of indoor and outdoor microenvironments during the course of a day, error is introduced in the estimation of PM risk coefficients using ambient, as opposed to personal, PM measurements.
- Total personal exposures to PM could vary from person to person, and even day to day, depending on the specific activities of each person. Separation of PM exposures into two components, ambient and nonambient PM, would reduce potential uncertainties in the analysis and interpretation of PM health effects data.
- Available data indicate that PM mass concentrations, especially fine PM, typically are distributed relatively uniformly in most metropolitan areas, thus reducing the potential for exposure misclassification because of spatial variability when a limited number of ambient PM monitors are used to represent population average ambient exposures in community time-series or long-term, cross-sectional epidemiological studies of PM.
- Even though the spatial variability of PM (in particular, $PM_{2.5}$) mass concentrations in urban environments seems to be small, the same conclusions drawn above regarding the influence of measurement errors may not necessarily hold for all PM components.
- There are important differences in the relationship of ambient PM concentrations with ambient PM exposures and with nonambient PM exposures. Various researchers have shown that ambient PM exposure is a function of ambient PM concentration and that concentrations of ambient PM are driven by meteorology, by changes in source emission rates, and in locations of emission sources relative to the measurement site. However, nonambient PM exposure is independent of ambient PM concentration because concentrations of nonambient PM are driven by the daily activities of people.
- Because personal exposures also include a contribution from ambient concentrations, the correlation between daily-average personal exposure and the daily-average ambient concentration increases as the number of subjects measured daily increases. An application of a Random Component Superposition model has shown that the contributions of ambient PM_{10} and indoor-generated PM_{10} to community mean exposure can be decoupled in modeling urban population exposure distributions.
- If linear nonthreshold models are assumed in time-series analysis of daily-average ambient PM concentrations and community health data, nonambient PM exposure is not expected to contribute to the relative risk estimates determined by regression of health responses on ambient PM concentration.
- Using the PTEAM study data, analysis of exposure measurement errors in the context of time-series epidemiology has shown that the error introduced by using ambient PM concentrations as a surrogate for ambient PM exposure biases the estimation of health risk coefficients low by the ratio of ambient PM exposure to ambient PM concentration (called the attenuation factor). However, the health risk coefficient determined using ambient PM concentrations provides the correct information on the change in health risks that would be produced by a change in ambient concentrations.

- Because sources and chemical composition of PM affecting personal exposures in different microenvironments vary by season, day of the week, and time of day, it is likely that some degree of misclassification of exposures to PM toxic agents of concern will be introduced when health-effects models use only daily-average mass measures such as PM₁₀ or PM_{2.5}. Because of the paucity of currently available data on many of these factors, at this point it is impossible to ascertain the significance of these more complex exposure misclassification problems in the interpretation of results from PM epidemiology.
- Exposure measurement errors may depend on particle size and composition. PM_{2.5} better reflects personal exposure to PM of outdoor origin than PM₁₀. Various indicators of ultrafine particle concentrations or other components of PM may also be useful exposure indicators for epidemiological studies.
- Seasonal or temporal variations in the measurement errors and their correlations between different PM concentration measures and co-pollutants (e.g., SO₂, CO, NO₂, O₃) could influence the error analysis results but are not likely to change the interpretation of current findings.
- Multipollutant personal exposure studies have suggested that ambient concentrations of gaseous co-pollutants serve as surrogates of personal exposures to particles rather than as confounders. The implications for epidemiology are discussed in Chapters 8 and 9.
- Ordinarily, PM exposure measurement errors are not expected to influence the interpretation of findings from either the community time-series or long-term epidemiologic studies that have used ambient concentration data if they include sufficient adjustments for seasonality and key personal and geographic confounders.
- In the context of long-term epidemiological studies, it appears that the errors introduced in estimating ambient PM indoor/ambient PM outdoor ratios have much more influence on the accuracy of the microenvironmental exposure estimation approach than do errors in estimating time spent in these microenvironments.
- To reduce exposure misclassification errors in PM epidemiology, conducting new cohort studies of sensitive populations with better real-time techniques for exposure monitoring and further speciation of indoor-generated, ambient, and personal PM mass are essential.
- Based on statistical modeling considerations, it is expected that existing PM exposure measurement errors or uncertainties most likely will reduce the statistical power of the PM health effects analysis, thus making it difficult to detect a true underlying association between the correct exposure metric and the health outcome studied.
- Although exposure measurement errors for fine particles are not expected to influence the interpretation of findings from either the community time-series or the long-term, cross-sectional epidemiological studies that have used ambient concentration data, they may underestimate the strength of the effect. Sufficient data are not available to evaluate the effect of exposure measurement error for other PM species or size fractions.

Key Findings

- Most people spend most of their time indoors where they are exposed to indoor-generated PM and ambient PM that has infiltrated indoors.
- Indoor-generated and ambient PM differ in sources, sizes, chemical composition, and toxicity.
- The ambient PM concentration and the indoor PM concentration can be measured by outdoor and indoor monitors. The total personal exposure can be measured by a personal exposure monitor carried by the person. However, the concentrations of indoor-generated PM and ambient PM that has infiltrated indoors and the related values of ambient and nonambient PM exposures must be estimated.
- From a regression of individual, daily values of total personal exposure on daily PM concentrations, the intercept gives the average nonambient PM exposure and the slope gives the average attenuation factor (the ratio of ambient PM exposure to ambient PM concentration).
- Similarly, from a regression of individual, daily values of indoor PM concentration on daily ambient concentrations, the intercept gives the average concentration of indoor-generated PM and the slope gives the average infiltration factor (concentrations of ambient PM that has infiltrated indoors/ambient PM concentration).
- The attenuation factor and the infiltration factor depend on the penetration coefficient, the fraction of ambient PM that penetrates through the walls, doors, windows, etc.; the deposition or removal rate, a measure of how rapidly PM within the indoor microenvironment is removed by deposition to surfaces or by filtration in a heating/cooling system; and the air exchange rate, a measure of how rapidly indoor air is replaced by outdoor air. The attenuation factor also depends on the fraction of time spent outdoors.
- The air exchange rate is an important variable for determining the concentration of ambient PM found indoors. It can be measured by release and measurement of an inert tracer gas indoors. The air exchange rate increases with the opening of windows or doors or the operation of window or attic fans. It also increases as the indoor/outdoor temperature difference increases. For closed homes, i.e., no open windows or doors, the air exchange rate does not appear to be a function of wind speed or direction.
- The penetration coefficient and the deposition rate can be estimated from measurements of outdoor and indoor concentrations under conditions when there are no indoor sources (nighttime or unoccupied home). These parameters are functions of particle size. The penetration coefficient is high and the deposition rate is low for accumulation mode particles (0.1 to 1.0 μm). The penetration coefficient is lower and the deposition rate is higher for ultrafine particles (< 0.1 μm) and coarse mode particles (> 1.0 μm). The attenuation factor and the infiltration factor are higher for particles in the accumulation mode than for ultrafine or coarse particles.

- The attenuation factor and the infiltration factor will vary with the air exchange rate and, therefore, will vary with season and housing characteristics. These factors will increase with increased opening of windows and doors. For closed homes, these factors will increase with an increase of the indoor/outdoor temperature difference but they do not appear to be affected by wind speed or direction.
- The regression technique is useful for finding average values of the attenuation factor and the nonambient exposure and possibly for estimating the distribution of individual, daily values of the nonambient PM exposure.
- Individual, daily values of the ambient PM exposure, the nonambient PM exposure, and the attenuation factor may be determined from individual, daily values of the total PM personal exposure and daily ambient PM concentrations by several techniques:
 - *Mass balance technique.* Direct measurement of the air exchange rate, measurement of the fraction of time spent outdoors by a diary of the subject's activity pattern, and use of the equilibrium mass balance equation for the attenuation factor with estimated values of the penetration coefficient and the deposition rate.
 - *Sulfate ratio technique.* Individual, daily values of the attenuation factor (for PM_{2.5}) will be given by individual, daily values of personal exposure to sulfate / the daily ambient sulfate concentration, provided that there are no indoor sources of sulfate and sulfate and PM_{2.5} have similar particle size distributions.
 - *Recursive technique.* Indoor-generated emissions, which tend to be episodic, can be removed from a continuous record of indoor PM concentration, allowing separation of indoor-generated PM and ambient PM that has infiltrated indoors.
- In pooled studies (different subjects measured on different days), individual, daily values of the total PM exposure are usually not well correlated with the daily ambient PM concentrations. In longitudinal studies (each subject measured for multiple days), individual, daily values of the total PM personal exposure and the daily ambient PM concentrations are found to be highly correlated for some, but not all, subjects.
- Only one study has reported estimated individual, daily values of ambient and nonambient PM exposure. Individual, daily values of the total PM personal exposure and the daily ambient PM concentrations were poorly correlated. However, individual, daily values of ambient PM exposure and the daily ambient PM concentrations were highly correlated. Individual daily values of ambient and nonambient PM exposure were not well correlated. Individual daily values of nonambient PM exposure and daily ambient PM concentrations were also not well correlated.

REFERENCES

- Abraham, J. L.; Siwinski, G.; Hunt, A. (2002) Ultrafine particulate exposures in indoor, outdoor, personal and mobile environments: effects of diesel, traffic, pottery kiln, cooking and HEPA filtration on micro-environmental particle number concentration. *Ann. Occup. Hyg.* 46(suppl. 1): 406-411.
- Abt, E.; Suh, H. H.; Allen, G.; Koutrakis, P. (2000a) Characterization of indoor particle sources: a study conducted in the metropolitan Boston area. *Environ. Health Perspect.* 108: 35-44.
- Abt, E.; Suh, H. H.; Catalano, P.; Koutrakis, P. (2000b) The relative contribution of outdoor and indoor particle sources to indoor concentrations. *Environ. Sci. Technol.* 34: 3579-3587.
- Adams, H. S.; Nieuwenhuijsen, M. J.; Colvile, R. N.; McMullen, M. A. S.; Khandelwal, P. (2001) Fine particle (PM_{2.5}) personal exposure levels in transport microenvironments, London, UK. *Sci. Total Environ.* 279: 29-44.
- Adgate, J. L.; Ramachandran, G.; Pratt, G. C.; Waller, L. A.; Sexton, K. (2002) Spatial and temporal variability in outdoor, indoor, and personal PM_{2.5} exposure. *Atmos. Environ.* 36: 3255-3265.
- Akland, G. G.; Hartwell, T. D.; Johnson, T. R.; Whitmore, R. W. (1985) Measuring human exposure to carbon monoxide in Washington, D.C., and Denver, Colorado, during the winter of 1982-1983. *Environ. Sci. Technol.* 19: 911-918.
- Al-Raheem, M. Y. A.; El-Desouky, M.; Mage, D. T.; Kollander, M.; Wallace, L. A.; Spengler, J. D. (2000) Personal exposures to PM and gaseous air pollutants at the end of the 1991 Kuwait oil fires. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.
- Alm, S.; Jantunen, M. J.; Vartiainen, M. (1999) Urban commuter exposure to particle matter and carbon monoxide inside an automobile. *J. Exposure Anal. Environ. Epidemiol.* 9: 237-244.
- Ames, R. B.; Hand, J. L.; Kreidenweis, S. M. (2000) Optical measurements of aerosol size distributions in Great Smoky Mountains National Park: dry aerosol characterization. *J. Air Waste Manage. Assoc.* 50: 665-676.
- Anuszewski, J.; Larson, T. V.; Koenig, J. Q. (1998) Simultaneous indoor and outdoor particle light-scattering measurements at nine homes using a portable nephelometer. *J. Exposure Anal. Environ. Epidemiol.* 8: 483-493.
- Babich, P.; Davey, M.; Allen, G.; Koutrakis, P. (2000) Method comparisons for particulate nitrate, elemental carbon, and PM_{2.5} mass in seven U.S. cities. *J. Air Waste Manage. Assoc.* 50: 1095-1105.
- Baek, S.-O.; Kim, Y.-S.; Perry, R. (1997) Indoor air quality in homes, offices and restaurants in Korean urban areas—indoor/outdoor relationships. *Atmos. Environ.* 31: 529-544.
- Bahadori, T. (1998) Human particulate exposure assessment: relationship between outdoor, indoor, and personal measurements (air pollution) [dissertation]. Boston, MA: Harvard University.
- Bohadana, A. B.; Massin, N.; Wild, P.; Toamain, J.-P.; Engel, S.; Goutet, P. (2000) Symptoms, airway responsiveness, and exposure to dust in beech and oak wood workers. *Occup. Environ. Med.* 57: 268-273.
- Brauer, M. (1995) Assessment of indoor aerosols with an integrating nephelometer. *J. Exposure Anal. Environ. Epidemiol.* 5: 45-56.
- Brauer, M.; Marnett, A. (1998) Restaurant smoking restrictions and environmental tobacco smoke exposure. *Am. J. Public Health* 88: 1834-1836.
- Brauer, M.; Koutrakis, P.; Spengler, J. D. (1989) Personal exposures to acidic aerosols and gases. *Environ. Sci. Technol.* 23: 1408-1412.
- Brauer, M.; Koutrakis, P.; Keeler, G. J.; Spengler, J. D. (1990) Indoor and outdoor concentrations of acidic aerosols and gases. In: *Indoor air '90: proceedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air*; July-August; Toronto, ON, Canada. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 447-452.
- Brauer, M.; Bartlett, K.; Regalado-Pineda, J.; Perez-Padilla, R. (1996) Assessment of particulate concentrations from domestic biomass combustion in rural Mexico. *Environ. Sci. Technol.* 30: 104-109.
- Brauer, M.; Hirtle, R. D.; Hall, A. C.; Yip, T. R. (1999) Monitoring personal fine particle exposure with a particle counter. *J. Exposure Anal. Environ. Epidemiol.* 9: 228-236.
- Brauer, M.; Hrubá, F.; Mihalíková, E.; Fabiánová, E.; Miskovic, P.; Plžíková, A.; Lendacká, M.; Vanderberg, J.; Cullen, A. (2000) Personal exposure to particles in Banská Bystrica, Slovakia. *J. Exposure Anal. Environ. Epidemiol.* 10: 478-487.

- Brown, K. G.; Paxton, M. B. (1998) Predictability of personal exposure to airborne particulate matter from ambient concentrations in four communities. In: Vostal, J. J., ed. Health effects of particulate matter in ambient air. Proceedings of an international conference; 1997; Prague, Czech Republic. Pittsburgh, PA: Air & Waste Management Association; pp. 333-350. (A&WMA publication VIP-80).
- Burke, J.; Zufall, M.; Özkaynak, H. (2001) A population exposure model for particulate matter: case study results for PM_{2.5} in Philadelphia, PA. *J. Exposure Anal. Environ. Epidemiol.* 11: 470-489.
- Burton, R. M.; Suh, H. H.; Koutrakis, P. (1996) Spatial variation in particulate concentrations within metropolitan Philadelphia. *Environ. Sci. Technol.* 30: 400-407.
- Buzorius, G.; Hämeri, K.; Pekkanen, J.; Kulmala, M. (1999) Spatial variation of aerosol number concentration in Helsinki city. *Atmos. Environ.* 33: 553-565.
- Byrne, M. A.; Lange, C.; Goddard, A. J. H.; Roed, J. (1992) Indoor aerosol deposition studies using neutron-activatable tracers. *J. Aerosol Sci.* 23(suppl. 1): S543-S546.
- Calder, K. L. (1957) A numerical analysis of the protection afforded by buildings against BW aerosol attack. Fort Detrick, MD: Office of the Deputy Commander for Scientific Activities; BWL tech. study no. 2.
- Carrer, P.; Alcini, D.; Cavallo, D. (1997) Daily personal exposure to air pollutants of office workers in Milano. In: Woods, et al. (eds.), Proceedings of Healthy Buildings '97, v.2, pp. 249-254; Washington, DC.
- Carrer, P.; Alcini, D.; Cavallo, D.; Bollini, D.; Ghalandar, R.; Lovato, L.; Ploia, E.; Vercelli, F.; Visigalli, F.; Maroni, M. (1998) Personal exposure to PM₁₀ of subjects living in Milan. In: Vostal, J. J., ed. Health effects of particulate matter in ambient air. Proceedings of an international conference; 1997; Prague, Czech Republic. Pittsburgh, PA: Air & Waste Management Association; pp. 359-365. (A&WMA publication VIP-80).
- Cha, Q.; Chen, Y.-Z.; Du, Y.-X. (1996) Influence of indoor air pollution on quality of outdoor air in city of Guangzhou, China. In: Indoor air '96: the 7th international conference on indoor air quality, volume I; July; Nagoya, Japan. Tokyo, Japan: Institute of Public Health; pp. 1131-1135.
- Chan, A. T. (2002) Indoor-outdoor relationships of particulate matter and nitrogen oxides under different outdoor meteorological conditions. *Atmos. Environ.* 36: 1543-1551.
- Chan, L. Y.; Lau, W. L.; Lee, S. C.; Chan, C. Y. (2002) Commuter exposure to particulate matter in public transportation modes in Hong Kong. *Atmos. Environ.* 36: 3363-3373.
- Chang, L.-T.; Koutrakis, P.; Catalano, P. J.; Suh, H. H. (2000) Hourly personal exposures to fine particles and gaseous pollutants—results from Baltimore, Maryland. *J. Air Waste Manage. Assoc.* 50: 1223-1235.
- Chang, L.-T.; Suh, H. H.; Wolfson, J. M.; Misra, K.; Allen, G. A.; Catalano, P. J.; Koutrakis, P. (2001) Laboratory and field evaluation of measurement methods for one-hour exposures to O₃, PM_{2.5}, and CO. *J. Air Waste Manage. Assoc.* 51: 1414-1422.
- Chao, C. Y. H.; Tung, T. C. (2001) An empirical model for outdoor contaminant transmission into residential buildings and experimental verification. *Atmos. Environ.* 35: 1585-1596.
- Chuang, J. C.; Callahan, P. J.; Lyu, C. W.; Wilson, N. K. (1999) Polycyclic aromatic hydrocarbon exposures of children in low-income families. *J. Exposure Anal. Environ. Epidemiol.* 2: 85-98.
- Clayton, C. A.; Perritt, R. L.; Pellizzari, E. D.; Thomas, K. W.; Whitmore, R. W.; Wallace, L. A.; Özkaynak, H.; Spengler, J. D. (1993) Particle total exposure assessment methodology (PTEAM) study: distributions of aerosol and elemental concentrations in personal, indoor, and outdoor air samples in a southern California community. *J. Exposure Anal. Environ. Epidemiol.* 3: 227-250.
- Clayton, C. A.; Pellizzari, E. D.; Rodes, C. E.; Mason, R. E.; Piper, L. L. (1999a) Estimating distributions of long-term particulate matter and manganese exposures for residents of Toronto, Canada. *Atmos. Environ.* 33: 2515-2526.
- Clayton, C. A.; Pellizzari, E. D.; Whitmore, R. W.; Perritt, R. L.; Quackenboss, J. J. (1999b) National human exposure assessment survey (NHEXAS): distributions and associations of lead, arsenic, and volatile organic compounds in EPA Region 5. *J. Exposure Anal. Environ. Epidemiol.* 9: 381-392.
- Conner, T. L.; Lonneman, W. A.; Seila, R. L. (1995) Transportation-related volatile hydrocarbon source profiles measured in Atlanta. *J. Air Waste Manage. Assoc.* 45: 383-394.
- Conner, T. L.; Norris, G. A.; Landis, M. S.; Williams, R. W. (2001) Chemical and physical characterization of indoor, outdoor, and personal samples collected in and around a retirement facility. In: Measurement of Toxic and Related Air Pollutants: proceedings of a specialty conference; September, 2000; Research Triangle Park, NC. Pittsburgh, PA: (A&WMA publication VIP-100/CD; pp. 678-700).
- Copeland, K. T.; Checkoway, H.; McMichael, A. J.; Holbrook, R. H. (1977) Bias due to misclassification in the estimation of relative risk. *Am. J. Epidemiol.* 105: 488-495.

- Crump, K. S. (2000) Manganese exposures in Toronto during use of the gasoline additive, methylcyclopentadienyl manganese tricarbonyl. *J. Exposure Anal. Environ. Epidemiol.* 10: 227-239.
- Daisey, J. M.; Mahanama, K. R. R.; Hodgson, A. T. (1998) Toxic volatile organic compounds in simulated environmental tobacco smoke: emission factors for exposure assessment. *J. Exposure Anal. Environ. Epidemiol.* 8: 313-334.
- Demokritou, P.; Kavouras, I. G.; Harrison, D.; Koutrakis, P. (2001) Development and evaluation of an impactor for a PM_{2.5} speciation sampler. *J. Air Waste Manage. Assoc.* 51: 514-523.
- Dockery, D. W.; Spengler, J. D. (1981) Indoor-outdoor relationships of respirable sulfates and particles. *Atmos. Environ.* 15: 335-343.
- Dockery, D. W.; Pope, C. A., III; Xu, X.; Spengler, J. D.; Ware, J. H.; Fay, M. E.; Ferris, B. G., Jr.; Speizer, F. E. (1993) An association between air pollution and mortality in six U.S. cities. *N. Engl. J. Med.* 329: 1753-1759.
- Dominici, F.; Zeger, S. L.; Samet, J. (2000) A measurement error model for time-series studies of air pollution and mortality. *Biostatistics* 1: 157-175.
- Donham, K. J.; Cumro, D.; Reynolds, S. J.; Merchant, J. A. (2000) Dose-response relationships between occupational aerosol exposures and cross-shift declines of lung function in poultry workers: recommendations for exposure limits. *J. Occup. Environ. Med.* 42: 260-269.
- Douce, D. S.; Clench, M. R.; Frost, B. (2001) Variations in the estimation of the contribution of environmental tobacco smoke (ETS) to respirable ($\leq 5\mu\text{m}$) indoor air particulates obtained by the use of different analytical methods. *J. Environ. Monit.* 3: 295-301.
- Duan, N. (1982) Models for human exposure to air pollution. *Environ. Int.* 8: 305-309.
- Duan, N. (1989) Estimation of microenvironment concentration distribution using integrated exposure measurements. In: Starks, T. H., ed. *Proceedings of the Research Planning Conference on Human Activity Patterns*. Las Vegas, NV: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; pp. 15-1 - 15-12; report no. EPA/600/4-89-004. Available from: NTIS, Springfield, VA; PB89-166599.
- Duan, N. (1991) Stochastic microenvironment models for air pollution exposure. *J. Exposure Anal. Environ. Epidemiol.* 1: 235-257.
- Dubowsky, S. D.; Wallace, L. A.; Buckley, T. J. (1999) The contribution of traffic to indoor concentrations of polycyclic aromatic hydrocarbons. *J. Exposure Anal. Environ. Epidemiol.* 9: 312-321.
- Ebelt, S. T.; Petkau, A. J.; Vedal, S.; Fisher, T. V.; Brauer, M. (2000) Exposure of chronic obstructive pulmonary disease patients to particulate matter: relationships between personal and ambient air concentrations. *J. Air Waste Manage. Assoc.* 50: 1081-1094.
- Esmen, N. A.; Hall, T. A. (2000) Theoretical investigation of the interrelationship between stationary and personal sampling in exposure estimation. *Appl. Occup. Environ. Hyg.* 15: 114-119.
- Evans, G. F.; Highsmith, R. V.; Sheldon, L. S.; Suggs, J. C.; Williams, R. W.; Zweidinger, R. B.; Creason, J. P.; Walsh, D.; Rodes, C. E.; Lawless, P. A. (2000) The 1999 Fresno particulate matter exposure studies: comparison of community, outdoor, and residential PM mass measurements. *J. Air Waste Manage. Assoc.* 50: 1887-1896.
- Ezzati, M.; Kammen, D. M. (2001) Quantifying the effects of exposure to indoor air pollution from biomass combustion on acute respiratory infections in developing countries. *Environ. Health Perspect.* 109: 481-488.
- Federal Register. (1986) Guidelines for estimating exposures. *F. R.* (September 24) 51: 34,042-34,054.
- Ferris, B. G., Jr.; Speizer, F. E.; Spengler, J. D.; Dockery, D.; Bishop, Y. M. M.; Wolfson, M.; Humble, C. (1979) Effects of sulfur oxides and respirable particles on human health: methodology and demography of populations in study. *Am. Rev. Respir. Dis.* 120: 767-779.
- Fischer, P. H.; Hoek, G.; Van Reeuwijk, H.; Briggs, D. J.; Lebret, E.; Van Wijnen, J. H.; Kingham, S.; Elliott, P. E. (2000) Traffic-related differences in outdoor and indoor concentrations of particles and volatile organic compounds in Amsterdam. *Atmos. Environ.* 34: 3713-3722.
- Fogh, C. L.; Byrne, M. A.; Roed, J.; Goddard, A. J. H. (1997) Size specific indoor aerosol deposition measurements and derived I/O concentrations ratios. *Atmos. Environ.* 31: 2193-2203.
- Freijer, J. I.; Bloemen, H. J. T. (2000) Modeling relationships between indoor and outdoor air quality. *J. Air Waste Manage. Assoc.* 50: 292-300.
- Frey, H. C.; Rhodes, D. S. (1996) Characterizing, simulating and analyzing variability and uncertainty: an illustration of methods using an air toxic emissions example. *Hum. Ecol. Risk Assess.* 2: 762-797.
- Geller, M. D.; Chang, M.; Sioutas, C.; Ostro, B. D.; Lipsett, M. J. (2002) Indoor/outdoor relationship and chemical composition of fine and coarse particles in the southern California deserts. *Atmos. Environ.* 36: 1099-1110.

- Georgopoulos, P. G.; Lioy, P. J. (1994) Conceptual and theoretical aspects of human exposure and dose assessment. *J. Exposure Anal. Environ. Epidemiol.* 4: 253-285.
- Gladden, B.; Rogan, W. J. (1979) Misclassification and the design of environmental studies. *Am. J. Epidemiol.* 109: 607-616.
- Glen, G.; Lakkadi, Y.; Tippet, J. A.; del Valle-Torres, M. (1997) Development of NERL/CHAD: the National Exposure Research Laboratory consolidated human activity database. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development; contract no. 68-D5-0049.
- Goldstein, B. R.; Tardiff, G.; Hoffnagle, G.; Kester, R. (1992) Valdez health study: summary report. Anchorage, AK: Alyeska Pipeline Service Company.
- Goswami, E.; Larson, T.; Lumley, T.; Sally, Liu, L.-J. (2002) Spatial characteristics of fine particulate matter: identifying representative monitoring locations in Seattle, Washington. *J. Air Waste Manage. Assoc.* 52: 324-333.
- Gotschi, T.; Oglesby, L.; Mathys, P.; Monn, C.; Manalis, N.; Koistinen, K.; Jantunen, M.; Hanninen, O.; Polanska, L.; Kunzli, N. (2002) Comparison of black smoke and PM_{2.5} levels in indoor and outdoor environments of four European cities. *Environ. Sci. Technol.* 36: 1191-1197.
- Hadwen, G. E.; McCarthy, J. F.; Womble, S. E.; Girman, J. R.; Brightman, H. S. (1997) Volatile organic compound concentrations in 41 office buildings in the continental United States. In: Woods, J. E.; Grimsrud, D. T.; Boschi, N., eds. *Proceedings of healthy buildings/indoor air quality international conference*; September; Bethesda, MD. Washington, DC: Healthy Buildings/IAQ. [Healthy Buildings - International Conference; v. 2, pp. 465-470.]
- Harrison, R. M.; Thornton, C. A.; Lawrence, R. G.; Mark, D.; Kinnorsley, R. P.; Ayres, J. G. (2002) Personal exposure monitoring of particulate matter, nitrogen dioxide, and carbon monoxide, including susceptible groups. *Occup. Environ. Med.* 59: 671-679.
- Hartwell, T. D.; Clayton, C. A.; Michie, R. M., Jr.; Whitmore, R. W.; Zelon, H. S.; Whitehurst, D. A.; Akland, G. G. (1984) Study of carbon monoxide exposures of residents of Washington, D.C. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; report no. EPA-600/D-84-177. Available from: NTIS, Springfield, VA; PB84-230051.
- Hayes, S. R.; Marshall, J. R. (1999) Designing optimal strategies to attain the new U.S. particulate matter standards: some initial concepts. *J. Air Waste Manage. Assoc.* 49: PM192-PM198.
- Henry, R. C.; Lewis, C. W.; Collins, J. F. (1994) Vehicle-related source compositions from ambient data: the GRACE/SAFER method. *Environ. Sci. Technol.* 28: 823-832.
- Hoek, G.; Fischer, P.; Van Den Brandt, P.; Goldbohm, S.; Brunekreef, B. (2001) Estimation of long-term average exposure to outdoor air pollution for a cohort study on mortality. *J. Exposure Anal. Environ. Epidemiol.* 11: 459-469.
- Hoek, G.; Brunekreef, B.; Goldbohm, S.; Fischer, P.; Van den Brandt, P. A. (2002) Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* 360: 1203-1209.
- Houseman, E. A.; Ryan, L.; Levy, J.; Richardson, D.; Spenger, J. D. (2002) Autocorrelation in real-time continuous monitoring of microenvironments. *J. Appl. Stat.* 29: 855-872.
- Howard-Reed, C.; Rea, A. W.; Zufall, M. J.; Burke, J. M.; Williams, R. W.; Suggs, J. C.; Sheldon, L. S.; Walsh, D.; Kwok, R. (2000) Use of a continuous nephelometer to measure personal exposure to particles during the U.S. Environmental Protection Agency Baltimore and Fresno panel studies. *J. Air Waste Manage. Assoc.* 50: 1125-1132.
- Howard-Reed, C.; Wallace, L. A.; Ott, W. R. (2002) The effect of opening windows on air change rates in two homes. *J. Air Waste Manage. Assoc.* 52: 147-159.
- Institute for Social Research. (1997) Child development supplement to the panel study of income dynamics. Ann Arbor, MI: University of Michigan. Available: www.isr.umich.edu/src/child-development/home.html [2001, January 31].
- Isukapalli, S. S.; Georgopoulos, P. G. (2000) Modeling of outdoor/indoor relationships of gas phase pollutants and particulate matter. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.
- Jang, M.; Kamens, R. M. (1999) Newly characterized products and composition of secondary aerosols from the reaction of α -pinene with ozone. *Atmos. Environ.* 33: 459-474.
- Janssen, N. (1998) Personal exposure to airborne particles: validity of outdoor concentrations as a measure of exposure in time series studies [dissertation]. Wageningen, Netherlands: Landbouwniversiteit Wageningen.

- Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1995) A relation between personal and ambient . *Epidemiology* 6(suppl.): S45.
- Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1997) Childhood exposure to PM₁₀: relation between personal, classroom, and outdoor concentrations. *Occup. Environ. Med.* 54: 888-894.
- Janssen, N. A. H.; Hoek, G.; Brunekreef, B.; Harssema, H.; Mensink, I.; Zuidhof, A. (1998a) Personal sampling of particles in adults: relation among personal, indoor, and outdoor air concentrations. *Am. J. Epidemiol.* 147: 537-547.
- Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1998b) Personal sampling of airborne particles: method performance and data quality. *J. Exposure Anal. Environ. Epidemiol.* 8: 37-49.
- Janssen, N. A. H.; Hoek, G.; Harssema, H.; Brunekreef, B. (1999a) Personal exposure to fine particles in children correlates closely with ambient fine particles. *Arch. Environ. Health* 54: 95-101.
- Janssen, N. A. H.; Hoek, G.; Brunekreef, B.; Harssema, H. (1999b) Mass concentration and elemental composition of PM₁₀ in classrooms. *Occup. Environ. Med.* 56: 482-487.
- Janssen, L. H. J. M.; Buringh, E.; van der Meulen, A.; van den Hout, K. D. (1999c) A method to estimate the distribution of various fractions of PM₁₀ in ambient air in the Netherlands. *Atmos. Environ.* 33: 3325-3334.
- Janssen, N. A. H.; De Hartog, J. J.; Hoek, G.; Brunekreef, B.; Lanki, T.; Timonen, K. L.; Pekkanen, J. (2000) Personal exposure to fine particulate matter in elderly subjects: relation between personal, indoor, and outdoor concentrations. *J. Air Waste Manage. Assoc.* 50: 1133-1143.
- Jantunen, M. J.; Hänninen, O.; Katsouyanni, K.; Knöppel, H.; Kuenzli, N.; Lebret, E.; Maroni, M.; Saarela, K.; Srám, R.; Zmirou, D. (1998) Air pollution exposure in European cities: the *EXPOLIS* study. *J. Exposure Anal. Environ. Epidemiol.* 8: 495-518.
- Jantunen, M. J.; Kousa, A.; Hanninen, O.; Koistinen, K.; Sram, R.; Lanki, T.; Liroy, P. J. (2000) Long term population PM exposures correlate well with outdoor air levels - short term individual exposures do not. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.
- Jenkins, R. A.; Palausky, M. A.; Counts, R. W.; Guerin, M. R.; Dindal, A. B.; Bayne, C. K. (1996a) Determination of personal exposure of non-smokers to environmental tobacco smoke in the United States. *Lung Cancer* 14(suppl. 1): S195-S213.
- Jenkins, R. A.; Palausky, A.; Counts, R. W.; Bayne, C. K.; Dindal, A. B.; Guerin, M. R. (1996b) Exposure to environmental tobacco smoke in sixteen cities in the United States as determined by personal breathing zone air sampling. *J. Exposure Anal. Environ. Epidemiol.* 6: 473-502.
- Jinsart, W.; Tamura, K.; Loetkamonwit, S.; Thepanondh, S.; Karita, K.; Yano, E. (2002) Roadside particulate air pollution in Bangkok. *J. Air Waste Manage. Assoc.* 52: 1102-1110.
- Johnson, T. (1984) A study of personal exposure to carbon monoxide in Denver, Colorado. Research Triangle Park, NC: U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory; report no. EPA-600/4-84-014.
- Johnson, T. (1989) Human activity patterns in Cincinnati, Ohio [final report]. Palo Alto, CA: Electric Power Research Institute; project no. RP940-06; report no. EN-6204.
- Johnson, T. R. (1995) Recent advances in the estimation of population exposure to mobile source pollutants. *J. Exposure Anal. Environ. Epidemiol.* 5: 551-571.
- Johnson, T.; Long, T.; Ollison, W. (2000) Prediction of hourly microenvironmental concentrations of fine particles based on measurements obtained from the Baltimore scripted activity study. *J. Exposure Anal. Environ. Epidemiol.* 10: 403-411.
- Jones, N. C.; Thornton, C. A.; Mark, D.; Harrison, R. M. (2000) Indoor/outdoor relationships of particulate matter in domestic homes with roadside, urban and rural locations. *Atmos. Environ.* 34: 2603-2612.
- Kamens, R.; Lee, C.-T.; Wiener, R.; Leith, D. (1991) A study to characterize indoor particles in three non-smoking homes. *Atmos. Environ. Part A* 25: 929-948.
- Keeler, G. J.; Dvonch, J. T.; Yip, F. Y.; Parker, E. A.; Israel, B. A.; Marsik, F. J.; Morishita, M.; Barres, J. A.; Robins, T. G.; Brakefield-Caldwell, W.; Sam, M. (2002) Assessment of personal and community-Level exposures to particulate matter among children with asthma in Detroit, Michigan, as part of Community Action Against Asthma (CAAA). *Environ. Health Perspect.* 110(suppl. 2): 173-181
- Keywood, M. D.; Ayers, G. P.; Gras, J. L.; Gillett, R. W.; Cohen, D. D. (1999) Relationships between size segregated mass concentration data and ultrafine particle number concentrations in urban areas. *Atmos. Environ.* 33: 2907-2913.

- Kingham, S.; Briggs, D.; Elliott, P.; Fischer, P.; Lebrecht, E. (2000) Spatial variations in the concentrations of traffic-related pollutants in indoor and outdoor air in Huddersfield, England. *Atmos. Environ.* 34: 905-916.
- Kinney, P. L.; Chillrud, S. N.; Ramstrom, S.; Ross, J.; Spengler, J. D. (2002) Exposures to multiple air toxics in New York City. *Environ. Health Perspect.* 110(suppl. 4): 539-546.
- Kleeman, M. J.; Cass, G. R. (1998) Source contributions to the size and composition distribution of urban particulate air pollution. *Atmos. Environ.* 32: 2803-2816.
- Klepeis, N. E. (1999) An introduction to the indirect exposure assessment approach: modeling human exposure using microenvironmental measurements and the recent national human activity pattern survey. *Environ. Health Perspect. Suppl.* 107(2): 365-374.
- Klepeis, N. E.; Ott, W. R.; Switzer, P. (1994) A Total Human Exposure Model (THEM) for Respirable Suspended Particles (RSP). Presented at: 87th annual meeting & exhibition of the Air & Waste Management Association; June; Cincinnati, OH. Pittsburgh, PA: Air & Waste Management Association; paper no. 94-WA75A.03. Available from: NTIS, Springfield, VA; PB94-197415.
- Klepeis, N. E.; Ott, W. R.; Switzer, P. (1995) Modeling the time series of respirable suspended particles and carbon monoxide from multiple smokers: validation in two public smoking lounges. Presented at: 88th annual meeting and exhibition of the Air & Waste Management Association; June; San Antonio, TX. Pittsburgh, PA: Air & Waste Management Association; paper no. 95-MPH.03.
- Klepeis, N. E.; Ott, W. R.; Switzer, P. (1996) A multiple-smoker model for predicting indoor air quality in public lounges. *Environ. Sci. Technol.* 30: 2813-2820.
- Klepeis, N. E.; Switzer, P.; Ott, W. R. (2000) Estimating exposure to environmental tobacco smoke particles using 24-hour time-activity recall diaries. Presented at: PM2000: particulate matter and health—the scientific basis for regulatory decision-making, specialty conference & exhibition; January; Charleston, SC. Pittsburgh, PA: Air & Waste Management Association.
- Koistinen, K. J.; Kousa, A.; Tenhola, V.; Hänninen, O.; Jantunen, M. J.; Oglesby, L.; Kuenzli, N.; Georgoulis, L. (1999) Fine particle (PM_{2.5}) measurement methodology, quality assurance procedures and pilot results of the *EXPOLIS* study. *J. Air Waste Manage. Assoc.* 49: 1212-1220.
- Koontz, M. D.; Rector, H. E. (1995) Estimation of distributions for residential air exchange rates: final report. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics; contract nos. 68-D9-0166 and 68-D3-0013.
- Koontz, M. D.; Niang, L. L. (1998) California population indoor exposure model (CPIEM): version 1.4F. Final report A933-157: user's guide. Sacramento, CA: California Air Resources Board; GEOMET report no. IE-2632.
- Korenstein, S.; Piazza, B. (2002) An exposure assessment of PM₁₀ from a major highway interchange: are children in nearby schools at risk? *J. Environ. Health* 65: 9-17.
- Kousa, A.; Oglesby, L.; Koistinen, K.; Künzli, N.; Jantunen, M. (2002) Exposure chain of urban air PM_{2.5}—associations between ambient fixed site, residential outdoor, indoor, workplace and personal exposures in four European cities in the *EXPOLIS*-study. *Atmos. Environ.* 36: 3031-3039.
- Koutrakis, P.; Briggs, S. L. K.; Leaderer, B. P. (1992) Source apportionment of indoor aerosols in Suffolk and Onondaga Counties, New York. *Environ. Sci. Technol.* 26: 521-527.
- Krewski, D.; Burnett, R. T.; Goldberg, M. S.; Hoover, K.; Siemiatycki, J.; Jerrett, M.; Abrahamowicz, M.; White, W. H. (2000) Reanalysis of the Harvard Six Cities study and the American Cancer Society study of particulate air pollution and mortality: a special report of the Institute's Particle Epidemiology Reanalysis Project. Cambridge, MA: Health Effects Institute.
- Lachenmyer, C.; Hidy, G. M. (2000) Urban measurements of outdoor-indoor PM_{2.5} concentrations and personal exposure in the deep south. Part I. Pilot study of mass concentrations for nonsmoking subjects. *Aerosol Sci. Technol.* 32: 34-51.
- Laden, F.; Neas, L. M.; Dockery, D. W.; Schwartz, J. (2000) Association of fine particulate matter from different sources with daily mortality in six U.S. cities. *Environ. Health Perspect.* 108: 941-947.
- Landis, M. S.; Norris, G. A.; Williams, R. W.; Weinstein, J. P. (2001) Personal exposures to PM_{2.5} mass and trace elements in Baltimore, MD, USA. *Atmos. Environ.* 35: 6511-6524.
- Lanki, T.; Alm, S.; Ruuskanen, J.; Janssen, N. A. H.; Jantunen, M.; Pekkanen, J. (2002) Photometrically measured continuous personal PM_{2.5} exposure: levels and correlation to a gravimetric method. *J. Exposure Anal. Environ. Epidemiol.* 12: 172-178.

- Leaderer, B. P.; Naeher, L.; Jankun, T.; Balenger, K.; Holford, T. R.; Toth, C.; Sullivan, J.; Wolfson, J. M.; Koutrakis, P. (1999) Indoor, outdoor, and regional summer and winter concentrations of PM₁₀, PM_{2.5}, SO₄²⁻, H⁺, NH₄⁺, NO₃⁻, NH₃, and nitrous acid in homes with and without kerosene space heaters. *Environ. Health Perspect.* 107: 223-231.
- Lee, S. C.; Chang, M. (1999) Indoor air quality investigations at five classrooms. *Indoor Air* 9: 134-138.
- Lena, T. S.; Ochieng, V.; Carter, M.; Holguin-Veras, J.; Kinney, P. L. (2002) Elemental carbon and PM_{2.5} levels in an urban community heavily impacted by truck traffic. *Environ. Health Perspect.* 110: 1009-1015.
- Lessler, J. T.; Kalsbeek, W. D. (1992) Nonsampling error in surveys. New York, NY: John Wiley & Sons, Inc.; Levy, J. I.; Dumyahn, T.; Spengler, J. D. (2002) Particulate matter and polycyclic aromatic hydrocarbon concentrations in indoor and outdoor microenvironments in Boston, Massachusetts. *J. Exposure Anal. Environ. Epidemiol.* 12: 104-114.
- Liao, D.; Creason, J.; Shy, C.; Williams, R.; Watts, R.; Zweidinger, R. (1999) Daily variation of particulate air pollution and poor cardiac autonomic control in the elderly. *Environ. Health Perspect.* 107: 521-525.
- Ligocki, M. P.; Liu, H. I. H.; Cass, G. R.; John, W. (1990) Measurements of particle deposition rates inside Southern California museums. *Aerosol Sci. Technol.* 13: 85-101.
- Lillquist, D. R.; Lee, J. S.; Ramsay, J. R.; Boucher, K. M.; Walton, Z. L.; Lyon, J. L. (1998) A comparison of indoor/outdoor PM₁₀ concentrations measured at three hospitals and a centrally located monitor in Utah. *Appl. Occup. Environ. Hyg.* 13: 409-415.
- Linn, W. S.; Gong, H., Jr.; Clark, K. W.; Anderson, K. R. (1999) Day-to-day particulate exposures and health changes in Los Angeles area residents with severe lung disease. *J. Air Waste Manage. Assoc.* 49: PM108-PM115.
- Lioy, P. J. (1990) Assessing total human exposure to contaminants: a multidisciplinary approach. *Environ. Sci. Technol.* 24: 938-945.
- Lioy, P. J.; Waldman, J. M.; Buckley, T.; Butler, J.; Pietarinen, C. (1990) The personal, indoor and outdoor concentrations of PM-10 measured in an industrial community during the winter. *Atmos. Environ. Part B* 24: 57-66.
- Lioy, P. J.; Wainman, T.; Zhang, J.; Goldsmith, S. (1999) Typical household vacuum cleaners: the collection efficiency and emissions characteristics for fine particles. *J. Air Waste Manage. Assoc.* 49: 200-206.
- Lippmann, M.; Ito, K.; Nádas, A.; Burnett, R. T. (2000) Association of particulate matter components with daily mortality and morbidity in urban populations. Cambridge, MA: Health Effects Institute; research report no. 95.
- Liu, D.-L.; Nazaroff, W. W. (2001) Modeling pollutant penetration across building envelopes. *Atmos. Environ.* 35: 4451-4462.
- Liu, L.-J. S.; Slaughter, J. C.; Larson, T. V. (2002) Comparison of light scattering devices and impactors for particulate measurements in indoor, outdoor, and personal environments. *Environ. Sci. Technol.* 36: 2977-2986.
- Long, C. M.; Suh, H. H.; Koutrakis, P. (2000) Characterization of indoor particle sources using continuous mass and size monitors. *J. Air Waste Manage. Assoc.* 50: 1236-1250.
- Long, C. M.; Suh, H. H.; Catalano, P. J.; Koutrakis, P. (2001a) Using time- and size-resolved particulate data to quantify indoor penetration and deposition behavior. *Environ. Sci. Technol.* 35: 2089-2099.
- Long, C. M.; Suh, H. H.; Kobzik, L.; Catalano, P. J.; Ning, Y. Y.; Koutrakis, P. (2001b) A pilot investigation of the relative toxicity of indoor and outdoor fine particles: *in-vitro* effects of endotoxin and other particulate properties. *Environ. Health Perspect.* 109: 1019-1026.
- Lurmann, F. W.; Korc, M. E. (1994) User's guide to the regional human exposure (REHEX) model. San Francisco, CA: Bay Area Air Quality Management District; draft report no. STI-93150-1414-DR.
- MacIntosh, D. L.; Xue, J.; Özkaynak, H.; Spengler, J. D.; Ryan, P. B. (1995) A population-based exposure model for benzene. *J. Exposure Anal. Environ. Epidemiol.* 5: 375-403.
- Magari, S. R.; Schwartz, J.; Williams, P. L.; Hauser, R.; Smith, T. J.; Christiani, D. C. (2002) The association between personal measurements of environmental exposure to particulates and heart rate variability. *Epidemiology* 13: 305-310.
- Mage, D. T. (1985) Concepts of human exposure assessment for airborne particulate matter. *Environ. Int.* 11: 407-412.
- Mage, D. T. (1998) How much PM of ambient origin are people exposed to? In: Measurement of toxic and related air pollutants. Volume II: proceedings of a specialty conference; September; Cary, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 666-681. (A&WMA publication VIP-85).

- Mage, D. T.; Ott, W. R. (1996) Accounting for nonuniform mixing and human exposure in indoor environments. In: Tichenor, B. A., ed. *Characterizing sources of indoor air pollution and related sink effects*. Philadelphia, PA: American Society for Testing and Materials; pp. 263-278.
- Mage, D.; Wilson, W.; Hasselblad, V.; Grant, L. (1999) Assessment of human exposure to ambient particulate matter. *J. Air Waste Manage. Assoc.* 49: 174-185.
- Mar, T. F.; Norris, G. A.; Koenig, J. Q.; Larson, T. V. (2000) Associations between air pollution and mortality in Phoenix, 1995-1997. *Environ. Health Perspect.* 108: 347-353.
- Marple, V. A.; Rubow, K. L.; Turner, W.; Spengler, J. D. (1987) Low flow rate sharp cut impactors for indoor air sampling: design and calibration. *JAPCA* 37: 1303-1307.
- McBride, S. J.; Ferro, A. R.; Ott, W. R.; Switzer, P.; Hildemann, L. M. (1999) Investigations of the proximity effect for pollutants in the indoor environment. *J. Exposure Anal. Environ. Epidemiol.* 9: 602-621.
- McCurdy, T. (1995) Estimating human exposure to selected motor vehicle pollutants using the NEM series of models: lessons to be learned. *J. Exposure Anal. Environ. Epidemiol.* 5: 533-550.
- McCurdy, T. (1997) Modeling the dose profile in human exposure assessments: ozone as an example. *Rev. Toxicol.* 1: 3-23.
- McCurdy, T. (2000) Conceptual basis for multi-route intake dose modeling using an energy expenditure approach. *J. Exposure Anal. Environ. Epidemiol.* 10: 86-97.
- McCurdy, T.; Glen, G.; Smith, L.; Lakkadi, Y. (2000) The National Exposure Research Laboratory's Consolidated Human Activity Database. *J. Exposure Anal. Environ. Epidemiol.* 10: 566-578.
- Mihlan, G. J.; Todd, L. A.; Truong, K. N. (2000) Assessment of occupational exposure patterns by frequency-domain analysis of time series data. *Appl. Occup. Environ. Hyg.* 15: 120-130.
- Miller, S. L.; Branoff, S.; Nazaroff, W. W. (1998) Exposure to toxic air contaminants in environmental tobacco smoke: an assessment for California based on personal monitoring data. *J. Exposure Anal. Environ. Epidemiol.* 8: 287-311.
- Monn, C. (2001) Exposure assessment of air pollutants: a review on spatial heterogeneity and indoor/outdoor/personal exposure to suspended particulate matter, nitrogen dioxide and ozone. *Atmos. Environ.* 35: 1-32.
- Monn, Ch.; Carabias, V.; Junker, M.; Waeber, R.; Karrer, M.; Wanner, H. U. (1997) Small-scale spatial variability of particulate matter <math><10\ \mu\text{m}</math> (PM_{10}) and nitrogen dioxide. *Atmos. Environ.* 31: 2243-2247.
- Morawska, L.; He, C.; Hitchins, J.; Gilbert, D.; Parappukaran, S. (2001) The relationship between indoor and outdoor airborne particles in the residential environment. *Atmos. Environ.* 35: 3463-3473.
- Muraleedharan, T. R.; Radojevic, M. (2000) Personal particle exposure monitoring using nephelometry during haze in Brunei. *Atmos. Environ.* 34: 2733-2738.
- Murray, D. M.; Burmaster, D. E. (1995) Residential air exchange rates in the United States: empirical and estimated parametric distributions by season and climatic region. *Risk Anal.* 15: 459-465.
- National Air Pollution Control Administration. (1969) *Air quality criteria for particulate matter*. Washington, DC: U.S. Department of Health, Education, and Welfare, Public Health Service; NAPCA publication no. AP-49. Available from: NTIS, Springfield, VA; PB-190251/BA.
- National Research Council. (1991) *Human exposure assessment for airborne pollutants: advances and opportunities*. Washington, DC: National Academy of Sciences.
- National Research Council. (1998) *Research priorities for airborne particulate matter. I. Immediate priorities and a long-range research portfolio*. Washington, DC: National Academy Press.
- National Research Council. (1999) *Research priorities for airborne particulate matter. II. Evaluating research progress and updating the portfolio*. Washington, DC: National Academy Press.
- Naumova, Y. Y.; Eisenreich, S. J.; Turpin, B. J.; Weisel, C. P.; Morandi, M. T.; Colome, S. D.; Totten, L. A.; Stock, T. H.; Winer, A. M.; Alimokhtari, S.; Kwon, J.; Shendell, D.; Jones, J.; Maberti, S.; Wall, S. J. (2002) Polycyclic aromatic hydrocarbons in the indoor and outdoor air of three cities in the U.S. *Environ. Sci. Technol.* 36: 2552-2559.
- Navidi, W.; Thomas, D.; Langholz, B.; Stram, D. (1999) *Statistical methods for epidemiologic studies of the health effects of air pollution*. Cambridge, MA: Health Effects Institute; research report no. 86.
- Nelson, P. R.; Conrad, F. W.; Kelly, S. P.; Maiolo, K. C.; Richardson, J. D.; Ogden, M. W. (1998) Composition of environmental tobacco smoke (ETS) from international cigarettes part II: nine country follow-up. *Environ. Int.* 24: 251-257.
- Nieuwenhuijsen, M. J.; Noderer, K. S.; Schenker, M. B.; Vallyathan, V.; Olenchock, S. (1999) Personal exposure to dust, endotoxin and crystalline silica in California agriculture. *Ann. Occup. Hyg.* 43: 35-42.

- Oglesby, L.; Künzli, N.; Rösli, M.; Braun-Fahrländer, C.; Mathys, P.; Stern, W.; Jantunen, M.; Kousa, A. (2000) Validity of ambient levels of fine particles as surrogate for personal exposure to outdoor air pollution—results of the European EXPOLIS-EAS study (Swiss Center Basel). *J. Air Waste Manage. Assoc.* 50: 1251-1261.
- Ostro, B. D.; Hurley, S.; Lipsett, M. J. (1999) Air pollution and daily mortality in the Coachella Valley, California: a study of PM₁₀ dominated by coarse particles. *Environ. Res.* 81: 231-238.
- Ostro, B. D.; Broadwin, R.; Lipsett, M. J. (2000) Coarse and fine particles and daily mortality in the Coachella Valley, California: a follow-up study. *J. Exposure Anal. Environ. Epidemiol.* 10: 412-419.
- Ott, W. R. (1984) Exposure estimates based on computer generated activity patterns. *J. Toxicol. Clin. Toxicol.* 21: 97-128.
- Ott, W. R. (1999) Mathematical models for predicting indoor air quality from smoking activity. *Environ. Health Perspect.* 107(suppl. 2): 375-381.
- Ott, W.; Thomas, J.; Mage, D.; Wallace, L. (1988) Validation of the simulation of human activity and pollutant exposure (SHAPE) model using paired days from the Denver, CO, carbon monoxide field study. *Atmos. Environ.* 22: 2101-2113.
- Ott, W. R.; Mage, D. T.; Thomas, J. (1992) Comparison of microenvironmental CO concentrations in two cities for human exposure modeling. *J. Exposure Anal. Environ. Epidemiol.* 2: 249-267.
- Ott, W. R.; Klepeis, N. E.; Switzer, P. (1995) Modeling environmental tobacco smoke in the home using transfer functions. Presented at: 88th annual meeting and exhibition of the Air & Waste Management Association; June; San Antonio, TX; paper no. 95-WP84B.03. Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research And Exposure Assessment Laboratory; report no. EPA/600/A-95/079. Available from: NTIS, Springfield, VA; PB95-225488.
- Ott, W.; Switzer, P.; Robinson, J. (1996) Particle concentrations inside a tavern before and after prohibition of smoking: evaluating the performance of an indoor air quality model. *J. Air Waste Manage. Assoc.* 46: 1120-1134.
- Ott, W.; Wallace, L.; Mage, D. (2000) Predicting particulate (PM₁₀) personal exposure distributions using a random component superposition (RCS) statistical model. *J. Air Waste Manage. Assoc.* 50: 1390-1406.
- Owen, M. K.; Ensor, D. S.; Sparks, L. E. (1992) Airborne particle sizes and sources found in indoor air. *Atmos. Environ. Part A* 26: 2149-2162.
- Özkaynak, H.; Spengler, J. (1996) The role of outdoor particulate matter in assessing total human exposure. In: Wilson, R.; Spengler, J., eds. *Particles in our air: concentrations and health effects*. Cambridge, MA: Harvard University Press.
- Özkaynak, H.; Thurston, G. D. (1987) Associations between 1980 U.S. mortality rates and alternative measures of airborne particle concentration. *Risk Anal.* 7: 449-461.
- Özkaynak, H.; Ryan, P. B.; Spengler, J. D.; Laird, N. M. (1986) Bias due to misclassification of personal exposures in epidemiologic studies of indoor and outdoor air pollution. In: Berglund, B.; Berglund, U.; Lindvall, T.; Spengler, J.; Sundell, J., eds. *Indoor air quality: papers from the third international conference on indoor air quality and climate; August 1984; Stockholm, Sweden*. *Environ. Int.* 12: 389-393.
- Özkaynak, H.; Spengler, J. D.; Ludwig, J. F.; Butler, D. A.; Clayton, C. A.; Pellizzari, E.; Wiener, R. W. (1990) Personal exposure to particulate matter: findings from the Particle Total Exposure Assessment Methodology (PTEAM) prepilot study. In: *Indoor air '90: proceedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air; July-August; Toronto, ON, Canada*. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 571-576.
- Özkaynak, H.; Spengler, J. D.; Xue, J.; Koutrakis, P.; Pellizzari, E. D.; Wallace, L. (1993) Sources and factors influencing personal and indoor exposures to particles, elements and nicotine: findings from the particle TEAM pilot study. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. *Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate, v. 3, combustion products, risk assessment, policies; July; Helsinki, Finland*. Helsinki, Finland: Indoor Air '93; pp. 457-462.
- Özkaynak, H.; Xue, J.; Spengler, J.; Wallace, L.; Pellizzari, E.; Jenkins, P. (1996a) Personal exposures to airborne particles and metals: results from the particle TEAM study in Riverside, California. *J. Exposure Anal. Environ. Epidemiol.* 6: 57-78.
- Özkaynak, H.; Xue, J.; Weker, R.; Butler, D.; Koutrakis, P.; Spengler, J. (1996b) The particle TEAM (PTEAM) study: analysis of the data: final report, volume III. Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/R-95/098. Available from: NTIS, Springfield, VA; PB97-102495.

- Patterson, E.; Eatough, D. J. (2000) Indoor/outdoor relationships for ambient PM_{2.5} and associated pollutants: epidemiological implications in Lindon, Utah. *J. Air Waste Manage. Assoc.* 50: 103-110.
- Pellizzari, E. D.; Mason, R. E.; Clayton, C. A.; Thomas, K. W.; Cooper, S.; Piper, L.; Rodes, C.; Goldberg, M.; Roberds, J.; Michael, L. (1998) Manganese exposure study (Toronto). Final report. Research Triangle Park, NC: Research Triangle Institute, Analytical and Chemical Sciences; report no. RTI/6312/02-01 F.
- Pellizzari, E. D.; Clayton, C. A.; Rodes, C. E.; Mason, R. E.; Piper, L. L.; Fort, B.; Pfeifer, G.; Lynam, D. (1999) Particulate matter and manganese exposures in Toronto, Canada. *Atmos. Environ.* 33: 721-734.
- Pellizzari, E. D.; Clayton, C. A.; Rodes, C. E.; Mason, R. E.; Piper, L. L.; Fort, B.; Pfeifer, G.; Lynam, D. (2001) Particulate matter and manganese exposures in Indianapolis, Indiana. *J. Exposure Anal. Environ. Epidemiol.* 11: 423-440.
- Phillips, K.; Howard, D. A.; Browne, D.; Lewsley, J. M. (1994) Assessment of personal exposures to environmental tobacco smoke in British nonsmokers. *Environ. Int.* 20: 693-712.
- Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G. (1996) Assessment of air quality in Stockholm by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. *Scand. J. Work Environ. Health* 22(suppl. 1): 1-24.
- Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G.; Huici, A. (1997a) Assessment of air quality in Barcelona by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. *Environ. Int.* 23: 173-196.
- Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1997b) Assessment of air quality in Turin by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. *Environ. Int.* 23: 851-871.
- Phillips, K.; Bentley, M. C.; Howard, D. A.; Alván, G. (1998a) Assessment of air quality in Paris by personal monitoring of nonsmokers for respirable suspended particles and environmental tobacco smoke. *Environ. Int.* 24: 405-425.
- Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1998b) Assessment of environmental tobacco smoke and respirable suspended particle exposures for nonsmokers in Lisbon by personal monitoring. *Environ. Int.* 24: 301-324.
- Phillips, K.; Howard, D. A.; Bentley, M. C.; Alván, G. (1999) Assessment of environmental tobacco smoke and respirable suspended particle exposures for nonsmokers in Basel by personal monitoring. *Atmos. Environ.* 33: 1889-1904.
- Pope, C. A., III; Thun, M. J.; Namboodiri, M. M.; Dockery, D. W.; Evans, J. S.; Speizer, F. E.; Heath, C. W., Jr. (1995) Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *Am. J. Respir. Crit. Care Med.* 151: 669-674.
- Pope, C. A., III; Hill, R. W.; Villegas, G. M. (1999) Particulate air pollution and daily mortality on Utah's Wasatch Front. *Environ. Health Perspect.* 107: 567-573.
- Praml, G.; Schierl, R. (2000) Dust exposure in Munich public transportation: a comprehensive 4-year survey in buses and trams. *Int. Arch. Occup. Environ. Health* 73: 209-214.
- Quintana, P. J. E.; Samimi, B. S.; Kleinman, M. T.; Liu, L.-J.; Soto, K.; Warner, G. Y.; Bufalino, C.; Valencia, J.; Francis, D.; Hovell, M. H.; Delfino, R. J. (2000) Evaluation of a real-time passive personal particle monitor in fixed site residential indoor and ambient measurements. *J. Exposure Anal. Environ. Epidemiol.* 10: 437-445.
- Quintana, P. J. E.; Valenzia, J. R.; Delfino, R. J.; Liu, L.-J. S. (2001) Monitoring of 1-min personal particulate matter exposures in relation to voice-recorded time-activity data. *Environ. Res.* 87: 199-213.
- Ramachandran, G.; Vincent, J. H. (1999) A Bayesian approach to retrospective exposure assessment. *Appl. Occup. Environ. Hyg.* 14: 547-557.
- Rea, A. W.; Zufall, M. J.; Williams, R. W.; Sheldon, L. S.; Howard-Reed, C. (2001) The influence of human activity patterns on personal PM exposure: a comparative analysis of filter-based and continuous particle measurements. *J. Air Waste Manage. Assoc.* 51: 1271-1279.
- Ren, P.; Jankun, T. M.; Leaderer, B. P. (1999) Comparisons of seasonal fungal prevalence in indoor and outdoor air and in house dusts of dwellings in one northeast American county. *J. Exposure Anal. Environ. Epidemiol.* 9: 560-568.
- Riediker, N.M.; Keller, S.; Wuthrich, B.; Koller, T.; Monn, C. (2000) Personal pollen exposure compared to stationary measurements. *J. Invest. Allergol. Clin. Immunol.* 10: 200-203.
- Riley, W. J.; McKone, T. E.; Lai, A. C. K.; Nazaroff, W. W. (2002) Indoor particulate matter of outdoor origin: importance of size-dependent removal mechanisms. *Environ. Sci. Technol.* 36: 200-207.

- Robinson, J. P.; Wiley, J. A.; Piazza, T.; Garrett, K.; Cirksena, K. (1991) Activity patterns of California residents and their implications for potential exposure to pollution (final report). Sacramento, CA: California Air Resources Board report no. CARB-A6-177-33.
- Robinson, J. P.; Switzer, P.; Ott, W. (1994) Microenvironmental factors related to Californians' potential exposure to environmental tobacco smoke (ETS). Research Triangle Park, NC: U.S. Environmental Protection Agency, Atmospheric Research and Exposure Assessment Laboratory; report no. EPA/600/R-94/116. Available from: NTIS, Springfield, VA; PB94-214814.
- Rodes, C.; Sheldon, L.; Whitaker, D.; Clayton, A.; Fitzgerald, K.; Flanagan, J.; DiGenova, F.; Hering, S.; Frazier, C. (1998) Measuring concentrations of selected air pollutants inside California vehicles [final report]. Sacramento, CA: California Environmental Protection Agency, Air Resources Board; contract no. 95-339.
- Rodes, C. E.; Lawless, P. A.; Evans, G. F.; Sheldon, L. S.; Williams, R. W.; Vette, A. F.; Creason, J. P.; Walsh, D. (2001) The relationships between personal PM exposures for elderly populations and indoor and outdoor concentrations for three retirement center scenarios. *J. Exposure Anal. Environ. Epidemiol.* 11: 103-115.
- Rojas-Bracho, L.; Suh, H. H.; Koutrakis, P. (2000) Relationships among personal, indoor, and outdoor fine and coarse particle concentrations for individuals with COPD. *J. Exposure Anal. Environ. Epidemiol.* 10: 294-306.
- Rojas-Bracho, L.; Suh, H. H.; Oyola, P.; Koutrakis, P. (2002) Measurements of children's exposures to particles and nitrogen dioxide in Santiago, Chile. *Sci. Total Environ.* 287: 249-264.
- Roorda-Knape, M. C.; Janssen, N. A. H.; De Hartog, J. J.; Van Vliet, P. H. N.; Harsssema, H.; Brunekreef, B. (1998) Air pollution from traffic in city districts near major motorways. *Atmos. Environ.* 32: 1921-1930.
- Rotko, T.; Oglesby, L.; Künzli, N.; Jantunen, M. J. (2000a) Population sampling in European air pollution exposure study, *EXPOLIS*: comparisons between the cities and representativeness of the samples. *J. Exposure Anal. Environ. Epidemiol.* 10: 355-364.
- Rotko, T.; Koistinen, K.; Hänninen, O.; Jantunen, M. (2000b) Sociodemographic descriptors of personal exposure to fine particles (PM_{2.5}) in *EXPOLIS* Helsinki. *J. Exposure Anal. Environ. Epidemiol.* 10: 385-393.
- Ryan, P. B. (1991) An overview of human exposure modeling. *J. Exposure Anal. Environ. Epidemiol.* 1: 453-474.
- Samet, J. M.; Dominici, F.; Zeger, S. L.; Schwartz, J.; Dockery, D. W. (2000) National morbidity, mortality, and air pollution study. Part I: methods and methodologic issues. Cambridge, MA: Health Effects Institute; research report no. 94.
- Santos-Burgoa, C.; Rojas-Bracho, L.; Rosas-Pérez, I.; Ramírez-Sánchez, A.; Sánchez-Rico, G.; Mejía-Hernández, S. (1998) Modelaje de exposición a partículas en población general y riesgo de enfermedad respiratoria [Particle exposure modeling in the general population and risk of respiratory disease]. *Gac. Med. Mex.* 134: 407-417.
- Sarnat, J. A.; Koutrakis, P.; Suh, H. H. (2000) Assessing the relationship between personal particulate and gaseous exposures of senior citizens living in Baltimore, MD. *J. Air Waste Manage. Assoc.* 50: 1184-1198.
- Sarnat, J. A.; Schwartz, J.; Catalano, P. J.; Suh, H. H. (2001) Gaseous pollutants in particulate matter epidemiology: confounders or surrogates? *Environ. Health Perspect.* 109: 1053-1061.
- Sarnat, J. A.; Long, C. M.; Koutrakis, P.; Coull, B. A.; Schwartz, J.; Suh, H. H. (2002) Using sulfur as a tracer of outdoor fine particulate matter. *Environ. Sci. Technol.* 36: 5305-5314.
- Scheff, P. A.; Wadden, R. (1979) Comparison of three methods of particulate measurement in Chicago air. *Atmos. Environ.* 13: 639-643.
- Schwartz, J.; Dockery, D. W.; Neas, L. M. (1996) Is daily mortality associated specifically with fine particles? *J. Air Waste Manage. Assoc.* 46: 927-939.
- Sexton, K.; Ryan, P. B. (1988) Assessment of human exposure to air pollution: methods, measurements, and models. In: Watson, A. Y.; Bates, R. R.; Kennedy, D., eds. *Air pollution, the automobile, and public health*. Washington, DC: National Academy Press; pp. 207-238.
- Sexton, K.; Spengler, J. D.; Treitman, R. D. (1984) Personal exposure to respirable particles: a case study in Waterbury, Vermont. *Atmos. Environ.* 18: 1385-1398.
- Sheldon, L.; Clayton, A.; Jones, B.; Keever, J.; Perritt, R.; Smith, D.; Whitaker, D.; Whitmore, R. (1992a) Indoor pollutant concentrations and exposures. Sacramento, CA: California Air Resources Board; report no. ARB/R-92/486.
- Sheldon, L.; Clayton, A.; Keever, J.; Perritt, R.; Whitaker, D. (1992b) PTEAM: monitoring of phthalates and PAHs in indoor and outdoor air samples in Riverside, California, volume II [final report]. Sacramento, CA: California Environmental Protection Agency, State Air Resources Board, Research Division; contract no. A933-144. Available from: NTIS, Springfield, VA; PB93-205649/XAB.

- Sheldon, L.; Whitaker, D.; Keever, J.; Clayton, A.; Perritt, R. (1993a) Phthalates and PAHs in indoor and outdoor air in a southern California community. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 109-114.
- Sheldon, L.; Clayton, A.; Keever, J.; Perritt, R.; Whitaker, D. (1993b) Indoor concentrations of polycyclic aromatic hydrocarbons in California residences. Sacramento, CA: Air Resources Board; ARB contract no. A033-132.
- Sheldon, L.; Clayton, A.; Perritt, R.; Whitaker, D. A.; Keever, J. (1993c) Indoor concentrations of polycyclic aromatic hydrocarbons in California residences and their relationship to combustion source use. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 29-34.
- Shy, C. M.; Kleinbaum, D. G.; Morgenstern, H. (1978) The effect of misclassification of exposure status in epidemiological studies of air pollution health effects. *Bull. N. Y. Acad. Med.* 54: 1155-1165.
- Siegmann, K.; Scherrer, L.; Siegmann, H. C. (1999) Physical and chemical properties of airborne nanoscale particles and how to measure the impact on human health. *J. Mol. Struct. (Theochem)* 458: 191-201.
- Sinclair, J. D.; Psota-Kelty, L. A.; Weschler, C. J. (1988) Indoor/outdoor ratios and indoor surface accumulations of ionic substances at Newark, New Jersey. *Atmos. Environ.* 22: 461-469.
- Sinclair, J. D.; Psota-Kelty, L. A.; Weschler, C. J.; Shields, H. C. (1990) Measurement and modeling of airborne concentrations and indoor surface accumulation rates of ionic substances at Neenah, Wisconsin. *Atmos. Environ. Part A* 24: 627-638.
- Sioutas, C.; Wang, P. Y.; Ferguson, S. T.; Koutrakis, P.; Mulik, J. D. (1996) Laboratory and field evaluation of an improved glass honeycomb denuder/filter pack sampler. *Atmos. Environ.* 30: 885-895.
- Sioutas, C.; Kim, S.; Chang, M.; Terrell, L. L.; Gong, H., Jr. (2000) Field evaluation of a modified DataRAM MIE scattering monitor for real-time PM_{2.5} mass concentration measurements. *Atmos. Environ.* 34: 4829-4838.
- Sørensen, M.; Daneshvar, B.; Hansen, M.; Dragsted, L. O.; Hertel, O.; Knudsen, L.; Loft, S. (2003) Personal PM_{2.5} exposure and markers of oxidative stress in blood. *Environ. Health Perspect.* 111: 161-165.
- Spengler, J. D.; Treitman, R. D.; Tosteson, T. D.; Mage, D. T.; Soczek, M. L. (1985) Personal exposures to respirable particulates and implications for air pollution epidemiology. *Environ. Sci. Technol.* 19: 700-707.
- Spengler, J. D.; Özkaynak, H.; Ludwig, J.; Allen, G.; Pellizzari, E. D.; Wiener, R. (1989) Personal exposures to particulate matter: instruments and methodologies P-TEAM. In: Measurement of toxic and related air pollutants: proceedings of the 1989 U.S. EPA/A&WMA international symposium; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 449-463. (A&WMA publication VIP-13).
- Spier, C. E.; Little, D. E.; Trim, S. C.; Johnson, T. R.; Linn, W. S.; Hackney, J. D. (1992) Activity patterns in elementary and high school students exposed to oxidant pollution. *J. Exposure Anal. Environ. Epidemiol.* 2: 277-293.
- Suggs, J. C.; Burton, R. M. (1983) Spatial characteristics of inhalable particles in the Philadelphia metropolitan area. *J. Air Pollut. Control Assoc.* 33: 688-691.
- Suh, H. H.; Spengler, J. D.; Koutrakis, P. (1992) Personal exposures to acid aerosols and ammonia. *Environ. Sci. Technol.* 26: 2507-2517.
- Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1993a) Indoor and outdoor acid aerosol and gas concentrations. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 257-262.
- Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1993b) Validation of personal exposure models for sulfate and aerosol strong acidity. *J. Air Waste Manage. Assoc.* 43: 845-850.
- Suh, H. H.; Koutrakis, P.; Spengler, J. D. (1994) The relationship between airborne acidity and ammonia in indoor environments. *J. Exposure Anal. Environ. Epidemiol.* 4: 1-23.
- Suh, H. H.; Allen, G. A.; Koutrakis, P.; Burton, R. M. (1995) Spatial variation in acidic sulfate and ammonia concentrations within metropolitan Philadelphia. *J. Air Waste Manage. Assoc.* 45: 442-452.
- Tamura, K.; Ando, M.; Sagai, M.; Matsumoto, Y. (1996a) Estimation of levels of personal exposure to suspended particulate matter and nitrogen dioxide in Tokyo. *Environ. Sci. (Tokyo)* 4: 37-51.
- Tamura, K.; Ando, M.; Miyazaki, T.; Funasaka, K.; Kuroda, K. (1996b) Indoor concentrations and personal exposure to particulate matter at roadside houses. In: Indoor air '96: the 7th international conference on indoor air quality, volume I; July; Nagoya, Japan. Tokyo, Japan: Institute of Public Health; pp. 599-602.

- Teschke, K.; Marion, S. A.; Vaughan, T. L.; Morgan, M. S.; Camp, J. (1999) Exposures to wood dust in U.S. industries and occupations, 1979 to 1997. *Am. J. Ind. Med.* 35: 581-589.
- Thatcher, T. L.; Layton, D. W. (1995) Deposition, resuspension, and penetration of particles within a residence. *Atmos. Environ.* 29: 1487-1497.
- Thatcher, T. L.; Lai, A. C. K.; Moreno-Jackson, R.; Sextro, R. G.; Nazaroff, W. W. (2002) Effects of room furnishings and air speed on particle deposition rates indoors. *Atmos. Environ.* 36: 1811-1819.
- Thomas, K. W.; Pellizzari, E. D.; Clayton, C. A.; Whitaker, D. A.; Shores, R. C.; Spengler, J.; Özkaynak, H.; Froehlich, S. E.; Wallace, L. A. (1993) Particle total exposure assessment methodology (PTEAM) 1990 study: method performance and data quality for personal, indoor, and outdoor monitoring. *J. Exposure Anal. Environ. Epidemiol.* 3: 203-226.
- Tsang, A. M.; Klepeis, N. E. (1996) Descriptive statistics tables from a detailed analysis of the national human activity pattern survey (NHAPS) data. Las Vegas, NV: U.S. Environmental Protection Agency, Office of Research and Development; report no. EPA/600/R-96/148.
- U.S. Environmental Protection Agency. (1982) Air quality criteria for particulate matter and sulfur oxides. Research Triangle Park, NC: Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office; EPA report no. EPA-600/8-82-029aF-cF. 3v. Available from: NTIS, Springfield, VA; PB84-156777.
- U.S. Environmental Protection Agency. (1992) Guidelines for exposure assessment. Washington, DC: Risk Assessment Forum, USEPA 600Z-92/001.
- U.S. Environmental Protection Agency. (1996) Air quality criteria for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment-RTP Office; report nos. EPA/600/P-95/001aF-cF. 3v.
- U.S. Environmental Protection Agency. (1998) Particulate matter research needs for human health risk assessment to support future reviews of the national ambient air quality standards for particulate matter. Research Triangle Park, NC: National Center for Environmental Assessment; report no. EPA/600/R-97/132F.
- U.S. Environmental Protection Agency. (2000) Aerometric Information Retrieval System (AIRS). Research Triangle Park, NC: Office of Air Quality Planning and Standards. Available: www.epa.gov/ttn/airs/ [2000, October 6].
- Väkevä, M.; Hämeri, K.; Kulmala, M.; Lahdes, R.; Ruuskanen, J.; Laitinen, T. (1999) Street level versus rooftop concentrations of submicron aerosol particles and gaseous pollutants in an urban street canyon. *Atmos. Environ.* 33: 1385-1397.
- Van Vliet, P.; Knape, M.; De Hartog, J.; Janssen, N.; Harssema, H.; Brunekreef, B. (1997) Motor vehicle exhaust and chronic respiratory symptoms in children living near freeways. *Environ. Res.* 74: 122-132.
- Vette, A. F.; Rea, A. W.; Lawless, P. A.; Rodes, C. E.; Evans, G.; Highsmith, V. R.; Sheldon, L. (2001) Characterization of indoor-outdoor aerosol concentration relationships during the Fresno PM exposure studies. *Aerosol Sci. Technol.* 34: 118-126.
- Wagoner, A. P.; Weiss, R. E. (1980) Comparison of fine particle mass concentration and light scattering extinction in ambient aerosol. *Atmos. Environ.* 14: 623-626.
- Wainman, T.; Zhang, J.; Weschler, C. J.; Lioy, P. J. (2000) Ozone and limonene in indoor air: a source of submicron particle exposure. *Environ. Health Perspect.* 108: 1139-1145.
- Waldman, J. M.; Liang, C. S. K. (1993) Penetration and neutralization of acid aerosols at institutional settings. In: Jantunen, M.; Kalliokoski, P.; Kukkonen, E.; Saarela, K.; Seppänen, O.; Vuorelma, H., eds. *Indoor air '93: proceedings of the 6th international conference on indoor air quality and climate. Volume 3: combustion products, risk assessment, policies; July; Helsinki, Finland. Helsinki, Finland: Indoor Air '93; pp. 263-268.*
- Waldman, J. M.; Liang, C. S. K.; Menon, P. (1990) Indoor exposures to acidic aerosols at child and elderly care facilities. In: *Indoor air '90: proceedings of the 5th international conference on indoor air quality and climate, volume 2, characteristics of indoor air; July-August; Toronto, ON, Canada. Ottawa, ON, Canada: International Conference on Indoor Air Quality and Climate, Inc.; pp. 607-612.*
- Waldman, J. M.; Lioy, P. J.; Greenberg, A.; Butler, J. P. (1991) Analysis of human exposure to benzo(a)pyrene via inhalation and food ingestion in the Total Human Environmental Exposure Study (THEES). *J. Exposure Anal. Environ. Epidemiol.* 1: 193-225.
- Wallace, L. (1996) Indoor particles: a review. *J. Air Waste Manage. Assoc.* 46: 98-126.
- Wallace, L. (2000a) Correlations of personal exposure to particles with outdoor air measurements: a review of recent studies. *Aerosol Sci. Technol.* 32: 15-25.
- Wallace, L. (2000b) Real-time monitoring of particles, PAH, and CO in an occupied townhouse. *Appl. Occup. Environ. Hyg.* 15: 39-47.
- Wallace, L.; Howard-Reed, C. (2002) Continuous monitoring of ultrafine, fine, and coarse particles in a residence for 18 months in 1999-2000. *J. Air Waste Manage. Assoc.* 52: 828-844.

- Wallace, L.; Quackenboss, J.; Rodes, C. (1997) Continuous measurements of particles, PAH, and CO in an occupied townhouse in Reston, VA. In: Measurement of toxic and related air pollutants. Volume II: proceedings of a specialty conference; April-May; Research Triangle Park, NC. Pittsburgh, PA: Air & Waste Management Association, pp. 860-871. (A&WMA publication VIP-74).
- Wallace, L. A.; Emmerich, S. J.; Howard-Reed, C. (2002) Continuous measurements of air change rates in an occupied house for 1 year: the effect of temperature, wind, fans, and windows. *J. Exposure Anal. Environ. Epidemiol.* 12: 296-306.
- Watson, J. G.; Cooper, J. A.; Huntzicker, J. J. (1984) The effective variance weighting for least squares calculations applied to the mass balance receptor model. *Atmos. Environ.* 18: 1347-1355.
- Watson, J. G.; Robinson, N. F.; Chow, J. C.; Henry, R. C.; Kim, B. M.; Pace, T. G.; Meyer, E. L.; Nguyen, Q. (1990) The USEPA/DRI chemical mass balance receptor model, CMB 7.0. *Environ. Software* 5: 38-49.
- Watson, J. G.; Chow, J. C.; Pace, T. G. (1991) Chemical mass balance. In: Hopke, P. K., ed. Receptor modeling for air quality management. Amsterdam, The Netherlands: Elsevier Science Publishers B.V.; pp. 83-116. (Vandeginste, B. G. M.; Kvalheim, O. M., eds. Data handling in science and technology, v. 7).
- Weschler, C. J.; Shields, H. C. (1999) Indoor ozone/terpene reactions as a source of indoor particles. *Atmos. Environ.* 33: 2301-2312.
- Wheeler, A. J.; Williams, I.; Beaumont, R. A.; Hamilton, R. S. (2000) Characterisation of particulate matter sampled during a study of children's personal exposure to airborne particulate matter in a UK urban environment. *Environ. Monit. Assess.* 65: 69-77.
- Whitby, K. T. (1978) The physical characteristics of sulfur aerosols. *Atmos. Environ.* 12: 135-159.
- White, W. H. (1998) Statistical considerations in the interpretation of size-resolved particulate mass data. *J. Air Waste Manage. Assoc.* 48: 454-458.
- Wiener, R. W. (1988) Measurement and evaluation of personal exposure to aerosols. In: Measurement of toxic and related air pollutants: proceedings of the 1988 EPA/APCA international symposium; May; Research Triangle Park, NC. Pittsburgh, PA: Air Pollution Control Association; pp. 84-88. (APCA publication VIP-10).
- Wiener, R. W. (1989) Particle total exposure assessment methodology—an overview of planning and accomplishments. In: Measurement of toxic and related air pollutants: proceedings of the 1989 U.S. EPA/A&WMA international symposium; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 442-448. (A&WMA publication VIP-13).
- Wiener, R. W.; Wallace, L.; Pahl, D.; Pellizzari, E.; Whittaker, D.; Spengler, J.; Özkaynak, H. (1990) Review of the particle TEAM 9 home field study. In: Measurement of toxic and related air pollutants: proceedings of the 1990 EPA/A&WMA international symposium, v. 2; May; Raleigh, NC. Pittsburgh, PA: Air & Waste Management Association; pp. 452-460. (A&WMA publication VIP-17).
- Wiley, J. A.; Robinson, J. P.; Piazza, T.; Garrett, K.; Cirkseña, K.; Cheng, Y.-T.; Martin, G. (1991a) Activity patterns of California residents. Final report. Sacramento, CA: California Air Resources Board; report no. ARB/R93/487. Available from: NTIS, Springfield, VA.; PB94-108719.
- Wiley, J. A.; Robinson, J. P.; Cheng, Y.-T.; Piazza, T.; Stork, L.; Pladsen, K. (1991b) Study of children's activity patterns: final report. Sacramento, CA: California Air Resources Board; report no. ARB-R-93/489.
- Williams, R. W.; Watts, R. R.; Stevens, R. K.; Stone, C. L.; Lewtas, J. (1999) Evaluation of a personal air sampler for twenty-four hour collection of fine particles and semivolatile organics. *J. Exposure Anal. Environ. Epidemiol.* 9: 158-166.
- Williams, R.; Suggs, J.; Zweidinger, R.; Evans, G.; Creason, J.; Kwok, R.; Rodes, C.; Lawless, P.; Sheldon, L. (2000a) The 1998 Baltimore Particulate Matter Epidemiology-Exposure Study: part 1. Comparison of ambient, residential outdoor, indoor and apartment particulate matter monitoring. *J. Exposure Anal. Environ. Epidemiol.* 10: 518-532.
- Williams, R.; Suggs, J.; Creason, J.; Rodes, C.; Lawless, P.; Kwok, R.; Zweidinger, R.; Sheldon, L. (2000b) The 1998 Baltimore particulate matter epidemiology-exposure study: part 2. Personal exposure assessment associated with an elderly study population. *J. Exposure Anal. Environ. Epidemiol.* 10: 533-543.
- Williams, R.; Creason, J.; Zweidinger, R.; Watts, R.; Sheldon, L.; Shy, C. (2000c) Indoor, outdoor, and personal exposure monitoring of particulate air pollution: the Baltimore elderly epidemiology-exposure pilot study. *Atmos. Environ.* 34: 4193-4204.
- Wilson, W. E.; Suh, H. H. (1997) Fine particles and coarse particles: concentration relationships relevant to epidemiologic studies. *J. Air Waste Manage. Assoc.* 47: 1238-1249.
- Wilson, W. E.; Mage, D. T.; Grant, L. D. (2000) Estimating separately personal exposure to ambient and nonambient particulate matter for epidemiology and risk assessment: why and how. *J. Air Waste Manage. Assoc.* 50: 1167-1183.

- Yakovleva, E.; Hopke, P. K.; Wallace, L. (1999) Receptor modeling assessment of particle total exposure assessment methodology data. *Environ. Sci. Technol.* 33: 3645-3652.
- Zartarian, V. G.; Ott, W. R.; Duan, N. (1997) A quantitative definition of exposure and related concepts. *J. Exposure Anal. Environ. Epidemiol.* 7: 411-437.
- Zeger, S. L.; Thomas, D.; Dominici, F.; Samet, J. M.; Schwartz, J.; Dockery, D.; Cohen, A. (2000) Exposure measurement error in time-series studies of air pollution: concepts and consequences. *Environ. Health Perspect.* 108: 419-426.
- Zhu, Y.; Hinds, W. C.; Kim, S.; Shen, S.; Sioutas, C. (2002) Study of ultrafine particles near a major highway with heavy-duty diesel traffic. *Atmos. Environ.* 36: 4323-4335.
- Zmirou, D.; Masclet, P.; Boudet, C.; Dor, F.; Déchenaux, J. (2000) Personal exposure to atmospheric polycyclic aromatic hydrocarbons in a general adult population and lung cancer risk assessment. *J. Occup. Environ. Med.* 42: 121-126.



United States
Environmental Protection
Agency

National Center for
Environmental Assessment
Research Triangle Park, NC 27711

Official Business
Penalty for Private Use
\$300

EPA/600/P-99/002aF
October 2004

Please make all necessary changes in the below label,
detach copy or copy, and return to the address in the upper
left-hand corner.

If you do not wish to receive these reports CHECK HERE ;
detach copy or copy, and return to the address in the
upper left-hand corner.

PRESORTED STANDARD
POSTAGE & FEES PAID
EPA
PERMIT No. G-35



United States
Environmental Protection
Agency

National Center for
Environmental Assessment
Research Triangle Park, NC 27711

Official Business
Penalty for Private Use
\$300

EPA/600/P-99/002aF
October 2004

Please make all necessary changes in the below label,
detach copy or copy, and return to the address in the upper
left-hand corner.

If you do not wish to receive these reports CHECK HERE ;
detach copy or copy, and return to the address in the
upper left-hand corner.

PRESORTED STANDARD
POSTAGE & FEES PAID
EPA
PERMIT No. G-35