

## Draft Guidance on Piroxicam

This draft guidance, once finalized, will represent the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

**Active ingredient:** Piroxicam

**Form/Route:** Capsule/Oral

**Recommended studies:** 2 studies

1. Type of study: Fasting  
Design: Single-dose, two-way crossover in-vivo  
Strength: 20 mg  
Subjects: Healthy males and nonpregnant females, general population.  
Additional Comments: Please refer to the Amiodarone Tablet Draft Guidance for additional information regarding long half-life drugs.  

---
2. Type of study: Fed  
Design: Single-dose, two-way crossover *in-vivo*  
Strength: 20 mg  
Subjects: Healthy males and nonpregnant females, general population.  
Additional comments: Please refer to Amantadine Hydrochloride Tablet Draft Guidance for additional information regarding fed studies.  

---

**Analytes to measure (in appropriate biological fluid):** Piroxicam in plasma

**Bioequivalence based on (90% CI):** Piroxicam

**Waiver request of in-vivo testing:** 10 mg based on (i) acceptable bioequivalence studies on the 20 mg strength, (ii) acceptable in-vitro dissolution testing of the 10 mg and 20 mg, and (iii) proportional similarity in the formulations of the 10 mg and 20 mg strengths.

### **Dissolution test method and sampling times:**

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products. Specifications will be determined upon review of the application.