



FENNEC PHARMA

September 2024 | Corporate Presentation

www.fennecpharma.com

Safe Harbor Statement





Except for historical information described in this press release, all other statements are forward-looking. Words such as “believe,” “anticipate,” “plan,” “expect,” “estimate,” “intend,” “may,” “will,” or the negative of those terms, and similar expressions, are intended to identify forward-looking statements. These forward-looking statements include statements about our business strategy, timeline and other goals, plans and prospects, including our commercialization plans respecting PEDMARK[®], the market opportunity for and market impact of PEDMARK[®], its potential impact on patients and anticipated benefits associated with its use, and potential access to further funding after the date of this release. Forward-looking statements are subject to certain risks and uncertainties inherent in the Company’s business that could cause actual results to vary, including the risks and uncertainties that regulatory and guideline developments may change, scientific data and/or manufacturing capabilities may not be sufficient to meet regulatory standards or receipt of required regulatory clearances or approvals, clinical results may not be replicated in actual patient settings, unforeseen global instability, including political instability, or instability from an outbreak of pandemic or contagious disease, such as the novel coronavirus (COVID-19), or surrounding the duration and severity of an outbreak, protection offered by the Company’s patents and patent applications may be challenged, invalidated or circumvented by its competitors, the available market for the Company’s products will not be as large as expected, the Company’s products will not be able to penetrate one or more targeted markets, revenues will not be sufficient to fund further development and clinical studies, our ability to obtain necessary capital when needed on acceptable terms or at all, the Company may not meet its future capital requirements in different countries and municipalities, and other risks detailed from time to time in the Company’s filings with the Securities and Exchange Commission including its Annual Report on Form 10-K for the year ended December 31, 2023. Fennec disclaims any obligation to update these forward-looking statements except as required by law.

For a more detailed discussion of related risk factors, please refer to our public filings available at www.sec.gov and www.sedar.com.

Fennec Pharmaceuticals is a Commercial Stage Biotechnology Company Dedicated to Improving the Lives of Children with Cancer



- **PEDMARK® is FDA-approved in the U.S.** to reduce the risk of ototoxicity or hearing loss associated with cisplatin in pediatric patients 1 month of age and older with localized, non-metastatic solid tumors.
 - Unique formulation of sodium thiosulfate specifically developed in pediatrics
 - U.S. commercial launch October 2022
- **In Europe:**
 - In March 2024, entered into exclusive licensing agreement with Norgine to commercialize PEDMARQSI (European brand name) in Europe, Australia, and New Zealand
 - \$43 million upfront with up to an additional \$230 million in milestone payments and tiered royalties up to the mid twenties
 - **10 YEARS E.U. exclusivity** with Pediatric-use Marketing Authorization (PUMA)
- **Orange Book Patents provide protection until 2039 and 7 years U.S. market exclusivity** with Orphan Drug Exclusivity



US packaging represented

PEDMARK® Registrational Trials Met Their Endpoints



Registrational Trials Design

SIOPEL 6: randomized, controlled, open-label study of 114 patients aged 1 month to 18 years receiving cisplatin-based chemotherapy for standard risk hepatoblastoma


COG ACCL0431: randomized, controlled, open-label study of 125 patients aged 1 to 18 years w/ solid tumors receiving a chemotherapy regimen for any stage disease that included a cumulative cisplatin dose of ≥ 200 mg/m²

Manageable Safety Profile

- The most common adverse reactions ($\geq 25\%$ with difference between arms of $>5\%$ compared to cisplatin alone) in SIOPEL6 are:
 - Vomiting
 - Nausea
 - Decreased hemoglobin
 - Hyponatremia
- The most common adverse reaction ($\geq 25\%$ with difference between arms of $>5\%$ compared to cisplatin alone) in COG ACCL0431 is:
 - Hypokalemia

Primary Endpoint: Hearing Loss

PEDMARK reduced the relative risk of hearing loss by **48%** compared with cisplatin alone*




N = 101 evaluable patients

● CIS ● CIS+STS

Brock et al, N Engl J Med 2018;378:2376-85.

PEDMARK reduced the relative risk of hearing loss by **49%** compared with cisplatin alone**



N = 104 evaluable patients

● CIS ● CIS+STS

Freyer et al, www.thelancet.com/oncology Vol 18 January 2017

*A total of 109 children were randomly assigned to receive cisplatin plus sodium thiosulfate (n=57) or cisplatin alone (n=52) and could be evaluated. The absolute hearing threshold was assessed in 101 children. PEDMARK's FDA-approved label included patients without data that were lost to follow up and assumed to have hearing loss.

**A total of 125 children were randomly assigned to receive either sodium thiosulfate (n=61) or observation (n=64). Of these, 104 participants were assessable for the primary endpoint. PEDMARK's FDA-approved label included patients without data that were lost to follow up and assumed to have hearing loss. The on-label reduction in relative risk of hearing loss with PEDMARK is 25% compared with cisplatin alone.

Please see Important Safety Information at the end of this presentation

Full Prescribing Information is available at www.PEDMARK.com

Cisplatin is the Standard of Care for Pediatric Solid Tumors



Cisplatin | Penicillin of Chemotherapy

- Interferes with DNA replication killing fast proliferating cells
- Administered as intravenous infusion in normal saline
 - For treatment of solid and hematological malignancies
 - Relatively short half-life
- First licensed in 1979
 - Introduced in pediatric patients in 1980s
 - It is on the WHO's List of Essential Medicines
 - High cure rates achieved in pediatric patients, in contrast to adults

Common Childhood Cancers Treated with Cisplatin

- Brain and CNS cancers
- Neuroblastoma
- Hepatoblastoma
- Osteosarcoma
- Germ cell tumors
- Retinoblastoma

Treatment plan depends on the individual cancer diagnosis, stage of disease and patient age

Platinum cancer drugs. Available at cisplatin.org Accessed September 7, 2022.
Robertson J, et al. Bull World Health Organ. 2016 Oct 1; 94(10): 735–742.
Ward et al. CA Cancer J Clin. 2014;64:83-103.



One of the unfortunate complications with cisplatin therapy is **ototoxicity**



**Actor Portrayal*



Cisplatin Results in High-Frequency Hearing Loss in Children



Common Clinical Presentation of Hearing Loss

- High frequency (≥ 4 kHz) sensorineural hearing loss^{1,2}
 - Bilateral (both ears)
 - Progressive
 - Irreversible
 - Can progress to include lower frequencies (< 4 kHz)³
- Can be accompanied by tinnitus³
- Prolonged retention of platinum may cause hearing loss progression after completion of therapy⁴
- Hearing aids may be necessary in up to 40%; and cochlear implants in an additional percentage of children affected³

Risk Factors for Ototoxicity^{1,2}


- Younger age (< 5 years of age)
- Cranial irradiation
- Total dose and duration of platinum agent
- Exposure to other ototoxic medications
- Pre-existing renal insufficiency
- Pre-exposure to therapies that impair hearing ability
- Genetic factors

Hearing loss that is serious enough for hearing aid use has been independently associated with declines in cognition and educational performance⁵

1. Weissbluth S et al. *Int J Pediatr Otorhinolaryngol.* 2018;111:174-179. 2. Paken J et al. *J Toxicol.* 2016;2016:1809394. 3. Langer T et al. *Trends in Pharmacological Sciences.* 2013;34:458-469. 4. Sprauten M. *J. Clin Oncol.* 2012;30:300-307. 5. Schreiber et al., *Neuro Oncol,* 2014;16(8):1129-36.



CIO Can Occur Early in Treatment - As Early As Cycles 1-21



- Ototoxicity is a cisplatin dose-limiting toxicity¹ meaning that efficacy of chemotherapy could be compromised due to ototoxicity management
- Effects can be seen as soon as the second or third dose of cisplatin
- Survivors are at risk of hearing deterioration years after completion of therapy²

Key: — Pre — Post 1st block — Post 2nd block — Post 3rd block — Post 4th block — 1st follow up

Audiogram indicates how loud a sound must be to hear it at a given frequency.

1. Langer T et al. Trends in Pharmacological Sciences. 2013;34:458-469. 2. Bertolini P et al, J Pediatric Hem Onc 2004;26:649-655.



Clinical Manifestations

Effects on growth and development

- Certain consonants (f/th/p/k/h/t) are inaudible, compromising speech recognition and comprehension in young children¹
- High frequency hearing loss affects recognition of plurals such as /s/ in 'ducks' and /z/ in 'girls', resulting in delayed language development³
- Speech perception in background noise is hindered, resulting in poorer school performance (e.g., literacy)^{1,4,5}
- Impaired perception of music and ambient noises, resulting in a poorer quality of life¹
- Delayed neurocognitive and psychosocial development¹

Hearing loss is associated with a lower IQ, phonetic decoding and reading comprehension⁶



1. Langer T et al. *Trends in Pharmacological Sciences*. 2013;34:458-469. 2. Bertolini P et al, *J Pediatric Hem Onc* 2004;26:649-655.
3. Brock PR et al. *J Clin Oncol* 2010;30:2408-2417. 4. Gurney JG et al. *Pediatrics*. 2007;e1229-e1236. 5. Crandell CC. *Ear & Hearing*. 1993; 14:210-217.
6. Hennegan, K, Silber A, Dehipawala S, Chithran K, Lockhart D. Poster PIH67. ISPOR Annual Meeting 2020.



The Impact of Ototoxicity is Profound

- The overall 5-year survival rate for children with localized, non-metastatic disease is 85% or greater, making the permanent and progressive impact of ototoxicity an important consideration¹, yet audiological follow up, today, is inconsistent
- Current interventions occur after hearing loss has occurred and do not restore normal hearing⁴



Nearly 1 in 5 children (18%) considered at-risk for hearing loss do not have hearing tests during follow-up²



More than half (57%) children do not have full audiological monitoring before, during, and after treatment²



More than half (55%) of children with hearing loss have not been documented to require hearing aids³


60% [and up to 90%] of children develop irreversible ototoxicity resulting in a devastating and life-long impact¹

1. ACS Key Statistics for Childhood Cancer <https://www.cancer.org/cancer/cancer-in-children/key-statistics.html>. 2. Clemens E, van den Heuvel-Eibrink MM, Mulder RL, et al.; Lancet Oncol. 2019;20(1):e29-e41. 3. Hennegan, K, Silber A, Dehipawala S, Chithran K, Lockhart D. Poster PIH67. ISPOR Annual Meeting 2020. 4. Landier W.; Cancer. June 2016; Vol. 122, No. 11: 1647-1658. 5. Langer T, Zehnhoff-Dinnesen A; Trends in Pharmacological Sciences. August 2013, Vol. 34, No. 8: 458-469.



Ototoxicity Can Have a Devastating Impact

Long term follow up of neuroblastoma survivors with hearing loss



- High risk for being **held back a grade** (37% vs. 3%)¹
- Twice the rate of parents reported **learning problems** with reading, math, attention and need for special education²
- **Poorer child-reported** school functioning

Even minimal hearing loss is damaging, resulting in compromised learning and language development¹

1. Bess et al., *Ear and Hearing*, 1998, 19:339-54. 2. Gurney et al., *Pediatrics*, 2007 120 (5):229-36 Minimum sensorineural hearing loss (MSHL).

Until Now, Intervention for Ototoxicity Was Not Preventative



Intervention occurs after hearing loss has been detected

Hearing Aids¹

- Do not block out background noise
- Unable to separate speech and noise in loud environments
- Don't allow distant sounds to be heard
- Generally replaced every 3-5 years²

Personal Frequency Modulation (FM Classroom Amplification)


- Patients with hearing loss as a result of cisplatin therapy are more likely to need hearing loss amplification technology
e.g., extended bandwidth hearing aids¹
- There is no data suggesting improvement in speech recognition with this technology³

Cochlear Implants¹

- A surgically implanted neuro-prosthetic device to provide a modified sense of sound for moderate to profound sensorineural hearing loss
- Could be unilateral or bilateral
- Lifelong commitment

Speech Rehabilitation³

- Speech reading and counseling on compensatory communication strategies are needed
- Counseling should include family members including parents and siblings



1. Landier W. *Cancer*. 2016;122:1647-1658.
2. <https://www.starkey.com/blog/2014/02/5-common-questions-about-hearing-aids> accessed Feb 18th 2020.
3. Paken et al, *Journal of Toxicology* 2016, 1809394 | Image: <https://pubs.asha.org/>



Until Now, Intervention for Ototoxicity Was Not Preventative

Intervention occurs after hearing loss has been detected

Hearing Aids¹

- Do not block out background noise
- Unable to separate speech and noise in loud environments
- Don't allow distant sounds to be heard
- Generally replaced every 3-5 years²

Personal Frequency Modulation (FM Classroom Amplification)


- Patients with hearing loss as a result of cisplatin therapy are more likely to need hearing loss amplification technology
e.g., extended bandwidth hearing aids¹
- There is no data suggesting improvement in speech recognition with this technology³

Cochlear Implants¹

- A surgically implanted neuro-prosthetic device to provide a modified sense of sound for moderate to profound sensorineural hearing loss
- Could be unilateral or bilateral
- Lifelong commitment

Speech Rehabilitation³

- Speech reading and counseling on compensatory communication strategies are needed
- Counseling should include family members including parents and siblings





1. Landier W. *Cancer*. 2016;122:1647-1658.
2. <https://www.starkey.com/blog/2014/02/5-common-questions-about-hearing-aids> accessed Feb 18th 2020.
3. Paken et al, *Journal of Toxicology* 2016, 1809394 | Image: <https://pubs.asha.org/>

PEDMARK® Value Proposition



- Only FDA- and EMA-approved treatment for CIO across pediatric solid tumors, based on two Phase 3 clinical studies
 - PEDMARK® and PEDMARQSI™ are indicated to prevent the risk of ototoxicity associated with cisplatin in pediatric patients 1 month of age and older with localized, non metastatic solid tumors
- Clear benefit with ~50% reduction in hearing loss
- Only commercial pharmaceutical company dedicated to CIO with no branded competitor in development
- Unique formulation developed for children with differentiated excipient profile with mild-to-moderate and manageable side effect profile
- Rapid infusion time
- NCCN AYA guidelines adoption Category 2A





Pediatric Oncology Incidence is Consistent in U.S. and EU

Annual incidence of pediatric solid tumor cases eligible for platinum-based therapy in both U.S. and EU markets*



~30% 1,462 Metastatic

~30% 1,710 Metastatic

~70% 3,554 Localized, non-metastatic

~70% 4,215 Localized, non-metastatic

PEDMARK® Market


*Sources: <http://accis.iarc.fr/results/2003/pdfs/summaryincidencetables.pdf> Accessed Feb 2022; Ward, E CA CANCER J CLIN 2014;64:83-103
Localized vs metastatic breakdown based on Qualitative Market Research Study Completed February 2018

U.S. Pediatric Oncology Landscape



Institutions

- ~500 target pediatric hospital centers including COG, NCI and NCCN institutions*
- ~80% of cisplatin use driven by key centers ~200 institutions



*COG: Children's Oncology Group; NCI: National Cancer Institute; NCCN: National Comprehensive Cancer Network



AYA Opportunity

Adolescent and young adult (AYA) cancer patients, those who are diagnosed with cancer for the first time at the age of 15–39 years, are recognized as a distinct population within the oncology community for spontaneous use due to the unique challenges they face throughout their disease trajectory.



At a Glance


Based on estimates of new cancer cases in 2023, 4.4% of all new cases will occur among ages 15 to 39. 85.8% of AYAs diagnosed with cancer will survive their cancer for 5 years after diagnosis.



New Cancer Cases, 2023

Estimated New Cancers Among AYAs in the U.S. in 2023	85,980
% of All New Cancer Cases at Any Age	4.4%


Common Types of New Cancers Among AYAs



Distribution based on age-adjusted rates of new cases. SEER 22, 2016–2020.

<https://seer.cancer.gov/statfacts/html/aya.html>

Rates of New Cases by Cancer Type and Sex



Age-adjusted rates of new cases per 100,000. SEER 22, 2016–2020.

[https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045\(18\)30858-1/fulltext](https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(18)30858-1/fulltext)

Childhood, adolescent, and young adult (CAYA) cancer survivors treated with platinum-based drugs, head or brain radiotherapy, or both have an increased risk of ototoxicity (hearing loss, tinnitus, or both).

PEDMARK® is FDA-approved to reduce the risk of ototoxicity or hearing loss associated with cisplatin in pediatric patients 1 month of age and older with localized, non-metastatic solid tumors.

NCCN Guidelines Category 2A – Significant Opportunity in Adolescents And Young Adults (AYA)



- The Adolescent And Young Adult (AYA) oncology patient is defined as an individual aged 15–39 years of age at the time of initial cancer diagnosis
- PEDMARK Label – Does NOT Have Age Restrictions
- Treatment Related Issues
 - AYA patients should be offered enrollment in open clinical trials for their specific disease when available and appropriate and supportive care should follow well-established guidelines such as those available at www.NCCN.org
- Toxicities
 - Ototoxicity – Consider sodium thiosulfate to prevent the risk of ototoxicity associated with cisplatin in pediatric patients with localized, non-metastatic solid tumors.

Note: All recommendations are category 2A unless otherwise indicated



There is only one PEDMARK®



NCCN Guidelines for Adolescent and Young Adult Oncology recommends sodium thiosulfate (PEDMARK) as a preventative treatment option to reduce CIO in pediatric patients with localized, non-metastatic solid tumors.¹

NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

PEDMARK is the first and only FDA-approved STS formulation for cisplatin-induced ototoxicity²

Source: NCCN Guidelines V3.2023 – AYA Oncology, 09.01.2023 <https://www.nccn.org/guidelines/guidelines-detail?category=4&id=1412>.

Commercial Strategy



PEDMARK® is the First and Only FDA- and EMA-Approved Agent to Lower the Risk of Cisplatin-Induced Hearing Loss in Children with Localized Solid Tumors



- Establish PEDMARK® as *the* necessary complementary agent when prescribing a cisplatin- based therapy for an identified patient with a localized solid tumor
 - Minimize barriers to access and rapid responses to questions
 - Establish Fennec as an optimal partner in pediatric oncology

Key Activities



Direct Promotion

- Digital materials
- Digital MD speaker bureau to engage pediatric oncologists, audiologists, nursing and pharmacists



Aligned Commercial & Medical Infrastructure

- Sales team originally focused on target pediatric facilities
- Expanded team with record of success in community oncology sales (October)



Commercial Partnerships


- Several contracts recently signed with group purchasing organizations
- Specialty pharmacy offering, home infusions, white bag delivery, and direct billing



Access & Patient Support

- 3PL and distribution network
- Patient access services HUB
- Strong support from advocacy groups

Ongoing Partnerships





A single source program for patients needing financial and product access support

Financial Support

- \$0 copay savings eligibility for patients with commercial or private insurance
- Copay assistance through independent charities for eligible Medicaid recipients
- The Fennec Patient Assistance Program for eligible patients without insurance

Patient & Product Support

- Fennec HEARS dedicated care coordinators available to:
 - Answer insurance questions about coverage for PEDMARK
 - Provide you with tips and resources for managing your child's treatment

Commercial Success With Institutions Via Multiple Stakeholders





Ongoing Partnerships

Advocacy Engagement



APHON Dinners Programs Completed or Planned



PEDMARK® (sodium thiosulfate injection): Reducing the Risk of Ototoxicity Associated With Cisplatin in Pediatric Patients 1 Month of Age and Older with Localized, Non-metastatic Solid Tumors

Limitations of Use: The safety and efficacy of PEDMARK have not been established when administered following cisplatin infusions longer than 6 hours. PEDMARK may not reduce the risk of ototoxicity when administered following longer cisplatin infusions, because irreversible ototoxicity may have already occurred.

Please see Important Safety Information on slide 4 and full Prescribing Information for PEDMARK.



This patient comfort kit is sponsored by Fennec Pharmaceuticals Inc., makers of PEDMARK.

To learn more, scan the QR code or visit pedmark.com.

PEDMARK® and Fennec® are registered trademarks of Fennec Pharmaceuticals Inc. ©2023 Fennec Pharmaceuticals Inc. All rights reserved. FEN-1500-v1 01/23



FENNEC | Capital Structure and Financial Information

Stock Listings Current	FENC – Nasdaq FRX – TSX, Canada
Shares Outstanding	27.3 Million
Cash and Cash Equivalents¹	USD \$43.1 Million
YTD Net Revenues¹	USD \$14.6 Million
2023 Cash Burn²	USD \$16.0 Million
Debt³	\$32.0 Million

INSTITUTIONAL OWNERSHIP⁴

Southpoint Capital	16%
Essetifin	16%
Sonic Fund	9%




1. As of June 30, 2024.

2. For the 12-month period ending December 31, 2023.

3. As of June 30, 2024.

4. As of most recent Schedule 13G or Schedule 13F filing by respective fund.

INDICATIONS AND IMPORTANT SAFETY INFORMATION



INDICATION & USAGE

PEDMARK (sodium thiosulfate injection) is indicated to reduce the risk of ototoxicity associated with cisplatin in pediatric patients 1 month of age and older with localized, non-metastatic solid tumors.

Limitations of Use

The safety and efficacy of PEDMARK have not been established when administered following cisplatin infusions longer than 6 hours. PEDMARK may not reduce the risk of ototoxicity when administered following longer cisplatin infusions, because irreversible ototoxicity may have already occurred.

IMPORTANT SAFETY INFORMATION

PEDMARK is contraindicated in patients with history of a severe hypersensitivity to sodium thiosulfate or any of its components.

Hypersensitivity reactions occurred in 8% to 13% of patients in clinical trials. Monitor patients for hypersensitivity reactions. Immediately discontinue PEDMARK and institute appropriate care if a hypersensitivity reaction occurs. Administer antihistamines or glucocorticoids (if appropriate) before each subsequent administration of PEDMARK. PEDMARK may contain sodium sulfite; patients with sulfite sensitivity may have hypersensitivity reactions, including anaphylactic symptoms and life-threatening or severe asthma episodes. Sulfite sensitivity is seen more frequently in people with asthma.

PEDMARK is not indicated for use in pediatric patients less than 1 month of age due to the increased risk of hypernatremia or in pediatric patients with metastatic cancers.

Hypernatremia occurred in 12% to 26% of patients in clinical trials, including a single Grade 3 case. Hypokalemia occurred in 15% to 27% of patients in clinical trials, with Grade 3 or 4 occurring in 9% to 27% of patients. Monitor serum sodium and potassium at baseline and as clinically indicated. Withhold PEDMARK in patients with baseline serum sodium greater than 145 mmol/L.

Monitor for signs and symptoms of hypernatremia and hypokalemia more closely if the glomerular filtration rate (GFR) falls below 60 mL/min/1.73m².

Administer antiemetics prior to each PEDMARK administration. Provide additional antiemetics and supportive care as appropriate.

The most common adverse reactions ($\geq 25\%$ with difference between arms of $> 5\%$ compared to cisplatin alone) in SIOPEL6 were vomiting, nausea, decreased hemoglobin, and hypernatremia. The most common adverse reaction ($\geq 25\%$ with difference between arms of $> 5\%$ compared to cisplatin alone) in COG ACCL0431 was hypokalemia.

Please see full Prescribing Information for PEDMARK at www.PEDMARK.com.



FENNEC PHARMA

**Actor Portrayal*

PEDMARK® and Fennec® are registered trademarks of Fennec Pharmaceuticals Inc.
Fennec HEARS™ is a trademark of Fennec Pharmaceuticals Inc.
©2023 Fennec Pharmaceuticals Inc. All rights reserved. FEN-1321-v9 November 2023